

9 **Highlights**

10 - The strategies used for the replacement of gluten functionality in pasta were reviewed

11 - The effects of treatments on raw-materials were examined

12 - The effects of processing conditions on starch properties and pasta quality were considered

13

14

15 **ABSTRACT:**

16 Defining and optimizing the technological process to improve the sensory and nutritional
17 characteristics of gluten-free (GF) products still represent a challenge for researchers and industry.
18 As regards pasta, several ingredients (modified starch, GF flours, additives) have been used as
19 alternatives to gluten in order to create a starchy network that can withstand the physical stresses of
20 cooking and impart firmness to the cooked product. Moreover, different variations of noodle-
21 making technology have been proposed to simplify the artisan process based on repeated heating
22 and cooling steps, which are difficult to control and monitor. This paper will overview how to
23 replace gluten functionality in GF pasta.

24

25 INTRODUCTION

26 The popularity of pasta is increasing worldwide, thanks to its convenience, palatability, long and
27 easy shelf-life, and, last but not least, its nutritional properties. In addition to the conventional pasta-
28 product made from durum wheat semolina, it is common to enrich pasta with some cereals (barley,
29 rye, etc.), pseudocereals (buckwheat, amaranth, quinoa), and legume flours (pea, chickpea, etc.), to
30 provide sources of fiber, minerals, antioxidants, and polyphenols. In the last few decades, a third
31 group of pasta-products, the gluten-free (GF), is being consumed not only by the growing number
32 of celiacs but also by others who wish to exclude gluten-based products from their diet for health
33 reasons. Moreover, as celiac disease can occur at any age, the production of good quality GF
34 products for people with a tradition consuming of wheat-based products is necessary as an
35 alternative. Currently, there is a broad variety of GF products available for celiacs made from rice,
36 corn, and other GF flours. Unfortunately, most of them exhibit poor cooking quality, particularly
37 when compared with their wheat counterparts (Hager, Zannini & Arendt, 2012; Lucisano, Cappa,
38 Fongaro & Mariotti, 2012). Moreover, many GF products are nutritionally inferior, i.e. poorer in
39 minerals and bio-components, to the wheat-based foods they are intended to replace. These findings
40 suggest that more attention should be paid to the nutritional and sensory quality of GF products. At
41 this regard, recently, the possibility of using green banana flour to produce pasta products with
42 bioactive compounds, such as resistant starch and phenolic acids, was also investigated by
43 Zandonadi et al. (2012). Although the demand for better-tasting, better-textured, and healthier GF
44 products offers great market opportunities for food manufacturers, the replacement of gluten
45 functionality still presents a major technological challenge. The degree of difficulty in producing
46 GF products is closely associated with the technological role of gluten in the food-system. Cookies,
47 whose texture mainly depends on sugar and fat to assure crispness and friability, are the easiest to
48 formulate without gluten because it plays a secondary role in their making and end-product quality
49 (Engleson & Atwell, 2008). The most challenging products to formulate and produce are GF bread

50 and pasta, as gluten is their architectural key. The few papers published in the last decade (FSTA
51 database) on GF pasta (about 20, excluding patents) indicates the difficulty of this task.

52 **GLUTEN FUNCTIONALITY IN PASTA FROM DURUM-WHEAT SEMOLINA**

53 Pasta is considered one of the simplest cereal-based products in terms of ingredients (only two:
54 semolina and water) and processing (a sequence of hydration, mixing, forming, and drying steps).
55 Both raw-material characteristics and processing conditions play a key role in determining the
56 quality of final pasta products (De Noni & Pagani, 2010). Protein quantity and quality have
57 received considerable attention as the most important factors affecting pasta properties (D'Egidio,
58 Mariani, Nardi, Novaro & Cubadda, 1990). A high protein content and a “strong” gluten (in terms
59 of its visco-elasticity) are required to process semolina into a suitable final pasta product with an
60 optimal cooking performance (D'Egidio, Mariani, Nardi, Novaro & Cubadda, 1990; Feillet &
61 Dexter, 1996). Microscopic observations have revealed that the gluten network in dried pasta is
62 more or less uniformly and regularly arranged around starch granules according to the quality of the
63 semolina used (Resmini & Pagani, 1983). On the contrary, starch in dried pasta is still in the form
64 of whole native granules, as in semolina. During cooking, starch and protein exhibit completely
65 different behaviors. The starch granules rapidly swell, tend to disperse, and become partly soluble.
66 While, proteins become completely insoluble and coagulate, creating a strengthened network, which
67 traps starch material (Resmini & Pagani, 1983). Starch gelatinisation and protein coagulation are
68 both competitive phenomena, occur at the same temperature and are influenced by water
69 availability (Pagani, 1986). The faster the formation of a continuous protein network, the more
70 limited the starch swelling, thus ensuring firm consistency and the absence of stickiness in pasta
71 (Resmini & Pagani, 1983). On the contrary, if the protein network lacks elasticity or its formation is
72 delayed, starch granules will easily swell, and part of the starchy material will pass into the cooking
73 water, resulting in a product characterized by stickiness and poor consistency (Resmini & Pagani,
74 1983).

75 **HOW TO REPLACE GLUTEN FUNCTIONALITY IN GLUTEN FREE PASTA**

76 While gluten proteins play a key role in conventional semolina pasta properties, starch is the
77 determining component in GF pasta only if it can re-organize the macromolecular structure in an
78 efficacious way giving a texture similar to that found in semolina products. Pasta companies can
79 adopt different approaches to reach this goal. In any case, starch has to assume a structuring role,
80 which is related to the tendency of its macromolecules to re-associate and interact after
81 gelatinization, resulting in newly organized structures that retard further starch swelling and
82 solubilisation during cooking. Despite this well known fact, few studies have dealt with starch
83 organization in GF pasta. In the late '80s, Mestres, Colonna & Buleon (1988) investigated the starch
84 network of GF noodles using DSC and X-rays, and found that new crystalline organizations were
85 formed as a consequence of starch retrogradation. Amylose-based structures were present in
86 retrograded form (B-type) and the good cooking behaviour of rice noodles was mainly attributed to
87 amylose networks. More recently, Marti, Seetharaman, & Pagani (2010) and Marti, Pagani &
88 Seetharaman (2011a,b) observed that the average molecular weight of amylose and amylopectin, as
89 well as their molecular organization within the granule, affected starch functionality and,
90 consequently, cooking performance.

91 Basically, in GF pasta, the role of gluten could be replaced by choosing suitable formulations and
92 recipes using heat-treated flours as the key-ingredients, or by adopting non-conventional pasta-
93 making processes to induce new rearrangements of starch macromolecules.

94 **GLUTEN-FREE PASTA FORMULATION**

95 The common ingredients in GF pasta are flour and/or starch from corn, rice, potato (or other
96 tubers), with the addition of protein, gums, and emulsifiers which may partially act as substitutes
97 for gluten. The diversity of GF raw materials help to increase the quantity and the quality of
98 products for celiacs.

99 Formulating GF pasta requires, firstly, a thorough knowledge of the component properties of GF
100 flours and starches. Then, appropriate additives may be selected to promote a cohesive mass in the
101 product.

102 ***THE PROPERTIES OF GLUTEN FREE STARCHY FLOURS***

103 The ideal starch for GF pasta products should have a marked tendency to retrograde: this property,
104 generally observed in high amylose cereals and pulses, assures good cooking behaviour in terms of
105 texture and low cooking loss, even after prolonged cooking (Lii & Chang, 1981; Bhattacharya, Zee
106 & Corke, 1999). Mung bean starch is considered one of the best raw material for producing high
107 quality starch spaghetti, due to its high amylose content and type C viscoamylogram pasting profile,
108 characterized by the absence of a peak and the presence of a constantly increasing viscosity during
109 heating and shearing, indicative of good hot-paste stability (Lii & Chang, 1991).

110 Today, GF flours are used more than starches, thus skipping the expensive stage of starch extraction
111 from the grains. Furthermore, from a technological point of view, the use of flours allows to exploit
112 the presence of interactions between starch and other components, such as proteins and lipids.

113 Despite scientific efforts to determine the physico-chemical properties of GF raw materials as they
114 relate to the final quality of noodles (Bhattacharya; Zee & Corke, 1999; Tam, Corke, Tan, Li &
115 Collado, 2004), the selection of raw-materials for GF pasta production is currently based solely on
116 checking for the absence of gluten, while neglecting the evaluation of starch characteristics of GF
117 flours. In fact, GF industries prefer using peculiar heat-treatments or additives for improving the
118 cooking behaviour of GF pasta products.

119 *Rice*

120 Rice (as flour or starch) is present in practically all GF products in the market. Frequently rice flour
121 is produced starting from broken grains which are removed during milling since they decrease the
122 commercial quality of whole grain rice.

123 Traditional rice noodles are made from long-grain rice flour with intermediate-to-high amylose
124 content (> 22 g/100 g), which plays a pivotal role in creating a starch network in rice noodles
125 (Kohlwey, Kendall & Mohindra, 1995). Several studies have assessed the quality of noodles made
126 from different rice varieties. On the basis of sensory evaluation, Sanchez (1975) found a highly
127 significant correlation between high amylose content and panel acceptability. Chen & Luh (1980)
128 reported that the swelling capacity of starch and amylose-amylopectin ratio are the two major
129 factors affecting rice noodle quality. Li & Luh (1980) noted that rice varieties with high amylose
130 content, low gelatinisation temperature, and hard gel consistency were best suited for making
131 noodles. These findings were confirmed some years later, when a good correlation between
132 physico-chemical properties and the texture of vermicelli was found (Bhattacharya; Zee & Corke,
133 1999).

134 Little attention has actually been paid to flour from brown rice, despite its high nutritional value
135 related to dietary fibre, phytic acid, vitamins E and B, and aminobutyric acid (GABA): these
136 components are present in relevant quantities in the bran layers and germ which are removed during
137 the polishing (or milling step) to obtain milled rice. Recently, Marti, Seetharaman & Pagani (2010)
138 prepared GF pasta from brown rice flour. The higher fibre content in brown rice was responsible for
139 a weakening of the starch network and consequently for the increase in cooking loss. At the same
140 time, the inclusion of fibre in the starch matrix partially reduced the extreme firmness and
141 springiness found in pasta from milled rice flour.

142 *Corn*

143 Amylose in corn noodles has also been indicated as the component accounting for their textural
144 integrity after cooking. Dexter & Matsuo (1979) showed that in corn blends, the lower the amylose
145 content, the lower the noodle cooking quality. However, corn starches with high amylose contents
146 ($>40\%$) don't provide a sufficient degree of gelatinisation during the heating process, limiting the
147 extent of the following starch retrogradation (Tam, Corke, Tan, Li & Collado, 2004). Corn starches

148 with amylose contents of around 26-28% were successfully used for bihon-type noodle production
149 (Tam, Corke, Tan, Li & Collado, 2004).

150 Mestres, Colonna, Alexandre & Matencio (1993) studied the effects of various heat-treatments
151 (drum-drying, extrusion-cooking, pasting with hot water, or steaming) on corn pasta properties. The
152 best cooking quality was observed using the drum-drying process, even if no reason was given.
153 Waniska *et al.* (1999) investigated the effects of several parameters on corn noodle quality.
154 Preheating the mixture of corn flour and water (43-45% moisture) at 90-95 °C was required to
155 successfully extrude noodles using a pasta-maker. Adding more water to noodle production resulted
156 in higher gelatinisation, which is associated with longer cooking times and lower cooking losses
157 (Waniska *et al.*, 1999).

158 *Sorghum*

159 The grain presents interesting characteristics from a nutritional standpoint, as it is a source of
160 protein, starch, and antioxidant compounds. For this reason, a potential novel use of sorghum could
161 be the manufacture of pasta products, in addition to or as a substitute for corn or rice flours in the
162 preparation of GF food.

163 Suhendro, Kunitz, McDonough, Rooney & Waniska (2000) investigated the effect of the cultivar,
164 flour particle size, and processing conditions on the cooking quality of noodles prepared from flour
165 of decorticated sorghum on laboratory scale. The fine flour preheated in a microwave oven and
166 dried using the two-stage method produced the best noodles with moderate dry matter loss. Noodles
167 from waxy sorghum proved to be of inferior quality compared to normal sorghum. Such noodles
168 were soft and sticky, with high losses during cooking, probably as a consequence of limited
169 retrogradation extent (Suhendro, Kunitz, McDonough, Rooney & Waniska, 2000).

170 Recently, flour from fermented sorghum was mixed to brown rice flour to prepare GF pasta (Pagani
171 *et al.*, 2010). The modification of the structural and physical properties promoted by fermentation
172 improved pasta quality with respect to the sample from unfermented sorghum.

173 *Pseudo-cereals*

174 Amaranth, quinoa, and buckwheat are becoming increasingly popular because they improve the
175 nutritional quality of GF products, in terms of high fibre, vitamins, minerals, and other bioactive
176 components (polyphenols, phytosterols, etc.) (Alvarez-Jubete, Arendt & Gallagher, 2010). Despite
177 the few published data on oat-enriched GF pasta, oat flour is not commonly used as ingredients for
178 GF formulations. In fact, a number of early studies produced conflicting results and most
179 gastroenterologists have been cautious and recommended avoidance of oats.

180 Good quality spaghetti were produced from blends of corn, soy, oat, and quinoa (5-15%) flours
181 (Caperuto, Amaya-Farfan & Camargo, 2001; Chillo *et al.*, 2009; Mastromatteo, Chillo, Iannetti,
182 Civica & Del Nobile, 2011). GF macaroni from blends of quinoa and rice flour obtained by
183 extrusion at 60 and 77 °C have also been successfully produced (Borges, Ramirez Acheri, Ramirez
184 Ascheri, Do Nascimento & Freitas, 2003). By blending buckwheat, amaranth, and quinoa in
185 different ratios by means of an experimental design (along with the addition of albumen, emulsifier,
186 and enzymes), Schoenlechner, Drausinger, Ottenschlaeger, Jurackova & Berghofer (2011)
187 improved the cooking quality of GF pasta. The best product was prepared from a combination of
188 amaranth, quinoa, and buckwheat (40:40:60), with 6% of egg white powder and 1.2% of emulsifier.
189 More recently, Cabrera-Chávez *et al.* (2012) prepared amaranth-supplemented GF pasta, observing
190 that the incorporation of amaranth to rice flour (25:75 ratio), combined with the cooking-extrusion
191 process, improved the nutritional quality of pasta, while maintaining good cooking behaviour.

192 ***THE USE OF ADDITIVES AND TEXTURING INGREDIENTS***

193 Pasta prepared only from non-gluten flour is generally considered to be inferior in textural quality
194 compared to semolina pasta: it does not tolerate overcooking, it is sticky, and, above all, it is
195 characterized by relevant cooking losses. Adding texturing ingredients can be a simple solution for
196 improving pasta cooking behaviour by decreasing these defects.

197 Hydrocolloids or gums are commonly used for their ability to make a gel in little quantities, provide
198 high consistency at room temperature, improve firmness, give body and mouthfeel to pasta. In
199 addition, because of their ability to bind water, gums can increase the rehydration rate of pasta
200 (Sozer, 2009). A wide range of hydrocolloids have been proposed: arabic gum, xanthan-gum, locust
201 bean gum, carboxymethylcellulose (CMC), etc.

202 Emulsifiers act as lubricants in the extrusion process and provide firmer consistency and a less
203 sticky surface, as they control starch swelling and leaching phenomena during cooking (Lai, 2002),
204 thereby improving the texture of the final product (Kaur, Singh & Singh., 2005; Charutigon,
205 Jitpupakdree, Namsreem & Rungsardthong, 2008).

206 Despite the several well-known positive effects of the addition of emulsifiers and hydrocolloids
207 (Huang, Knight, & Goad 2001; Lai, 2002; Singh, Raina, Bawa & Saxena, 2004; Kaur, Singh &
208 Singh., 2005; Chillo, Laverse, Falcone & Del Nobile, 2007; Charutigon, Jitpupakdree, Namsreem &
209 Rungsardthong, 2008; Sozer, 2009) consumers often associate their presence in GF pasta to an
210 “artificial” food. Consequently, the use of proteins as structuring building ingredients represents an
211 interesting alternative for producing GF pasta, not to mention its positive nutritional effects
212 (Thompson, 2009). In this regard, recent studies found an improvement in pasta texture when egg
213 and milk proteins were used in GF formulations (Chillo *et al.*, 2009; Sozer, 2009; Schoenlechner,
214 Drausinger, Ottenschlaeger, Jurackova & Berghofer, 2011).

215 **THE OPTIMIZATION OF GF PASTA-MAKING PROCESS**

216 Up to now, GF pasta made from solely GF flour has usually been prepared in one of two ways. The
217 first approach focuses on the use of heat-treated flours, in which starch is already mostly
218 gelatinized. Here, the pre-treated flour can be formed into pasta by the continuous extrusion press
219 commonly used in durum wheat semolina pasta-making. In the second technological approach
220 (extrusion-cooking process), native flour is treated with steam and extruded at high temperatures
221 (more than 100°C) for promoting starch gelatinization directly inside the extruder-cooker. Marti,

222 Caramanico, Bottega & Pagani (2012) applied both these processes to native rice flour, without
223 additives or structuring ingredients. Because a regular and continuous protein network was lacking,
224 starch polymers were less efficaciously entrapped in the rice matrix, resulting in a product with high
225 cooking losses (10g/100g), two-three times higher than those of pasta from durum wheat semolina.
226 As regards the texture, pasta prepared from pre-gelatinized flour (Pasta A) exhibited higher
227 firmness compared to that of pasta from extrusion-cooking of native flour, using a single-screw
228 extruder (Pasta B). The ultrastructure images reported in Figure 1 highlighted differences in starch
229 arrangement inside the two products. At the beginning of cooking, Pasta A showed a compact and
230 homogeneous matrix (Figure 1a). On the contrary, the mere immersion of Pasta B in hot water
231 induced a great disruption of surface structure (Figure 1b), accounting for the high water absorption
232 (91 g/100g and 78g/100g, by Pasta B and Pasta A, respectively) and the low firmness (190 N and
233 310 N for Pasta B and Pasta A, respectively). In addition, the extrusion-cooking of native rice was
234 not efficacious in creating a continuous and smoothed starchy matrix, since some aggregates are
235 still recognizable (Figure 1c).

236 Recently, Chillo *et al.* (2010) investigated the effect of the repeated extrusion steps (at temperatures
237 below 46°C) on the sensory characteristics of GF spaghetti. This processing promoted the formation
238 of a compact structure in the dried product. But, the application of shear stress without the
239 combination of high temperature was not efficacious in promoting starch gelatinisation, and thus
240 there was no improvement in the sensory quality of the cooked pasta.

241 Careful selection of processing conditions is the starting point for promoting new starch
242 arrangements in GF raw materials to assure good cooking behaviour and effective structure, not
243 only for the texture but also for nutritional properties in terms of enzyme accessibility and starch
244 digestibility.

245 ***FROM TRADITIONAL NOODLE-MAKING PROCESS TO THE CURRENT***
246 ***TECHNOLOGIES***

247 GF pasta-making is still based on ancient but still-in-use processes for making Oriental starch
248 noodles. As the main ingredient of GF raw-materials, starch plays a key role in noodle production.
249 Non-gluten noodle-technology is mainly based on dough heating and cooling operations, that
250 exploit two phenomena: firstly starch gelatinisation and, then, its retrogradation. The greater the
251 degree of starch gelatinisation, the better the cooking quality. On the contrary, a slight starch
252 swelling is related to pasta disruption during cooking due to the lack of a continuous network of
253 retrograded starch (Pagani, 1986). For this reason the traditional noodle-making process suggests
254 heat-treatments at high temperatures (90-95 °C) during extrusion, which may be repeated several
255 times (Tan, Li & Tan, 2009). During the cooling steps, new and spontaneous starch crystallization
256 occurs, resulting in a translucent, vitreous, and consistent product. These modifications promote a
257 loss of starch granular structure during gelatinisation, and an extensive reticular and fibrillar
258 network after cooling (Resmini & Pagani, 1983).

259 Even if the highly reticulated starch network can account for the good cooking quality of the
260 artisanal pasta, traditional Oriental noodle-technology is difficult to transfer to an industrial scale.
261 Controlling gelatinisation and retrogradation phenomena is hard and requires many hours of work
262 and high amounts of energy and water to heat and cool the dough. Moreover, the size or the
263 diameter of the product is a critical factor: the thin layer of noodles (diameter of 0.68-0.78 mm) is
264 essential in decreasing the sensory perception of extreme hardness and springiness in a product
265 characterized by a strong degree of retrogradation.

266 Considering all these disadvantages, the use of pre-heated flour or extrusion-cooking processed
267 flour bypasses the steps of the discontinuous process (steaming, cooking in boiling water, and
268 cooling), thus simplifying traditional noodle making technology.

269 ***EXTRUSION-COOKING PROCESS***

270 Extrusion-cooking is one of the most suitable technologies for GF pasta-making. Extrusion-cooking
271 consists of using high temperature for a relatively short time, and is commonly used for producing

272 several food items (pre-gelatinized starch, snacks, ready-to-eat breakfast cereals, etc.). The main
273 phenomenon associated with the extrusion-cooking used and exploited in GF pasta-making is again
274 starch gelatinisation. In fact, starch granule organisation is disrupted to render it digestible and to
275 produce a malleable product. In other words, the crystalline starch macromolecules are converted
276 into a more amorphous material, as recently reported by Wolf (2010).

277 Tsao (1976) was one of the first authors to apply extrusion-cooking to make rice spaghetti. More
278 recently, the suitability of pea starch and pea flour for pasta-making using a twin-screw cooking-
279 extruder was investigated (Wang *et al.*, 1999; Vasanthan & Li, 2003). Pasta obtained by extrusion-
280 cooking exhibited superior firmness, flavour, and texture after cooking, compared to pasta-products
281 prepared from the same flour using a conventional extruder (Wang *et al.*, 1999). Extrusion-cooking
282 has been successfully used for pasta production from corn (Budelli & Fontanesi, 2007; Merayo,
283 Gonzalez, Drago, Torres & De Greef, 2011; Giménez *et al.*, 2013). The GF flour was first heat-
284 treated in an extruder by contact with a heated wall and/or steam injection, and then, extruded,
285 formed and shaped, and finally dried. A certain degree of cooking has to be reached so as to obtain
286 pasta with good cooking characteristics and resistance to overcooking (Merayo, Gonzalez, Drago,
287 Torres & De Greef, 2011; Giménez *et al.*, 2013).

288 **THE USE OF PRE-TREATED FLOURS**

289 The use of pre-treated flours, whereby starch is disorganized by pre-cooking it in a separate plant
290 before pasta-making, is one of the processes currently used to prepare GF pasta. In this regard,
291 several heat-treatments have been proposed and each of them specifically affects starch properties
292 (Table 1). Physical treatments have also been applied to starches to alter their native
293 physiochemical properties in order to meet various industrial needs (Zavareze, Storck, de Castro,
294 Schirmer & Dias, 2010). Understanding the nature of the changes could help determine the choice
295 of efficacious treatments on starch for the GF pasta sector.

296 Annealing (ANN), consisting in the treatment of starch in excess of water (more than 40%) at a
297 temperature below gelatinisation (for rice 50-60°C), and heat-moisture treatment (HMT; treatment
298 at low moisture and high temperatures, 100-120°C for rice) are hydrothermal processes often used
299 in modifying the native physiochemical properties of starch (Jacobs & Delcour, 1998). Both ANN
300 and HMT increase starch crystallinity, granule rigidity, and polymer chain associations (Jacobs &
301 Delcour, 1998; Tester & Debon, 2000). These particular hydrothermal treatments suppress granule
302 swelling, retard gelatinization, and increase starch paste stability (Hoover & Vasanthan, 1994;
303 Horndok & Noomhorm, 2007), thus improving cooking behaviour and texture properties of rice
304 noodles (Yoenyongbuddhagal & Noomhorm, 2002; Horndok & Noomhorm, 2007). In addition,
305 Cham & Suwannaporn (2010) optimized hydrothermal treatment conditions to obtain rice noodles
306 exhibiting different cooking qualities. ANN is suitable for preparing fresh rice noodles that require
307 a soft texture, whereas HMT is more appropriate for semi-dried and dried noodles characterized by
308 high tensile strength and gel hardness.

309 Despite the improvements associated with the use of ANN and HMT flours, the use of pre-
310 gelatinized flour is generally considered a cheaper approach for improving rice noodle quality.
311 Raina, Singh, Bawa & Saxena, (2005) reported that the textural quality of both uncooked and
312 cooked pasta improved significantly when pre-gelatinized rice flour was used. Moreover, the
313 intensity of flour pre-gelatinisation plays a very important role in imparting a desirable noodle
314 texture. Although gelatinisation is required to produce the binding effect during extrusion, excessive
315 gelatinisation may cause extremely high extrusion pressures (Juliano & Sakurai 1985). More
316 recently, Yalcin & Basman (2008a) investigated the effect of the gelatinisation level of rice flour on
317 noodle cooking behaviour. Samples obtained with 25% gelatinisation level exhibited lower cooking
318 loss and better tolerance during cooking compared to samples prepared with 15, 20, or 30%
319 gelatinisation level. Other works noted that the effects of gelatinisation extent on the final product
320 depended on the cereal variety and processing conditions used. In the case of corn, noodles
321 containing 80% gelatinised corn flour exhibited the best cooking and sensory properties (Yalcin &

322 Basman, 2008b). The hydration level and the time-temperature conditions of the pre-gelatinisation
323 process significantly affected the pasta-making process and the cooking quality of rice pasta (Lai,
324 2002). Low hydration level (400g water/kg flour) and steaming for short times and low
325 temperatures (85°C for 10 min) resulted in the formation of rice dough that easily extruded into
326 pasta. On the contrary, rice dough prepared using a high hydration level and high gelatinisation was
327 too viscous to be extruded (Lai, 2002). More recently, Marti (2010) reported that the substitution of
328 50% rice flour with pre-gelatinized flour improved the quality of the pasta. These authors supposed
329 that the pre-gelatinized flour may have acted as a binder, re-polymerizing into a network around the
330 starch granules of rice flour during the extrusion step, because of the different gelatinization
331 temperatures of the two flours, thereby increasing their tolerance to cooking stress, as Pagani (1986)
332 suggested too.

333 Pre-gelatinized flours is currently preferred by GF pasta companies also because it can be used in
334 the same conventional press for semolina pasta. However, some technological know-how has to be
335 used. In GF pasta production, the amount of water added to the pre-gelatinized flour has to
336 be calculated by taking into account the higher water affinity of this raw-material. Generally, the final
337 moisture of dough could amount to 40% of the mixture (Marti, Seetharaman & Pagani, 2010),
338 higher than in semolina dough (approximately 30% moisture; Dalbon, Grivon & Pagani, 1996).

339 Recently, Grugni, Mazzini, Viazzo & Viazzo (2009) patented the use of parboiled rice in GF pasta
340 production. Parboiling, carried out on paddy rice, promotes changes in the physicochemical,
341 nutritional, and sensory properties of the kernel: starch gelatinizes, part of the vitamins and minerals
342 migrate towards the endosperm, and a lipid-amylose complex is formed, restricting starch swelling
343 and amylose leaching during cooking (Bhattacharya, 2004). These modifications on starch
344 organization are responsible for decreasing stickiness and increasing hardness of the cooked kernels
345 (Bhattacharya, 2004). Marti, Seetharaman & Pagani (2010) demonstrated that the use of flour from
346 parboiled rice promoted the formation of a new macromolecular structure, resulting in good texture

347 after cooking, also according to the parboiling conditions (steeping temperature) (Marti,
348 Seetharaman & Pagani, 2010; Marti, 2010). Further, the strong interactions of amylopectin and/or
349 amylose promoted by the extrusion-cooking process suggested that the amylopectin matrix is likely
350 a combination of amylose and amylopectin chains. By using parboiled rice flour, the traditional
351 recipe for producing pasta (flour and water) are maintained, while additives (such as modified
352 starches, gums, mono and diglycerides of fatty acids, etc.) are avoided.

353 **CONCLUSIONS**

354 Despite the great efforts made in the last few decades to produce GF pasta with sensory
355 characteristics similar to durum wheat products, the GF pasta currently on the market is still far
356 from what the consumer is looking for. Moreover, little information is available regarding starch
357 arrangements that can guarantee good quality cooking. In fact, up-to-now only a few works have
358 investigated the molecular starch organizations induced by different treatments and how these
359 impact on pasta cooking behaviour. Most studies adopt an empiric approach: varying ingredients
360 and processing conditions rather than understanding the macromolecule organization associated
361 with good or poor cooking quality. Moreover, most of the few studies published refer to laboratory
362 scale pasta-making, neglecting its transfer to an industrial scale. Understanding the relationship
363 between starch structure and processing conditions will help the industry re-formulate and develop
364 products with the desired texture as well as improved nutritional and digestive properties.

365 **References**

- 366 Alvarez-Jubete, L., Arendt, E.K., & Gallagher, E. (2010). Nutritive value of pseudocereals and their
367 increasing use as functional gluten-free ingredients. *Trends in Food Science and Technology*, 21,
368 106-113.
- 369 Bhattacharya, M., Zee, S.Y., & Corke, H. (1999). Physicochemical Properties Related to quality of
370 rice noodles. *Cereal Chemistry*, 76, 861-867.
- 371 Bhattacharya, K. R. (2004). Parboiling of rice. In E.T. Champagne, (Eds.), *Rice: Chemistry and*
372 *Technology* (pp. 329-404). St. Paul: The American Association of Cereal Chemists.

- 373 Borges, J.T. da S., Ramirez Acheri, J. L., Ramirez Ascheri, D., Do Nascimento, R. E., & Freitas, A.
374 S. (2003). Cooking properties and physico-chemical characterization of precooked macaroni of
375 whole quinoa (*Chenopodium quinoa* Willd) flour and rice (*Oryza sativa* L.) flour by extrusion
376 cooking. *Boletim Centro de Pesquisa de Processamento de Alimentos*, 21, 303-322.
- 377 Budelli, A, & Fontanesi, M. (2007). Gluten-free pasta and dough, use of the dough and process for
378 preparing same. EP 1749450 A1.
- 379 Cabrera-Chávez, F., Calderon de la Barca, A., Islas-Rubio, A. R., Marti, A., Marengo, M., Pagani,
380 M.A., Bonomi, F., & Iametti, S. (2012). Molecular rearrangements in extrusion processes for the
381 production of amaranth-enriched, gluten-free rice pasta. *Food Science and Technology*, 47, 421-
382 426.
- 383 Caperuto, L.C., Amaya-Farfan, J., & Camargo, C.R. (2001). Performance of quinoa (*Chenopodium*
384 quinoa Willd) flour in the manufacture of gluten-free spaghetti. *Journal of Food and Agriculture*,
385 81, 95-101.
- 386 Cham, S., & Suwannaporn, P. (2010). Effect of hydrothermal treatment of rice flour on various rice
387 noodles quality. *Journal of Cereal Science*, 51, 284-291.
- 388 ~~Chan, P.S.K., Chen, J., Ettelaie, R., Law, Z., Alevisopoulos, S., Day, E., & Smith, S. (2007). Study
389 of the shear and extensional rheology of casein, waxy maize starch and their mixtures. *Food*
390 *Hydrocolloids*, 21, 716-725.~~
- 391 Charutigon, C., Jitpupakdree, J., Namsreem P., & Rungsardthong, V. (2008). Effects of processing
392 conditions and the use of modified starch and monoglyceride on some properties of extruded rice
393 vermicelli. *Food Science and Technology*, 41, 642-651.
- 394 Chen, S.Y., & Luh, B.S. (1980). Chemical, physical and cooking properties of rice vermicelli. In
395 Proceeding of Rice Technology. Working Group, 18th. Texas A&M University, College Station,
396 TX, USA.
- 397 Chillo, S., Laverse, J., Falcone, P.M., & Del Nobile, M.A. (2007). Effect of carboxymethylcellulose
398 and pregelatinized corn starch on the quality of amaranthus spaghetti. *Journal of Food*
399 *Engineering*, 83, 492-500.
- 400 Chillo, S., Civica, V., Iannetti, M., Suriano, N., Mastromatteo, M., & Del Nobile, M.A. (2009).
401 Properties of quinoa and oat spaghetti loaded with carboxymethylcellulose sodium salt and
402 pregelatinized starch as structuring agents. *Carbohydrate Polymers*, 78, 932-937.
- 403 Chillo S, Civica V., Iannetti, M., Mastromatteo, M., Suriano, N., & Del Nobile, M.A. (2010).
404 Influence of repeated extrusions on some properties of non-conventional spaghetti. *Journal of Food*
405 *Engineering*, 100, 329-335.
- 406 Dalbon, G., Grivon, D., & Pagani, M.A. (1996). Continuous manufacturing process. In J.E. Kruger,
407 R.R. Matsuo, & J.W. Dick (Eds.), *Pasta and Noodle Technology* (pp. 13-58). St. Paul: The
408 American Association of Cereal Chemists.
- 409 D'Egidio, M.G., Mariani, B.M., Nardi, S., Novaro, P. and Cubadda, R. (1990). Chemical and
410 technological variables and their relationships: A predictive equation for pasta cooking quality.
411 *Cereal Chemistry*, 67, 275-281.
412

- 413 Dexter, J.E., & Matsuo, R.R. (1979). Effect of starch on pasta dough rheology and spaghetti cooking
414 quality. *Cereal Chemistry*, 56, 190-195.
- 415 De Noni, I., & Pagani, M.A. (2010). Cooking properties and heat damage of dried pasta as
416 influenced by raw material characteristics and processing conditions. *Critical Review in Food
417 Science and Nutrition*, 50, 465-472.
- 418 Dias, A.R.G., Zavarese, E. da R., Spier, F., de Castro, L.A.S., & Gutkoski, L.C. (2010). Effects of
419 annealing on the physicochemical properties and enzymatic susceptibility of rice starches with
420 different amylase contents. *Food Chemistry*, 123, 711-719.
- 421 Engleson, J., & Atwell, B. (2008). Gluten-free product development. *Cereal Foods World*, 53, 180-
422 184.
- 423 Feillet, P. & Dexter, J.E. (1996). Quality requirements of durum wheat for semolina milling and
424 pasta production. In J.E. Kruger, R.R. Matsuo, & J.W. Dick (Eds.), *Pasta and Noodle Technology*
425 (pp. 95-131). St. Paul: The American Association of Cereal Chemists.
- 426 Giménez, M.A., Gonzalez, R.J., Wagner, J., Torres, R., lobo, M.O., Samman, N.C. (2013). Effect
427 of extrusion conditions on physicochemical and sensorial properties of corn-broad beans (*Vicia
428 faba*) spaghetti type pasta. *Food Chemistry*, 136, 538-545.
- 429 Grugni, G. Mazzini, F., Viazzo, G., & Viazzo, N. (2009). Process for the production of rice pasta
430 and product obtained thereby. EP 2110026 A1.
- 431 Hagenimana, A., Ding, X., & Fang, T. (2006). Evaluation of rice flour modified by extrusion
432 cooking. *Journal of Cereal Science*, 43, 38-46.
- 433 Hager, A.S., Zannini, E., & Arendt, E., K. (2012). Gluten-free pasta: advances in research and
434 commercialization. *Cereal Foods World*, 57, 225-229.
- 435 Hoover, R., & Vasanthan, T. (1994). Effect of heat-moisture treatment on the structure and
436 physicochemical properties of cereal, legume, and tuber starches. *Carbohydrate Research*, 252, 33-
437 53.
- 438 Horndok, R., & Noomhorm, A. (2007). Hydrothermal treatments of rice starch for improvement of
439 rice noodle quality. *Food Science and Technology*, 40, 1723-1731.
- 440 Huang, J.C., Knight, S., & Goad, C. (2001). Model prediction for sensory attributes of non-gluten
441 pasta. *Journal of Food Quality*, 24, 495-511.
- 442 Jacobs, H., Eerlingen, C., Clauwaert, W., & Delcour, J.A. (1995). Influence of annealing on the
443 pasting properties of starch from varying botanical sources. *Cereal Chemistry*, 72, 480-487.
- 444 Jacobs, H., & Delcour, J.A. (1998) Hydrothermal modifications of granular starch, with retention of
445 the granular structure: a review. *Journal of Agriculture and Food Chemistry*, 46, 2895-2905.
- 446 Juliano, B.O., & Sakurai, J. (1985). Miscellaneous products. In B.O. Juliano (Eds), *Rice: chemistry
447 and technology* (pp. 569-618). St. Paul: The American Association of Cereal Chemists.
- 448 Kaur, L., Singh, J., & Singh, N. (2005). Effect of glycerol monostearate on the physico-chemical,
449 thermal, rheological and noodle making properties of corn and potato starches. *Food Hydrocolloids*,
450 19, 839-849.

- 451 Kohlwey, D.E., Kendall, J.H., & Mohindra, R.B. (1995). Using the physical properties of rice as a
452 guide to formulation. *Cereal Foods World*, 40, 728-732.
- 453 Lai, H.M. (2002). Effects of rice properties and emulsifiers on the quality of rice pasta. *Journal of*
454 *Science and Food Agriculture*, 82, 203-216.
- 455 Lai, H.M., & Cheng, H.H. (2004). Properties of pregelatinized rice flour made by hot air or gum
456 puffing. *International Journal of Food Science and Technology*, 39, 201-212.
- 457 Li, C.F., & Luh B.S. (1980). Rice snack foods. In B.S. Luh (Eds), *Rice: Production and Utilization*
458 (pp. 690-711). Westport: Avi.
- 459 Lii, C.Y., & Chang, S.M. (1981). Characterization of red bean (*Phaseolus radiatus* var. aureal)
460 starch and its noodles quality. *Journal of Food Science*, 46, 78-81.
- 461 Lii, C.Y., & Chang, Y.H. (1991). Study of starch in Taiwan. *Food Review International*, 7, 185-
462 203.
- 463 Lu, S., Chen, C.Y., & Lip, C.Y. (1996). Gel-chromatography fractionation and thermal
464 characterization of rice starch affected by hydrothermal treatment. *Cereal Chemistry*, 73, 5-11.
- 465 Lucisano, M., Cappa, C., Fongaro, L., Mariotti, M. (2012). Characterisation of gluten-free pasta
466 through conventional and innovative methods: evaluation of the cooking behaviour. *Journal of*
467 *Cereal Science*, 56, 667-675.
- 468 Marti, A., Seetharaman, K., & Pagani, M.A. (2010). Rice-based pasta: a comparison between
469 conventional pasta-making and extrusion- cooking. *Journal of Cereal Science*, 52, 404-409.
- 470 Marti, A. (2010). Physical approaches to improve the quality of gluten-free pasta. PhD Dissertation,
471 University of Milan.
- 472 Marti, A., Pagani, M.A., & Seetharaman, K. (2011a). Understanding starch organisation in gluten-
473 free pasta from rice flour. *Carbohydrate Polymers*, 84, 1069-1084.
- 474 Marti, A., Pagani, M.A., & Seetharaman, K. (2011b). Characterizing starch structure in a gluten-
475 free pasta by using iodine vapor as a tool. *Starch/Starke*, 63, 241-244.
- 476 Marti, A., Caramanico, R., Bottega, G., & Pagani, M.A. (2012). Effect of treatments on rice flour
477 and coking behaviour of rice pasta. *Cereal Foods World*, 57-4A.
- 478 Mastromatteo, M., Chillo, S., Iannetti, M., Civica, V., & Del Nobile, M.A. (2011). Formulation
479 optimisation of gluten-free functional spaghetti based on quinoa, maize and soy flours.
480 *International Journal of Food Science & Technology*, 46, 1201-1208.
- 481 Merayo, Y. A., Gonzalez, R. J., Drago, S. L., Torres, R. L., & De Greef, D. M. (2011). Extrusion
482 conditions and zea mays endosperm hardness affecting gluten-free spaghetti quality. *International*
483 *Journal of Food Science and Technology* 46, 2321-2328.
- 484 Mestres, C., Colonna, P., & Buleon, A. (1988). Characteristics of starch networks within rice flour
485 noodles and mungbean starch vermicelli. *Journal of Food Science*, 53, 1809-1812.
- 486 Mestres, C., Colonna, P., Alexandre, M.C., & Matencio, F. (1993). Comparison of various
487 processes for making maize pasta. *Journal of Cereal Science*, 17, 277-290.

- 488 Nakorn. K.N., Tongdang, T., & Sirivongpaisal, P. (2009). Crystallinity and rheological properties
489 of pregelatinized rice starches differing in amylose content. *Starch/Starke*, *61*, 101-108.
- 490 Pagani, M.A. (1986). Pasta products from non-conventional raw materials. In C. Mercier, & C.
491 Cantarelli (Eds), *Pasta and extruded products* (pp. 52-68). London: Elsevier Applied Science.
- 492 Pagani, M.A., Bonomi, F., Bottega, G., Casiraghi, M.C., Elkhalfa, A.E.O., & Iametti, S. (2010).
493 Characterizing sorghum-based pasta: a multi-disciplinary approach. Proceedings in 2nd Second
494 International Symposium on Gluten-Free Cereal Products and Beverages, June 8–11, Tampere,
495 Finland.
- 496 Raina, C.S., Singh, S., Bawa, A.S., & Saxena, D.C. (2005). Textural characteristics of pasta made
497 from rice flour supplemented with proteins and hydrocolloids. *Journal of Texture Studies*, *36*, 402-
498 420.
- 499 Resmini, P., & Pagani, M.A. (1983). Ultrastructure studies of pasta: a review. *Food Microstructure*,
500 *2*, 1-12.
- 501 Sanchez, P.C. (1975). Varietal influence on the quality of Phillipines rice. *Food Technology*, *25*, 51-
502 52.
- 503 Schoenlechner, R., Drausinger, J., Ottenschlaeger, V., Jurackova, K., & Berghofer, E. (2011).
504 Functional properties of gluten-free pasta produced from amaranth, quinoa and buckwheat. *Plant*
505 *for Foods and Human Nutrition*, *65*, 339-349.
- 506 Singh, S., Raina, C.S., Bawa, A.S., & Saxena, D.C. (2004). Sweet potato-based pasta product:
507 optimization of ingredient levels using response surface methodology. *International Journal of*
508 *Food Science and Technology*, *39*, 191-200.
- 509 Sozer, N. (2009). Reological properties of rice pasta dough supplemented with proteins and gums.
510 *Food Hydrocolloids*, *23*, 849-855.
- 511 Suhendro, E.L., Kunetz, C.F, McDonough, C.M., Rooney, L.W. & Waniska, R.D. (2000). Cooking
512 characteristics and quality of noodles from food sorghum. *Cereal Chemistry*, *77*, 96-100.
- 513 Tam, L.M., Corke, H, Tan, W.T., Li, J., & Collado, L.S. (2004). Production of Bihon-type noodles
514 from maize starch differing in amylose content. *Cereal Chemistry*, *81*, 475-480.
- 515 Tan, H.Z., Li, Z.G., & Tan, B. (2009). Starch noodles: history, classification, materials, processing,
516 structure, nutrition, quality evaluating and improving. *Food Research International*, *42*, 551-576.
- 517 Tester, R.F., & Debon, S.J.J. (2000). Annealing of starch - a review. *International Journal of*
518 *Biological Macromolecules*, *27*, 1-12.
- 519 Thompson, T. (2009). The nutritional quality of gluten-free foods. In E. Gallagher (Eds), *Gluten-*
520 *free food science and technology* (pp. 42-51). Oxford: Blackwell Publishing Ltd.
- 521 Tsao, T., Beetner, G., Lorenz, K., & Frey, A. (1976). Extrusion processing of instant rice spaghetti.
522 *Lebensmittel-Wissenschaft and Technologie*, *9*, 96-98.
- 523 Vasanthan, T., & Li, J.H. (2003). Hypochlorite oxidation of field pea starch and its suitability for
524 noodle making using an extrusion cooker. *Food Research International*, *36*, 381-386.

- 525 Wang, N., Bhirud, P.R., Sosulski, F.W., & Tyler, R.T. (1999). Pasta-like product from pea flour by
526 twin-screw extrusion. *Journal of Food Science*, 64, 671-678.
- 527 Waniska, R.D., Yi, T., Lu, J., Xue-Ping, L., Xu, W., & Lin, H. (1999). Effects of preheating
528 temperature, moisture, and sodium metabisulfite content on quality of noodles prepared from maize
529 flour or meal. *Food Science and Technology International*, 5, 339-346.
- 530 Wolf, B. (2010). Polysaccharide functionality through extrusion processing. *Current Opinion in*
531 *Colloid and Interface Science*, 15, 50-54.
- 532 Yalcin, S., & Basman, A. (2008a). Effects of gelatinisation level, gum and transglutaminase on the
533 quality characteristics of rice noodle. *International Journal of Food Science and Technology*, 43,
534 1637-1644.
- 535 Yalcin, S., & Basman, A. (2008b). Quality characteristics of corn noodles containing gelatinized
536 starch, transglutaminase and gum. *Journal of Food Quality*, 31, 465-479.
- 537 Yoenyongbuddhagal, S., & Noomhorm, A. (2002). Effect of physicochemical properties of high-
538 amylose thai rice flours on vermicelli quality. *Cereal Chemistry*, 79, 481-485.
- 539 Zandonadi, R.P., Botelho, R.B.A., Gandolfi, L., Ginani, J.S., Montenegro, F.M., Pratesi, R. (2012).
540 Green banana pasta: an alternative for gluten-free diets. *Journal of the Academy of Nutrition and*
541 *Dietetics*, 112, 1068-1072.
- 542 Zavareze, E. da R., Storck, C.R., de Castro, L.A.S., Schirmer, M.A., & Dias, A.R.G. (2010). Effect
543 of heat-moisture treatment on rice starch of varying amylose content. *Food Chemistry*, 121, 358-
544 365.