

ORIGINAL ARTICLE

Variation in *PCSK9* and *HMGCR* and Risk of Cardiovascular Disease and Diabetes

Brian A. Ference, M.D., Jennifer G. Robinson, M.D., M.P.H.,
Robert D. Brook, M.D., Alberico L. Catapano, Ph.D., M. John Chapman, Ph.D.,
David R. Neff, D.O., Szilard Voros, M.D., Robert P. Giugliano, M.D.,
George Davey Smith, M.D., D.Sc., Sergio Fazio, M.D., Ph.D.,
and Marc S. Sabatine, M.D., M.P.H.

ABSTRACT

BACKGROUND

Pharmacologic inhibitors of proprotein convertase subtilisin–kexin type 9 (*PCSK9*) are being evaluated in clinical trials for the treatment of cardiovascular disease. The effect of lowering low-density lipoprotein (LDL) cholesterol levels by inhibiting *PCSK9* on the risk of cardiovascular events or diabetes is unknown.

METHODS

We used genetic scores consisting of independently inherited variants in the genes encoding *PCSK9* and 3-hydroxy-3-methylglutaryl–coenzyme A reductase (*HMGCR*; the target of statins) as instruments to randomly assign 112,772 participants from 14 studies, with 14,120 cardiovascular events and 10,635 cases of diabetes, to groups according to the number of LDL cholesterol–lowering alleles that they had inherited. We compared the effects of lower LDL cholesterol levels that were mediated by variants in *PCSK9*, *HMGCR*, or both on the risk of cardiovascular events and the risk of diabetes.

RESULTS

Variants in *PCSK9* and *HMGCR* were associated with nearly identical protective effects on the risk of cardiovascular events per decrease of 10 mg per deciliter (0.26 mmol per liter) in the LDL cholesterol level: odds ratio for cardiovascular events, 0.81 (95% confidence interval [CI], 0.74 to 0.89) for *PCSK9* and 0.81 (95% CI, 0.72 to 0.90) for *HMGCR*. Variants in these two genes were also associated with very similar effects on the risk of diabetes: odds ratio for each 10 mg per deciliter decrease in LDL cholesterol, 1.11 (95% CI, 1.04 to 1.19) for *PCSK9* and 1.13 (95% CI, 1.06 to 1.20) for *HMGCR*. The increased risk of diabetes was limited to persons with impaired fasting glucose levels for both scores and was lower in magnitude than the protective effect against cardiovascular events. When present together, *PCSK9* and *HMGCR* variants had additive effects on the risk of both cardiovascular events and diabetes.

CONCLUSIONS

In this study, variants in *PCSK9* had approximately the same effect as variants in *HMGCR* on the risk of cardiovascular events and diabetes per unit decrease in the LDL cholesterol level. The effects of these variants were independent and additive. (Funded by the Medical Research Council and the National Heart, Lung, and Blood Institute.)

From the Division of Cardiovascular Medicine, Wayne State University School of Medicine, Detroit (B.A.F.), the Division of Cardiovascular Medicine, University of Michigan Medical School, Ann Arbor (R.D.B.), and Michigan State University, East Lansing (D.R.N.) — all in Michigan; the Departments of Epidemiology and Medicine, College of Public Health, University of Iowa, Iowa City (J.G.R.); the Department of Pharmacological and Biomolecular Sciences, University of Milan and MultiMedica Istituto di Ricovero e Cura a Carattere Scientifico, Milan (A.L.C.); INSERM, Pitié–Salpêtrière University Hospital, Paris (M.J.C.); the Global Genomics Group, Richmond, VA (S.V.); the Thrombolysis in Myocardial Infarction Study Group, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston (R.P.G., M.S.S.); the Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom (G.D.S.); and the Center for Preventive Cardiology, Knight Cardiovascular Institute, Oregon Health and Science University, Portland (S.F.). Address reprint requests to Dr. Ference at the Division of Cardiovascular Medicine, Wayne State University School of Medicine, UHC, 4H-34, Detroit, MI 48202, or at bference@med.wayne.edu.

N Engl J Med 2016;375:2144–53.

DOI: 10.1056/NEJMoa1604304

Copyright © 2016 Massachusetts Medical Society.

MONOCLONAL ANTIBODIES AND OTHER therapies that inhibit proprotein convertase subtilisin–kexin type 9 (PCSK9) have been shown to reduce low-density lipoprotein (LDL) cholesterol levels by approximately 50 to 60% in several randomized trials.^{1–8} Whether lowering LDL cholesterol levels by inhibiting PCSK9 will reduce the risk of cardiovascular events and, like statins, also increase the risk of new-onset diabetes is unknown.⁹

Exploratory and post hoc analyses of randomized trials have suggested that lowering LDL cholesterol levels by approximately 70 mg per deciliter (1.81 mmol per liter) with a PCSK9 inhibitor may reduce the risk of major cardiovascular events by up to 50%.^{10,11} However, these trials included a total of fewer than 120 cardiovascular events, and data on the risk of new-onset diabetes are very limited. Four much larger and longer-term cardiovascular outcome trials are currently ongoing and will provide a much more robust estimate of the clinical effect of PCSK9 inhibitors.^{12–15}

In anticipation of the results of those trials, we have used a “mendelian randomization” approach to test the effect of LDL cholesterol–lowering variants in *PCSK9* on the risk of cardiovascular events and diabetes. The results of a mendelian randomization study can be interpreted as follows: if a genetic variant (e.g., in *PCSK9*) is associated with an exposure of interest (e.g., LDL cholesterol levels) that is observationally associated with the outcome under study (e.g., coronary heart disease), then the observed association between the exposure and the outcome is likely to be causal if the variant is also associated with the outcome. If not, the observed association between the exposure and the outcome is likely to be noncausal. (For more information, see the Methods section in the Supplementary Appendix, available with the full text of this article at NEJM.org.)

Because PCSK9 inhibitors are designed to recapitulate the phenotype of loss-of-function mutations,^{16,17} we used the presence of LDL cholesterol–lowering variants in *PCSK9* to estimate the biologic effect of inhibiting PCSK9 on both the risk of cardiovascular events and the risk of diabetes. We constructed genetic scores that mimic the effect of PCSK9 inhibitors and the effect of statins (which target 3-hydroxy-3-methylglutaryl–coenzyme A reductase [HMGCR]) and

compared the effect of these scores on the risk of cardiovascular disease and the risk of diabetes to make inferences about the potential clinical benefit and safety of treatment with a PCSK9 inhibitor as compared with treatment with a statin.

METHODS

STUDY POPULATION

The study included 112,772 participants (with 14,120 cardiovascular events and 10,635 cases of diabetes) from 14 prospective cohort or case–control studies who had provided written informed consent for genetic studies and for whom individual participant-level data were available as part of the Database of Genotypes and Phenotypes program of the National Center for Biotechnology Information.¹⁸ A description of the included studies and the genotyping platforms that were used in each study is provided in Table S1 in the Supplementary Appendix.

GENETIC INSTRUMENTS

We constructed genetic scores for *PCSK9* and *HMGCR* by combining all variants within 100 kb on either side of each gene that were associated with LDL cholesterol levels at a genomewide level of significance ($P < 5.0 \times 10^{-8}$) as reported by the Global Lipids Genetics Consortium and that were in low linkage disequilibrium ($r^2 < 0.2$) with all other variants included in the score.^{19,20} For each variant, we defined the exposure allele as the allele associated with lower LDL cholesterol levels.¹⁹ For each study participant, we calculated a weighted *PCSK9* genetic score and a weighted *HMGCR* score by adding the number of LDL cholesterol–lowering alleles that the person had inherited at each variant that was included in either score, weighted by the effect of each variant on LDL cholesterol levels measured in milligrams per deciliter.¹⁹

STUDY DESIGN

We dichotomized each genetic score and used this instrument to divide participants into two groups of approximately equal size on the basis of whether their genetic score was above the median or below the median (Fig. S1 in the Supplementary Appendix). Because each variant that was included in either genetic score is inherited approximately randomly at the time of con-

ception²¹ and is inherited approximately independently of other variants included in the score owing to low linkage disequilibrium, the number of LDL cholesterol–lowering alleles that a person inherits in either score should also be random. Therefore, assignment to each group with the use of this instrument should be random. To evaluate the dose–response relationship, we divided participants into four groups on the basis of the quartile value of their genetic score. To compare the separate and combined effect of variants in *PCSK9* and *HMGCR*, we conducted a 2×2 factorial analysis (Fig. S1 in the Supplementary Appendix).

STUDY OUTCOMES

The primary cardiovascular outcome for the study was a composite of the first occurrence of myocardial infarction or death from coronary heart disease. Key secondary cardiovascular outcomes were major coronary events (defined as the first occurrence of myocardial infarction, coronary revascularization, or death from coronary heart disease) and major vascular events (defined as the first occurrence of a major coronary event or stroke). The primary safety outcome was diabetes, defined as either a glycated hemoglobin level greater than 6.5% or treatment with a glucose-lowering medication. Key secondary safety outcomes were fasting plasma glucose level, weight, and body-mass index. To increase the statistical power of the analyses, we combined prevalent and incident outcome events under the assumption that all events occur incident to a genetic exposure.

STATISTICAL ANALYSIS

We assessed whether the assignment to each group was indeed random by comparing the baseline characteristics of the participants in each group. We measured the difference in LDL cholesterol level between groups using linear regression and compared the risk of cardiovascular events or diabetes using logistic-regression analyses that were adjusted for age and sex. To compare the effect of different variants or genetic scores on the risk of cardiovascular events and diabetes, we adjusted each effect size estimate for a standard decrement of 10 mg per deciliter (0.26 mmol per liter) in the LDL cholesterol level using the usual ratio of effect estimates method (for details, see the Methods section in the Supplementary Appendix).

All analyses were performed separately in each of the included studies and then combined across studies in a fixed-effects inverse-variance-weighted meta-analysis to produce summary estimates of effect. To minimize the potential for bias with respect to population stratification, separate analyses were performed for each included ancestral group before being combined.

In a test of replication, we compared the effect of lower LDL cholesterol levels on the risk of coronary heart disease mediated by the *PCSK9* and *HMGCR* genetic scores in up to 62,240 case patients and 127,299 controls without such disease who were enrolled in the Coronary Artery Disease Genomewide Replication and Meta-Analysis plus the Coronary Artery Disease (CARDIoGRAMplusC4D) consortium studies and in up to 86,196 participants of European descent (with 22,669 cases of diabetes) who were enrolled in the Diabetes Genetics Replication and Meta-Analysis (DIAGRAM) consortium studies (Table S2 in the Supplementary Appendix).^{22–24} Pleiotropy was assessed with the use of mendelian randomization–Egger regression.²⁵ All analyses were performed with the use of Stata 12 software or Golden Helix SNP & Variation Suite software (version 8.1.4).²⁶ A detailed description of the methods is provided in the Supplementary Appendix.

RESULTS

PARTICIPANT CHARACTERISTICS

The weighted mean age of the study participants was 59.9 years. The participants had weighted mean cholesterol values as follows: LDL cholesterol, 129.9 mg per deciliter (3.36 mmol per liter); high-density lipoprotein (HDL) cholesterol, 52.3 mg per deciliter (1.35 mmol per liter); and non-HDL cholesterol, 155.3 mg per deciliter (4.02 mmol per liter) (Table S3 in the Supplementary Appendix). Seven variants were included in the *PCSK9* genetic score and six variants in the *HMGCR* genetic score (Tables S4 through S7 in the Supplementary Appendix). There were no significant differences in any nonlipid baseline characteristics between the groups being compared, thus showing that assignment to each group was indeed random (Table 1).

CARDIOVASCULAR EVENTS

As expected, participants in the group with higher *PCSK9* genetic scores had a lower mean LDL

Table 1. Baseline Characteristics of the Participants, According to PCSK9 Genetic Score.*

Characteristic	Below Median Score (N=57,064)	Above Median Score (N=55,708)	P Value
Lipids (mg/dl)			
LDL cholesterol	132.6±35.2	128.4±35.4	5.6×10 ⁻¹⁶
HDL cholesterol	52.4±15.6	52.9±15.8	5.4×10 ⁻⁵
Triglycerides			
Median	121.4	116.1	6.8×10 ⁻¹⁰
Interquartile range	82–164	79–158	
Non-HDL cholesterol	157.6±37.5	153.1±38.2	1.8×10 ⁻¹⁶
Nonlipid characteristics			
Age (yr)	61.3±7.2	61.4±7.2	0.24
Female sex (%)	58.2	58.1	0.68
Blood pressure (mm Hg)			
Systolic	127.7±17.5	127.8±17.2	0.43
Diastolic	74.9±9.9	75.0±10.3	0.36
Weight (kg)	76.9±16.7	76.9±16.2	0.75
Body-mass index†	27.5±5.3	27.7±5.0	0.17
Ever smoked (%)	54.1	54.3	0.28

* Plus–minus values are means ±SD. The weighted PCSK9 genetic score was calculated for each participant by adding the number of low-density lipoprotein (LDL) cholesterol–lowering alleles that the person had inherited at each variant that was included in the score, weighted by the effect of each variant on LDL cholesterol levels measured in milligrams per deciliter. Values in the table represent weighted mean values of the baseline characteristics for the entire study sample (for age and sex) or from the prospective cohort studies (for all other variables) in either group, after combining study-specific estimates in an inverse-variance–weighted meta-analysis. To convert the values for LDL, high-density lipoprotein (HDL), and non-HDL cholesterol to millimoles per liter, multiply by 0.02586. To convert the values for triglycerides to millimoles per liter, multiply by 0.01129.

† The body-mass index is the weight in kilograms divided by the square of the height in meters.

cholesterol level than those in the group with lower PCSK9 scores (difference, –4.2 mg per deciliter [–0.11 mmol per liter]; $P=5.6\times 10^{-16}$), as well as a lower mean level of non-HDL cholesterol (difference, –4.5 mg per deciliter [–0.12 mmol per liter]; $P=1.8\times 10^{-16}$), a lower median level of triglycerides (difference, –5.3 mg per deciliter [–0.06 mmol per liter]; $P=6.8\times 10^{-10}$), and a higher mean level of HDL cholesterol (difference, 0.5 mg per deciliter [0.01 mmol per liter]; $P=5.4\times 10^{-5}$) (Table 1). Participants in the group with higher PCSK9 genetic scores had an 8.4% lower risk of myocardial infarction or death from coronary heart disease (odds ratio, 0.92; 95% confidence interval [CI], 0.88 to 0.95), as well as similarly lower risks of major coronary events, major vascular events, myocardial infarction, and death from coronary heart disease (Fig. S2 in the Supplementary Appendix). In dose–response analyses, increasing PCSK9 scores were associated with a

stepwise decrease in LDL cholesterol levels and a corresponding stepwise decrease in the risk of myocardial infarction or death from coronary heart disease (Fig. 1A). Indeed, when the effects of each PCSK9 score and the individual PCSK9 variants included in these scores were plotted, there was a dose-dependent log-linear association between PCSK9-mediated lower LDL cholesterol levels and the risk of myocardial infarction or death from coronary heart disease (Fig. S3 in the Supplementary Appendix). The effect of the PCSK9 score on the risk of myocardial infarction or death from coronary heart disease was similar in all subgroups studied (Fig. S4 in the Supplementary Appendix).

In similar analyses using the HMGCR genetic score, participants in the group with higher HMGCR scores had a mean LDL cholesterol level that was lower by 3.2 mg per deciliter (0.08 mmol per liter) than participants with lower





Figure 1. Effect of PCSK9 and HMGR Genetic Scores on the Risk of Myocardial Infarction or Death from Coronary Heart Disease.

For each study participant, we calculated a weighted PCSK9 genetic score and a weighted HMGR score by adding the number of low-density lipoprotein (LDL) cholesterol-lowering alleles that the person had inherited at each variant that was included in either score, weighted by the effect of each variant on LDL cholesterol levels measured in milligrams per deciliter. A total of 10,401 primary cardiovascular events (myocardial infarction or death from coronary heart disease [CHD]) were included in the analysis. Of these events, 5508 (1528 prevalent myocardial infarctions and 3980 incident myocardial infarctions or deaths from CHD) occurred in the prospective cohort studies and 4893 in the case-control studies. Across the included studies, the median weighted PCSK9 score was 12.7 (range, 0 to 24.5), and the median weighted HMGR score was 16.8 (range, 4.1 to 26.7). Boxes represent point estimates of effect. Lines represent 95% confidence intervals (CIs).

HMGR scores ($P=2.9 \times 10^{-15}$) and a 6.6% lower risk of myocardial infarction or death from coronary heart disease (odds ratio, 0.93; 95% CI, 0.90 to 0.97) (Fig. 1B). As with the PCSK9 score, the HMGR score had a very consistent effect on each of the secondary outcomes and a similar effect in all subgroups studied.

After adjustment for a standard decrement of 10 mg per deciliter in the LDL cholesterol level, PCSK9 variants were associated with an 18.9% decrease in the risk of myocardial infarction or death from coronary heart disease (odds ratio, 0.81; 95% CI, 0.74 to 0.89) and HMGR variants were associated with a nearly identical 19.1%



decrease in risk (odds ratio, 0.81; 95% CI, 0.72 to 0.90) (Fig. 1C). The effects of the *PCSK9* and *HMGCR* scores were very similar for all of the cardiovascular outcomes studied (Fig. S5 in the Supplementary Appendix). In the 2x2 factorial analysis, the *PCSK9* and *HMGCR* genetic scores had additive effects on LDL cholesterol and the corresponding risk of cardiovascular events (Fig. 2A).


In external validation analyses involving up to 62,240 case patients with coronary heart disease and 127,299 controls without such disease, the *PCSK9* genetic score (odds ratio, 0.84; 95% CI, 0.80 to 0.88) and *HMGCR* genetic score (odds ratio, 0.84; 95% CI, 0.81 to 0.89) had nearly identical associations with the risk of coronary heart disease per decrease of 10 mg per deciliter in the LDL cholesterol level (Figs. S6, S7, and S8 in the Supplementary Appendix). Mendelian randomization–Egger analyses showed that the effect of both *PCSK9* and *HMGCR* variants on the risk of cardiovascular disease was due entirely to their LDL cholesterol–lowering effect, with no evidence for any significant pleiotropic effects (Figs. S9 and S10 in the Supplementary Appendix).

RISK OF DIABETES

Participants in the group with higher *PCSK9* scores had a 6.1% higher risk of diabetes than

those in the group with lower *PCSK9* scores (odds ratio, 1.06; 95% CI, 1.02 to 1.11). After adjustment for a standard decrement of 10 mg per deciliter in the LDL cholesterol level, *PCSK9* variants were associated with an 11.2% increase in the risk of diabetes (odds ratio, 1.11; 95% CI, 1.04 to 1.19). This effect was very similar to the 12.7% increase in the risk of diabetes per 10 mg per deciliter decrease in the LDL cholesterol level that was associated with *HMGCR* variants (odds ratio, 1.13; 95% CI, 1.06 to 1.20) (Fig. 3). Both the *PCSK9* and *HMGCR* genetic scores appeared to have dose-dependent effects on the risk of diabetes. When present together in the 2x2 factorial analysis, the *PCSK9* and *HMGCR* variants had additive effects on the risk of diabetes (Fig. 2B).

Among persons without prevalent diabetes at baseline, neither the *PCSK9* genetic score nor the *HMGCR* genetic score was significantly associated with baseline fasting plasma glucose levels (difference in plasma glucose level for each 10 mg per deciliter decrease in LDL cholesterol level for *PCSK9* score, 0.26 mg per deciliter [0.01 mmol per liter]; $P=0.33$; difference for *HMGCR* score, 0.42 mg per deciliter [0.02 mmol per liter]; $P=0.10$). Among persons with impaired fasting glucose levels at baseline (≥ 100 mg per deciliter [5.6 mmol per liter]), both the *PCSK9* and the *HMGCR* genetic scores were associated with a



higher risk of incident diabetes per decrease of 10 mg per deciliter in the LDL cholesterol level (odds ratio for PCSK9 score, 1.22; 95% CI, 1.03 to 1.45; odds ratio for HMGCR score, 1.19; 95% CI, 1.00 to 1.41). By contrast, among persons with normal fasting glucose levels at baseline, neither the PCSK9 nor the HMGCR genetic score was associated with an increased risk of incident diabetes (odds ratio for PCSK9 score, 0.99; 95% CI, 0.84 to 1.17; odds ratio for HMGCR score, 1.04; 95% CI, 0.89 to 1.22) (Fig. 4).

REPLICATION AND ADDITIONAL ANALYSES

In additional analyses designed to evaluate the role of LDL receptor–mediated pathways as a

potential common mechanism by which PCSK9 and HMGCR inhibition may increase the risk of diabetes, a genetic score consisting of variants in the gene encoding the LDL receptor (*LDLR*) had a very similar effect on the risk of diabetes per unit decrease in the LDL cholesterol level as compared with the PCSK9 and HMGCR genetic scores (Fig. S11 in the Supplementary Appendix).

In a test of replication involving up to 86,196 participants of European descent (with 22,669 cases of diabetes) who were enrolled in the DIAGRAM consortium studies, the PCSK9 and HMGCR genetic scores had very similar effects on the risk of diabetes per decrease of 10 mg per deciliter in the LDL cholesterol level (odds ratio




Figure 4. Effect of *PCSK9* and *HMGCR* Scores on the Risk of Incident Diabetes.

A total of 6295 incident cases of diabetes occurred during follow-up in the prospective cohort studies. After the exclusion of participants with prevalent diabetes, baseline fasting plasma glucose levels were available for 31,077 participants. The main analysis included all the participants after the exclusion of 4340 participants with prevalent diabetes; the subgroup analysis that was stratified according to fasting plasma glucose level included the 31,077 participants without prevalent diabetes for whom baseline fasting plasma glucose levels were available. Boxes represent point estimates of effect. Lines represent 95% CIs.

for *PCSK9* score, 1.08; 95% CI, 1.02 to 1.15; odds ratio for *HMGCR* score, 1.08; 95% CI, 1.02 to 1.14) (Figs. S12, S13, and S14 in the Supplementary Appendix). In additional analyses involving up to 133,100 persons enrolled in the Meta-Analyses of Glucose and Insulin-Related Traits Consortium studies,²⁷ the *PCSK9* and *HMGCR* genetic scores had very similar effects on plasma glucose levels 2 hours after an oral glucose challenge (effect of *PCSK9* score per 10 mg per deciliter decrease in LDL cholesterol, 1.55 mg per deciliter; 95% CI, 0.46 to 2.63 [0.09 mmol per liter; 95% CI, 0.03 to 0.15]; effect of *HMGCR* score per 10 mg per deciliter decrease in LDL cholesterol, 1.44 mg per deciliter; 95% CI, 0.11 to 2.77 [0.08 mmol per liter; 95% CI, 0.01 to 0.15]). These effects were quantitatively similar to the effect of the *PCSK9* 46L loss-of-function allele alone on both the risk of diabetes (odds ratio, 1.09; 95% CI, 0.98 to 1.23) and on plasma glucose levels 2 hours after an oral glucose challenge (effect per 46L allele, 1.51 mg per deciliter; 95% CI, 0.00 to 3.02 [0.08 mmol per liter; 95% CI, 0.00 to 0.17]).

Unlike the *HMGCR* genetic score, the *PCSK9* genetic score was not associated with weight, body-mass index, waist circumference, or waist-to-hip ratio in up to 339,224 participants enrolled in the Genetic Investigation of Anthropometric Traits consortium studies (Table S8 in the Supplementary Appendix).²⁸

DISCUSSION

We found that genetic variants that mimic the effect of *PCSK9* inhibitors had remarkably similar effects on the risk of cardiovascular events and the risk of diabetes as compared with variants that mimic the effect of statins when measured per unit change in the LDL cholesterol level. Furthermore, we found that when variants that mimic the effect of *PCSK9* inhibitors and statins were present together, they had independent and additive effects on the risk of both cardiovascular events and diabetes.

Our finding that *PCSK9* and *HMGCR* variants were associated with approximately the same effect on the risk of cardiovascular disease per unit decrease in the LDL cholesterol level suggests that treatment with a *PCSK9* inhibitor should reduce the risk of cardiovascular events by approximately the same amount as treatment with a statin. Therefore, treatment with a *PCSK9* inhibitor, used either alone or in combination with a statin, should reduce the risk of cardiovascular events by approximately 20% per decrease of 1.0 mmol per liter (39 mg per deciliter) in the LDL cholesterol level.²⁹

Our finding that variants in *PCSK9* and *HMGCR* were associated with very similar effects on the risk of diabetes per unit decrease in the LDL cholesterol level implies that, like statins,

PCSK9 inhibitors may also increase the risk of new-onset diabetes. However, the increased risk of diabetes that was associated with both *PCSK9* and *HMGCR* variants appeared to be confined to persons with impaired fasting glucose levels. Therefore, as with statins, any potential increased risk of new-onset diabetes during treatment with a PCSK9 inhibitor is likely to be confined to persons with impaired fasting glucose levels.

Although variants that mimic the effect of PCSK9 inhibitors and statins were associated with an increased risk of diabetes, the corresponding proportional reduction in cardiovascular risk was much greater than the increased risk of diabetes. Therefore, as with statins, the reduction in cardiovascular risk with PCSK9 inhibitors should far exceed any potential increased risk of diabetes. Furthermore, we found that both *PCSK9* and *HMGCR* variants were associated with a decreased risk of cardiovascular events among persons with diabetes and those without diabetes. This finding is consistent with the results of meta-analyses showing that statins are associated with the same proportional reduction in the risk of cardiovascular events in persons with diabetes as in those without diabetes.³⁰ Therefore, like statins, PCSK9 inhibitors should reduce the risk of cardiovascular events equally well among persons with diabetes and those without diabetes.

The mechanism by which *PCSK9* and *HMGCR* variants increase the risk of diabetes is unclear. However, it is unlikely to be mediated by weight gain because unlike *HMGCR* variants, *PCSK9* variants are not associated with obesity or its subphenotypes, such as weight, body-mass index, or waist circumference. Instead, the mechanism may involve an LDL receptor–mediated pathway. We found that each set of gene-specific variants in *PCSK9*, *HMGCR*, and *LDLR* had a very similar effect as the other sets on the risk of diabetes per unit decrease in the LDL cholesterol level. This finding is consistent with the fact that both PCSK9 and HMGCR inhibitors ultimately reduce plasma LDL cholesterol levels by increasing the density of LDL receptors.³¹ It is also consistent with the observation that persons with familial hypercholesterolemia appear to have a lower prevalence of diabetes than unaffected relatives.³²

The genetic evidence suggests that PCSK9 and HMGCR inhibition, possibly acting through an LDL receptor–mediated pathway, may cause mildly impaired glucose tolerance (as suggested by higher plasma glucose levels 2 hours after an

oral glucose challenge) without materially increasing fasting glucose levels, which may then lead to an increased likelihood of incident diabetes among persons who have impaired fasting glucose levels. This conclusion is consistent with data from the Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin (JUPITER), in which treatment with rosuvastatin was associated with an increased risk of diabetes but not an increase in fasting plasma glucose levels, and virtually all of the increased risk of diabetes occurred among persons with impaired fasting glucose levels.³³

Our study has limitations. Lifelong exposure to decreased levels of LDL cholesterol that are mediated by genetic variants is associated with much greater reductions in the risk of cardiovascular disease per unit decrease in the LDL cholesterol level than is short-term pharmacologic treatment.³⁴ Therefore, the effect of *PCSK9* variants on the risk of cardiovascular events (and probably diabetes) that was estimated is likely to be quantitatively much larger than the effect of treatment with a PCSK9 inhibitor observed in the ongoing outcome trials as measured according to the unit decrease in the LDL cholesterol level. However, having first established that variants that mimic the effect of PCSK9 inhibitors and statins have biologically equivalent effects on the risk of cardiovascular events and diabetes per unit decrease in the LDL cholesterol level, we believe it is reasonable to anticipate that PCSK9 inhibitors and statins are likely to have therapeutically equivalent effects on the risk of cardiovascular events and diabetes per unit decrease in the LDL cholesterol level.^{35,36} In addition, it is important to note that monoclonal antibodies bind extracellular PCSK9 and therefore may not have the same biologic effect as *PCSK9* variants that lower LDL cholesterol levels.

In conclusion, we found that variants in *PCSK9* and *HMGCR* were associated with approximately the same effects on the risk of cardiovascular events and very similar effects on the risk of diabetes per unit decrease in the LDL cholesterol level. We also found that these effects were independent and additive.

Supported by a grant (MC_UU_12013/1, to Dr. Davey Smith) from the Medical Research Council and a grant (R01HL132985, to Dr. Fazio) from the National Heart, Lung, and Blood Institute. Funders for the various studies that are discussed in this article are listed in the Supplementary Appendix.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

REFERENCES

- Roth EM, McKenney JM, Hanotin C, Asset G, Stein EA. Atorvastatin with or without an antibody to PCSK9 in primary hypercholesterolemia. *N Engl J Med* 2012; 367:1891-900.
- Giugliano RP, Desai NR, Kohli P, et al. Efficacy, safety, and tolerability of a monoclonal antibody to proprotein convertase subtilisin/kexin type 9 in combination with a statin in patients with hypercholesterolemia (LAPLACE-TIMI 57): a randomised, placebo-controlled, dose-ranging, phase 2 study. *Lancet* 2012;380:2007-17.
- Koren MJ, Lundqvist P, Bolognese M, et al. Anti-PCSK9 monotherapy for hypercholesterolemia: the MENDEL-2 randomised, controlled phase III clinical trial of evolocumab. *J Am Coll Cardiol* 2014;63:2531-40.
- Robinson JG, Nedergaard BS, Rogers WJ, et al. Effect of evolocumab or ezetimibe added to moderate- or high-intensity statin therapy on LDL-C lowering in patients with hypercholesterolemia: the LAPLACE-2 randomized clinical trial. *JAMA* 2014;311:1870-82.
- Koren MJ, Giugliano RP, Raal FJ, et al. Efficacy and safety of longer-term administration of evolocumab (AMG 145) in patients with hypercholesterolemia: 52-week results from the Open-Label Study of Long-Term Evaluation Against LDL-C (OSLER) randomized trial. *Circulation* 2014;129:234-43.
- Blom DJ, Hala T, Bolognese M, et al. A 52-week placebo-controlled trial of evolocumab in hyperlipidemia. *N Engl J Med* 2014;370:1809-19.
- Kereiakes DJ, Robinson JG, Cannon CP, et al. Efficacy and safety of the proprotein convertase subtilisin/kexin type 9 inhibitor alirocumab among high cardiovascular risk patients on maximally tolerated statin therapy: the ODYSSEY COMBO I study. *Am Heart J* 2015;169(6):906-915.e13.
- Cannon CP, Cariou B, Blom D, et al. Efficacy and safety of alirocumab in high cardiovascular risk patients with inadequately controlled hypercholesterolemia on maximally tolerated doses of statins: the ODYSSEY COMBO II randomized controlled trial. *Eur Heart J* 2015;36:1186-94.
- Swerdlow DI, Preiss D, Kuchenbaecker KB, et al. HMG-coenzyme A reductase inhibition, type 2 diabetes, and bodyweight: evidence from genetic analysis and randomised trials. *Lancet* 2015;385:351-61.
- Sabatine MS, Giugliano RP, Wiviott SD, et al. Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med* 2015;372:1500-9.
- Robinson JG, Farnier M, Krempf M, et al. Efficacy and safety of alirocumab in reducing lipids and cardiovascular events. *N Engl J Med* 2015;372:1489-99.
- Schwartz GG, Bessac L, Berdan LG, et al. Effect of alirocumab, a monoclonal antibody to PCSK9, on long-term cardiovascular outcomes following acute coronary syndromes: rationale and design of the ODYSSEY outcomes trial. *Am Heart J* 2014;168:682-9.
- Sabatine MS, Giugliano RP, Keech A, et al. Rationale and design of the Further cardiovascular Outcomes Research with PCSK9 Inhibition in subjects with Elevated Risk trial. *Am Heart J* 2016;173:94-101.
- ClinicalTrials.gov. The evaluation of bococizumab (PF-04950615;RN316) in reducing the occurrence of major cardiovascular events in high risk subjects (SPIRE-1) (<https://clinicaltrials.gov/ct2/show/NCT01975376>).
- ClinicalTrials.gov. The evaluation of bococizumab (PF-04950615; RN316) in reducing the occurrence of major cardiovascular events in high risk subjects (SPIRE-2) (<https://www.clinicaltrials.gov/ct2/results?term=NCT01975389&Search=Search>).
- Cohen J, Pertsemlidis A, Kotowski IK, Graham R, Garcia CK, Hobbs HH. Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. *Nat Genet* 2005;37:161-5.
- Cohen JC, Boerwinkle E, Mosley TH Jr, Hobbs HH. Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 2006;354:1264-72.
- Mailman MD, Feolo M, Jin Y, et al. The NCBI dbGaP database of genotypes and phenotypes. *Nat Genet* 2007;39:1181-6.
- The Global Lipids Genetics Consortium. Discovery and refinement of loci associated with lipid levels. *Nat Genet* 2013;45:1274-83.
- Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PI. SNAP: a Web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 2008;24:2938-9.
- Lawlor DA, Harbord RM, Sterne JA, Timpson N, Davey Smith G. Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Stat Med* 2008;27:1133-63.
- The CARDIoGRAMplusC4D Consortium. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet* 2013;45:25-33.
- Nikpay M, Goel A, Won HH, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* 2015; 47:1121-30.
- Morris AP, Voight BF, Teslovich TM, et al. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet* 2012;44:981-90.
- Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015;44:512-25.
- SNP & Variation Suite, (version 8.1.4). Bozeman, MT: Golden Helix (http://www.goldenhelix.com/SNP_Variation/index.html).
- Scott RA, Lagou V, Welch RP, et al. Large-scale association analyses identify new loci influencing glyemic traits and provide insight into the underlying biological pathways. *Nat Genet* 2012;44:991-1005.
- Locke AE, Kahali B, Berndt SI, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015;518:197-206.
- The Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010;376:1670-81.
- The Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy of cholesterol-lowering therapy in 18,686 people with diabetes in 14 randomised trials of statins: a meta-analysis. *Lancet* 2008;371:117-25.
- Tavori H, Fan D, Blakemore JL, et al. Serum proprotein convertase subtilisin/kexin type 9 and cell surface low-density lipoprotein receptor: evidence for a reciprocal regulation. *Circulation* 2013;127:2403-13.
- Besseling J, Kastelein JJ, Defesche JC, Hutten BA, Hovingh GK. Association between familial hypercholesterolemia and prevalence of type 2 diabetes mellitus. *JAMA* 2015;313:1029-36.
- Ridker PM, Pradhan A, MacFadyen JG, Libby P, Glynn RJ. Cardiovascular benefits and diabetes risks of statin therapy in primary prevention: an analysis from the JUPITER trial. *Lancet* 2012;380:565-71.
- Ference BA, Yoo W, Alesh I, et al. Effect of long-term exposure to lower low-density lipoprotein cholesterol beginning early in life on the risk of coronary heart disease: a Mendelian randomization analysis. *J Am Coll Cardiol* 2012;60:2631-9.
- Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;372:2387-97.
- Ference BA, Majeed F, Penumetcha R, Flack JM, Brook RD. Effect of naturally random allocation to lower low-density lipoprotein cholesterol on the risk of coronary heart disease mediated by polymorphisms in NPC1L1, HMGCR, or both: a 2x2 factorial Mendelian randomization study. *J Am Coll Cardiol* 2015;65:1552-61.

Copyright © 2016 Massachusetts Medical Society.