

Accepted Manuscript

External Validation Of Equations To Estimate Resting Energy Expenditure In 14952 Adults With Overweight And Obesity And 1948 Adults With Normal Weight From Italy

Giorgio Bedogni, Simona Bertoli, Alessandro Leone, Ramona de Amicis, Elisa Lucchetti, Fiorenza Agosti, Nicoletta Marazzi, Alberto Battezzati, Alessandro Sartorio



PII: S0261-5614(17)31409-7

DOI: [10.1016/j.clnu.2017.11.011](https://doi.org/10.1016/j.clnu.2017.11.011)

Reference: YCLNU 3308

To appear in: *Clinical Nutrition*

Received Date: 7 February 2017

Revised Date: 1 July 2017

Accepted Date: 17 November 2017

Please cite this article as: Bedogni G, Bertoli S, Leone A, de Amicis R, Lucchetti E, Agosti F, Marazzi N, Battezzati A, Sartorio A, External Validation Of Equations To Estimate Resting Energy Expenditure In 14952 Adults With Overweight And Obesity And 1948 Adults With Normal Weight From Italy, *Clinical Nutrition* (2017), doi: 10.1016/j.clnu.2017.11.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **EXTERNAL VALIDATION OF EQUATIONS TO ESTIMATE RESTING ENERGY**
2 **EXPENDITURE IN 14952 ADULTS WITH OVERWEIGHT AND OBESITY AND 1948**
3 **ADULTS WITH NORMAL WEIGHT FROM ITALY**

4
5 Giorgio BEDOGNI ^{1,2*}, Simona BERTOLI ^{1*}, Alessandro LEONE ¹, Ramona DE AMICIS ¹,
6 Elisa LUCCHETTI ³, Fiorenza AGOSTI ³, Nicoletta MARAZZI ³, Alberto BATTEZZATI ^{1**},
7 Alessandro SARTORIO ^{3,4**}

8
9 ¹ International Center for the Assessment of Nutritional Status (ICANS), University of Milan,
10 Milan, Italy.

11
12 ² Clinical Epidemiology Unit, Liver Research Center, Basovizza, Trieste, Italy.

13
14 ³ Istituto Auxologico Italiano, IRCCS, Laboratorio Sperimentale Ricerche Auxo-
15 endocrinologiche, Milan and Verbania, Italy.

16
17 ⁴ Istituto Auxologico Italiano, IRCCS, Divisione di Auxologia, Verbania, Italy.

18
19 * These authors share first authorship; ** these authors share last authorship.

20
21 **Keywords:** overweight, obesity, adults, resting energy expenditure, indirect calorimetry,
22 prediction equations.

23
24 **Running title:** External validation of REE equations.

25 **Funding:** The study was supported by internal grants from the International Center for the
26 Assessment of Nutritional Status and the Italian Institute of Auxology (Progetti di Ricerca
27 Corrente).

28

29 **Send correspondence to:** Dr Giorgio Bedogni, International Center for the Assessment of
30 Nutritional Status (ICANS), Department of Food, Environmental and Nutritional Sciences
31 (DEFENS), University of Milan, Via Botticelli 21, 20133 Milano, Italy. Phone: +39-02-5031
32 6643; Fax: +39-02-50216077; E-mail: giorgiobedogni@gmail.com.

33 **ABSTRACT**

34

35 **Background & aims:** We cross-validated 28 equations to estimate resting energy
36 expenditure (REE) in a very large sample of adults with overweight or obesity.

37

38 **Methods:** 14952 Caucasian men and women with overweight or obesity and 1498 with
39 normal weight were studied. REE was measured using indirect calorimetry and estimated
40 using two meta-regression equations and 26 other equations. The correct classification
41 fraction (CCF) was defined as the fraction of subjects whose estimated REE was within
42 10% of measured REE.

43

44 **Results:** The highest CCF was 79%, 80%, 72%, 64%, and 63% in subjects with normal
45 weight, overweight, class 1 obesity, class 2 obesity, and class 3 obesity, respectively. The
46 Henry weight and height and Mifflin equations performed equally well with CCFs of 77% vs.
47 77% for subjects with normal weight, 80% vs. 80% for those with overweight, 72% vs. 72%
48 for those with class 1 obesity, 64% vs. 63% for those with class 2 obesity, and 61% vs.
49 60% for those with class 3 obesity. The Sabounchi meta-regression equations offered an
50 improvement over the above equations only for class 3 obesity (63%).

51

52 **Conclusions:** The accuracy of REE equations decreases with increasing values of body
53 mass index. The Henry weight & height and Mifflin equations are similarly accurate and
54 the Sabounchi equations offer an improvement only in subjects with class 3 obesity.

55 **ABBREVIATIONS**

56

57 BMI = body mass index

58 eREE = estimated resting energy expenditure

59 FFM = fat-free mass

60 FM = fat mass

61 Ht = height

62 ICANS = International Center for the Assessment of Nutritional Status

63 IQR = interquartile range

64 mREE = measured resting energy expenditure

65 NIH = National Institutes of Health

66 REE = resting energy expenditure

67 RQ = respiratory quotient

68 TEE = total energy expenditure

69 Wt = weight

70 **INTRODUCTION**

71

72 An evaluation of individual energy expenditure is important to deliver effective weight loss
73 programs. Total energy expenditure (TEE) is most commonly calculated from measured
74 (mREE) or estimated (eREE) resting energy expenditure (REE) using a constant
75 correction for the thermic effect of food and a variable correction for physical activity (1).

76

77 As reviewed by Madden et al. (2), REE ($\text{kcal}\cdot\text{day}^{-1}$) is higher in subjects with than in those
78 without obesity. This is explained by the expansion of fat-free mass (FFM) that
79 accompanies the expansion of fat mass (FM) in most subjects with obesity, with the
80 exception of those with genetic obesities such as the Prader-Willi syndrome (3). However,
81 REE standardized on body weight ($\text{kcal}\cdot\text{day}^{-1}\cdot\text{kg}^{-1}$) is lower in obesity because FM, which
82 contributes to REE much less than FFM, accounts for most of the weight of subjects with
83 obesity. Body weight is included in most prediction equations because it explains the
84 greatest portion of REE variability (1). Mostly because the REE-weight relationship differs
85 in subjects with and without obesity, population-specific equations are considered to be
86 needed for subjects with obesity (2).

87

88 Sabounchi et al. (4) have recently developed REE meta-regression equations for 20
89 population groups by pooling the algorithms produced by 47 studies. The 20 population
90 groups are defined on the basis of race, sex and age and the coefficients of the meta-
91 regression equations are weighted averages of the same coefficients across the available
92 equations for a given population. The attractiveness of the Sabounchi equations lies in the
93 fact that the aggregation of different studies is expected to provide more generalizable
94 estimates. The Sabounchi equations have presently undergone external validation only in

95 a small sample of 30 subjects with values of body mass index (BMI) ranging from 19 to 39
96 $\text{kg}\cdot\text{m}^{-2}$ (5).

97

98 Madden et al. (2) have recently performed a systematic review of the equations used to
99 estimate REE in adults with overweight and obesity. They evaluated the accuracy of 28
100 equations that had been cross-validated in external populations. Equations based on
101 simple anthropometric and demographic characteristics were chosen so that they could be
102 easily employed in clinical practice. The conclusion of the systematic review of Madden et
103 al. (2) was that no single equation provided accurate estimates of REE in adults with
104 overweight and obesity.

105

106 The aim of the present study was to externally validate the meta-regression equations of
107 Sabounchi et al. (4) and those systematically reviewed by Madden et al. (2) in subjects
108 with overweight or obesity using subjects with normal weight as comparator.

109

110 **MATERIALS AND METHODS**

111

112 **Study design**

113

114 We retrospectively collected the data of consecutive Caucasian men and women followed
115 between January 2009 and June 2017 at the International Center for the Assessment of
116 Nutritional Status (ICANS, Milan, Italy) and at the Italian Institute of Auxology (Verbania,
117 Italy). The REE of the subjects with overweight and obesity was measured at the inception
118 of a weight-loss program at both Centers. The REE of the subjects with normal weight was
119 measured only at ICANS, which offers weight-maintaining and nutrition counseling
120 programs also for subjects with normal weight. The inclusion criteria were: 1) age \geq 18

121 years; 2) BMI $\geq 18.5 \text{ kg}\cdot\text{m}^{-2}$ and; 3) availability of REE. The exclusion criteria were: 1)
122 syndromic obesity (6); 2) dysthyroidism; 3) use of drugs known to affect energy
123 expenditure (e.g. levothyroxine) and; 4) respiratory quotient (RQ) < 0.67 or > 1.3 (7). The
124 study protocol was approved by the Ethical Committee of the Italian Institute of Auxology.

125

126 **Anthropometric assessment**

127

128 Weight and height were measured following international guidelines (8). BMI was
129 calculated as weight (kg) \cdot height (m) $^{-2}$ and classified as normal weight ($18.5 \leq \text{BMI} \leq 24.9$
130 $\text{kg}\cdot\text{m}^{-2}$), overweight ($25.0 \leq \text{BMI} \leq 29.9 \text{ kg}\cdot\text{m}^{-2}$), class 1 obesity ($30.0 \leq \text{BMI} \leq 34.9 \text{ kg}\cdot\text{m}^{-2}$),
131 class 2 obesity ($35.0 \leq \text{BMI} \leq 39.9 \text{ kg}\cdot\text{m}^{-2}$), and class 3 obesity ($\text{BMI} \geq 40.0 \text{ kg}\cdot\text{m}^{-2}$) (9).

132

133 **REE measurement**

134

135 In both study centers, REE was measured between 8:00 and 10:00 AM in thermo-neutral
136 conditions using an open-circuit indirect calorimeter equipped with a canopy (Vmax 29,
137 Sensor Medics, Yorba Linda, CA). Each indirect calorimeter underwent an ethanol burning
138 test at least one time per year during the study period. The gas analyzers were calibrated
139 before each test using a reference gas mixture made of 15% O₂ and 5% CO₂. The
140 subjects were in the fasting state from at least 8 hours, were not smoking from at least 1
141 hour, and waited at least 30 minutes in the sitting position before undergoing REE
142 measurement. REE was measured in the supine position for at least 30 minutes, including
143 an acclimation period of 10 minutes. The data relative to the acclimation period were
144 discarded. The steady state was defined as at least 5 minutes with less than 5% variation
145 in RQ, less than 10% variation in O₂ consumption, and less than 10% variation in minute
146 ventilation (7). After the steady state was reached, O₂ consumption and CO₂ production

147 were recorded at intervals of one minute for at least 20 minutes and averaged over the
148 whole measurement period. REE was calculated from O₂ consumption and CO₂
149 production using Weir's equation (10).

150

151 **REE estimation**

152

153 REE was estimated using 2 of the 20 Sabouchi meta-regression equations (4) and 26 of
154 the 28 equations systematically reviewed by Madden (2).

155

156 The two Sabouchi equations employed for the present study are the so-called S1
157 equations: 1) REE (kcal·day⁻¹) = 10.2·weight (kg) + 3.09·height (cm) - 3.09·age (years) +
158 301 for women and, 2) REE (kcal·day⁻¹) = 10.4·weight (kg) + 3.19·height (cm) - 3.10·age
159 (years) + 522 for men. These are the Sabouchi weight and height equations applicable to
160 white men and women aged ≥ 18 years and thus to our study subjects (4). Although the
161 equations contributing the most weight to the Sabouchi meta-regression equations were
162 developed at the Italian Institute of Auxology (on a sample of subjects different from that
163 enrolled for the present study) (11), other algorithms were taken into account by the
164 Sabouchi equations (12–14). Moreover, 53% of the present subjects were enrolled at
165 ICANS, which was not involved in the development of the Italian Institute of Auxology REE
166 equations (11). Thus, we considered the Sabouchi equations suitable for our purpose of
167 externally validating REE equations.

168

169 Two of the 28 equations reviewed by Madden et al. (2) had been developed at the Italian
170 Institute of Auxology (on a sample of subjects different from that enrolled for the present
171 study) and were therefore not considered suitable for the present study aimed at validating

172 externally developed equations (15, 16). All the remaining 26 equations (13, 14, 17–21,
173 21–36) were evaluated in the present study.

174

175 **Statistical analysis**

176

177 Most continuous variables were not Gaussian-distributed and all are reported as median
178 (50th percentile) and interquartile range (IQR, 25th and 75th percentiles). Categorical
179 variables are reported as the number and proportion of subjects with the characteristic of
180 interest. Bland-Altman plots of the absolute bias (eREE - mREE) vs. the average bias
181 $[(eREE + mREE) / 2]$ and of the percent bias $[(eREE - mREE) / mREE]$ vs. the average
182 bias were used to investigate the presence of proportional bias (37). The correct
183 classification fraction (CCF) of an equation was defined as the fraction of subjects whose
184 eREE was within 10% of mREE (2). Not unexpectedly (37), proportional bias was detected
185 for almost all equations using both absolute and percent bias (data not shown). Because
186 of this fact and of our primary interest in the CCF of the equations (2), the Bland-Altman
187 limits of agreement were not computed (37). Statistical analysis was performed using
188 Stata 14.2 (Stata Corporation, College Station, TX, USA).

189

190 **RESULTS**

191

192 **Table 1** gives the anthropometric measurements, the mREE and the eREEs of the 16900
193 studied subjects.

194

195 **Table 1 here**

196

197 The median (IQR) age of the subjects was 48 (37;57) years and 72.7% of them were
198 women (**Table 1**). 11.5% of the subjects had a normal weight, 20.9% were overweight,
199 20.5% had class 1 obesity, 20.3% had class 2 obesity, and 26.8% had class 3 obesity
200 (**Table 1**).

201

202 **Table 2** gives the median (IQR) percent bias of the REE equations stratified by BMI class.
203 Using this criterion, the best equation is that with the median bias nearest to 0 and the
204 narrowest IQR.

205

206 **Table 2 here**

207

208 The median percent bias of the REE equations is also plotted in **Figure 1**. Using this
209 criterion, the best equation is that with the dot nearest to the 0 value of the Y-axis.

210

211 **Figure 1 here**

212

213 **Table 3** gives the CCF, i.e. the proportion of subjects whose eREE was within 10% of
214 mREE. Using this criterion, the best equation is that with the highest CCF. This criterion is
215 more useful than the median (IQR) bias to evaluate the applicability of the REE equations
216 at the individual level (2).

217

218 **Table 3 here**

219

220 The CCF is also plotted in **Figure 2**. According to this criterion, the best equation is that
221 with the dot corresponding to the highest value on the Y-axis. Looking at **Figure 2**, it can
222 be clearly seen that, moving from subjects with normal weight to those with class 3 obesity,

223 the CCF of all equations decreases substantially (from 79% to 63% under the best case
224 scenario).

225

226 **Figure 2 here**

227

228 Among the subjects with normal weight, the highest CCF was associated with the Henry
229 weight (Wt) equation (79%, 95% confidence interval 77 to 81%) followed by the Huang
230 (78%, 76% to 80%), Sabounchi (78%, 76% to 80%), and Mifflin equations (77%, 76% to
231 79%) (**Table 3 and Figure 2**).

232

233 Among the subjects with overweight, the highest CCF was associated with the Henry
234 weight and height (Wt & Ht) (80%, 95% confidence interval 78% to 81%) and the Mifflin
235 equations (80%, 95% confidence interval 78% to 81%) followed by the Huang (78%, 77%
236 to 80%), Henry Wt (78%, 76% to 79%), and Sabounchi equations (77%, 76% to 79%)
237 (**Table 3 and Figure 2**).

238

239 Among the subjects with class 1 obesity, the highest CCF was associated with the Mifflin
240 equation (72%, 95% confidence interval 71% to 74%) and the Henry Wt & ht equations
241 (72%, 71% to 74), followed by the Huang (71%, 69% to 72%), and Sabounchi (70%, 69 to
242 72%) equations (**Table 3 and Figure 2**).

243

244 Among the subjects with class 2 obesity, the highest CCF was associated with the Huang
245 equation (65%, 95% confidence interval 64% to 67%) followed by the Sabounchi (64%,
246 63% to 66%), Henry Wt & ht (64%, 62 to 66%), and Mifflin equations (63%, 61% to 65%)
247 (**Table 3 and Figure 2**).

248

249 Lastly, among the subjects with class 3 obesity, the highest CCF was associated with the
250 Huang equation (63%, 95% confidence interval 62 to 65%), followed by the Sabouchi
251 (63%, 61 to 64%), Roza (61%, 59 to 62%), Henry Wt & ht (61%, 59 to 62%), and Mifflin
252 (60%, 59 to 61%) equations (**Table 3** and **Figure 2**).

253

254 **DISCUSSION**

255

256 In the largest study performed so far on Caucasian adults with overweight and obesity, we
257 evaluated the accuracy of two of the 20 REE meta-regression equations of Sabouchi et
258 al. (4) and 26 (13, 14, 17–21, 21–36) of the 28 REE equations systematically reviewed by
259 Madden et al. (2).

260

261 In agreement with Madden et al. (2), we found that the Henry Wt & Ht and the Mifflin
262 equations gave similarly accurate predictions of REE. The CCFs for the Mifflin and the
263 WHO equations were better than those obtained in a previous study performed at the
264 Italian Institute of Auxology (11). The greater accuracy of the WHO and Mifflin equations in
265 the present study may be partly explained by a different case-mix of subjects. 53% of the
266 subjects were in fact contributed by ICANS and the remaining 47% were not involved in
267 the previous study performed at the Italian Institute of Auxology (11). The Sabouchi
268 equation performed better than the Henry Wt & Ht and Mifflin equations only in subjects
269 with class 3 obesity. The Sabouchi equation was however paralleled by the Huang
270 equation, which showed also similar or slightly better CCFs for subjects with normal-weight,
271 overweight, class 1 and class 2 obesity. (It is to be noted that the Huang equation is one of
272 those used by Sabouchi to develop the meta-regression equations). It is noteworthy that
273 there was not a clear winner among the REE equations within any given BMI class (**Table**
274 **3**) and that an equation developed in the general population, i.e. the Henry Wt & Ht

275 equation, had the same accuracy of one specifically developed in obese subjects, i.e. the
276 Mifflin equation (**Table 3**).

277

278 The main strength of the present study is the very large number of enrolled subjects (N =
279 19600) and their balanced distribution within the classes of overweight (N = 3524), degree
280 1 obesity (N = 3464), degree 2 obesity (N = 3429), and degree 3 obesity (N = 4535).

281 Another strength of the present study is that REE was measured using the same
282 instrumentation and protocol at the two study Centers. This is expected to reduce the
283 variability of the bias attributable to the application of the reference method, i.e. indirect
284 calorimetry. Another strength of the present study is the use of a comparator group of
285 subjects with normal weight (N = 1948). We believe that the present study adds
286 substantially to the available data, which were collected mostly on subjects with
287 overweight or class 1 obesity (2).

288

289 The present study has nonetheless two clear limitations. The first limitation is that we
290 studied only Caucasian subjects. Non-Caucasian individuals account for less than 2% of
291 the subjects presently followed at our Centers. The number of non-Caucasian subjects
292 available during the time frame of the study was too low to allow a precise estimate of the
293 bias of the REE equations, especially because stratification on BMI was needed (Tables 2
294 and 3) (2). The second limitation is that our findings may not extend to the general
295 population. This is possibly true also for the subjects with normal weight, because the fact
296 that they sought professional help to maintain their weight and/or ameliorate their diet is
297 likely to select an health-conscious sector of the population. However, if one considers the
298 50th (34.3 kg·m⁻²) and 75th (40.3 kg·m⁻²) percentiles of BMI of our study subjects, it should
299 be clear that subjects with such degree of obesity can be adequately studied only at
300 specialized centers such as ICANS and the Italian Institute of Auxology.

301 The very high number of studied subjects allowed us to obtain precise estimates of the
302 CCF. Because of such precision, we can confidently state that, in our study sample, the
303 Henry Wt & Ht and Mifflin equations perform equally well with a CCF of 77% vs. 77%
304 among subjects with normal weight, 80% vs. 80% among subjects with overweight, 72%
305 vs. 72% among subjects with class 1 obesity, 64% vs. 63% among subjects with class 2
306 obesity, and 60% vs. 60% among subjects with class 3 obesity and that the Sabouchi
307 equations offers an improvement over these equations only in class 3 obesity (CCF =
308 63%).

309
310 The most interesting finding of the present study is that, if one chooses the most accurate
311 equation for a given BMI class, the CCF decreases from 79% among subjects with normal
312 weight and 80% among subjects with overweight to 72% among subjects with class 1
313 obesity to 64% among subjects with class 2 obesity to 63% among subjects with class 3
314 obesity (**Table 3 and Figure 2**). Thus, the accuracy of REE equations decreases
315 substantially with increasing BMI. This has important practical implications as the higher is
316 the BMI of the subject, the higher is the possibility of having her/his REE misclassified with
317 the currently employed REE equations independently of the fact that they were developed
318 in overweight and obese subjects.

319
320 In conclusion, the accuracy of REE equations decreases with increasing BMI. The Henry
321 Wt & Ht and Mifflin equations are similarly accurate to estimate the REE of subjects with
322 overweight and obesity. The Sabouchi equations are more accurate than these equations
323 only in subjects with class 3 obesity.

324

325 **ACKNOWLEDGMENTS**

326

327 We wish to thank the ICANS research staff and especially the dietitians Chiara Caporali,
328 Giulia De Carlo, Valentina Giustizieri, Chiara Lessa, Lidia Lewandowski, and Diana Osio
329 for their help in the collection of the data used in the present study.

330

331 **AUTHORS' CONTRIBUTIONS**

332

333 Study design: Giorgio Bedogni, Simona Bertoli, Alberto Battezzati, Alessandro Sartorio.

334

335 Data collection: Alessandro Leone, Ramona De Amicis, Elisa Lucchetti, Fiorenza Agosti,
336 Nicoletta Marazzi.

337

338 Manuscript writing: Giorgio Bedogni, Simona Bertoli, Alberto Battezzati, Alessandro
339 Sartorio.

340

341 Data management: Alessandro Leone, Giorgio Bedogni.

342

343 Statistical analysis: Giorgio Bedogni.

344 **FIGURE CAPTIONS AND LEGENDS**

345

346 **Figure 1** – Dot chart showing the median percent bias of the REE equations. The best
347 equation is that with the dot nearest to the 0 value of the Y-axis.

348

349 **Figure 2** - Dot chart showing the correct classification fraction of the REE equations. The
350 best equation is that with the dot corresponding to the highest value on the Y-axis.

| | Women | Men | Total |
|--|------------------|------------------|------------------|
| | N = 12281 | N = 4619 | N = 16900 |
| Center | | | |
| Italian Institute of Auxology | 5782 (47.1%) | 2230 (48.3%) | 8012 (47.4%) |
| International Center for the Assessment of Nutritional Status | 6499 (52.9%) | 2389 (51.7%) | 8888 (52.6%) |
| Age (years) | 48 (37;57) | 48 (38;57) | 48 (37;57) |
| Weight (kg) | 87 (72;102) | 105 (91;121) | 92 (76;108) |
| Height (m) | 1.60 (1.55;1.65) | 1.74 (1.70;1.78) | 1.63 (1.57;1.70) |
| BMI (kg·m ⁻²) | 34.0 (27.6;40.5) | 34.8 (30.0;40.3) | 34.3 (28.3;40.4) |
| BMI classification (NIH) | | | |
| Normal weight | 1724 (14.0%) | 224 (4.8%) | 1948 (11.5%) |
| Overweight | 2595 (21.1%) | 929 (20.1%) | 3524 (20.9%) |
| Class 1 obesity | 2268 (18.5%) | 1196 (25.9%) | 3464 (20.5%) |
| Class 2 obesity | 2369 (19.3%) | 1060 (22.9%) | 3429 (20.3%) |
| Class 3 obesity | 3325 (27.1%) | 1210 (26.2%) | 4535 (26.8%) |
| mREE indirect calorimetry (kcal·day ⁻¹) | 1506 (1346;1711) | 1923 (1725;2200) | 1609 (1403;1865) |
| mREE indirect calorimetry (kcal·day ⁻¹ ·kg weight ⁻¹) | 18 (16;20) | 19 (17;20) | 18 (16;20) |
| eREE Bernstein 1983 (14) (kcal·day ⁻¹) | 1279 (1172;1400) | 1618 (1442;1834) | 1344 (1204;1514) |
| eREE De Lorenzo 2001 (35) (kcal·day ⁻¹) | 1743 (1561;1954) | 1844 (1686;2047) | 1773 (1595;1981) |
| eREE de Luis 2006 (34) (kcal·day ⁻¹) | 1626 (1484;1798) | 1796 (1645;1986) | 1674 (1523;1854) |
| eREE Fredrix 1990 (33) (kcal·day ⁻¹) | 1727 (1572;1916) | 2125 (1959;2332) | 1835 (1631;2067) |
| eREE Ganpule 2007 (32) (kcal·day ⁻¹) | 1626 (1465;1814) | 2043 (1878;2251) | 1739 (1524;1970) |
| eREE Harris 1919 (31) (kcal·day ⁻¹) | 1552 (1416;1709) | 2048 (1843;2299) | 1651 (1464;1890) |
| eREE Henry 2005 (Wt) (30) (kcal·day ⁻¹) | 1526 (1379;1695) | 2048 (1848;2305) | 1635 (1431;1889) |
| eREE Henry 2005 (Wt & Ht) (30) (kcal·day ⁻¹) | 1488 (1368;1630) | 1979 (1809;2196) | 1582 (1410;1819) |
| eREE Huang 2004 (13) (kcal·day ⁻¹) | 1500 (1358;1660) | 1996 (1856;2175) | 1614 (1409;1866) |
| eREE Ireton-Jones 1989 (29) (kcal·day ⁻¹) | 1878 (1654;2140) | 2262 (2004;2595) | 1971 (1717;2285) |
| eREE Kleiber 1932 (28) (kcal·day ⁻¹) | 1538 (1403;1699) | 1806 (1652;1999) | 1610 (1446;1802) |
| eREE Korth 2007 (27) (kcal·day ⁻¹) | 1561 (1418;1731) | 2121 (1967;2311) | 1681 (1473;1970) |
| eREE Livingston 2005 (26) (kcal·day ⁻¹) | 1482 (1352;1623) | 1503 (1385;1642) | 1488 (1361;1628) |
| eREE Mifflin 1990 (25) (kcal·day ⁻¹) | 1465 (1322;1635) | 1898 (1744;2086) | 1573 (1375;1803) |

| | | | |
|--|------------------|------------------|------------------|
| eREE Muller 2004 (24) (kcal·day ⁻¹) | 1572 (1409;1752) | 2015 (1857;2214) | 1691 (1468;1918) |
| eREE Muller 2004 (BMI) (24) (kcal·day ⁻¹) * | 1604 (1461;1774) | 2042 (1881;2253) | 1725 (1517;1959) |
| eREE Owen 1986; 1987 (23, 36) (kcal·day ⁻¹) | 1417 (1310;1528) | 1948 (1805;2114) | 1499 (1349;1745) |
| eREE Roza 1984 (22) (kcal·day ⁻¹) | 1533 (1402;1687) | 2050 (1854;2294) | 1633 (1449;1879) |
| eREE Schofield 1985 (Wt) (21) (kcal·day ⁻¹) | 1539 (1411;1688) | 2039 (1857;2262) | 1634 (1456;1888) |
| eREE Schofield 1985 (Wt & Ht) (21) (kcal·day ⁻¹) | 1550 (1414;1705) | 2029 (1842;2259) | 1646 (1460;1908) |
| eREE Siervo 2003 (20) (kcal·day ⁻¹) | 1539 (1367;1716) | 1747 (1586;1935) | 1600 (1418;1784) |
| eREE Tabata 2012 (18) (kcal·day ⁻¹) | 1863 (1542;2194) | 2253 (1952;2604) | 1978 (1638;2321) |
| eREE Weijs 2010 (19) (kcal·day ⁻¹) | 1663 (1462;1886) | 2119 (1922;2360) | 1790 (1533;2049) |
| eREE WHO 1985 (Wt) (17) (kcal·day ⁻¹) | 1579 (1440;1734) | 2068 (1888;2286) | 1678 (1488;1927) |
| eREE WHO 1985 (Wt & Ht) (17) (kcal·day ⁻¹) | 1568 (1435;1723) | 2062 (1873;2290) | 1665 (1480;1910) |
| eREE Sabounchi (S1) 2013 (4) (kcal·day ⁻¹) | 1528 (1383;1692) | 2012 (1867;2200) | 1643 (1435;1889) |

351 *Not available for the 1948 subjects with normal weight

352

353 **Table 1** – Measurements of the study subjects. Continuous variables are reported as
 354 median (50th percentile) and interquartile range (25th and 75th percentiles). Categorical
 355 variables are reported as number and proportion. Abbreviations: BMI = body mass index;
 356 eREE = estimated resting energy expenditure; Ht = height; mREE = measured resting
 357 energy expenditure; NIH = National Institutes of Health; Wt = weight.

358

| | Normal weight | Overweight | Class 1 obesity | Class 2 obesity | Class 3 obesity |
|-------------------------------|---------------|---------------|-----------------|-----------------|-----------------|
| | N = 1948 | N = 3524 | N = 3464 | N = 3429 | N = 4535 |
| Bernstein 1983 (14) | -14 (-19;-9) | -15 (-19;-10) | -16 (-21;-10) | -16 (-22;-10) | -17 (-23;-11) |
| De Lorenzo 2001 (35) | 13 (6;19) | 11 (2;19) | 9 (-1;19) | 10 (-2;20) | 11 (-1;21) |
| de Luis 2006 (34) | 9 (3;17) | 6 (-2;13) | 3 (-6;11) | 2 (-7;12) | 2 (-7;11) |
| Fredrix 1990 (33) | 17 (11;24) | 15 (9;21) | 13 (6;21) | 12 (3;20) | 11 (2;19) |
| Ganpule 2007 (32) | 7 (1;13) | 8 (2;13) | 8 (1;14) | 7 (-1;14) | 6 (-2;15) |
| Harris 1919 (31) | 4 (-1;10) | 4 (-1;10) | 4 (-2;10) | 3 (-4;11) | 2 (-6;10) |
| Henry 2005 (Wt) (30) | 1 (-5;6) | 2 (-3;8) | 3 (-3;10) | 3 (-5;11) | 2 (-6;11) |
| Henry 2005 (Wt & Ht) (30) | 2 (-4;8) | 1 (-4;7) | 1 (-5;7) | -1 (-8;6) | -3 (-11;5) |
| Huang 2004 (13) | 0 (-6;6) | 1 (-4;7) | 1 (-5;8) | 0 (-7;7) | -1 (-9;6) |
| Ireton-Jones 1989 (29) | 31 (22;39) | 37 (29;46) | 5 (-4;15) | 15 (4;26) | 26 (15;40) |
| Kleiber 1932 (28) | 5 (-1;11) | 3 (-3;9) | 0 (-7;7) | -2 (-10;7) | -5 (-13;4) |
| Korth 2007 (27) | 7 (1;14) | 7 (1;13) | 6 (0;13) | 4 (-4;12) | 2 (-6;10) |
| Livingston 2005 (26) | -2 (-9;4) | -4 (-14;2) | -7 (-19;1) | -8 (-20;1) | -10 (-20;-1) |
| Mifflin 1990 (25) | -1 (-7;4) | -1 (-7;4) | -2 (-8;4) | -3 (-11;4) | -4 (-12;3) |
| Muller 2004 (24) | 2 (-3;8) | 4 (-1;9) | 5 (-1;11) | 4 (-3;11) | 4 (-4;12) |
| Muller 2004 (BMI) (24) | Not available | 4 (-1;9) | 3 (-3;10) | 4 (-4;11) | 4 (-4;12) |
| Owen 1986; 1987 (23, 36) | -4 (-9;3) | -2 (-8;4) | -3 (-9;4) | -6 (-13;2) | -8 (-15;0) |
| Roza 1984 (22) | 4 (-2;10) | 4 (-2;9) | 3 (-3;10) | 2 (-5;9) | 1 (-7;9) |
| Schofield 1985 (Wt) (21) | 4 (-1;10) | 5 (-1;10) | 4 (-3;11) | 2 (-6;11) | 1 (-7;10) |
| Schofield 1985 (Wt & Ht) (21) | 7 (0;15) | 6 (-1;12) | 4 (-3;11) | 2 (-6;11) | 0 (-8;9) |
| Siervo 2003 (20) | -4 (-10;2) | -3 (-9;4) | -2 (-9;5) | -1 (-10;7) | 0 (-9;9) |
| Tabata 2012 (18) | 1 (-5;8) | 11 (5;18) | 19 (11;27) | 25 (15;35) | 33 (22;44) |
| Weijs 2010 (19) | 3 (-2;9) | 8 (2;14) | 10 (4;17) | 11 (3;19) | 12 (4;21) |
| WHO 1985 (Wt) (17) | 6 (0;12) | 6 (1;12) | 6 (-1;13) | 5 (-3;13) | 4 (-4;12) |
| WHO 1985 (Wt & Ht) (17) | 6 (0;12) | 7 (1;12) | 5 (-1;13) | 4 (-4;12) | 2 (-6;11) |
| Sabounchi (S1) 2013 (4) | 2 (-4;7) | 3 (-2;8) | 3 (-3;9) | 1 (-6;9) | 0 (-7;8) |

359

360 **Table 2** – Percent bias of the REE equations. Variables are reported as median (50th
361 percentile) and interquartile range (25th and 75th percentiles).

| | Normal weight | Overweight | Class 1 obesity | Class 2 obesity | Class 3 obesity |
|-------------------------------|---------------|------------|-----------------|-----------------|-----------------|
| | N = 1948 | N = 3524 | N = 3464 | N = 3429 | N = 4535 |
| Bernstein 1983 (14) | 28% | 24% | 25% | 24% | 23% |
| De Lorenzo 2001 (35) | 36% | 42% | 43% | 39% | 38% |
| de Luis 2006 (34) | 49% | 58% | 57% | 53% | 54% |
| Fredrix 1990 (33) | 23% | 30% | 37% | 41% | 43% |
| Ganpule 2007 (32) | 64% | 61% | 56% | 53% | 53% |
| Harris 1919 (31) | 72% | 73% | 67% | 61% | 60% |
| Henry 2005 (Wt) (30) | 79% | 78% | 68% | 60% | 58% |
| Henry 2005 (Wt & Ht) (30) | 77% | 80% | 72% | 64% | 61% |
| Huang 2004 (13) | 78% | 78% | 71% | 65% | 63% |
| Ireton-Jones 1989 (29) | 5% | 1% | 52% | 34% | 16% |
| Kleiber 1932 (28) | 68% | 72% | 66% | 58% | 54% |
| Korth 2007 (27) | 61% | 60% | 58% | 59% | 60% |
| Livingston 2005 (26) | 72% | 62% | 50% | 47% | 45% |
| Mifflin 1990 (25) | 77% | 80% | 72% | 63% | 60% |
| Muller 2004 (24) | 78% | 75% | 66% | 60% | 59% |
| Muller 2004 (BMI) (24) | Not available | 75% | 69% | 61% | 58% |
| Owen 1986; 1987 (23, 36) | 71% | 72% | 66% | 55% | 50% |
| Roza 1984 (22) | 74% | 76% | 69% | 63% | 61% |
| Schofield 1985 (Wt) (21) | 71% | 71% | 65% | 59% | 58% |
| Schofield 1985 (Wt & Ht) (21) | 60% | 65% | 62% | 57% | 56% |
| Siervo 2003 (20) | 69% | 69% | 64% | 58% | 56% |
| Tabata 2012 (18) | 71% | 45% | 21% | 15% | 8% |
| Weijs 2010 (19) | 72% | 60% | 48% | 43% | 39% |
| WHO 1985 (Wt) (17) | 68% | 65% | 61% | 56% | 56% |
| WHO 1985 (Wt & Ht) (17) | 66% | 64% | 61% | 57% | 58% |
| Sabounchi (S1) 2013 (4) | 78% | 77% | 70% | 64% | 63% |

363 **Table 3** – Correct classification fraction of the REE equations, i.e. proportion of subjects
364 whose estimated resting energy expenditure was within 10% of measured resting energy
365 expenditure.

366

ACCEPTED MANUSCRIPT

367 REFERENCES

368


- 369 1. Shetty P. Energy requirements of adults. *Public Health Nutr* 2005;8:994-1009.
- 370 2. Madden AM, Mulrooney HM, Shah S. Estimation of energy expenditure using
371 prediction equations in overweight and obese adults: a systematic review. *J Hum*
372 *Nutr Diet* 2016;29:458-476.
- 373 3. Bedogni G, Grugni G, Tringali G, Agosti F, Sartorio A. Assessment of fat-free mass
374 from bioelectrical impedance analysis in obese women with Prader-Willi syndrome.
375 *Ann Hum Biol* 2015;42:538-542.
- 376 4. Sabounchi NS, Rahmandad H, Ammerman A. Best-fitting prediction equations for
377 basal metabolic rate: informing obesity interventions in diverse populations. *Int J*
378 *Obes (Lond)* 2013;37:1364-1370.
- 379 5. Flack KD, Siders WA, Johnson L, Roemmich JN. Cross-Validation of Resting
380 Metabolic Rate Prediction Equations. *J Acad Nutr Diet* 2016
- 381 6. Kaur Y, de Souza RJ, Gibson WT, Meyre D. A systematic review of genetic
382 syndromes with obesity. *Obes Rev* 2017;18:603-634.
- 383 7. McClave SA, Lowen CC, Kleber MJ, McConnell JW, Jung LY, Goldsmith LJ. Clinical
384 use of the respiratory quotient obtained from indirect calorimetry. *JPEN J Parenter*
385 *Enteral Nutr* 2003;27:21-26.
- 386 8. Lohman TG, Roche AF. *Anthropometric standardization reference manual*. Human
387 Kinetics Books: Champaign IL, 1988.
- 388 9. National Heart Lung and Blood Institute, North American Association for the Study of
389 Obesity. *The practical guide: identification, evaluation, and treatment of overweight*
390 *and obesity in adults*. National Heart, Lung, and Blood Institute: Bethesda, MD, 2000,
- 391 10. Weir JB. New methods for calculating metabolic rate with special reference to protein
392 metabolism. 1949. *Nutrition* 1990;6:213-221.

- 393 11. Lazzer S, Bedogni G, Lafortuna CL et al. Relationship between basal metabolic rate,
394 gender, age, and body composition in 8,780 white obese subjects. *Obesity*
395 2010;18:71-78.
- 396 12. Javed F, He Q, Davidson LE et al. Brain and high metabolic rate organ mass:
397 contributions to resting energy expenditure beyond fat-free mass. *Am J Clin Nutr*
398 2010;91:907-912.
- 399 13. Huang KC, Kormas N, Steinbeck K, Loughnan G, Caterson ID. Resting metabolic
400 rate in severely obese diabetic and nondiabetic subjects. *Obes Res* 2004;12:840-845.
- 401 14. Bernstein RS, Thornton JC, Yang MU et al. Prediction of the resting metabolic rate in
402 obese patients. *Am J Clin Nutr* 1983;37:595-602.
- 403 15. Lazzer S, Agosti F, Silvestri P, Derumeaux-Burel H, Sartorio A. Prediction of resting
404 energy expenditure in severely obese Italian women. *J Endocrinol Invest* 2007;30:20-
405 27.
- 406 16. Lazzer S, Agosti F, Resnik M, Marazzi N, Mornati D, Sartorio A. Prediction of resting
407 energy expenditure in severely obese Italian males. *J Endocrinol Invest* 2007;30:754-
408 761.
- 409 17. Joint FAO/WHO/UNU Expert Consultation on Energy and Protein Requirements.
410 *Energy and protein requirements: report of a Joint FAO WHO UNU Expert*
411 *Consultation*. World Health Organization: Geneva, 1985.
- 412 18. Tabata I, Ebine N, Kawashima Y et al. Dietary Reference Intakes for Japanese 2010:
413 Energy. *J Nutr Sci Vitaminol* 2012;59:S26-S35.
- 414 19. Weijs PJ, Vansant GA. Validity of predictive equations for resting energy expenditure
415 in Belgian normal weight to morbid obese women. *Clin Nutr* 2010;29:347-351.
- 416 20. Siervo M, Boschi V, Falconi C. Which REE prediction equation should we use in
417 normal-weight, overweight and obese women. *Clin Nutr* 2003;22:193-204.
- 418 21. Schofield WN. Predicting basal metabolic rate, new standards and review of previous


- 419 work. *Hum Nutr Clin Nutr* 1985;39 Suppl 1:5-41.
- 420 22. Roza AM, Shizgal HM. The Harris Benedict equation reevaluated: resting energy
421 requirements and the body cell mass. *Am J Clin Nutr* 1984;40:168-182.
- 422 23. Owen OE, Kaval E, Owen RS et al. A reappraisal of caloric requirements in healthy
423 women. *Am J Clin Nutr* 1986;44:1-19.
- 424 24. Muller MJ, Bosy-Westphal A, Klaus S et al. World Health Organization equations
425 have shortcomings for predicting resting energy expenditure in persons from a
426 modern, affluent population: generation of a new reference standard from a
427 retrospective analysis of a German database of resting energy expenditure. *Am J*
428 *Clin Nutr* 2004;80:1379-1390.
- 429 25. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive
430 equation for resting energy expenditure in healthy individuals. *Am J Clin Nutr*
431 1990;51:241-247.
- 432 26. Livingston EH, Kohlstadt I. Simplified resting metabolic rate-predicting formulas for
433 normal-sized and obese individuals. *Obes Res* 2005;13:1255-1262.
- 434 27. Korth O, Bosy-Westphal A, Zschoche P, Gluer CC, Heller M, Muller MJ. Influence of
435 methods used in body composition analysis on the prediction of resting energy
436 expenditure. *Eur J Clin Nutr* 2007;61:582-589.
- 437 28. Kleiber M. Body size and metabolism. *Hilgardia* 1932;11:315-353.
- 438 29. Ireton-Jones CS. Evaluation of energy expenditures in obese patients. *Nutr Clin Pract*
439 1989;4:127-129.
- 440 30. Henry CJ. Basal metabolic rate studies in humans: measurement and development of
441 new equations. *Public Health Nutr* 2005;8:1133-1152.
- 442 31. Harris J. *A biometric study of basal metabolism in man*. Carnegie institution of
443 Washington, 1919.
- 444 32. Ganpule AA, Tanaka S, Ishikawa-Takata K, Tabata I. Interindividual variability in

- 445 sleeping metabolic rate in Japanese subjects. *Eur J Clin Nutr* 2007;61:1256-1261.
- 446 33. Fredrix EW, Soeters PB, Deerenberg IM, Kester AD, von Meyenfeldt MF, Saris WH.
447 Resting and sleeping energy expenditure in the elderly. *Eur J Clin Nutr* 1990;44:741-
448 747.
- 449 34. de Luis DA, Aller R, Izaola O, Romero E. Prediction equation of resting energy
450 expenditure in an adult Spanish population of obese adult population. *Ann Nutr*
451 *Metab* 2006;50:193-196.
- 452 35. De Lorenzo A, Tagliabue A, Andreoli A, Testolin G, Comelli M, Deurenberg P.
453 Measured and predicted resting metabolic rate in Italian males and females, aged 18-
454 59 y. *Eur J Clin Nutr* 2001;55:208-214.
- 455 36. Owen OE, Holup JL, D'Alessio DA et al. A reappraisal of the caloric requirements of
456 men. *Am J Clin Nutr* 1987;46:875-885.
- 457 37. Agostoni C, Edefonti A, Calderini E et al. Accuracy of Prediction Formulae for the
458 Assessment of Resting Energy Expenditure in Hospitalized Children. *J Pediatr*
459 *Gastroenterol Nutr* 2016
460


Normal weight




Overweight




Class 1 obesity




Class 2 obesity




Class 3 obesity




Normal weight




Overweight



Class 1 obesity



Class 2 obesity



Class 3 obesity

