NMCD

Nutrition, Metabolism & Cardiovascular Diseases

An International Journal on Diabetes,
Atherosclerosis and Human Nutrition

Volume 18 Supplement 1 November 2008 ISSN 0939-4753

Official Journal of

- the Italian Society of Diabetology (SID)
- the Italian Society for the Study of Atherosclerosis (SISA)
- the Italian Society of Human Nutrition (SINU)

Abstracts of the XXII National Congress of the
Italian Society for the Study of Atherosclerosis (SISA)
19 - 22 November 2008, Rome, Italy

ABSTRACTS

Nutrition, Metabolism & Cardiovascular Diseases

www.elsevier.com/locate/nmcd

XXII National Congress of the Italian Society for the Study of Atherosclerosis (SISA), 19–22 November, Rome, Italy

1 DIAGNOSTIC AND CLINICAL ASPECTS OF CHILDREN AFFECTED BY PRIMARY DYSLIPIDEMIA

F. Abello, V. Baracco, O. Guardamagna. *Department of Pediatrics, University of Turin, Italy*

E-mail: progetto.child@unito.it

Objective: Dyslipidemia is one of the leading causes of death and cardiovascular morbidity in western countries. Some of these genetic disorders can be diagnosed in childhood through the evaluation of lipoprotein profile and DNA testing. Guidelines suggest the utility of an early selective screening in children/adolescents with a family history of dyslipidemia and/or premature cardiovascular disease. Aim of this study was to carry out the definite diagnosis of primitive lipoprotein disorders in children/adolescents who had been referred to our Lipid Clinic.

Methods: we studied 288 children (age 10 ± 3.37 years) with a family history of dyslipidemia and/or premature cardiovascular disease. On the basis of the first level investigation (TC, HDL-C, triglycerides) results, subjects were divided in 3 phenotype groups including IIa, IIb, IV and a further hypocholesterolemia group. Subsequently second level (apoB and apoA-I) and third level [Lipoprotein(a), ApoE gene polymorphism, DNA-based test and phytosterols dosage] investigations were performed.

Results: on the basis of the first level analysis, patients were so divided: 76.7% IIa, 15.3% IIb, 6.9% IV, 1% hypocholesterolemia. We reached a definite diagnosis of primitive dyslipidemia in 95% of tested patients through second and third level investigations. 30.9% children resulted affected by Familial Hypercholesterolemia (FH), 30.6% by Familial Combined Hyperlipidaemia (FCHL), 14.1% by dominant hypercholesterolemia, 13.9% by Hyperlipoproteinemia(a), 3.9% by Familial Hypertriglyceridemia (FHTG), 1% by Hypocholesterolemia due to mutations in ApoB and PCSK9 genes, 0.6% by Hyperchylomicronemia attributable to LPL and ApoC-II defects and 0.3% by Phytosterolemia.

Conclusion: this study remarks the importance of an early diagnosis of primitive lipoprotein disorders. When the suspect of primary dyslipidemia is supported by first level analysis it is advisable performing second and third level investigations in order to reach a definite diagnosis and check the cardiovascular risk. Combining the proband atherogenic profile and family history allow to establish an adequate and personalized treatment approach, based on lifestyle intervention and, if needed, dietary supplements and drugs since childhood/adolescence.

2 EFFECT OF PROBUCOL DERIVATE SUCCINOBUCOL (AGI-1067) ON CELLULAR CHOLESTEROL EFFLUX PATHWAYS

M.P. Adorni¹, R. Stocker², F. Bernini¹, E. Favari¹. ¹Dept. of Pharmacological and Biological Sciences and Applied Chemistry, University of Parma, Italy, ²Center for Vascular Research, School of Medical Sciences, University of New South Wales, Australia

E-mail: mariapia.adorni@nemo.unipr.it

Succinobucol (AGI-1067) is a metabolically stable monosuccinate ester derivative of probucol. In addition to its antioxidant effects, AGI-1067 modestly lowers lipid and HDL concentrations in animals compared to probucol, and it inhibits inflammation. Since probucol selectively inhibits ABCA1-mediated efflux of cellular cholesterol, the aim of this study was to compare the activity of AGI-1067 to probucol to modulate the cholesterol efflux process through the different pathways. Like probucol, AGI-1067 had no effect on either the passive diffusion of cholesterol in mouse peritoneal macrophages (MPM) or on SRBI-mediated cholesterol efflux in Fu5AH rat hepatoma cells. The ABCA1-mediated efflux of cholesterol to the specific acceptor apolipoprotein A-I was measured in MPM stimulated with an LXR-RXR agonist. A dose response curve showed AGI-1067 to be a stronger inhibitor of ABCA1-mediated efflux compared to probucol (99% and 24.5% respectively at 5 μ M). A similar result was obtained when MPM were treated with cAMP to enhance ABCA1 expression. Also, AGI-1067, but not probucol, inhibited efflux of cholesterol to HDL in MPM treated with an LXR-RXR agonist. Therefore, despite its reduced ability to decrease HDL levels in vivo, AGI-1067 is more active than probucol in inhibiting ABCA1-mediated cholesterol efflux and,

different to probucol, AGI-1067 seems to be less selective for ABCA1 pathway as it also inhibits ABCG1-mediated cholesterol efflux.

3 ARTERIAL STIFFNESS IN OBESE PATIENTS WITHOUT CARDIOVASCULAR RISK FACTORS

V. Alaimo, G. Cardella, E. Fertitta, M. Maggiore, L. Catalano, A. Falletta, C.M. Barbagallo, M.R. Averna. *Department of Clinical Medicine and Emerging Disease, University of Palermo, Italy*

E-mail: carlob@unipa.it

Despite relationship between obesity and cardiovascular disease are widely defined, mechanisms underlying vascular damage in obesity are less clear. In particular it is debated if weight gain might represent an independent condition able to predict atherosclerosis and clinical event or if other conditions could play a significant role. To furnish further insight in this topic, in this study we analyzed arterial stiffness parameters, as early marker of vascular damage, in a group of obese patients without cardiovascular risk factors. For this reason we selected a group of 12 non-smoking obese patients (BMI > 30 kg/m²), without cardiovascular disease, hypertension, diabetes or glucose intolerance (fasting glycemia < 110 mg/dL) or dyslipidemias. Exclusion criteria was also the presence of renal or liver disease, neoplasms or other conditions or drugs known to affect lipid metabolism. As controls we selected a group of 12 non obese healthy subjects with the same inclusion and exclusion criteria. In all subjects a fasting blood sample has been drawn for plasma lipid analysis and the analysis of arterial pulse wave has been done by applanation tonometry (SphygmoCor SPT-301), which is a non-invasive, bedside technology able to determine the Augmentation index (AI) and the velocity (PWV) of pulse wave. We found that obese subjects were significantly older and had systolic and diastolic blood pressure, total, LDL- and non-HDL cholesterol levels higher and HDL-cholesterol levels lower than controls. No significant difference was found for AI or PWV between obese patients and controls and no correlation was found between BMI and AI or PWV in all subjects. However AI was significantly positively correlated with age, triglyceride and total, LDL- and HDL-cholesterol levels, whereas PWV was positively correlated with diastolic blood pressure and negatively with HDL-cholesterol levels. In conclusion our study has preliminary demonstrated that obese patients without cardiovascular risk factors have no significant abnormality of arterial stiffness. These data, wherever further confirmed, reinforce the importance of early preventive tools, before the development of cardiovascular risk factors associated to obesity.

4 CHRONIC USE OF LIGHT OR REGULAR CIGARETTES AND FLOW MEDIATED DILATION

M. Amato¹, S. Castelnuovo², B. Frigerio², A. Ravani¹, E. Tremoli^{1,2}, C.R. Sirtori², D. Baldassarre^{1,2}. ¹Cardiologico "Monzino" Center, IRCCS, Milan and ²Dept of Pharmacological Sciences University Milan, Milan, Italy

E-mail: mauro.amato@ccfm.it

Objective: to compare the effect of "light" and "regular" cigarettes on brachial artery (BA) flow mediated dilation (FMD).

Methods: BA-FMD was evaluated in 45 light-cigarette consumers, in 45 regular-cigarettes consumers and in 45 never-smokers. The groups were matched for age, sex and status of hypercholesterolemia. FMD was calculated either considering the percent change of BA versus the "at-rest" diameter (FMDvs.at-rest) or considering the percent change versus the BA diameter measured in the last 60 seconds of cuff inflation (FMDvs.last-60"). Cigarettes were defined as "light" or "regular" according to the concomitant presence of tar and nicotine above or below 9 and 0.8 mg, respectively.

Results: In never-smokers, BA diameter increased during cuff inflation (change: 0.004 ± 0.012 ; $p = 0.023$). In smokers no changes in BA diameter during cuff inflation were observed (change: -0.001 ± 0.014 ; $p = 0.6$). In smokers, FMD was lower than in never smokers both considering the FMDvs.at-rest ($3.4 \pm 3.0\%$ vs. $6.3 \pm 3.2\%$; $p < 0.0001$) and the FMDvs.last-60" (3.5 ± 3.4 vs. 5.2 ± 3.3 respectively; $p = 0.007$). In the whole group, both FMDvs.at-rest and FMDvs.last-60" were inversely associated with pack-years (both p-trend < 0.0001). FMDvs.at-rest of both light ($2.78 \pm 3.13\%$) or regular ($4.03 \pm 2.93\%$) cigarette consumers were reduced with respect to

never-smokers ($6.26 \pm 3.21\%$) ($p < 0.0004$ and $p = 0.002$, respectively) but no differences were observed when the FMDvs.at-rest of light- and regular-cigarette consumers were compared ($p = 0.173$). Almost identical results were obtained when FMDvs.last-60'' were considered.

Conclusion: Light and regular cigarettes have the same detrimental effect of FMD, thus light cigarette consumption cannot be considered as a good alternative to smoking cessation.

Funding: Researches described were supported in part by Philip Morris USA Inc. and Philip Morris International.

5 CYTOKINE/CHEMOKINE DIFFERENTIATION PATTERN IN PATIENTS WITH ST-ELEVATION MYOCARDIAL INFARCTION (STEMI) ASSOCIATED WITH HIGH LEVELS OF CIRCULATING IL-6: EIGHTEEN CYTOKINES/CHEMOKINES ANALYZED SIMULTANEOUSLY WITH THE FLEX-SET CAPTURE BEAD ASSAY (CBA)

E. Ammirati, N. Cristell, V. Vecchio, A. Palini, A. Paganoni, A. Monello, C. Cannistraci, A. Durante, M. Banfi, D. Cianflone, A. Maseri. *Università Vita-Salute San Raffaele, Milano, Italy*
E-mail: durante.alessandro@gmail.com

Background: Preliminary analysis of the population enrolled in the international, case-controlled First Acute Myocardial Infarction (FAMI) study showed a 3-fold increase in circulating IL-6 in STEMI patients in comparison to controls.

Objectives: We assessed the possible clustering of patients' behavior by means of multiple biomarkers' measurements, in patients with very high levels or low levels of IL-6.

Materials and Methods: Eighteen cytokine/chemokines (IL-2, IL-4, IL-7, IL-1 β , IL-10, VEGF, FAS-L, GM-CSF, TNF α , IL-12p70, INF γ , IL-8, IP-10, MIP-1 α , MIP-1 β , eotaxin, MCP-1 and MIG) were analysed simultaneously with the Flex-Set CBA. Assays were performed on a total of 308 serum samples obtained within 6 hours from the onset of chest pain as follows: 109 patients belonging to the fourth interquartile (IQ) of IL-6 levels (IL6^{high} STEMI with IL-6 median 17.1 pg/ml, iq 11.9–29.8 pg/ml), 96 patients belonging to the first IQ of IL-6 levels (IL6^{low} STEMI with IL-6 median 3.7; 2.7–4.6 pg/ml) and 103 controls (IL-6 median 3.1; 2.3–5.2 pg/ml).

Results: The IL6^{high} STEMI versus IL6^{low} STEMI group showed increased levels of 6 out of 18 analyzed cytokines: IL-10 (5.3 vs 2.4 pg/ml; $p < 0.001$), IL-8 (10.5 vs 7.2 pg/ml; $p < 0.001$), MIP-1 α (4.4 vs 3.5 pg/ml; $p < 0.001$), MIP-1 β (86 vs 63 pg/ml; $p < 0.05$), MIG (160 vs 81 pg/ml; $p < 0.001$), MCP-1 (100 vs 68 pg/ml; $p < 0.001$). Similar levels of all cytokines were found in IL6^{low} STEMI and controls with the exception of IL-10 (2.4 vs 2.0 pg/ml; $p < 0.05$) which was increased in IL6^{low} STEMI, and MCP-1 (68 vs 92 pg/ml; $p < 0.05$) which was decreased in IL6^{low} STEMI. Independence analysis of cytokine distribution, taking into account the 6 cytokines and chemokines which were significantly different and the levels of IL-6 and C-reactive protein, showed a distinct cytokine patterns in the IL6^{low} and IL6^{high} STEMI groups ($P < 0.05$).

Conclusions: The multi-cytokine approach allowed us to detect an inflammatory pattern in IL6^{high} STEMI patients, consistent with the hypothesis that some patients with STEMI have a multifaceted inflammatory component, identified by high levels of circulating IL-6. This subgroup of patients might benefit by immune-modulating therapy.

6 CYTOKINE AND RECEPTOR PATTERN OF ACTIVATED VERSUS REGULATORY T CELLS IN CORONARY ARTERY DISEASES

E. Ammirati, A. Durante, V. Vecchio, A. Palini, M. Banfi, M. De Metrio, A. Monello, A.C. Vermi, A.A. Manfredi, D. Cianflone, A. Maseri. *Università Vita-Salute San Raffaele, Milano, Italy*
E-mail: durante.alessandro@gmail.com

Background: Regulatory T lymphocytes (Treg) play a key role in maintaining self tolerance and in suppression of pathological immune response. In atherosclerosis an unbalance between activated and regulatory T-cell activity might play a role in the development, evolution and instability of plaques. In murine models a reduction in Treg results in increased atherogenesis.

Aims of the study: The aim of the study is to evaluate the balance between circulating activated and regulatory T cells in various manifestations of coronary artery diseases (CAD) and the receptor patterns of these cells. The cytokine profile in its activation (IL-6)/regulation (IL-10) balance as expressed by changes in IL-6 and IL-10 was also evaluated.

Materials and Methods: Treg percentage, identified by CD3+CD4dimCD25high panel within the CD3+CD4+ T-cell compartment, was assessed in the study by means of 4- and 5-colour flow cytometry with conjugate antibodies (CD3, CD4, CD25, CD69 and CCR5 the 5-colour panel; CD3, CD4, CCR5 and HLA-DR the 4-colour panel). Circulating levels of cytokines were assessed by the CBA-Flexset technology. We studied four groups of patients: 49 healthy controls with no evidence of coronary disease, 36 patients with Chronic Stable Angina (CSA), 34 patients with Unstable Angina/Non ST Elevation Myocardial Infarction (NSTEMI) and 29 patients with ST Elevation Myocardial Infarction (STEMI). The state of activation of CD4+ T cells was determined by measuring two different markers: CD69, a marker of early activation, and HLA-DR, a marker of late activation. The reliability of the Treg percentage was assessed by two different measurements at 1 and 3 months in patients and controls, respectively. All values are expressed as medians (25%-75% pct).

Results: The percentage of Treg was stable over time. Treg percentage was significantly increased in STEMI (6.2, 5.4–7.5) vs controls (5.4, 4.6–6.2), vs CSA (5.3, 4.5–6.4) and vs NSTEMI (4.7, 4.4–5.7); while in NSTEMI a not significant trend towards an overall reduction in Treg number was observed. The expression of CD69 was higher in STEMI (1.0, 0.7–1.8) vs NSTEMI (0.5, 0.7–1.4), $p < 0.05$. A significantly increased expression of HLA-DR in NSTEMI (9.2, 5.7–12.2) and STEMI (5.3, 2.4–6.5) was observed vs controls (2.4, 1.7–4.0) and vs CSA (3.2, 1.5–6.8), $p < 0.0001$. The balance between activation and regulation, expressed as the ratio of Treg and activation markers was altered in CAD groups showing an increased activation of T-cell response. Moreover a decreased CCR5 Treg expression was observed in STEMI (20.7, 13.5–32.8) vs controls (31.3, 23.3–39.2), $p < 0.05$; These data seem to indicate a decreased tissutal migration ability since CCR5 represents a chemokine receptor for Treg homing. We observed a 2-fold increase in IL-10 levels in STEMI (4.1 pg/ml, 2.4–12.7) vs controls (2.0 pg/ml, 1.6–2.2) and vs NSTEMI (2.0 pg/ml, 1.7–2.9), both $p < 0.001$. IL-6 showed an increased serum concentration both in NSTEMI (6.9, 4.5–16.5) and STEMI (6.7, 4.2–24.9) vs controls (3.1, 2.5–4.5), $p < 0.001$ for all. IL-10/IL-6 ratio, used to determine the cytokine suppression/activation balance, was decreased in NSTEMI (0.7, 0.3–1.0) vs controls (0.6, 0.4–1.2), vs CSA (0.5, 0.3–0.8) and vs STEMI (0.7, 0.3–1.0), $p < 0.001$ for all.

Conclusions: This study showed a pro-inflammatory unbalance both in T-cell response and in cytokine network in all CAD groups as compared to controls. These findings could help understanding a possible protective role of Treg in patients with various manifestations of coronary artery diseases. The variability we observed so far may reflect the complexity of the pro- and anti-inflammatory network which still needs to be explored further.

7 EXPANSION OF T-CELL RECEPTOR γ^{dim} EFFECTOR T CELLS IN ACUTE CORONARY SYNDROMES

E. Ammirati, A.C. Vermi, D. Cianflone, M. Banfi, C. Foglieni, C.L. Gorman, A. Durante, A. Monello, A. Maseri, A.P. Cope, C. Monaco. *Università Vita-Salute San Raffaele, Milano, Italy*
E-mail: durante.alessandro@gmail.com

Objective: The T-cell receptor zeta (TCR ζ)-chain is a master sensor and regulator of lymphocyte responses. Loss of TCR ζ -chain expression has been documented during infectious and inflammatory diseases and defines a population of effector T cells (TCR ζ^{dim} cells) that migrate to inflamed tissues. We assessed the expression and functional correlates of circulating TCR ζ^{dim} cells in coronary artery disease.

Methods and Results: We examined the expression of TCR ζ -chain by flow cytometry in 140 subjects. Increased peripheral blood CD4⁺TCR ζ^{dim} cells were found in patients with acute coronary syndromes (ACS, $n = 66$; median 5.3%, inter-quartile 2.6–9.1% of total CD4⁺ cells; $p < 0.0001$) compared to chronic stable angina (CSA, $n = 32$; 1.6%; 1.0–4.1%) and controls ($n = 42$; 1.5%; 0.5–2.9%). Such increase was significantly greater in ACS patients with elevated levels of C-reactive protein (CRP ≥ 2 mg/l: 7.7, 3.8–11.3%, $n = 41$) compared to patients with ACS and low CRP levels (< 2 mg/l: 3.3; 1.7–7.7%, $n = 25$), $p = 0.003$, and it persisted after the acute event. Moreover, TCR ζ^{dim} cells were also more represented in ACS compared to CSA and controls within CD8⁺T cell subset (4.0%, 2.5–7.5% vs 1.2%, 0.8–2.1% vs 0.6%, 0.2–1.9%; $p < 0.0001$), NK subset (5.1%, 2.4–7.5% vs 1.6%, 0.9–2.7% vs 0.5%, 1.3–3.8%; $p = 0.001$) and CD4⁺CD28null subset (67.2, 39.4–82.9% vs 29.2, 23.0–46.9% vs 20.9, 11.1–33.1%; $p < 0.0001$). Finally, CD4⁺ and CD8⁺TCR ζ^{dim} cells isolated from ACS displayed an enhanced transendothelial migratory capacity.

Conclusions: TCR ζ^{dim} cells, an effector T-cell subset with trans-endothelial migratory ability, are increased in ACS, and may be implicated in coronary instability.

8 THE METABOLIC SYNDROME DOES NOT ADD TO CAROTID ATHEROSCLEROSIS BEYOND THAT EXPECTED BY RISK FACTOR COUNTING OR RISK SCORING

D. Baldassarre^{1,2}, S. Castelnuovo², B. Frigerio², M. Amato¹, A. Ravani¹, E. Tremoli^{1,2}, J.P. Werba¹. ¹Cardiologic Monzino Center IRCCS, Milan; ²Dept of Pharmacological Sciences, Milan University, Italy
E-mail: damiano.baldassarre@unimi.it

Objective: The aim of this study was to assess if the Metabolic Syndrome (MS) has any add-on effect on subclinical atherosclerosis beyond that expected by risk factors (RF) counting or risk scoring.

Methods: Intima-media thickness (IMT) of carotid arteries was assessed by using B-mode ultrasound in 1805 patients (56 $\pm$ 13 y; 52% women) attending a cardiovascular prevention program. Patients with (cases) or without (controls) MS according to NCEP ATP III criteria were 1:1 matched for sex, age and either the number of conventional RF (Analysis 1) or the Framingham risk score (Analysis 2). For Analysis 1 not more than 2 components of the MS were accepted as RF in the control group.

Results: Case:control matches were 211 for Analysis 1 and 244 for Analysis 2. No significant differences in carotid IMTmean and carotid IMTmax were found between cases and controls in both analyses (Analysis 1: IMTmean 1.03 ± 0.38 vs 1.07 ± 0.37 ; IMTmax 1.90 ± 0.96 vs 1.95 ± 0.90 , cases and controls, respectively; Analysis 2: IMTmean 1.03 ± 0.36 vs 1.01 ± 0.33 ; IMTmax 1.91 ± 0.94 vs 1.83 ± 0.81 , cases and controls, respectively; all $p > 0.1$).

Conclusions: According to our results the metabolic syndrome does not add to the extent of carotid subclinical atherosclerosis beyond that expected by RF