

BAKER'S YEAST HYDROGENATION OF CARBONYL ACTIVATED DOUBLE BONDS.
 ENANTIOSELECTIVE SYNTHESIS OF THE (S)-FORM OF THE DIHYDROTERPENEDIOL SECRETED BY
DANAUS CHRYSIPPUS AND OF A PHEROMONE OF CALLOSOBRUCHUS CHINENSIS L.

Paola GRAMATICA,* Giuseppe GIARDINA, Giovanna SPERANZA,[†] and Paolo MANITTO*

Dipartimento di Chimica Organica e Industriale, Università di Milano and Centro
 per lo Studio delle Sostanze Organiche Naturali del CNR, Via Venezian 21, 20133-
 Milano, Italy

[†]Centro Studi Maria Branca, Via E. Porro 1, 20158-Milano, Italy


The (S)-forms of (E)-3,7-dimethyl-2-octene-1,8-diol (secreted by
 male danaid butterflies) and of (E)-3,7-dimethyl-2-octene-1,8-dioic
 acid (a pheromone of the azuki bean weevil) were synthesized via
 stereoselective hydrogenation by baker's yeast of the carbonyl
 activated double bond of achiral precursors.

(E)-3,7-Dimethyl-2-octene-1,8-diol (dihydroterpenediol, 1) was isolated by
 Meinwald et al. as one of the major components of the hairpencil secretion of male
 danaid butterflies (Danaus chrysippus, commonly called "the African Monarch");¹⁾
 its absolute configuration, however, remained unknown. To our knowledge, no report
 has so far been published on the preparation of 1 in one enantiomerically pure
 form,²⁾ although a number of syntheses have been performed giving racemic 1.^{1,3)}

In this paper we report two independent syntheses of (S)-1, both based on the
 stereoselective reduction by baker's yeast of a carbonyl activated double bond.

Previous findings that α -methyl- α,β -unsaturated aldehydes (alcohols or acetals)
 are reduced by Saccharomyces cerevisiae to give the corresponding α -methyl saturat-
 ed alcohols⁴⁾ prompted us to utilize such a biological hydrogenation in order to
 obtain enantiomerically pure (S)-1.⁵⁾ from 8-acetoxy-2,6-dimethylocta-2(E),6(E)-di-
 enal (4). This compound was in turn prepared from geraniol (2), an achiral and
 readily available material, through the acetylation (Ac₂O in dry Py) and subse-
 quent oxidation of the resulting 3 with SeO₂ in EtOH⁶⁾ (36% overall yield from
2).^{7,8)} The protection of the allylic hydroxyl group in 4 was retained to avoid
 the hydrogenation of the adjacent double bond, which occurs in the yeast catalysed
 conversion of geraniol (2) into (R)-(+)-citronellol (9).⁹⁾

Reduction of 4 by baker's yeast (3 g educt, 1 kg yeast, 3 l H₂O, 3 d at r.t.),
 followed by ether extraction, afforded a crude product, which was then treated
 with alcoholic KOH (5%) for 4 h at 40 °C. After flash chromatography (silica gel-
 AgNO₃ 5%; hexane:EtOAc 1:2 as eluent) (S)-1¹⁰⁾ was isolated in 20% yield.¹¹⁾
 $[\alpha]_D^{25} -9.1^\circ$ (c 4.6, CHCl₃); e.e. > 97% as estimated by a comparison of its rotatory
 power with that of (S)-1 prepared from 7 as described below (see Scheme 1).



Scheme 1.

Methyl-(S)-8-hydroxy-3,7-dimethyl-2(E)-octenoate (7) was obtained by baker's yeast hydrogenation of 6¹²⁾ (fermentation conditions as mentioned above; ether extraction; flash chromatography silica gel-AgNO₃ 10%, hexane:EtOAc 1:1 as eluent; 35% yield):¹¹⁾ $[\alpha]_D^{21}$ -10.1° (c 2.4, CHCl₃); IR (liquid film): 3420, 1715, 1650 cm⁻¹; ¹H-NMR (CDCl₃, δ, TMS): 0.9 (2H, d, J=6 Hz, 7-CH₃), 1.0-2.2 (7H, m, CH₂ and CH), 2.18 (3H, d, J=1 Hz, 3-CH₃), 2.28 (1H, s, OH), 3.44 (2H, d, J=6 Hz, CH₂OH), 3.65 (3H, s, OCH₃), 5.65 (1H, q, J=1 Hz, =CH); ¹³C-NMR (CDCl₃, δ, TMS): 16.4 (7-CH₃), 18.6 (3-CH₃), 24.8 (C-5), 32.7 (C-6), 35.6 (C-7), 41.0 (C-4), 50.6 (OCH₃), 68.0 (C-8), 115.1 (C-2), 159.9 (C-3), 167.0 (C-1).


The conversion (PDC oxidation in DMF¹³⁾ followed by HCl-hydrolysis of the mono-methyl ester) of **7** into the diacid (S)-**8** (65% yield, $[\alpha]_D^{21} +11.2^\circ$ (c 0.66, CHCl₃)),¹⁴⁾ whose absolute configuration and optical activity were interrelated by Mori *et al.*¹⁵⁾ through unambiguous asymmetric synthesis and accurate estimation of the e.e. of the product, clearly defines the (S)-configuration and the enantiomeric excess (> 97%) of **7**.


LiAlH₄ reduction of **7** gave (S)-**1** (82% yield; $[\alpha]_D^{25} -8.9^\circ$ (c 1.55, CHCl₃)).¹⁶⁾

The diacid **8** is also known as callosobruchusic acid in that it is a component of the copulation release pheromone (erectin) of the azuki bean weevil (*Callosobruchus chinensis* L.).¹⁷⁾ The absolute configuration of the naturally occurring **8** is still unknown. Although both enantiomers of **8** were found to be biologically active, the potency of the (S)-form appeared higher than that of its antipode and significantly closer to that of the natural pheromone.¹⁵⁾

We thank Dr. Paola Gambini for technical assistance. We gratefully acknowledge the National Research Council (CNR), Rome - Progetto Finalizzato "Chimica Fine e Secondaria", and Curt Georgi Imes (Sesto S.Giovanni, Italy) for financial supports.

References

- 1) J. Meinwald, W.R. Thompson, T. Eisner, and D.F. Owen, *Tetrahedron Lett.*, **1971**, 3485.
- 2) The only attempt to obtain optically active **1** by asymmetric hydroboration of:
 with isopinocampheylborane (from B₂H₆ and (-)-α-pinene) afforded a product having $[\alpha]_D^{20} +2.2^\circ$, which appears to be the (R)-isomer (e.e. ca. 23% from the data of the present paper): M. Julia and J.N. Verpeaux, *Tetrahedron*, **39**, 3289 (1983).
- 3) a) J.P. Morizur, G. Bidan, and J. Kossanyi, *Tetrahedron Lett.*, **1975**, 4167; b) G. Bidan, J. Kossanyi, V. Meyer, and J.P. Morizur, *Tetrahedron*, **33**, 2193 (1977); c) T. Fujisawa, T. Sato, T. Kawara, and A. Noda, *Tetrahedron Lett.*, **23**, 3193 (1982).
- 4) a) C. Fuganti, D. Ghiringhelli, and P. Grasselli, *J. Chem. Soc., Chem. Commun.*, **1975**, 846; b) M.G.W. Leuenberger, W. Boguth, R. Barner, M. Schmid, and R. Zell, *Helv. Chim. Acta*, **62**, 455 (1979); c) C. Fuganti and P. Grasselli, *J. Chem. Soc., Chem. Commun.*, **1979**, 995; d) P. Gramatica, B.M. Ranzi, and P. Manitto, *Bioorg. Chem.*, **10**, 22 (1981); e) P. Gramatica, P. Manitto, and L. Poli, results to be published concerning the microbial conversion of:



- 5) It must be pointed out that, in all cases examined from a stereochemical point of view,^{4b,c,e)} the hydrogenation of α-methyl-α,β-unsaturated aldehydes by baker's yeast was shown to give the (S)-configuration at C-α of the resulting alcohols.

- 6) This reaction is known to be stereoselective: U.T. Bhalerao and H. Rapoport, *J. Am. Chem. Soc.*, **93**, 4835 (1971).
- 7) All compounds reported gave satisfactory spectroscopic and analytical data.
- 8) 4: IR (liquid film): 1730, 1680, 1640 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3 , δ , TMS): 1.8 (6H, s, $2\text{CH}_3\text{-C=}$), 2.1 (3H, s, CH_3CO), 2.0-2.8 (4H, m, CH_2), 4.6 (2H, d, $J=6$ Hz, $\text{CH}_2\text{-OAc}$), 5.4 (1H, t, $J=6$ Hz, 7-H), 6.45 (1H, t, $J=6$ Hz, 3-H), 9.4 (1H, s, (E)-CHO); $^{13}\text{C-NMR}$ (CDCl_3 , δ , TMS): 9.1 (2- CH_3), 16.3 (6- CH_3), 20.8 (CH_3CO), 27.0 (C-4), 37.7 (C-5), 60.9 (C-8), 119.7 (C-7), 139.6 (C-2), 140.0 (C-6), 152.8 (C-3), 170.4 (CH_3CO), 194.4 (C-1).
- 9) P. Gramatica, P. Manitto, B.M. Ranzi, A. Delbianco, and M. Francavilla, *Experientia*, **38**, 775 (1982).
- 10) (S)-1: IR and $^1\text{H-NMR}$ spectra were identical with those reported in Refs.1, 3b,c for racemic 1. $^{13}\text{C-NMR}$ (CDCl_3 , δ , TMS): 16.1 (3- CH_3), 16.5 (7- CH_3), 24.9 (C-5), 32.6 (C-6), 35.5 (C-7), 39.6 (C-4), 60.8 (C-1), 67.9 (C-8), 123.5 (C-2), 138.9 (C-3). Its physiological activities remain to be clarified.
- 11) The rather low yields are likely due to an extensive degradation of the starting material.
- 12) Methyl geraniate was converted into 6 by SeO_2 oxidation according to Ref.6 (32% yield): IR (liquid film): 1715, 1680, 1640 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3 , δ , TMS): 1.8 (3H, d, $J=1$ Hz, 7- CH_3), 2.2 (3H, d, $J=1$ Hz, 3- CH_3), 2.0-2.8 (4H, m, 2CH_2), 3.75 (3H, s, OCH_3), 5.8 (1H, q, $J=1$ Hz, 2-H), 6.45 (1H, tq, $J=6$ and 1 Hz, 6-H), 9.4 (1H, s, (E)-CHO). $^{13}\text{C-NMR}$ (CDCl_3 , δ , TMS): 9.23 (7- CH_3), 18.6 (3- CH_3), 26.7 (C-5), 39.0 (C-4), 50.7 (OCH_3), 116.1 (C-2), 139.9 (C-7), 151.7 (C-6), 157.4 (C-3), 166.5 (C-1), 194.4 (C-8).
- 13) E.J. Corey and G. Schmidt, *Tetrahedron Lett.*, **1979**, 399.
- 14) Lit,¹⁵⁾ $[\alpha]_D^{21} +11.4^\circ$ (c 0.10, CHCl_3), calcd. assuming 100% e.e. Analytical and spectroscopic data of (S)-8 were identical with those reported by Mori.¹⁵⁾
- 15) K. Mori, T. Ito, K. Tanaka, H. Honda, and I. Yamamoto, *Tetrahedron*, **39**, 2303 (1983).
- 16) It may be pointed out that rotations observed for the (S)-form of 1, 7 (levo) and 8 (dextro) are in agreement with the empirical rules relating the sign of rotation at the sodium D line to the absolute configuration in the homologous series $\text{Me-(CH}_2)_n\text{-CHMe-CH}_2\text{OH}$ and $\text{Me-(CH}_2)_n\text{-CHMe-CO}_2\text{H}$ where $n \geq 1$; cf.: W. Kline and J. Buckingham, "Atlas of Stereochemistry," 2nd ed, Chapman and Hall, London (1978), Vol.1 pp. 64-65; P.A. Levene and A. Rothen, *J. Org. Chem.*, **1**, 76 (1936).
- 17) K. Tanaka, K. Ohsawa, H. Honda, and I. Yamamoto, *J. Pesticide Sci.*, **6**, 75 (1981).
- 18) E.J. Corey, N.W. Gilman, and B.E. Ganem, *J. Am. Chem. Soc.*, **90**, 5616 (1968).

(Received May 7, 1985)