



CERN-EP-2018-210
LHCb-PAPER-2018-030
12 December 2018

Search for lepton-flavour-violating decays of Higgs-like bosons

LHCb collaboration[†]

Abstract

A search is presented for a Higgs-like boson with mass in the range 45 to 195 GeV/c^2 decaying into a muon and a tau lepton. The dataset consists of proton-proton interactions at a centre-of-mass energy of 8 TeV, collected by the LHCb experiment, corresponding to an integrated luminosity of 2 fb^{-1} . The tau leptons are reconstructed in both leptonic and hadronic decay channels. An upper limit on the production cross-section multiplied by the branching fraction at 95% confidence level is set and ranges from 22 pb for a boson mass of 45 GeV/c^2 to 4 pb for a mass of 195 GeV/c^2 .

Published in Eur. Phys. J. C78 (2018) 1008

© 2018 CERN for the benefit of the LHCb collaboration. CC-BY-4.0 licence.

[†]Authors are listed at the end of this paper.

1 Introduction

Decays mediated by charged-lepton flavour-violating (CLFV) processes are forbidden in the Standard Model (SM). Their observation would be a clear sign for physics beyond the SM. Such processes are predicted by several theoretical models [1–8], in particular those based on an effective theory with relaxed renormalisability requirements [9], supersymmetric models [10–14], composite Higgs models [15, 16], Randall-Sundrum models [17, 18], and non-abelian flavour symmetry models [19]. Nonetheless, no evidence for CLFV effects has been reported to date.

The LEP experiments set stringent limits on the CLFV decay of the Z boson [20–23]. In the presence of CLFV couplings, the decays to $e^\pm\mu^\mp$, $e^\pm\tau^\mp$ and $\mu^\mp\tau^\mp$ could be mediated by a Higgs boson. At LEP2, limits on the cross-section of the $e^+e^- \rightarrow e^\pm\mu^\mp$, $e^+e^- \rightarrow e^\pm\tau^\mp$ and $e^+e^- \rightarrow \mu^\pm\tau^\mp$ processes were obtained by the OPAL collaboration for centre-of-mass energies (\sqrt{s}) ranging from 192 to 209 GeV [24]. These constraints can be translated into limits on the Higgs CLFV decay branching fraction [9, 25], which are on the order of 10^{-8} for a SM Higgs decay into an electron and muon [25]. Recent searches for the $H \rightarrow \mu^\pm\tau^\mp$ decay have been performed by the CMS [26] and ATLAS [27] collaborations for the Higgs boson with $m_H = 125 \text{ GeV}/c^2$. Upper limits on the branching fraction $\mathcal{B}(H \rightarrow \mu^\pm\tau^\mp)$ have been placed by the two collaborations at 0.25% and 1.85%, respectively.

The possible existence of low-mass Higgs-like bosons is a feature of models like the two-Higgs-doublet models (2HDM) [28]. Searches for such particles have been performed by the ATLAS [29] and CMS [30] collaborations in the ditau decay mode. Another scenario is that of a hidden gauge sector [31, 32]. In this context, the BaBar and Belle collaborations have performed searches for a resonance with a mass below $10 \text{ GeV}/c^2$ [33, 34]. The LHCb collaboration has recently published the results of a search for dark photons decaying into the dimuon channel, placing a stringent limit for the production of a dimuon in the mass range from 10.6 to 70 GeV/c^2 [35].

The LHCb detector probes the forward rapidity region which is only partially covered by the other LHC experiments, and triggers on particles with low transverse momenta (p_T), allowing the experiment to explore relatively small boson masses. In this paper a search for CLFV decays into a muon and a tau lepton of a Higgs-like boson with a mass ranging from 45 to 195 GeV/c^2 is presented, using proton-proton collision data collected at $\sqrt{s} = 8 \text{ TeV}$. The Higgs-like boson is assumed to be produced by gluon-fusion, similarly to the main production mechanism of the SM Higgs boson at LHC [36].¹ The analysis is separated into four channels depending on the final state of the τ lepton decay: (i) single muon $\tau^- \rightarrow \mu^-\bar{\nu}_\mu\nu_\tau$, (ii) single electron $\tau^- \rightarrow e^-\bar{\nu}_e\nu_\tau$, (iii) single charged hadron $\tau^- \rightarrow \pi^-(\pi^0)\nu_\tau$, and (iv) three charged hadrons $\tau^- \rightarrow \pi^-\pi^-\pi^+(\pi^0)\nu_\tau$. They are denoted as τ_μ , τ_e , τ_{h1} , and τ_{h3} respectively. The main sources of background are $Z \rightarrow \tau^+\tau^-$ decays,² heavy flavour production from QCD processes (“QCD” in the following) and electroweak boson production accompanied by jets (“ Vj ”). This analysis utilizes reconstruction techniques and results obtained from the $Z \rightarrow \tau^+\tau^-$ measurement by the LHCb collaboration [37].

¹The remaining Higgs production modes (*e.g.*, $\sim 10\%$ from Vector-Boson Fusion) are neglected in this study.

²Throughout this note, Z implies Z/γ^* , *i.e.* includes contributions from Z boson production, virtual photon production, and also their interference.

2 Detector and simulation description

The LHCb detector [38, 39] is a single-arm forward spectrometer covering the $2 < \eta < 5$ pseudorapidity range, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream of the magnet. The tracking system provides a measurement of the momentum of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at $200\text{ GeV}/c$. The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T)\mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/c . Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad (SPD) and preshower detectors (PS), an electromagnetic calorimeter (ECAL) and a hadronic calorimeter (HCAL). Muons are identified by a system composed of five stations of alternating layers of iron and multiwire proportional chambers.

Simulated data samples are used to calculate the efficiency for selecting signal processes, to estimate the residual background level, and to produce templates for the fit used to determine the signal yield. For this analysis, the simulation is validated primarily by comparing $Z \rightarrow l^+l^-$ decays in simulation and data. The Higgs boson is generated assuming a gluon-fusion process, and with mass values from 45 to $195\text{ GeV}/c^2$ in steps of $10\text{ GeV}/c^2$, using PYTHIA 8 [40] with a specific LHCb configuration [41]. The parton density functions (PDF) are taken from the CTEQ6L set [42]. Decays of hadronic particles are described by EVTGEN [43], in which final-state radiation is generated using PHOTOS [44]. The interaction of the particles with the detector and its response are implemented using the GEANT4 toolkit [45] as described in Ref. [46]. Samples of $H \rightarrow \mu^\pm\tau^\mp$ decays generated at next-to-leading order precision by POWHEG-BOX [47–50] with the PDF set MMHT2014nlo68c1 [51] are used for the signal acceptance determination.

3 Signal selection

This analysis uses data corresponding to a total integrated luminosity of $1976 \pm 23\text{ pb}^{-1}$ [52]. The data collected uses a trigger system consisting of a hardware stage followed by a software stage. The hardware trigger requires a muon track identified by matching hits in the muon stations, as well as a global event cut (GEC) requiring the hit multiplicity in the SPD to be less than 600. The software trigger selects muons or electrons with a minimum p_T of $15\text{ GeV}/c$.

The $H \rightarrow \mu^\pm\tau^\mp$ candidates are identified and reconstructed into the four channels: $\mu\tau_e$, $\mu\tau_{h1}$, $\mu\tau_{h3}$ and $\mu\tau_\mu$. The τ_{h3} candidates are reconstructed from the combination of three charged hadrons from a secondary vertex (SV). The $\mu^\pm\tau^\mp$ candidates are required to be compatible with originating from a common PV. The muon track and the tracks used to reconstruct the tau candidate must be in the geometrical region $2.0 < \eta < 4.5$. Electron candidates are chosen amongst tracks failing the muon identification criteria and falling into the acceptance of the PS, ECAL, and HCAL sub-detectors. A large energy deposit, E , in the PS, ECAL, but not in HCAL is required, satisfying: $E_{\text{PS}} > 50\text{ MeV}$, $E_{\text{ECAL}}/p > 0.1$,

and $E_{\text{HCAL}}/p < 0.05$, where p is the reconstructed momentum of the electron candidate, after recovering the energy of the bremsstrahlung photons [53]. Charged hadrons are required to be in the HCAL acceptance, to deposit an energy E_{HCAL} with $E_{\text{HCAL}}/p > 0.05$, and to fail the muon identification criteria. The pion mass is assigned to all charged hadrons.

The selection criteria need to be optimised over the m_H range used in this analysis, from 45 to 195 GeV/c^2 . Three different sets of selection criteria are considered, dubbed L-selection, C-selection, and H-selection. The C-selection is similar to that used for the analysis of $Z \rightarrow \tau^+ \tau^-$ decays [37]; as such, it is optimised for $m_H \sim m_Z$. The L-selection and H-selection are optimised for the m_H regions below and above the Z mass respectively. All selection sets are applied in parallel to compute background estimation and exclusion limits. Subsequently, for each m_H hypothesis, the chosen selection is that of L-, C-, or H-selection which provides the smallest expected signal limit, allowing precise separation between adjacent mass regions. As expected, it is found that the C-selection is optimal for a boson mass of 75 and 85 GeV/c^2 . Below and above that range the best upper limits are obtained from the L- and H-selections, respectively. In the following discussion the requirements are applied identically for all decay channels and selection sets unless stated otherwise.

The tau candidates are selected with $p_T > 5 \text{ GeV}/c$ for τ_e, τ_μ , and $p_T > 10 \text{ GeV}/c$ for τ_{h1} . For the τ_{h3} candidate, the charged hadrons are required to have $p_T > 1 \text{ GeV}/c$ and one of them with $p_T > 6 \text{ GeV}/c$. They are combined to form the tau candidates, which are required to have $p_T > 12 \text{ GeV}/c$ and an invariant mass in the range 0.7 to 1.5 GeV/c^2 . In the H-selection, the tau candidates must have p_T in excess of 20 GeV/c . This requirement is not applied in the $\mu\tau_\mu$ channel as it favours the selection of $Z \rightarrow \mu^+ \mu^-$ background. The muon from $H \rightarrow \mu^\pm \tau^\mp$ decay is expected to have a relatively large p_T , thus the selection requires the muon p_T to be greater than 20 GeV/c , 30 GeV/c , and 40 GeV/c in the L-, C-, and H-selections, respectively. A tighter requirement of 50 GeV/c is applied for the muon in the $\mu\tau_\mu$ channel in the H-selection due to the $Z \rightarrow \mu^+ \mu^-$ background. Additionally, for the $\mu\tau_e$ channel, the contribution from $W/Z \rightarrow e + \text{jet}$ background is suppressed by requiring the transverse momentum of the muon to be larger than that of the τ_e candidate.

The relatively large lifetime of the τ lepton is used to suppress prompt background. For the τ_{h3} candidate, a SV is reconstructed. A correction to the visible invariant mass, m , computed from the three-track combination, is obtained by exploiting the direction of flight defined from the PV to the SV. The relation used is $m_{\text{corr}} = \sqrt{m^2 + p^2 \sin^2 \theta} + p \sin \theta$, where θ is the angle between the momentum of the τ_{h3} candidate, and its flight direction. The m_{corr} value is required to not exceed 3 GeV/c^2 . A time-of-flight variable is also computed from the distance of flight and the partially reconstructed momentum of the τ lepton, and a minimum value of 30 fs is required. The m_{corr} and time-of-flight requirements together retain 80% of the signal, while rejecting about 75% of the QCD background. For tau decay channels with a single charged particle, it is not possible to reconstruct a SV, and a selection on the particle IP is applied. A threshold of $\text{IP} > 10 \mu\text{m}$ selects 85% of the τ_e and τ_{h1} candidates, and rejects about 50% of the Vj background. The threshold is increased to 50 μm for τ_μ candidates, in order to suppress $Z \rightarrow \mu^+ \mu^-$ background. The prompt muon instead is selected by requiring IP less than 50 μm , allowing up to 50% rejection of QCD and $Z \rightarrow \tau^+ \tau^-$ backgrounds.

The two leptons from the Higgs decay should be approximately back-to-back in the plane transverse to the beam. The absolute difference in azimuthal angle of muon and

tau candidates is required to be greater than 2.7 radians. This rejects 50% of the Vj background. The transverse momentum asymmetry of the two particles, defined as $A_{p_T} = |p_{T1} - p_{T2}|/(p_{T1} + p_{T2})$, can be used to effectively suppress various background processes. The background from the Vj processes is suppressed by up to 60% for the $\mu\tau_h$ channel by requiring $A_{p_T} < 0.4$ (0.5) in the L-selection (S-selection), because of the large p_T imbalance between the high- p_T muon from the vector boson and a hadron from a jet. For the $\mu\tau_e$ channel, the worse momentum resolution increases the average A_{p_T} value, hence a softer selection $A_{p_T} < 0.6$ is used to preserve efficiency. On the contrary, for the $\mu\tau_\mu$ channel, a tighter cut is applied to suppress the dominant background from $Z \rightarrow \mu^+\mu^-$ decays. By requiring $A_{p_T} > 0.3$ (0.4) in the L-selection and C-selection (H-selection), such background is reduced by 80%, while the signal decreases to 70%.

The two leptons from the Higgs decay are required to be isolated from other charged particles. Two particle-isolation variables are defined as $I_{p_T} = (\vec{p}_{\text{cone}})_T$ and $\hat{I}_{p_T} = p_T / (\vec{p} + \vec{p}_{\text{cone}})_T$ where \vec{p} is the momentum of the lepton candidate, the subscript T denotes the component in the transverse plane, and \vec{p}_{cone} is the sum of the momenta of all charged tracks within a distance $R_{\eta\phi} = 0.5$ in the (η, ϕ) plane around the lepton candidate. The isolation requirement $\hat{I}_{p_T} > 0.9$ is applied to the muon and tau candidates for all decay channels and selection sets, and retain 70% of the signal candidates while rejecting 90% of QCD events. In addition, a cut $I_{p_T} < 2 \text{ GeV}/c$ is applied in the L-selection to both candidates, as the lower p_T reduces the background rejection power of the \hat{I}_{p_T} variable.

The selection criteria common or specific to each selection set and decay channel are summarised in Table 1. The signal selection efficiencies are found to vary from 10 to 50%. Due to the kinematic selection, the decay channels are mutually exclusive and just one $\mu^\pm\tau^\mp$ candidate per event is found.

4 Background estimation

Several background processes are considered: $Z \rightarrow \tau^+\tau^-$, $Z \rightarrow l^+l^-$ ($l = e, \mu$), QCD, Vj , double bosons production (VV), $t\bar{t}$, and $Z \rightarrow b\bar{b}$. All backgrounds except $Z \rightarrow \tau^+\tau^-$ are estimated following the procedures described in Ref. [37]. The expected yields can be found in Table 2. The corresponding invariant-mass distributions compared with candidates observed in the data are shown in Fig. 1. For illustration, examples of $H \rightarrow \mu^\pm\tau^\mp$ distributions from simulation are also superimposed.

The $Z \rightarrow \tau^+\tau^-$ background is estimated from the cross-section measured by the LHCb collaboration [37] where the reconstruction efficiency is determined from data, and the acceptance and selection efficiency are obtained from simulation. The estimated background includes a small amount of cross-feed from different final states of the tau decay, as determined from simulation. The $Z \rightarrow \mu^+\mu^-$ background is dominant in the $\mu\tau_\mu$ channel. The corresponding invariant-mass distribution is obtained from simulation and normalised to data in the Z peak region, from 80 to 100 GeV/c^2 . In order to suppress the potential presence of signal in this region, the muons are required to be promptly produced. For other channels, the $Z \rightarrow l^+l^-$ decay becomes a background source in case a lepton is misidentified. This contribution is computed from the $Z \rightarrow l^+l^-$ in data, and weighted by the particle misidentification probability obtained from simulation.

The QCD and Vj backgrounds are inferred from data using the same criteria as for the signal but selecting same-sign $\mu^\pm\tau^\pm$ candidates. Their amounts are determined by a fit to

Table 1: Requirements for each decay channel and selection set.

Selection set	Variable	$\mu\tau_e$	$\mu\tau_{h1}$	$\mu\tau_{h3}$	$\mu\tau_\mu$
All	$p_T(\tau) [\text{GeV}/c]$	> 5	> 10	> 12	> 5
	$p_T(\tau_{h3}^{\text{prong}1}) [\text{GeV}/c]$	—	—	> 1	—
	$p_T(\tau_{h3}^{\text{prong}2}) [\text{GeV}/c]$	—	—	> 1	—
	$p_T(\tau_{h3}^{\text{prong}3}) [\text{GeV}/c]$	—	—	> 6	—
	$p_T(\mu) - p_T(\tau) [\text{GeV}/c]$	> 0	—	—	—
	$m(\tau_{h3}) [\text{GeV}/c^2]$	—	—	0.7–1.5	—
	$m_{\text{corr}}(\tau_{h3}) [\text{GeV}/c^2]$	—	—	> 3	—
	Time-of-flight (τ_{h3}) [fs]	—	—	> 30	—
	IP(τ) [μm]	> 10	> 10	—	> 50
	IP(μ) [μm]	< 50	< 50	< 50	< 50
	$\Delta\phi$ [rad]	> 2.7	> 2.7	> 2.7	> 2.7
	$\hat{I}_{p_T}(\tau)$	> 0.9	> 0.9	> 0.9	> 0.9
	$\hat{I}_{p_T}(\mu)$	> 0.9	> 0.9	> 0.9	> 0.9
L-selection	$p_T(\mu) [\text{GeV}/c]$	> 20	> 20	> 20	> 20
	A_{p_T}	< 0.6	< 0.4	—	> 0.3
	$I_{p_T}(\tau) [\text{GeV}/c]$	< 2	< 2	< 2	< 2
	$I_{p_T}(\mu) [\text{GeV}/c]$	< 2	< 2	< 2	< 2
C-selection	$p_T(\mu) [\text{GeV}/c]$	> 30	> 30	> 30	> 30
	A_{p_T}	—	< 0.5	—	> 0.3
H-selection	$p_T(\tau) [\text{GeV}/c]$	> 20	> 20	> 20	—
	$p_T(\mu) [\text{GeV}/c]$	> 40	> 40	> 40	> 50
	A_{p_T}	—	—	—	> 0.4

the distribution of $p_T(\mu) - p_T(\tau)$, with templates representing each of them. The template for the QCD component is obtained from data requiring an anti-isolation $\hat{I}_{p_T} < 0.6$ selection. The distribution obtained from simulation is used for the Vj component. Factors are subsequently applied for the correction of the relative yield of opposite-sign to same-sign candidates. For the QCD background the number of anti-isolated opposite-sign candidates found in data is used in the calculation of the correction factor, where it is found to be close to unity. The factors are found consistent with the simulation. The factors for the Vj component are taken from simulation, and are in general larger than unity (1.3 for $\mu\tau_e$ up to 3.1 for $\mu\tau_{h1}$, for the L-selection). The minor contributions from VV , $t\bar{t}$, and $Z \rightarrow b\bar{b}$ processes are estimated from simulation.

5 Results

The signal cross-section multiplied by the branching fraction is given by

$$\sigma(gg \rightarrow H \rightarrow \mu^\pm \tau^\mp) = N_{\text{sig}} / (\mathcal{L} \cdot \mathcal{B}(\tau \rightarrow X) \cdot \varepsilon), \quad (1)$$

where N_{sig} is the signal yield obtained from the fit procedure described below, \mathcal{L} the total integrated luminosity, $\mathcal{B}(\tau \rightarrow X)$ the tau branching fraction, and ε the detection efficiency. The latter is the product of acceptance, reconstruction, and offline selection efficiencies.




Figure 1: Invariant-mass distributions for the $\mu^\pm\tau^\mp$ candidates for the four decay channels (from top to bottom: $\mu\tau_e$, $\mu\tau_{h1}$, $\mu\tau_{h3}$, $\mu\tau_\mu$) and the three selections (from left to right: L-selection, C-selection, H-selection). The distribution of candidates observed (black points) is compared with backgrounds (filled colour, stacked), and with signal hypothesis (cyan). The signal is normalised to \sqrt{N} , with N the total number of candidates in the corresponding data histogram.

These efficiencies are obtained from simulated samples and data for each decay channel and selection set, following the methods developed for the $Z \rightarrow \tau^+\tau^-$ measurement [37]. The acceptance obtained from the POWHEG-BOX generator is identical for the $\mu\tau_e$, $\mu\tau_{h3}$, and $\mu\tau_\mu$ channels, varying from 1.0% for $m_H = 195 \text{ GeV}/c^2$ to 3.2% for $m_H = 75 \text{ GeV}/c^2$. The reconstruction efficiency, which is the product of contributions from trigger, tracking, and particle identification, is in the range 40–70%, but only about 15% in the case of the $\mu\tau_{h3}$ channel because of the limited tracking efficiency for the low-momentum hadrons. With the exception of the $\mu\tau_\mu$ channel, the selection efficiency is 18–30% in the L-selection, and 24–49% in the C-selection and H-selection. In the case of the $\mu\tau_\mu$ channel, the tighter selection on the muon p_T and impact parameter reduces the selection efficiency to 10–15%.

Table 2: Expected number of background candidates from each component, total background with uncertainty, and number of observed candidates with statistical uncertainty, from each decay channel and selection set.

Selection set	Process	$\mu\tau_e$	$\mu\tau_{h1}$	$\mu\tau_{h3}$	$\mu\tau_\mu$
L-selection	$Z \rightarrow \tau^+\tau^-$	371.1 ± 26.0	681.7 ± 47.1	135.1 ± 11.7	137.4 ± 9.5
	$Z \rightarrow l^+l^-$	8.2 ± 1.6	4.0 ± 1.8	—	155.3 ± 5.0
	QCD	67.5 ± 10.6	463.6 ± 5.4	93.1 ± 10.9	19.4 ± 5.5
	Vj	14.5 ± 10.3	143.2 ± 58.6	40.1 ± 15.8	10.7 ± 5.8
	VV	3.4 ± 0.3	0.9 ± 0.2	0.3 ± 0.1	0.3 ± 0.1
	$t\bar{t}$	1.7 ± 0.1	1.3 ± 0.1	0.7 ± 0.1	1.3 ± 0.2
	$Z \rightarrow b\bar{b}$	0.2 ± 0.2	0.2 ± 0.2	0.1 ± 0.1	0.2 ± 0.2
Total background		466.6 ± 28.0	1294.9 ± 75.5	269.4 ± 20.3	324.5 ± 12.5
Observed		472.0 ± 21.7	1284.0 ± 35.8	240.0 ± 15.5	344.0 ± 18.5
C-selection	$Z \rightarrow \tau^+\tau^-$	200.0 ± 14.3	288.1 ± 20.2	61.3 ± 5.5	71.7 ± 5.2
	$Z \rightarrow l^+l^-$	8.0 ± 1.7	4.3 ± 1.8	—	126.7 ± 4.5
	QCD	10.0 ± 14.0	137.9 ± 14.0	29.9 ± 9.0	6.1 ± 3.6
	Vj	48.3 ± 17.2	242.9 ± 25.3	30.8 ± 17.6	7.9 ± 4.7
	VV	3.4 ± 0.3	1.5 ± 0.2	0.3 ± 0.1	0.3 ± 0.1
	$t\bar{t}$	2.5 ± 0.1	1.6 ± 0.1	0.7 ± 0.1	1.5 ± 0.2
	$Z \rightarrow b\bar{b}$	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
Total background		272.3 ± 17.8	676.4 ± 35.2	123.1 ± 15.0	214.3 ± 8.1
Observed		296.0 ± 17.2	679.0 ± 26.1	123.0 ± 11.1	235.0 ± 15.3
H-selection	$Z \rightarrow \tau^+\tau^-$	13.7 ± 1.8	18.4 ± 1.6	8.9 ± 1.1	2.2 ± 0.4
	$Z \rightarrow l^+l^-$	4.7 ± 1.1	2.5 ± 1.1	—	33.7 ± 2.3
	QCD	—	15.8 ± 6.3	9.7 ± 5.1	—
	Vj	3.5 ± 2.6	142.6 ± 26.0	18.6 ± 16.5	7.8 ± 4.0
	VV	1.7 ± 0.2	1.0 ± 0.2	0.1 ± 0.1	0.2 ± 0.1
	$t\bar{t}$	1.2 ± 0.1	0.9 ± 0.1	0.4 ± 0.1	0.8 ± 0.1
	$Z \rightarrow b\bar{b}$	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
Total background		24.9 ± 3.4	181.2 ± 26.7	37.8 ± 13.6	44.7 ± 4.6
Observed		27.0 ± 5.2	184.0 ± 13.6	37.0 ± 6.1	39.0 ± 6.2

The systematic uncertainties are summarised in Table 3. The uncertainty on the acceptance receives contributions from the gluon PDF uncertainty, as well as from factorization and renormalisation scales. The uncertainties on the reconstruction and selection efficiencies are estimated from simulation and are calibrated using data as described in Ref. [37]. The uncertainty associated with the invariant-mass shape is handled by selecting the weakest expected limits among the different choices of distribution (kernel estimation and histograms with different bin widths are used). The uncertainties on the integrated luminosity and acceptance are fully correlated among channels, while only a partial correlation is found for the reconstruction efficiency uncertainties. All the other uncertainties are taken as uncorrelated.

The signal yield is determined from a simultaneous extended likelihood fit of the binned

Table 3: Relative systematic uncertainties (in %) on the normalisation factors in the cross-section calculation. When the uncertainty depends on m_H a range is indicated.

	$\mu\tau_e$	$\mu\tau_{h1}$	$\mu\tau_{h3}$	$\mu\tau_\mu$
Luminosity	1.16	1.16	1.16	1.16
Tau branching fraction	0.22	0.18	0.48	0.23
PDF	2.6–7.1	3.5–7.2	2.6–7.3	3.0–7.9
Scales	0.9–1.9	0.8–1.7	0.9–1.7	0.9–1.9
Reconstruction efficiency	1.8–3.6	1.9–5.4	3.3–7.1	1.5–3.3
Selection efficiency	2.5–6.0	1.9–4.1	4.0–9.3	3.8–8.5




Figure 2: Cross-section times branching fraction 95% CL limits for the $H \rightarrow \mu^\pm \tau^\mp$ decay as a function of m_H , from the simultaneous fit. The observed limits from individual channels are also shown.

invariant-mass distributions of the $\mu\tau$ candidates. The distributions for signal are obtained from simulation, while distributions of the different background sources are obtained using the method described in Sect. 4. The amount of each background component as well as other terms in Eq. (1) containing uncertainties are treated as nuisance parameters and are constrained to a Gaussian distribution with mean and standard deviation corresponding to the expected value and its uncertainty, respectively.

The fit results for all m_H values are compatible with a null signal, hence cross-section upper limits are computed. The exclusion limits of $\sigma(gg \rightarrow H \rightarrow \mu^\pm \tau^\mp)$ defined at 95% confidence level are obtained from the CL_s method [54]. As mentioned before, for each mass hypothesis the selection considered is that providing the smallest expected limit. The $\sigma(gg \rightarrow H \rightarrow \mu^\pm \tau^\mp)$ exclusion limits are shown in Fig. 2, ranging from 22 pb for $m_H = 45 \text{ GeV}/c^2$ to 4 pb for $m_H = 195 \text{ GeV}/c^2$. In the particular case of $m_H = 125 \text{ GeV}/c^2$, using the production cross-section from Ref. [55] gives a best fit for the branching fraction of $\mathcal{B}(H \rightarrow \mu^\pm \tau^\mp) = -2^{+14}_{-12}\%$ and an observed exclusion limit $\mathcal{B}(H \rightarrow \mu^\pm \tau^\mp) < 26\%$. The corresponding exclusion limit on the Yukawa coupling is $\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 1.7 \times 10^{-2}$, assuming the decay width $\Gamma_{\text{SM}} = 4.1 \text{ MeV}/c^2$ [56].

6 Conclusion

A search for Higgs-like bosons decaying via a lepton-flavour-violating process $H \rightarrow \mu^\pm \tau^\mp$ in pp collisions at $\sqrt{s} = 8$ TeV is presented, with the tau lepton reconstructed in leptonic and hadronic decay modes. No signal has been found. The upper bound on the cross-section multiplied by the branching fraction, at 95% confidence level, ranges from 22 pb for a boson mass of $45\text{ GeV}/c^2$, to 4 pb for $195\text{ GeV}/c^2$. The search provides information complementary to the ATLAS and CMS collaborations.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom); Laboratory Directed Research and Development program of LANL (USA).

References

- [1] M. Blanke *et al.*, *$\Delta F = 2$ observables and fine-tuning in a warped extra dimension with custodial protection*, *JHEP* **03** (2009) 001, [arXiv:0809.1073](https://arxiv.org/abs/0809.1073).
- [2] G. F. Giudice and O. Lebedev, *Higgs-dependent Yukawa couplings*, *Phys. Lett. B* **665** (2008) 79, [arXiv:0804.1753](https://arxiv.org/abs/0804.1753).
- [3] J. A. Aguilar-Saavedra, *A minimal set of top-Higgs anomalous couplings*, *Nucl. Phys. B* **821** (2009) 215, [arXiv:0904.2387](https://arxiv.org/abs/0904.2387).
- [4] M. E. Albrecht *et al.*, *Electroweak and flavour structure of a warped extra dimension with custodial protection*, *JHEP* **09** (2009) 064, [arXiv:0903.2415](https://arxiv.org/abs/0903.2415).
- [5] A. Goudelis, O. Lebedev, and J.-h. Park, *Higgs-induced lepton flavour violation*, *Phys. Lett. B* **707** (2012) 369, [arXiv:1111.1715](https://arxiv.org/abs/1111.1715).

- [6] D. McKeen, M. Pospelov, and A. Ritz, *Modified Higgs branching ratios versus CP and lepton flavour violation*, Phys. Rev. D **86** (2012) 113004, [arXiv:1208.4597](#).
- [7] E. Arganda, A. M. Curiel, M. J. Herrero, and D. Temes, *Lepton flavour violating Higgs boson decays from massive seesaw neutrinos*, Phys. Rev. D **71** (2005) 035011, [arXiv:hep-ph/0407302](#).
- [8] E. Arganda, M. J. Herrero, X. Marcano, and C. Weiland, *Imprints of massive inverse seesaw model neutrinos in lepton flavour violating Higgs boson decays*, Phys. Rev. D **91** (2015) 015001, [arXiv:1405.4300](#).
- [9] R. Harnik, J. Kopp, and J. Zupan, *Flavour violating Higgs decays*, JHEP **03** (2013) 026, [arXiv:1209.1397](#).
- [10] J. D. Bjorken and S. Weinberg, *A mechanism for nonconservation of muon number*, Phys. Rev. Lett. **38** (1977) 622.
- [11] J. L. Diaz-Cruz and J. J. Toscano, *Lepton flavour violating decays of Higgs bosons beyond the standard model*, Phys. Rev. D **62** (2000) 116005, [arXiv:hep-ph/9910233](#).
- [12] T. Han and D. Marfatia, *$H \rightarrow \mu\tau$ at hadron colliders*, Phys. Rev. Lett. **86** (2001) 1442, [arXiv:hep-ph/0008141](#).
- [13] A. Arhrib, Y. Cheng, and O. C. W. Kong, *Comprehensive analysis on lepton flavour violating Higgs boson to $\mu^\mp\tau^\pm$ decay in supersymmetry without R parity*, Phys. Rev. D **87** (2013) 015025, [arXiv:1210.8241](#).
- [14] M. Arana-Catania, E. Arganda, and M. J. Herrero, *Non-decoupling SUSY in LFV Higgs decays: a window to new physics at the LHC*, JHEP **09** (2013) 160, Erratum *ibid.* **10** (2015) 192, [arXiv:1304.3371](#).
- [15] K. Agashe and R. Contino, *Composite Higgs-mediated flavor-changing neutral current*, Phys. Rev. D **80** (2009) 075016, [arXiv:0906.1542](#).
- [16] A. Azatov, M. Toharia, and L. Zhu, *Higgs mediated flavor-changing neutral current's in warped extra dimensions*, Phys. Rev. D **80** (2009) 035016, [arXiv:0906.1990](#).
- [17] G. Perez and L. Randall, *Natural neutrino masses and mixings from warped geometry*, JHEP **01** (2009) 077, [arXiv:0805.4652](#).
- [18] S. Casagrande *et al.*, *Flavour physics in the Randall-Sundrum model: I. Theoretical setup and electroweak precision tests*, JHEP **10** (2008) 094, [arXiv:0807.4937](#).
- [19] H. Ishimori *et al.*, *Non-Abelian discrete symmetries in particle physics*, Prog. Theor. Phys. Suppl. **183** (2010) 1, [arXiv:1003.3552](#).
- [20] ALEPH collaboration, D. Decamp *et al.*, *Searches for new particles in Z decays using the ALEPH detector*, Phys. Rept. **216** (1992) 253.
- [21] DELPHI collaboration, P. Abreu *et al.*, *A search for lepton flavour violation in Z decays*, Phys. Lett. B **298** (1993) 247.

- [22] L3 collaboration, O. Adriani *et al.*, *Search for lepton flavour violation in Z decays*, *Phys. Lett. B* **316** (1993) 427.
- [23] OPAL collaboration, M. Z. Akrawy *et al.*, *A Search for lepton flavour violation in Z decays*, *Phys. Lett. B* **254** (1991) 293.
- [24] OPAL collaboration, G. Abbiendi *et al.*, *Search for lepton flavour violation in e^+e^- collisions at $\sqrt{s} = 189\text{--}209$ GeV*, *Phys. Lett. B* **519** (2001) 23, [arXiv:hep-ex/0109011](https://arxiv.org/abs/hep-ex/0109011).
- [25] G. Blankenburg, J. Ellis, and G. Isidori, *Flavour-changing decays of a 125 GeV Higgs-like particle*, *Phys. Lett. B* **712** (2012) 386, [arXiv:1202.5704](https://arxiv.org/abs/1202.5704).
- [26] CMS collaboration, A. M. Sirunyan *et al.*, *Search for lepton flavour violating decays of the Higgs boson to $\mu\tau$ and $e\tau$ in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2018) 001, [arXiv:1712.07173](https://arxiv.org/abs/1712.07173).
- [27] ATLAS collaboration, G. Aad *et al.*, *Search for lepton-flavour-violating $H \rightarrow \mu\tau$ decays of the Higgs boson with the ATLAS detector*, *JHEP* **11** (2015) 211, [arXiv:1508.03372](https://arxiv.org/abs/1508.03372).
- [28] J. F. Gunion and H. E. Haber, *The CP-conserving two-Higgs doublet model: The Approach to the decoupling limit*, *Phys. Rev. D* **67** (2003) 075019, [arXiv:hep-ph/0207010](https://arxiv.org/abs/hep-ph/0207010).
- [29] ATLAS collaboration, G. Aad *et al.*, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *JHEP* **02** (2013) 095, [arXiv:1211.6956](https://arxiv.org/abs/1211.6956).
- [30] CMS collaboration, S. Chatrchyan *et al.*, *Search for neutral Higgs bosons decaying to tau pairs in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **713** (2012) 68, [arXiv:1202.4083](https://arxiv.org/abs/1202.4083).
- [31] J. Jaeckel and A. Ringwald, *The low-energy frontier of particle physics*, *Ann. Rev. Nucl. Part. Sci.* **60** (2010) 405, [arXiv:1002.0329](https://arxiv.org/abs/1002.0329).
- [32] M. Baumgart *et al.*, *Non-Abelian dark sectors and their collider signatures*, *JHEP* **04** (2009) 014, [arXiv:0901.0283](https://arxiv.org/abs/0901.0283).
- [33] BaBar collaboration, J. P. Lees *et al.*, *Search for low-mass dark-sector Higgs bosons*, *Phys. Rev. Lett.* **108** (2012) 211801, [arXiv:1202.1313](https://arxiv.org/abs/1202.1313).
- [34] Belle collaboration, I. Jaegle, *Search for the 'Dark Photon' and the 'Dark Higgs' at Belle*, *Nucl. Phys. Proc. Suppl.* **234** (2013) 33, [arXiv:1211.1403](https://arxiv.org/abs/1211.1403).
- [35] LHCb collaboration, R. Aaij *et al.*, *Search for dark photons produced in 13 TeV pp collisions*, *Phys. Rev. Lett.* **120** (2018) 061801, [arXiv:1710.02867](https://arxiv.org/abs/1710.02867).
- [36] H. M. Georgi, S. L. Glashow, M. E. Machacek, and D. V. Nanopoulos, *Higgs bosons from two-gluon annihilation in proton-proton collisions*, *Phys. Rev. Lett.* **40** (1978) 692.

- [37] LHCb collaboration, R. Aaij *et al.*, *Measurement of $Z \rightarrow \tau^+ \tau^-$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV*, **JHEP** **09** (2018) 159, [arXiv:1806.05008](https://arxiv.org/abs/1806.05008).
- [38] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, **JINST** **3** (2008) S08005.
- [39] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, **Int. J. Mod. Phys. A** **30** (2015) 1530022, [arXiv:1412.6352](https://arxiv.org/abs/1412.6352).
- [40] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, **Comput. Phys. Commun.** **178** (2008) 852, [arXiv:0710.3820](https://arxiv.org/abs/0710.3820); T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, **JHEP** **05** (2006) 026, [arXiv:hep-ph/0603175](https://arxiv.org/abs/hep-ph/0603175).
- [41] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, **J. Phys. Conf. Ser.** **331** (2011) 032047.
- [42] J. Pumplin *et al.*, *New generation of parton distributions with uncertainties from global QCD analysis*, **JHEP** **07** (2002) 012, [arXiv:hep-ph/0201195](https://arxiv.org/abs/hep-ph/0201195).
- [43] D. J. Lange, *The EvtGen particle decay simulation package*, **Nucl. Instrum. Meth. A** **462** (2001) 152.
- [44] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, **Eur. Phys. J. C** **45** (2006) 97, [arXiv:hep-ph/0506026](https://arxiv.org/abs/hep-ph/0506026).
- [45] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, **IEEE Trans. Nucl. Sci.** **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, **Nucl. Instrum. Meth. A** **506** (2003) 250.
- [46] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, **J. Phys. Conf. Ser.** **331** (2011) 032023.
- [47] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, **JHEP** **11** (2004) 040, [arXiv:hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146).
- [48] S. Frixione, P. Nason, and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, **JHEP** **11** (2007) 070, [arXiv:0709.2092](https://arxiv.org/abs/0709.2092).
- [49] S. Alioli, P. Nason, C. Oleari, and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, **JHEP** **06** (2010) 043, [arXiv:1002.2581](https://arxiv.org/abs/1002.2581).
- [50] S. Alioli, P. Nason, C. Oleari, and E. Re, *NLO Higgs boson production via gluon fusion matched with shower in POWHEG*, **JHEP** **04** (2009) 002, [arXiv:0812.0578](https://arxiv.org/abs/0812.0578).
- [51] L. A. Harland-Lang, A. D. Martin, P. Motylinski, and R. S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, **Eur. Phys. J. C** **75** (2015) 204, [arXiv:1412.3989](https://arxiv.org/abs/1412.3989).
- [52] LHCb collaboration, R. Aaij *et al.*, *Precision luminosity measurements at LHCb*, **JINST** **9** (2014) P12005, [arXiv:1410.0149](https://arxiv.org/abs/1410.0149).

- [53] F. Machefer, *LHCb calorimeters and muon system lepton identification*, in *AIP Conference Proceedings*, AIP, 2004. doi: [10.1063/1.1807309](https://doi.org/10.1063/1.1807309).
- [54] A. L. Read, *Presentation of search results: The CL_s technique*, *J. Phys. G* **28** (2002) 2693.
- [55] LHC Higgs Cross Section Working Group, J. R. Andersen *et al.*, *Handbook of LHC Higgs cross sections: 3. Higgs properties*, tech. rep., CERN, Jul, 2013. doi: [CERN-2013-004](https://cds.cern.ch/record/1442313).
- [56] A. Denner *et al.*, *Standard Model Higgs-boson branching ratios with uncertainties*, *Eur. Phys. J. C* **71** (2011) 1753, [arXiv:1107.5909](https://arxiv.org/abs/1107.5909).

LHCb collaboration

R. Aaij²⁷, C. Abellán Beteta⁴⁴, B. Adeva⁴¹, M. Adinolfi⁴⁸, C.A. Aidala⁷³, Z. Ajaltouni⁵, S. Akar⁵⁹, P. Albicocco¹⁸, J. Albrecht¹⁰, F. Alessio⁴², M. Alexander⁵³, A. Alfonso Albero⁴⁰, G. Alkhazov³³, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁴¹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An³, L. Anderlini¹⁷, G. Andreassi⁴³, M. Andreotti^{16,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, F. Archilli²⁷, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁹, M. Artuso⁶¹, K. Arzymatow³⁷, E. Aslanides⁶, M. Atzeni⁴⁴, B. Audurier²², S. Bachmann¹², J.J. Back⁵⁰, S. Baker⁵⁵, V. Balagura^{7,b}, W. Baldini¹⁶, A. Baranov³⁷, R.J. Barlow⁵⁶, S. Barsuk⁷, W. Barter⁵⁶, F. Baryshnikov⁷⁰, V. Batozskaya³¹, B. Batsukh⁶¹, V. Battista⁴³, A. Bay⁴³, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, A. Beiter⁶¹, L.J. Bel²⁷, S. Belin²², N. Belyi⁶³, V. Bellee⁴³, N. Belloli^{20,i}, K. Belous³⁹, I. Belyaev^{34,42}, E. Ben-Haim⁸, G. Bencivenni¹⁸, S. Benson²⁷, S. Beranek⁹, A. Berezhnoy³⁵, R. Bernet⁴⁴, D. Berninghoff¹², E. Bertholet⁸, A. Bertolin²³, C. Betancourt⁴⁴, F. Betti^{15,42}, M.O. Bettler⁴⁹, M. van Beuzekom²⁷, Ia. Bezshyiko⁴⁴, S. Bhasin⁴⁸, J. Bhom²⁹, S. Bifani⁴⁷, P. Billoir⁸, A. Birnkraut¹⁰, A. Bizzeti^{17,u}, M. Bjørn⁵⁷, M.P. Blago⁴², T. Blake⁵⁰, F. Blanc⁴³, S. Blusk⁶¹, D. Bobulska⁵³, V. Bocci²⁶, O. Boente Garcia⁴¹, T. Boettcher⁵⁸, A. Bondar^{38,w}, N. Bondar³³, S. Borghi^{56,42}, M. Borisyak³⁷, M. Borsato⁴¹, F. Bossu⁷, M. Bouabdil⁹, T.J.V. Bowcock⁵⁴, C. Bozzi^{16,42}, S. Braun¹², M. Brodski⁴², J. Brodzicka²⁹, A. Brossa Gonzalo⁵⁰, D. Brundu²², E. Buchanan⁴⁸, A. Buonaura⁴⁴, C. Burr⁵⁶, A. Bursche²², J. Buytaert⁴², W. Byczynski⁴², S. Cadeddu²², H. Cai⁶⁴, R. Calabrese^{16,g}, R. Calladine⁴⁷, M. Calvi^{20,i}, M. Calvo Gomez^{40,m}, A. Camboni^{40,m}, P. Campana¹⁸, D.H. Campora Perez⁴², L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni²⁵, R. Cardinale^{19,h}, A. Cardini²², P. Carniti^{20,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina²⁰, M. Cattaneo⁴², G. Cavallero^{19,h}, R. Cenci^{24,p}, D. Chamont⁷, M.G. Chapman⁴⁸, M. Charles⁸, Ph. Charpentier⁴², G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, V. Chekalina³⁷, C. Chen³, S. Chen²², S.-G. Chitic⁴², V. Chobanova⁴¹, M. Chrzaszcz⁴², A. Chubykin³³, P. Ciambrone¹⁸, X. Cid Vidal⁴¹, G. Ciezarek⁴², P.E.L. Clarke⁵², M. Clemencic⁴², H.V. Cliff⁴⁹, J. Closier⁴², V. Coco⁴², J.A.B. Coelho⁷, J. Cogan⁶, E. Cogneras⁵, L. Cojocariu³², P. Collins⁴², T. Colombo⁴², A. Comerma-Montells¹², A. Contu²², G. Coombs⁴², S. Coquereau⁴⁰, G. Corti⁴², M. Corvo^{16,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴², G.A. Cowan⁵², D.C. Craik⁵⁸, A. Crocombe⁵⁰, M. Cruz Torres¹, R. Currie⁵², C. D'Ambrosio⁴², F. Da Cunha Marinho², C.L. Da Silva⁷⁴, E. Dall'Occo²⁷, J. Dalseno⁴⁸, A. Danilina³⁴, A. Davis³, O. De Aguiar Francisco⁴², K. De Bruyn⁴², S. De Capua⁵⁶, M. De Cian⁴³, J.M. De Miranda¹, L. De Paula², M. De Serio^{14,d}, P. De Simone¹⁸, C.T. Dean⁵³, D. Decamp⁴, L. Del Buono⁸, B. Delaney⁴⁹, H.-P. Dembinski¹¹, M. Demmer¹⁰, A. Dendek³⁰, D. Derkach³⁷, O. Deschamps⁵, F. Desse⁷, F. Dettori⁵⁴, B. Dey⁶⁵, A. Di Canto⁴², P. Di Nezza¹⁸, S. Didenko⁷⁰, H. Dijkstra⁴², F. Dordei⁴², M. Dorigo^{42,y}, A. Dosil Suárez⁴¹, L. Douglas⁵³, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour²⁷, G. Dujany⁸, P. Durante⁴², J.M. Durham⁷⁴, D. Dutta⁵⁶, R. Dzhelyadin³⁹, M. Dziewiecki¹², A. Dziurda²⁹, A. Dzyuba³³, S. Easo⁵¹, U. Egede⁵⁵, V. Egorychev³⁴, S. Eidelman^{38,w}, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, S. Ely⁶¹, A. Ene³², S. Escher⁹, S. Esen²⁷, T. Evans⁵⁹, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, D. Fazzini^{20,42,i}, L. Federici²⁵, P. Fernandez Declara⁴², A. Fernandez Prieto⁴¹, F. Ferrari¹⁵, L. Ferreira Lopes⁴³, F. Ferreira Rodrigues², M. Ferro-Luzzi⁴², S. Filippov³⁶, R.A. Fini¹⁴, M. Fiorini^{16,g}, M. Firlej³⁰, C. Fitzpatrick⁴³, T. Fiutowski³⁰, F. Fleuret^{7,b}, M. Fontana^{22,42}, F. Fontanelli^{19,h}, R. Forty⁴², V. Franco Lima⁵⁴, M. Frank⁴², C. Frei⁴², J. Fu^{21,q}, W. Funk⁴², C. Färber⁴², M. Féo Pereira Rivello Carvalho²⁷, E. Gabriel⁵², A. Gallas Torreira⁴¹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², Y. Gan³, M. Gandelman², P. Gandini²¹, Y. Gao³, L.M. Garcia Martin⁷², B. Garcia Plana⁴¹, J. García Pardiñas⁴⁴, J. Garra Tico⁴⁹, L. Garrido⁴⁰, D. Gascon⁴⁰, C. Gaspar⁴², L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck⁵⁶, M. Gersabeck⁵⁶, T. Gershon⁵⁰, D. Gerstel⁶, Ph. Ghez⁴, S. Gianì⁴³, V. Gibson⁴⁹, O.G. Girard⁴³, L. Giubega³²,

K. Gizzdov⁵², V.V. Gligorov⁸, D. Golubkov³⁴, A. Golutvin^{55,70}, A. Gomes^{1,a}, I.V. Gorelov³⁵,
 C. Gotti^{20,i}, E. Govorkova²⁷, J.P. Grabowski¹², R. Graciani Diaz⁴⁰, L.A. Granado Cardoso⁴²,
 E. Graugés⁴⁰, E. Graverini⁴⁴, G. Graziani¹⁷, A. Grecu³², R. Greim²⁷, P. Griffith²², L. Grillo⁵⁶,
 L. Gruber⁴², B.R. Gruberg Cazon⁵⁷, O. Grünberg⁶⁷, C. Gu³, E. Gushchin³⁶, Yu. Guz^{39,42},
 T. Gys⁴², C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴³, C. Haen⁴²,
 S.C. Haines⁴⁹, B. Hamilton⁶⁰, X. Han¹², T.H. Hancock⁵⁷, S. Hansmann-Menzemer¹²,
 N. Harnew⁵⁷, S.T. Harnew⁴⁸, T. Harrison⁵⁴, C. Hasse⁴², M. Hatch⁴², J. He⁶³, M. Hecker⁵⁵,
 K. Heinicke¹⁰, A. Heister¹⁰, K. Hennessy⁵⁴, L. Henry⁷², E. van Herwijnen⁴², M. Heß⁶⁷,
 A. Hicheur², R. Hidalgo Charman⁵⁶, D. Hill⁵⁷, M. Hilton⁵⁶, P.H. Hopchev⁴³, W. Hu⁶⁵,
 W. Huang⁶³, Z.C. Huard⁵⁹, W. Hulsbergen²⁷, T. Humair⁵⁵, M. Hushchyn³⁷, D. Hutchcroft⁵⁴,
 D. Hynds²⁷, P. Ibis¹⁰, M. Idzik³⁰, P. Ilten⁴⁷, K. Ivshin³³, R. Jacobsson⁴², J. Jalocha⁵⁷,
 E. Jans²⁷, A. Jawahery⁶⁰, F. Jiang³, M. John⁵⁷, D. Johnson⁴², C.R. Jones⁴⁹, C. Joram⁴²,
 B. Jost⁴², N. Jurik⁵⁷, S. Kandybei⁴⁵, M. Karacson⁴², J.M. Kariuki⁴⁸, S. Karodia⁵³, N. Kazeev³⁷,
 M. Kecke¹², F. Keizer⁴⁹, M. Kelsey⁶¹, M. Kenzie⁴⁹, T. Ketel²⁸, E. Khairullin³⁷, B. Khanji⁴²,
 C. Khurewathanakul⁴³, K.E. Kim⁶¹, T. Kirn⁹, S. Klaver¹⁸, K. Klimaszewski³¹, T. Klimkovich¹¹,
 S. Koliiev⁴⁶, M. Kolpin¹², R. Kopecna¹², P. Koppenburg²⁷, I. Kostiuk²⁷, S. Kotriakhova³³,
 M. Kozeiha⁵, L. Kravchuk³⁶, M. Kreps⁵⁰, F. Kress⁵⁵, P. Krokovny^{38,w}, W. Krupa³⁰,
 W. Krzemien³¹, W. Kucewicz^{29,l}, M. Kucharczyk²⁹, V. Kudryavtsev^{38,w}, A.K. Kuonen⁴³,
 T. Kvaratskheliya^{34,42}, D. Lacarrere⁴², G. Lafferty⁵⁶, A. Lai²², D. Lancierini⁴⁴, G. Lanfranchi¹⁸,
 C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷, R. Le Gac⁶, A. Leflat³⁵, J. Lefrançois⁷,
 R. Lefèvre⁵, F. Lemaitre⁴², O. Leroy⁶, T. Lesiak²⁹, B. Leverington¹², P.-R. Li⁶³, T. Li³, Z. Li⁶¹,
 X. Liang⁶¹, T. Likhomanenko⁶⁹, R. Lindner⁴², F. Lionetto⁴⁴, V. Lisovskyi⁷, X. Liu³, D. Loh⁵⁰,
 A. Loi²², I. Longstaff⁵³, J.H. Lopes², G.H. Lovell⁴⁹, D. Lucchesi^{23,o}, M. Lucio Martinez⁴¹,
 A. Lupato²³, E. Luppi^{16,g}, O. Lupton⁴², A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷, F. Maciuc³²,
 V. Macko⁴³, P. Mackowiak¹⁰, S. Maddrell-Mander⁴⁸, O. Maev^{33,42}, K. Maguire⁵⁶,
 D. Maisuzenko³³, M.W. Majewski³⁰, S. Malde⁵⁷, B. Malecki²⁹, A. Malinin⁶⁹, T. Maltsev^{38,w},
 G. Manca^{22,f}, G. Mancinelli⁶, D. Marangotto^{21,q}, J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵,
 C. Marin Benito⁷, M. Marinangeli⁴³, P. Marino⁴³, J. Marks¹², P.J. Marshall⁵⁴, G. Martellotti²⁶,
 M. Martin⁶, M. Martinelli⁴², D. Martinez Santos⁴¹, F. Martinez Vidal⁷², A. Massafferri¹,
 M. Materok⁹, R. Matev⁴², A. Mathad⁵⁰, Z. Mathe⁴², C. Matteuzzi²⁰, A. Mauri⁴⁴, E. Maurice^{7,b},
 B. Maurin⁴³, A. Mazurov⁴⁷, M. McCann^{55,42}, A. McNab⁵⁶, R. McNulty¹³, J.V. Mead⁵⁴,
 B. Meadows⁵⁹, C. Meaux⁶, F. Meier¹⁰, N. Meinert⁶⁷, D. Melnychuk³¹, M. Merk²⁷, A. Merli^{21,q},
 E. Michelin²³, D.A. Milanes⁶⁶, E. Millard⁵⁰, M.-N. Minard⁴, L. Minzoni^{16,g}, D.S. Mitzel¹²,
 A. Mogini⁸, J. Molina Rodriguez^{1,z}, T. Mombächer¹⁰, I.A. Monroy⁶⁶, S. Monteil⁵,
 M. Morandin²³, G. Morello¹⁸, M.J. Morello^{24,t}, O. Morgunova⁶⁹, J. Moron³⁰, A.B. Morris⁶,
 R. Mountain⁶¹, F. Muheim⁵², M. Mulder²⁷, C.H. Murphy⁵⁷, D. Murray⁵⁶, A. Mödden¹⁰,
 D. Müller⁴², J. Müller¹⁰, K. Müller⁴⁴, V. Müller¹⁰, P. Naik⁴⁸, T. Nakada⁴³, R. Nandakumar⁵¹,
 A. Nandi⁵⁷, T. Nanut⁴³, I. Nasteva², M. Needham⁵², N. Neri²¹, S. Neubert¹², N. Neufeld⁴²,
 M. Neuner¹², T.D. Nguyen⁴³, C. Nguyen-Mau^{43,n}, S. Nieswand⁹, R. Niet¹⁰, N. Nikitin³⁵,
 A. Nogay⁶⁹, N.S. Nolte⁴², D.P. O'Hanlon¹⁵, A. Oblakowska-Mucha³⁰, V. Obraztsov³⁹,
 S. Ogilvy¹⁸, R. Oldeman^{22,f}, C.J.G. Onderwater⁶⁸, A. Ossowska²⁹, J.M. Otalora Goicochea²,
 P. Owen⁴⁴, A. Oyanguren⁷², P.R. Pais⁴³, T. Pajero^{24,t}, A. Palano¹⁴, M. Palutan^{18,42},
 G. Panshin⁷¹, A. Papanestis⁵¹, M. Pappagallo⁵², L.L. Pappalardo^{16,g}, W. Parker⁶⁰, C. Parkes⁵⁶,
 G. Passaleva^{17,42}, A. Pastore¹⁴, M. Patel⁵⁵, C. Patrignani^{15,e}, A. Pearce⁴², A. Pellegrino²⁷,
 G. Penso²⁶, M. Pepe Altarelli⁴², S. Perazzini⁴², D. Pereima³⁴, P. Perret⁵, L. Pescatore⁴³,
 K. Petridis⁴⁸, A. Petrolini^{19,h}, A. Petrov⁶⁹, S. Petrucci⁵², M. Petruzzo^{21,q}, B. Pietrzyk⁴,
 G. Pietrzyk⁴³, M. Pikies²⁹, M. Pili⁵⁷, D. Pinci²⁶, J. Pinzino⁴², F. Pisani⁴², A. Piucci¹²,
 V. Placinta³², S. Playfer⁵², J. Plews⁴⁷, M. Plo Casasus⁴¹, F. Polci⁸, M. Poli Lener¹⁸,
 A. Poluektov⁵⁰, N. Polukhina^{70,c}, I. Polyakov⁶¹, E. Polycarpo², G.J. Pomery⁴⁸, S. Ponce⁴²,
 A. Popov³⁹, D. Popov^{47,11}, S. Poslavskii³⁹, C. Potterat², E. Price⁴⁸, J. Prisciandaro⁴¹,

C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴⁴, H. Pullen⁵⁷, G. Punzi^{24,p}, W. Qian⁶³, J. Qin⁶³, R. Quagliani⁸, B. Quintana⁵, B. Rachwal³⁰, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas⁴¹, M.S. Rangel², F. Ratnikov^{37,x}, G. Raven²⁸, M. Ravonel Salzgeber⁴², M. Reboud⁴, F. Redi⁴³, S. Reichert¹⁰, A.C. dos Reis¹, F. Reiss⁸, C. Remon Alepuz⁷², Z. Ren³, V. Renaudin⁷, S. Ricciardi⁵¹, S. Richards⁴⁸, K. Rinnert⁵⁴, P. Robbe⁷, A. Robert⁸, A.B. Rodrigues⁴³, E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁶, M. Roehrken⁴², S. Roiser⁴², A. Rollings⁵⁷, V. Romanovskiy³⁹, A. Romero Vidal⁴¹, M. Rotondo¹⁸, M.S. Rudolph⁶¹, T. Ruf⁴², J. Ruiz Vidal⁷², J.J. Saborido Silva⁴¹, N. Sagidova³³, B. Saitta^{22,f}, V. Salustino Guimaraes⁶², C. Sanchez Gras²⁷, C. Sanchez Mayordomo⁷², B. Sanmartin Sedes⁴¹, R. Santacesaria²⁶, C. Santamarina Rios⁴¹, M. Santimaria¹⁸, E. Santovetti^{25,j}, G. Sarpis⁵⁶, A. Sarti^{18,k}, C. Satriano^{26,s}, A. Satta²⁵, M. Saur⁶³, D. Savrina^{34,35}, S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁵³, H. Schindler⁴², M. Schmelling¹¹, T. Schmelzer¹⁰, B. Schmidt⁴², O. Schneider⁴³, A. Schopper⁴², H.F. Schreiner⁵⁹, M. Schubiger⁴³, M.H. Schune⁷, R. Schwemmer⁴², B. Sciascia¹⁸, A. Sciubba^{26,k}, A. Semennikov³⁴, E.S. Sepulveda⁸, A. Sergi^{47,42}, N. Serra⁴⁴, J. Serrano⁶, L. Sestini²³, A. Seuthe¹⁰, P. Seyfert⁴², M. Shapkin³⁹, Y. Shcheglov^{33,†}, T. Shears⁵⁴, L. Shekhtman^{38,w}, V. Shevchenko⁶⁹, E. Shmanin⁷⁰, B.G. Siddi¹⁶, R. Silva Coutinho⁴⁴, L. Silva de Oliveira², G. Simi^{23,o}, S. Simone^{14,d}, N. Skidmore¹², T. Skwarnicki⁶¹, M.W. Slater⁴⁷, J.G. Smeaton⁴⁹, E. Smith⁹, I.T. Smith⁵², M. Smith⁵⁵, M. Soares¹⁵, l. Soares Lavra¹, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³, B. Souza De Paula², B. Spaan¹⁰, E. Spadaro Norella^{21,q}, P. Spradlin⁵³, F. Stagni⁴², M. Stahl¹², S. Stahl⁴², P. Steffko⁴³, S. Stefkova⁵⁵, O. Steinkamp⁴⁴, S. Stemmle¹², O. Stenyakin³⁹, M. Stepanova³³, H. Stevens¹⁰, A. Stocchi⁷, S. Stone⁶¹, B. Storaci⁴⁴, S. Stracka²⁴, M.E. Stramaglia⁴³, M. Stratificiuc³², U. Straumann⁴⁴, S. Strokov⁷¹, J. Sun³, L. Sun⁶⁴, K. Swientek³⁰, T. Szumlak³⁰, M. Szymanski⁶³, S. T'Jampens⁴, Z. Tang³, A. Tayduganov⁶, T. Tekampe¹⁰, G. Tellarini¹⁶, F. Teubert⁴², E. Thomas⁴², J. van Tilburg²⁷, M.J. Tilley⁵⁵, V. Tisserand⁵, M. Tobin³⁰, S. Tolk⁴², L. Tomassetti^{16,g}, D. Tonelli²⁴, D.Y. Tou⁸, R. Tourinho Jadallah Aoude¹, E. Tournefier⁴, M. Trail⁵³, M.T. Tran⁴³, A. Trisovic⁴⁹, A. Tsaregorodtsev⁶, G. Tuci^{24,p}, A. Tully⁴⁹, N. Tuning^{27,42}, A. Ukleja³¹, A. Usachov⁷, A. Ustyuzhanin³⁷, U. Uwer¹², A. Vagner⁷¹, V. Vagnoni¹⁵, A. Valassi⁴², S. Valat⁴², G. Valenti¹⁵, R. Vazquez Gomez⁴², P. Vazquez Regueiro⁴¹, S. Vecchi¹⁶, M. van Veghel²⁷, J.J. Velthuis⁴⁸, M. Veltri^{17,r}, G. Veneziano⁵⁷, A. Venkateswaran⁶¹, T.A. Verlage⁹, M. Vernet⁵, M. Veronesi²⁷, N.V. Veronika¹³, M. Vesterinen⁵⁷, J.V. Viana Barbosa⁴², D. Vieira⁶³, M. Vieites Diaz⁴¹, H. Viemann⁶⁷, X. Vilasis-Cardona^{40,m}, A. Vitkovskiy²⁷, M. Vitti⁴⁹, V. Volkov³⁵, A. Vollhardt⁴⁴, D. Vom Bruch⁸, B. Voneki⁴², A. Vorobyev³³, V. Vorobyev^{38,w}, J.A. de Vries²⁷, C. Vázquez Sierra²⁷, R. Waldi⁶⁷, J. Walsh²⁴, J. Wang⁶¹, M. Wang³, Y. Wang⁶⁵, Z. Wang⁴⁴, D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷, D. Websdale⁵⁵, A. Weiden⁴⁴, C. Weisser⁵⁸, M. Whitehead⁹, J. Wicht⁵⁰, G. Wilkinson⁵⁷, M. Wilkinson⁶¹, I. Williams⁴⁹, M.R.J. Williams⁵⁶, M. Williams⁵⁸, T. Williams⁴⁷, F.F. Wilson^{51,42}, J. Wimberley⁶⁰, M. Winn⁷, J. Wishahi¹⁰, W. Wislicki³¹, M. Witek²⁹, G. Wormser⁷, S.A. Wotton⁴⁹, K. Wyllie⁴², D. Xiao⁶⁵, Y. Xie⁶⁵, A. Xu³, M. Xu⁶⁵, Q. Xu⁶³, Z. Xu³, Z. Xu⁴, Z. Yang³, Z. Yang⁶⁰, Y. Yao⁶¹, L.E. Yeomans⁵⁴, H. Yin⁶⁵, J. Yu^{65,ab}, X. Yuan⁶¹, O. Yushchenko³⁹, K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, D. Zhang⁶⁵, L. Zhang³, W.C. Zhang^{3,aa}, Y. Zhang⁷, A. Zhelezov¹², Y. Zheng⁶³, X. Zhu³, V. Zhukov^{9,35}, J.B. Zonneveld⁵², S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

⁷LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

⁸LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

- ⁹*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁰*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹¹*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹²*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ¹³*School of Physics, University College Dublin, Dublin, Ireland*
- ¹⁴*INFN Sezione di Bari, Bari, Italy*
- ¹⁵*INFN Sezione di Bologna, Bologna, Italy*
- ¹⁶*INFN Sezione di Ferrara, Ferrara, Italy*
- ¹⁷*INFN Sezione di Firenze, Firenze, Italy*
- ¹⁸*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ¹⁹*INFN Sezione di Genova, Genova, Italy*
- ²⁰*INFN Sezione di Milano-Bicocca, Milano, Italy*
- ²¹*INFN Sezione di Milano, Milano, Italy*
- ²²*INFN Sezione di Cagliari, Monserrato, Italy*
- ²³*INFN Sezione di Padova, Padova, Italy*
- ²⁴*INFN Sezione di Pisa, Pisa, Italy*
- ²⁵*INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ²⁶*INFN Sezione di Roma La Sapienza, Roma, Italy*
- ²⁷*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ²⁸*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ²⁹*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ³⁰*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³¹*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³²*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³³*Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- ³⁴*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- ³⁵*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³⁶*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*
- ³⁷*Yandex School of Data Analysis, Moscow, Russia*
- ³⁸*Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- ³⁹*Institute for High Energy Physics (IHEP), Protvino, Russia*
- ⁴⁰*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴¹*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴²*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴³*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁴*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁵*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷*University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵²*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵*Imperial College London, London, United Kingdom*
- ⁵⁶*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷*Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸*Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁹*University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁰*University of Maryland, College Park, MD, United States*
- ⁶¹*Syracuse University, Syracuse, NY, United States*

- ⁶²*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to* ²
- ⁶³*University of Chinese Academy of Sciences, Beijing, China, associated to* ³
- ⁶⁴*School of Physics and Technology, Wuhan University, Wuhan, China, associated to* ³
- ⁶⁵*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to* ³
- ⁶⁶*Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to* ⁸
- ⁶⁷*Institut für Physik, Universität Rostock, Rostock, Germany, associated to* ¹²
- ⁶⁸*Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to* ²⁷
- ⁶⁹*National Research Centre Kurchatov Institute, Moscow, Russia, associated to* ³⁴
- ⁷⁰*National University of Science and Technology "MISIS", Moscow, Russia, associated to* ³⁴
- ⁷¹*National Research Tomsk Polytechnic University, Tomsk, Russia, associated to* ³⁴
- ⁷²*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to* ⁴⁰
- ⁷³*University of Michigan, Ann Arbor, United States, associated to* ⁶¹
- ⁷⁴*Los Alamos National Laboratory (LANL), Los Alamos, United States, associated to* ⁶¹

^a*Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

^b*Laboratoire Leprince-Ringuet, Palaiseau, France*

^c*P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*

^d*Università di Bari, Bari, Italy*

^e*Università di Bologna, Bologna, Italy*

^f*Università di Cagliari, Cagliari, Italy*

^g*Università di Ferrara, Ferrara, Italy*

^h*Università di Genova, Genova, Italy*

ⁱ*Università di Milano Bicocca, Milano, Italy*

^j*Università di Roma Tor Vergata, Roma, Italy*

^k*Università di Roma La Sapienza, Roma, Italy*

^l*AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*

^m*LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*

ⁿ*Hanoi University of Science, Hanoi, Vietnam*

^o*Università di Padova, Padova, Italy*

^p*Università di Pisa, Pisa, Italy*

^q*Università degli Studi di Milano, Milano, Italy*

^r*Università di Urbino, Urbino, Italy*

^s*Università della Basilicata, Potenza, Italy*

^t*Scuola Normale Superiore, Pisa, Italy*

^u*Università di Modena e Reggio Emilia, Modena, Italy*

^v*MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*

^w*Novosibirsk State University, Novosibirsk, Russia*

^x*National Research University Higher School of Economics, Moscow, Russia*

^y*Sezione INFN di Trieste, Trieste, Italy*

^z*Escuela Agrícola Panamericana, San Antonio de Oriente, Honduras*

^{aa}*School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China*

^{ab}*Physics and Micro Electronic College, Hunan University, Changsha City, China*

[†]*Deceased*