

The archaeology and ethnoarchaeology of rain-fed cultivation in arid and hyper-arid North Africa

Carla Lancelotti^{1,*}, Stefano Biagetti^{1,2}, Andrea Zerboni³,
Donatella Usai⁴ & Marco Madella^{1,2,5}



Rain-fed cultivation in drylands—especially in arid and hyper-arid areas—is often considered to play a minor role in human subsistence. Drawing upon the results of ethnoarchaeological research in North Africa, this paper reviews non-irrigated agricultural practices in the absence of anthropogenic water-harvesting structures, and presents a proposal for how such practices can be identified in the drylands of the past. An improved understanding of the long-term development of rain-fed cultivation augments our knowledge of past land-use strategies and can inform future models of sustainable agriculture in some of the world’s driest regions.

Keywords: Sahara, Al Khiday, rain-fed cultivation, drylands, resilience, phytoliths

Introduction

Today, drylands cover approximately 40 per cent of the world’s land area and host around 40 per cent of the world’s population. The United Nations Environment Programme defines drylands as tropical and temperate areas with an aridity index of less than 0.65, including hyper-arid drylands or deserts with an aridity index below 0.05 (Safariel & Adeel 2005: 626). These areas are especially sensitive to climate change, shifts in precipitation patterns

¹ CaSEs Research Group, Department of Humanities, Universitat Pompeu Fabra, c/Trias Fargas 25–27, Barcelona 08005, Spain

² School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, 1 Jan Smuts Avenue, Braamfontein 2000, Johannesburg, South Africa

³ Dipartimento di Scienze della Terra ‘A. Desio’, Università degli Studi di Milano, via Mangiagalli 34, Milan, Italy

⁴ Centro Studi Sudanesi e Sub-Sahariani, Strada di Cannizzano 128/d, Treviso 31100, Italy

⁵ Catalan Institution for Research and Advanced Studies, Passeig Lluís Companys 23, Barcelona 08010, Spain

* Author for correspondence (Email: carla.lancelotti@upf.edu)

and increases in extreme climatic and meteorological events (Huang *et al.* 2017). Nonetheless, drylands have been inhabited for thousands of years, and are today home to societies that display a wide array of adaptive behaviours. Such practices, evidenced by traditional ecological knowledge, are key to the resilience of these communities.

In this article, we present a review of rain-fed agriculture in the absence of water-harvesting structures in the arid and hyper-arid lands of North Africa; farming in the Sahara is a surprisingly common practice even though it is omitted from most maps of current and past land use. On the basis of historical records and our field observations, we argue that this type of rain-fed agriculture in arid and hyper-arid lands is an overlooked practice that holds huge potential for designing effective strategies to cope with current climate-driven desertification. Further, we suggest that rain-fed cultivation without water-harvesting techniques might have played a so-far unrecognised role in the development of food production in arid areas in the past, even where rainfall is normally considered too scarce for crop cultivation.

In Old World contexts, considerable research has been directed to the study of the origin of agriculture, focusing especially on the Near East—one of the cradles of domestication—and crops such as wheat and barley that provide the foundation for many modern agricultural systems. It is clear, however, that past human groups also experimented with a diverse assemblage of crops, some of which (e.g. millets) are well adapted to areas with water constraints. Recent work by Winchell *et al.* (2018) has traced the domestication trajectories of African crops, including sorghum (*Sorghum bicolor* (L.) Moench) and pearl millet (*Pennisetum glaucum* (L.) R.Br). Past cultivation practices related to these crops, including how water was managed, however, are still under study.

In this article, we first provide an overview of rain-fed cultivation of cereals in the arid and hyper-arid lands of North Africa, based upon the study of ethnographies from the nineteenth and twentieth centuries, as well as a few more recent observations. We then present the results from Al Khiday, Sudan, where a multidisciplinary pilot investigation on rain-fed crops was carried out in 2015–2016. This case study has driven the development of a larger research project (RAINDROPS), featuring an experimental methodology aimed at identifying crop-water availability and management directly from archaeobotanical remains.

A review of rain-fed cultivation in arid and hyper-arid North Africa

In arid lands, irrigation, a permanent water source, or some form of rainwater-harvesting is usually deemed necessary to support cultivation. There are modern examples, however, that demonstrate the presence, even in hyper-arid areas, of successful rain-fed cultivation systems. For the purposes of this article, we employ the following definitions:

- a) Irrigated cultivation: water is provided to crops at regular intervals throughout the growing season by human intervention.
- b) *Décrue* cultivation: water is provided by natural inundation, typically from major river systems (floodplain cultivation).
- c) Rain-fed cultivation: water is provided by rainfall alone (directly or as run-off); cultivation occurs far from any permanent water sources (e.g. rivers, wells, lakes or cisterns) and without any water harvesting.

This last type, rain-fed cultivation in arid and hyper-arid environments, has received comparatively little research attention as it is often considered to play only a minor role in human livelihoods. Although the harvest yielded by this type of cultivation is lower compared to irrigated or *décrue* cultivation, its benefits as an adaptive strategy in drylands can be important. Indeed, rain-fed crops, such as millets and sorghum, may offer a considerable contribution to a predominantly pastoral economy, as they provide carbohydrates and a high percentage of daily protein and minerals (Gulia *et al.* 2007). Such crops are currently identified as ‘future smart foods’ (Li & Siddique 2018), as they have the potential to offer better options than traditional crops, such as wheat/barley or corn, for mitigating the effects of environmental and, especially, climate change, and to supply affordable and sustainable dietary diversity.

North Africa is dominated by the Sahara, which is often considered to be agriculturally ‘inactive’ (Rockström & Falkenmark 2015). Indeed, land-use models and maps of North Africa generally characterise the Sahara and its margins as a huge empty area that stretches from the northern edge of the Sahel up to the Mediterranean coast and the Atlas Mountains (Rockström & Falkenmark 2015). Similarly, modern technical reports and studies on dryland food production tend to regard the Sahara as being devoid of cultivation (Portmann *et al.* 2010).

The perceived dominance of pastoralism, together with the notion that hot deserts are unsuitable for growing crops, seems to have obscured the role that cultivation has played in the past. While therefore not surprising that the extent of rain-fed modern and ancient agriculture in the Sahara is poorly understood, observations published over the past two centuries indicate the practice of crop cultivation in different parts of the Sahara (Figure 1). In central Mauritania, for example, where rainfall averages 170mm/year, rain-fed crops were observed in the early 1960s, and so-called pastoral communities were reportedly growing cereals without irrigation (Toupet 1963).

Millet was grown in the nineteenth century in the Ahaggar Mountains, where rainfall varies between 0 and 100mm/year (Duveyrier 1864: 372). The Tuareg peoples inhabiting the valleys of the Ahaggar used a specific word (*tawgest*) to designate non-irrigated plots of cultivated land (Nicolaisens & Nicolaisens 1997: 251). Similarly, in the Air massif, where average precipitation is 120mm/year, Rodd (1926: 133) observed that rain-fed cereals could be grown most years. The engagement of nomadic Tuareg groups with rain-fed cultivation is further confirmed by Nicolaisen and Nicolaisen (1997: 250–51), who observed rain-fed wheat and millet/sorghum fields in the Tassili n’Ajjer in the early 1950s. This last report is particularly surprising, given the very low (and mostly uneven) rainfall in this area, which ranges between 0 and 40mm/year. Cultivation has also been recorded in the Tadrart Acacus massif in south-west Libya (di Lernia *et al.* 2012) and around the city of Ghat (Bourbon del Monte di Santa Maria 1912), where annual rainfall is less than 20mm. In the Tibesti Mountains of south-east Libya, close to the Guezendi area, rain-fed cereal plots were observed in areas with less than 50mm/year of precipitation (Desio 1942). These examples clearly show that the cultivation of rain-fed crops complemented pastoral and foraged resources in the Sahara in the recent past.




Figure 1. CGIAR-CSI aridity index and the localities mentioned in the text. The aridity index takes into account the values of precipitation, temperature and/or potential evapo-transpiration. The cases of rain-fed agriculture in the Sahara fall within the 'hyper-arid' (Ghat, Guezendou, Tassili, Tadrart Acacus), or 'arid' (Al Khiday, Moudjeria, and Air) class—well beyond the threshold for the supposed suitability of rain-fed agriculture. Map produced in QGIS by the authors; baseline data (Aridity Index values) obtained from <https://cgiarcsi.community/data/global-aridity-and-pet-database/> (accessed 10 July 2019) (Zomer et al. 2007, 2008).




Figure 2. GoogleEarth view of RFA15_01 and RFA15_02/04: a) the study area; b) RFA15_01, imagery taken in November 2014, with plough marks clearly visible; c) RFA15_02/04, imagery taken in September 2012 showing plots of cultivated lands; the crops growing form a pattern of coarsely parallel, vertical lines, clearly bounded by uncultivated land (one of the field's boundaries is indicated by the arrow) (GIS elaboration by the authors using QGIS).

The Al Khiday case study

In 2015–2016, we conducted ethnoarchaeological surveys in the area west of Al Khiday in central Sudan (Figure 2), with the aim of investigating the late prehistoric and historical landscape of the region, as well as the traditional ecological knowledge of contemporary populations. Today, the area is categorised as arid by both the Koppen-Geigen and CGIAR-CSI classifications (Zomer *et al.* 2007, 2008), with an aridity index of 0.0765, a yearly average rainfall of approximately 100mm (concentrated in July, August and partially September) and widespread sandy aridisols (i.e. soils typical of arid and hot environments, characterised by low organic content and colonised by drought- and salt-tolerant vegetation). Donatella Usai has been conducting archaeological investigations in this region since 2001 (Usai 2018). Upon joining her project in 2015, we noted the presence of many regularly spaced, rows of tufts of vegetation, located around 30km inland from the Nile, in an area devoid of farms, wells and permanent rivers (Figure 3). The following year, systematic fieldwork led to the identification of several rain-fed fields of pearl millet, located between 15 and 30km west of the White Nile.

Millets, and specifically pearl millet, are key crops in drylands due to their short growing season and their high productivity in arid, high-temperature conditions. As both the entire plant (before the grains mature) and the by-products from grain-processing can also be used



Figure 3. View of recently harvested fields in the Al Khiday area in December 2016. The upper photograph shows a general view, where the margins of the fields are clearly marked in straight lines by the wild grass cover. The lower photograph shows a recently harvested plot, with tufts of millets still visible in regular rows (photographs by the authors).

as fodder, millets are well suited for mixed agro-pastoral systems. Pearl millet is a critical West Africa crop, and the Sahel zone south of the Sahara is an important area for its domestication (D'Andrea & Casey 2002; Winchell *et al.* 2018). It is particularly suitable for rain-fed cultivation in hyper-arid areas, as it is well adapted to soil with low fertility and to high temperatures, and is able to grow with as little as 40mm of water per annum.

Desktop and field research

Our analysis of time-series imagery (GoogleEarth™), combined with rainfall data for the last 20 years, highlights that 70mm of rain in July/August is sufficient for pearl millet cultivation over extremely large expanses of land (up to 120ha per field system; Figure 2).

We conducted ethnographic interviews with local farmers, who revealed that the cultivated areas are divided into plots (Figure 3) owned by different families from neighbouring villages. After the late spring/early summer rains, the fields are hoed to clear wild grasses while the soil is still wet. Farmers then walk the fields along parallel lines, making holes and planting

seeds approximately every metre. The sown fields are then left unattended until the harvest in September, at which time the farmers use iron sickles to cut the plants just above the crown. Grains are then threshed in the fields, and the by-products are left for the domestic livestock to graze. Although no productivity data are currently available for this type of arid or hyper-arid rain-fed agriculture, a preliminary evaluation—based on plant density and average single-plant seed production—gives a value of approximately 0.1t of millet per hectare. These figures are clearly much lower than the production in dry sub-humid and semi-arid regions of the world, where rain-fed agriculture yields vary between 0.5 and 2t/ha (Rockström & Falkenmark 2015). In spite of the low yield, however, these farmers in central Sudan are able to produce twice the quantity of millet necessary for the families' annual needs, leaving part of the crop to be sold at local markets after having set aside enough seed for the next planting season. What is surprising about these rain-fed fields in Sudan is that, in wetter years (with a rainfall of approximately 100mm, for example), a double cropping of millet, along with a cash crop, such as hibiscus (*Hibiscus* sp.) can be produced.

Phytolith analysis: preliminary results of a pilot project

In 2015, we conducted a pilot project at Al Khiday to check for the presence and preservation of phytoliths in modern deposits associated with rain-fed cultivation and to confirm the morphological ratios of phytoliths in relation to water availability, as highlighted by previous studies (Jenkins *et al.* 2016). Phytoliths are silica bodies produced in plant tissues. Their inorganic nature means that they are far less affected by pre- and post-depositional issues than organic plant material, and hence, their preservation is ubiquitous (Piperno 2006). Two different field systems were sampled: two small ephemeral fields (RFA15_02 and RFA15_04; Figure 2) that had been cultivated the previous year, and an abandoned field (RFA15_01; Figure 2b) that had been cultivated for over 10 years, before being abandoned during the last 2–3 years. According to local informants, this latter field was mechanically ploughed but relied only on rain for water. Comparative samples were collected from inside and outside the furrows and holes used for planting (Table 1). Phytoliths were extracted and analysed according to standard laboratory protocols employed at the Laboratory for Environmental Archaeology at the Universitat Pompeu Fabra (Lancelotti *et al.* 2017).

All the analysed samples yielded well-preserved phytoliths, with little or no evidence of chemical or mechanical damage, and several different morphotypes. Phytolith concentration is generally low and seems dependent on the intensity of use (i.e. both how long and also the different ways in which these fields were cultivated), as it is generally higher in RFA15_01 than in RFA15_02 and 04. The number of morphotypes identified, however, is virtually the same in both cases, thus indicating good representativeness of the original vegetation cover (Madella & Lancelotti 2012). Another notable aspect is the relationship between two of the main types recovered: bulliforms and trichomes. Samples from RFA15_02 and 04 yielded proportionally more bulliforms (average = 68) and fewer trichomes (average = 42) than those from RFA15_01 (Figure 4). These two morphotypes are considered indicative of environmental conditions in C4 grasses (Olsen *et al.* 2013). Specifically, plants grown in mesic (more humid conditions) tend to have more bulliforms and fewer trichomes than plant grown in xeric (drier) conditions. Our results suggest that the plants from the less

Table 1. Phytolith data; summary results of the phytolith analysis performed at the three sampling locations.

Field	Trench	Sample	Concentration g(AIF)	Morphotypes (n)	Bulliforms	Trichomes	Ratio bull: tric
RFA 15_01	1	Outside	32 427	13	20	93	0.22
		Outside	79 389	14	5	74	0.07
		Inside	177 932	12	14	59	0.24
		Inside	150 907	19	18	58	0.31
RFA 15_02	1	Inside	593	19	109	54	2.02
		Outside	459	18	84	34	2.47
	4	Top	507	12	44	53	0.83
		Bottom	1399	12	39	18	2.17
RFA 15_04	2	Inside	253	12	78	60	1.30
		Outside	1576	17	55	33	1.67




Figure 4. Phytolith analysis. Proportions of bulliforms (upper-left photograph) and trichomes (lower-right photograph) identified at the three sampling locations (photographs by the authors).

intensely cultivated area and farther from water sources (RFA15_02/04) grew in more humid conditions than those in the ploughed field. This may be a result of the techniques used for working the soil in preparation for sowing, with the intensive mechanical ploughing of

RFA15_01 depleting the soil of moisture, whereas the soil in RFA15_02 and 04 conserved more humidity. These preliminary results confirm the viability of the methodology and encourage the further exploration of this line of research.

Exploring the past: a methodological proposal

The results of the pilot project show the potential of phytoliths as a proxy to study crop-water management practices in drylands. The methodology, however, needs to be further tested and improved in order to obtain significant results. We present a proposal on how this research should develop.

State of the art

Research on past crops in North Africa—especially for the drier areas—is still very limited. During the Early and Middle Holocene (10 000–5000 BC), patchy, savannah-like vegetation supported small communities of hunter-gatherers that exploited a range of wild millet-type grasses (Fuller 2005), including attempts at opportunistic cultivation without domestication (Mercuri *et al.* 2018). These populations had certainly adopted livestock by 6000–5000 BC (Marshall & Hildebrand 2002; Fuller 2005). The shift from the more humid ‘Green Sahara’ (Holocene climatic optimum) to the present-day conditions occurred by the end of the Middle Holocene, and it is generally marked by relevant socio-economic transformations (Gasse 2000; Kuper & Kröpelin 2006). After 3500 BC, when the climate conditions became drier, some of the nomadic pastoral groups seem to have adopted cultivation in selected Saharan locations (Cremaschi & Zerboni 2009). Yet, the role that rain-fed agriculture may have played in such a critical transition has not been explored, and very few data are available on the earliest agricultural experiments in the Sahara. Past studies have emphasised the contemporaneous adoption of more drought-tolerant livestock, such as the dromedary and goat, in parallel with the progressive abandonment of cattle (Clarke *et al.* 2016).

Research on rain-fed practices in the Sahara and an understanding of their chronological depth is urgently needed in order to address the role of cultivation in the resilience of human societies living in unfavourable climatic settings. Notwithstanding the importance that rain-fed agriculture could have had in the Sahara and North Africa during the Late Holocene, the practice has only occasionally formed part of the current discourse on strategies of human adaptation. This is probably due to the challenges in detecting direct evidence for rain-fed cultivation in the archaeological record. The identification of ancient fields, for example, is extremely difficult, even using techniques such as geoarchaeology or molecular footprints (Motuzaitė-Matuzeviciute *et al.* 2016), and fields are usually inferred from the presence of related technology, such as ploughs and terracing. Furthermore, the different types of water management are also difficult to discern, and currently there are few proxies able to identify and differentiate such practices clearly. Determining water availability directly from archaeobotanical material offers a possible solution.

Previous efforts to address past water-management practices have explored charred macrobotanical remains (Jenkins *et al.* 2016), have analysed weed assemblages and performed isotopic studies of $\delta^{13}\text{C}$. All of these analyses have demonstrated good potential for assessing

The archaeology and ethnoarchaeology of rain-fed cultivation in arid and hyper-arid North Africa

water-management practices in C3 plants, such as trees, legumes and winter cereals (Styring *et al.* 2016). Approaches exclusively based on charred remains, however, can prove problematic in drylands, where they are often poorly preserved. Consequently, the analysis of plant microfossils, such as phytoliths, has been explored as an alternative (Madella *et al.* 2009; Jenkins *et al.* 2016). Morphology—assessed through indices based on morphotype proportions—and isotopic analyses have been explored to identify different water regimes from archaeobotanical samples (McClaran & Umlauf 2000; Webb & Longstaffe 2003; Madella *et al.* 2009; Hodson 2016; Jenkins *et al.* 2016).

RAINDROPS: an innovative approach

RAINDROPS aims to develop an innovative and reliable methodology for the identification of crop-water availability and management based on archaeobotanical remains. Highly controlled data on phytolith ratios and carbon, oxygen and silicon isotopes from macro- and micro-remains from experimental fields are being validated with ethnographic evidence, before being applied to selected key archaeological case studies in Africa and Asia.

Crops grown in experimental fields at the International Crop Research Institute for the Semi-Arid Tropics—under rigorously controlled conditions—form the basis of genotypic variation in phytolith production and isotopic fractionation for sorghum, finger millet (*Eleusine coracana* Gaertn.) and pearl millet landraces. To strengthen further the comparison between the experimental isotopic data and archaeological samples, the effect of charring on isotopic ratios is also being assessed.

Phytolith concentration and ratios between genetically *vs* environmentally controlled morphotypes analysed from experimentally grown plants are being tested to confirm that the observed correlation with plant-water availability identified in C3 plants holds true for C4 species. Simultaneously, oxygen and silicon isotopes are assessed for their potential use as proxies for evaluating crop-water availability and soil conditions. Pilot projects have anticipated the possible benefits of such an approach and have laid a strong foundation for the analysis of these oxygen and silicon isotopes in phytoliths (Chapligin *et al.* 2011). These studies indicate that the degree of silica ¹⁸O enrichment in leaves can be related to water loss through transpiration and relative humidity, and that silicon isotopes are sensitive to environmental conditions and changes (Webb & Longstaffe 2003; Opfergelt *et al.* 2006; Ding *et al.* 2008; Leng *et al.* 2009).

Ethnographic data and traditional ecological knowledge

The same set of analyses is being performed on crops grown locally using traditional methods (no chemical fertilisers and pesticides) in three study areas: Sudan, Ethiopia and Pakistan. At the same time, structured and unstructured interviews are being conducted with a number of local farmers—according to established ethnographic protocols—specifically, to assess water-management practices, as well as to collect traditional ecological knowledge on cultivation systems. These interviews target quantifiable data on cultivation methods, such as tilling, weeding, manuring and ploughing methods, as well as information on decision-making

related to where and when cultivation is carried out and how these processes are affected by climate variability.

The three case studies (Sudan, Ethiopia and Pakistan) were selected because of their geographical and environmental locations, which represent biophysical hotspot areas in terms of their significance in the long-term human adaptation to drylands, as well as for the presence of preliminary results—such as for Al Khiday, with the ethnoarchaeological work presented here. Furthermore, each site is located in one of the three ecological areas that broadly define the concept of drylands (United Nations Environment Programme 1997): arid to hyper-arid (Al Khiday, Sudan), dry sub-humid (Mezber, Ethiopia) and semi-arid (Harappa, Pakistan), thereby covering the geo-climatic variability of drylands. Water-management practices at these sites have been hypothesised on the basis of local geographic or geomorphological features, economy and degree of social organisation (Giosan *et al.* 2012; C. D’Andrea *pers. comm.*), although no test for these hypotheses has ever been provided. All of the sites are archaeologically well studied, with abundant palaeoenvironmental and palaeoclimatic evidence and long occupation sequences—factors that will allow for the study of the long-term dynamics of human-environment interaction (Terwilliger *et al.* 2011; Giosan *et al.* 2012; D’Andrea *et al.* 2015; Williams *et al.* 2015; Iacumin *et al.* 2016). Lastly, the three sites are located in areas where traditional agriculture is still practised, and where a rich source of traditional ecological knowledge exists. RAINDROPS is both recording and utilising this information as part of its data analyses.

Concluding remarks

Desertification has been internationally recognised as one of the main challenges of sustainable development, and in 1994, the United Nations Convention to Combat Desertification (UNCCD) was created. One of the aims of the UNCCD is to put science, technology and traditional knowledge at the forefront in mitigating desertification. In this perspective, environmental archaeology and ethnoarchaeology can play a fundamental role in understanding the dynamics of long-term strategies of human adaptation and resilience in drylands. Archaeologically, considerable research has been devoted to the study of the origins of agriculture—focusing especially on the Near East—and crops such as wheat and barley that provide the foundation for modern agricultural systems in the Old World. It is clear, however, that past human groups also experimented with a diverse assemblage of crops, some of which, such as millets, are well adapted to areas with water constraints. Such crops constitute one of the most important sources for future biodiverse and sustainable agriculture in many parts of the world. Furthermore, drylands agriculture is becoming highly relevant in development studies, due to the impact of desertification on food security for a great proportion of the human population. Our current work in Sudan highlights the importance of rain-fed cultivation in drylands, and our approach will provide the research community with an innovative and reliable methodology for understanding past water-management practices for C4 crops. This will extend the chronology of traditional ecological knowledge in key low-precipitation areas, contributing long-term data to explore resilience and to improve model-building for future policies.

Acknowledgements

We would like to thank all those who helped during fieldwork, particularly Mungida Khaled Magzoub and Ahmed Altayeb, as well as the Director General of the National Corporation of Antiquities and Museums (Sudan), Abdelrhman Ali Mohamed, for granting the licence. Funding is provided by the Italian Ministry for Foreign Affairs (MAE-DGSP VI). RAINDROPS has received funding from the European Union's Horizon 2020 research and innovation programme (ERC-Stg-2017) under grant agreement 759800. Additional funding for fieldwork was provided by the University of Milano. This work is also part of the PAGES LandCover6k effort and the INQUA International Focus Group HoLa (Holocene Global Landuse). Fieldwork in Al Khiday has been conducted in the area licensed to Donatella Usai of the Center for Sudanese and Sub-Saharan Studies. The licence includes permits for exporting sediments for analysis, for which clearance has also been obtained by the National Corporation for Antiquities and Museums in Khartoum. Oral consent has been obtained from the people interviewed during ethnoarchaeological surveys. No personal or sensitive data have been collected during, or stored after, these interviews. Universitas Pompeu Fabra health and safety measures have been followed during both field and laboratory work. C.L., S.B. and M.M. are members of CaSEs (UPF), an 'Excellence Research Group' of the Catalan Agency for Research (AGAUR SGR-1417 and SGR-0212) and an associated unit (unidad asociada) of the Institució Milà i Fontanals of the Spanish National Research Council (IMF-CSIC).

References

- BOURBON DEL MONTE SANTA MARIA, G. 1912. *L'oasi di Ghat e sue adiacenze, Comando del Corpo di Stato Maggiore (Ufficio Coloniale)*. Citta' di Castello: Tipografia dell'Unione Arti Grafiche.
- CHAPLIGIN, B., M.J. LENG, E. WEBB, A. ALEXANDRE, J.P. DODD, A. IJIRI & F.J. LONGSTAFFE. 2011. Inter-laboratory comparison of oxygen isotope compositions from biogenic silica. *Geochimica et Cosmochimica Acta* 75: 7242–56. <https://doi.org/10.1016/j.gca.2011.08.011>
- CLARKE, J. *et al.* 2016. Climatic changes and social transformations in the Near East and North Africa during the 'long' 4th millennium BC: a comparative study of environmental and archaeological evidence. *Quaternary Science Review* 136: 96–121. <https://doi.org/10.1016/j.quascirev.2015.10.003>
- CREMASCHI, M. & A. ZERBONI. 2009. Early to Middle Holocene landscape exploitation in a drying environment: two case studies compared from the Central Sahara (SW Fezzan, Libya). *Comptes Rendus Géoscience* 341: 689–702. <https://doi.org/10.1016/j.crte.2009.05.001>
- D'ANDREA, A.C. & J. CASEY. 2002. Pearl millet and Kintampo subsistence. *African Archaeological Review* 19: 147–73. <https://doi.org/10.1023/A:1016518919072>
- D'ANDREA, A.C., A.G. FAHMY, L. PERRY, M.P. RICHARDS, L. DARCUS, M. TOFFOLO & A.E. ATTIA EL SHAFAY. 2015. Ancient agricultural economy in the Horn of Africa: new evidence from grinding stones and stable isotopes. *Proceedings of the IWAAS, Società dei Naturalisti e Matematici di Modena* 144: 155–57.
- DESIO, A. 1942. *Il Sahara Italiano. Il Tibesti nord-orientale: Reale Società Geografica Italiana*. Roma: Società Italiana Arti Grafiche.
- DI LERNIA, S., I. MASSAMBA N'SIALA & A. ZERBONI. 2012. 'Saharan waterscapes': Traditional knowledge and historical depth of water management in the Akakus Mts. (SW Libya), in L. Mol & T. Sternberg (ed.) *Changing deserts: integrating people and their environment*: 101–28. Cambridge: White Horse.
- DING, T.P., J.X. ZHOU, D.F. WAN, Z.Y. CHEN, C.Y. WANG & F. ZHANG. 2008. Silicon isotope fractionation in bamboo and its significance to the biogeochemical cycle of silicon. *Geochimica et Cosmochimica Acta* 72: 1381–95. <https://doi.org/10.1016/j.gca.2008.01.008>
- DUVEYRIER, H. 1864. *L'exploration du Sahara. Les touaregs du Nord*. Paris: Challamel.
- FULLER, D.Q. 2005. Farming: Stone Age farmers of the savanna, in K. Shillington (ed.) *Encyclopedia of African history*: 521–23. New York: Fitzroy Dearborn.
- GASSE, F. 2000. Hydrological changes in the African tropics since the Last Glacial Maximum. *Quaternary Science Reviews* 19: 189–211. [https://doi.org/10.1016/S0277-3791\(99\)00061-X](https://doi.org/10.1016/S0277-3791(99)00061-X)
- GIOSAN, L. *et al.* 2012. Fluvial landscapes of the Harappan civilization. *Proceedings of the National Academy of Sciences of the USA* 109: E1688–94. <https://doi.org/10.1073/pnas.1112743109>
- GULLIA, S.K., J.P. WILSON, J. CARTER & B.P. SINGH. 2007. Progress in grain pearl millet research and

- market development, in J. Janick & A. Whipkey (ed.) *Issues in new crops and new uses*: 196–203. Alexandria (VA): ASHS.
- HODSON, M.J. 2016. The development of phytoliths in plants and its influence on their chemistry and isotopic composition. Implications for palaeoecology and archaeology. *Journal of Archaeological Science* 68: 62–69. <https://doi.org/10.1016/j.jas.2015.09.002>
- HUANG, J. et al. 2017. Dryland climate change: recent progress and challenges. *Reviews of Geophysics* 55: 719–78. <https://doi.org/10.1002/2016RG000550>
- IACUMIN, P., A. DI MATTEO, D. USAI, S. SALVADORI & G. VENTURELLI. 2016. Stable isotope study on ancient populations of central Sudan: insights on their diet and environment. *American Journal of Physical Anthropology* 160: 498–518. <https://doi.org/10.1002/ajpa.22987>
- JENKINS, E., K. JAMJOUR, S. NUIMAT, R. STAFFORD, S. NORTCLIFF & S. MITHEN. 2016. Identifying ancient water availability through phytolith analysis: an experimental approach. *Journal of Archaeological Science* 73: 82–93. <https://doi.org/10.1016/j.jas.2016.07.006>
- KUPER, R. & S. KRÖPELIN. 2006. Climate-controlled Holocene occupation in the Sahara: motor of Africa's evolution. *Science* 313: 803–807. <https://doi.org/10.1126/science.1130989>
- LANCELOTTI, C., J. RUIZ-PÉREZ & J.J. GARCÍA-GRANERO. 2017. Investigating fuel and fireplaces with a combination of phytoliths and multi-element analysis: an ethnographic experiment. *Vegetation History and Archaeobotany* 26: 75–83. <https://doi.org/10.1007/s00334-016-0574-y>
- LENG, M.J., G.E. SWANN, M.J. HODSON, J.J. TYLER, S.V. PATWARDHAN & H.J. SLOANE. 2009. The potential use of silicon isotope composition of biogenic silica as a proxy for environmental change. *Silicon* 1: 65–77. <https://doi.org/10.1007/s12633-009-9014-2>
- LI, X. & K.H.M. SIDDIQUE. 2018. *Future smart food —rediscovering hidden treasures of neglected and underutilized species for zero hunger in Asia*. Bangkok: Food and Agriculture Organisation of the United Nations. <https://doi.org/10.18356/23b5f7ab-en>
- MADELLA, M. & C. LANCELOTTI. 2012. Taphonomy and phytoliths: a user manual. *Quaternary International* 275: 76–83. <https://doi.org/10.1016/j.quaint.2011.09.008>
- MADELLA, M., M.K. JONES, P. ECHLIN, A. POWERS-JONES & M. MOORE. 2009. Plant water availability and analytical microscopy of phytoliths: implications for ancient irrigation in arid zones. *Quaternary International* 193: 32–40. <https://doi.org/10.1016/j.quaint.2007.06.012>
- MARSHALL, F. & E. HILDEBRAND. 2002. Cattle before crops: the beginnings of food production in Africa. *Journal of World Prehistory* 16: 99–144. <https://doi.org/10.1023/A:1019954903395>
- MCCCLARAN, M.P. & M. UMLAUF. 2000. Desert grassland dynamics estimated from carbon isotopes in grass phytoliths and soil organic matter. *Journal of Vegetation Science* 11: 71–76. <https://doi.org/10.2307/3236777>
- MERCURI, A.M., R. FORNACIARI, M. GALLINARO, S. VANIN & S. DI LERNIA. 2018. Plant behaviour from human imprints and the cultivation of wild cereals in Holocene Sahara. *Nature Plants* 4: 71–81. <https://doi.org/10.1038/s41477-017-0098-1>
- MOTUZAITE-MATUZEVICIUTE, G., J. JACOB, S. TELIZHENKO & M.K. JONES. 2016. Miliacin in palaeosols from an Early Iron Age in Ukraine reveal in situ cultivation of broomcorn millet. *Archaeological and Anthropological Sciences* 8: 43–50. <https://doi.org/10.1007/s12520-013-0142-7>
- NICOLAISEN, J. & I. NICOLAISEN. 1997. *The pastoral Tuareg: ecology, culture, and society*. Copenhagen: Thames & Hudson.
- OLSEN, J.T., K.L. CAUDLE, L.C. JOHNSON, S.G. BAER & B.R. MARICLE. 2013. Environmental and genetic variation in leaf anatomy among populations of *Andropogon gerardii* (Poaceae) along a precipitation gradient. *American Journal of Botany* 100: 1957–68. <https://doi.org/10.3732/ajb.1200628>
- OPFERGELT, S., D. CARDINAL, C. HENRIET, L. ANDRÉ & B. DELVAUX. 2006. Silicon isotope fractionation between plant parts in banana: *in situ vs. in vitro*. *Journal of Geochemical Exploration* 88: 224–27. <https://doi.org/10.1016/j.gexplo.2005.08.044>
- PIPERNO, D.R. 2006. *Phytoliths: a comprehensive guide for archaeologists and paleoecologists*. Lanham (MD): Altamira.
- PORTMANN, F.T., S. SIEBERT & P. DÖLL. 2010. MIRCA2000—global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and

- hydrological modeling. *Global Biogeochemical Cycles* 24: 1–24.
<https://doi.org/10.1029/2008GB003435>
- ROCKSTRÖM, J. & M. FALKENMARK. 2015. Agriculture: increase water harvesting in Africa. *Nature* 519: 283–85.
<https://doi.org/10.1038/519283a>
- RODD, F.R. 1926. *People of the veil*. London: Macmillan.
- SAFRIEL, U. & Z. ADEEL. 2005. Drylands systems, in R. Hassan, R. Sholes & N. Ash (ed.) *Ecosystems and human well-being: current state and trends*: 625–62. Washington, D.C.: Island.
- STYRING, A.K., M. ATER, Y. HMIMSA, R. FRASER, H. MILLER, R. NEEF, J.A. PEARSON & A. BOGAARD. 2016. Disentangling the effect of farming practice from aridity on crop stable isotope values: a present-day model from Morocco and its application to early farming sites in the Eastern Mediterranean. *The Anthropocene Review* 3: 2–22.
<https://doi.org/10.1177/2053019616630762>
- TERWILLIGER, V.J., Z. ESHETU, M. ALEXANDRE, Y. HUANG, M. UMER & T. GEBRU. 2011. Local variation in climate and land use during the time of the major kingdoms of the Tigray Plateau of Ethiopia and Eritrea. *Catena* 85: 130–43.
<https://doi.org/10.1016/j.catena.2010.08.003>
- TOUPET, C. 1963. L'évolution de la nomadisation en Mauritanie sahelienne, in C. Bataillon (ed.) *Nomades et nomadisme au Sahara: recherches sur la zone aride XIX*: 67–79. Munich: R. Oldenbourg.
- United Nations Environment Programme. 1997. *World atlas of desertification*. London: United Nations Environment Programme.
- USAI, D. 2018. Prehistory in central Sudan, in M. Honegger (ed.) *Nubian archaeology in the XXIst century. Proceedings of the Thirteenth International Conference for Nubian Studies, Neuchatel, 1–6 September 2014*: 3–18. Leuven: Peeters.
- WEBB, E.A. & F.J. LONGSTAFFE. 2003. Climatic influences on the oxygen isotopic composition of biogenic silica in prairie grass. *Geochimica et Cosmochimica Acta* 66: 1891–1904.
[https://doi.org/10.1016/S0016-7037\(02\)00822-0](https://doi.org/10.1016/S0016-7037(02)00822-0)
- WILLIAMS, M.A.J., D. USAI, S. SALVATORI, F.M. WILLIAMS, A. ZERBONI, L. MARITAN & V. LINSELE. 2015. Late Quaternary environments and prehistoric occupation in the lower White Nile valley, central Sudan. *Quaternary Science Reviews* 130: 72–88.
<https://doi.org/10.1016/j.quascirev.2015.03.007>
- WINCHELL, F., M. BRASS, A. MANZO, A. BELDADOS, V. PERNA, C. MURPHY, C. STEVENS & D.Q. FULLER. 2018. On the origins and dissemination of domesticated sorghum and pearl millet across Africa and into India: a view from the Butana Group of the Far Eastern Sahel. *African Archaeological Review* 35: 483–505.
<https://doi.org/10.1007/s10437-018-9314-2>
- ZOMER, R.J., D.H. BOSSIO, A. TRABUCCO, L. YUANJIE, D.C. GUPTA & P.S. VIRENDRA. 2007. *Trees and water: smallholder agroforestry on irrigated lands in northern India*. Colombo: International Water Management Institute.
- ZOMER, R.J., A. TRABUCCO, D.A. BOSSIO, O. VAN STRAATEN & L. VERCHOT. 2008. Climate change mitigation: a spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, Ecosystems and Environment* 126: 67–80.
<https://doi.org/10.1016/j.agee.2008.01.014>

Received: 27 November 2018; Revised: 27 February 2019; Accepted: 11 March 2019