



## W307

## Genetic Variability Study in a Wide Germplasm of Domesticated Peach Through High Throughput Genotyping

Date: Saturday, January 12, 2013

Time: 9:20 AM

Room: Pacific Salon 3

Diego Micheletti , IRTA- Centre of Research in Agricultural Genomics (CRAG), Barcelona, Spain

Valeria Aramini , CRA - Fruit Tree Research Center, Roma, Italy

Pere Arus , IRTA- Centre of Research in Agricultural Genomics (CRAG), Barcelona, Spain

Elisa Banchi , IASMA, Foundation Edmund Mach, San Michele all'Adige, Italy

Teresa Barreneche , INRA, UMR 1332 de Biologie du Fruit et Pathologie, Villenave d'Ornon, France

Daniele Bassi , University of Milano, Milano, Italy

Elisabeth Dirlewanger , INRA, UMR 1332 Biologie du Fruit et Pathologie, Villenave d'Ornon, France

Zhongshan Gao , Zhejiang University. Department of Horticulture, Hangzhou, China

Laura Gazza , CRA - Fruit Tree Research Center, Roma, Italy

Patrick Lambert , INRA. UR1052, Génét. et Amélior. des Fruits et Légumes, Montfavet, France

François Laurens , IRHS-INRA, Angers, France

Xiongwei Li , Zhejiang University. Department of Horticulture, Hangzhou, China

Thierry Pascal , INRA. UR1052, Génét. et Amélior. des Fruits et Légumes, Montfavet, France

Bénédicte Quilot-Turion , INRA. UR1052, Génét. et Amélior. des Fruits et Légumes, Montfavet, France

Laura Rossini , University of Milan, Milano, Italy

Michela Troggio , FEM-IASMA, San Michele all'Adige, Italy

Ignazio Verde , CRA - Fruit Tree Research Center, Roma, Italy

Maria Jose Aranzana , IRTA- Centre of Research in Agricultural Genomics (CRAG), Barcelona, Spain

Peach (*Prunus persica* (L.) Batsch) is one of the most economically important fruit crops in temperate areas. Classical fruit tree breeding is generally slow and inefficient. Molecular markers could improve its efficiency but, although nowadays many Mendelian traits are mapped in peach and SSR markers have been found to be linked to some of the key major genes, its use in breeding programs is still limited. Main reasons for that are insufficient linkage between the markers and the genes and the lack of markers suitable for medium-high degree of multiplexing. To address this limitation, about 1,300 peach cultivars were genotyped with the 9K peach SNP chip (Verde et al. 2012) in the frame of FruitBreedomics project. This germplasm was chosen to be representative of the genetic diversity present in five germplasm collection in Europe and in China. Out of the 8144SNPs present in the chip, about 4300 were positively genotyped and used for the further analysis. The average number of heterozygous loci in the genotyped accessions was 1186 (spanning from 13 to 2775). The preliminary results of the population structure reveal three main subpopulations and the presence of high number of admixed individuals. LD seems to decay at distance longer than ca. 1 Mb. These results will be instrumental for implementing LD-based mapping of QTLs and genes in peach.

[Back to: Fruit/Nuts](#)

[<< Previous Abstract](#) | [Next Abstract >>](#)

[Home/Search](#)[Browse by Day](#)[Browse by Type](#)[Author Index](#)[Exhibitor Index](#)[Poster Categories](#)[Personal Scheduler](#)

## Meeting Information

**When:**

January 11 - 16, 2013

**Where:**

San Diego, CA