USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-Annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 11 or above). (Note that this document uses screenshots from Adobe Reader DC.)
The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab (right-hand panel or under the Tools menu).

This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:

1. **Replace (Ins)** Tool – for replacing text.

Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it:

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.

2. **Strikethrough (Del)** Tool – for deleting text.

Strikes a red line through text that is to be deleted.

How to use it:

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.

experimental data if available. For ORFs to be had to meet all of the following criteria:

1. Small size (35-250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus with another coding feature; over both ends; or ORF containing a tRNA.

3. **Commenting** Tool – for highlighting a section to be changed to bold or italic or for general comments.

Use these 2 tools to highlight the text where a comment is then made.

How to use it:

- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.

4. **Insert** Tool – for inserting missing text at specific points in the text.

Marks an insertion point in the text and opens up a text box where comments can be entered.

How to use it:

- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.

5. **Attach File** Tool – for inserting large amounts of text or replacement figures.

Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it:

- Click on .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.

chondrial preparator
ative damage injury
e extent of membra
i, malondialdehyde (TBARS) formation. I
used by high perform

6. **Add stamp** Tool – for approving a proof if no corrections are required.

Inserts a selected stamp onto an appropriate place in the proof.

How to use it:

- Click on .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic*, *Sign Here*, *Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

or the business cycle, starting with the
. on perfect competition, constant ret
production. In this environment goods
extra profits and hence a return of market
he market is not perfect competition. The
etermined by the model. The New-Key
otaki (1987), has introduced produc
general equilibrium models with nomin
ed and sunk costs. Most of this literat

APPROVED

Drawing tools available on comment ribbon

How to use it:

- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select *Open Pop-up Note*.
- Type any text in the red box that appears.

7. **Drawing Markups** Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:

AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query References	Query	Remarks
Q1	Please confirm that given names (blue) and surnames/family names (vermilion) have been identified and spelled correctly.	Confirmed
Q2	Editor comment DG1: Edited to meet format requirements – Is this OK?	OK
Q3	Correct supplier information?	No, correct supplier information is: Ethicon, Guaynabo, Puerto Rico

ORIGINAL ARTICLE

Lip-to-nose flap for nasal plane reconstruction in dogs: A cadaveric and in vivo feasibility study

Lavinia E. Chiti DVM | Vincenzo Montinaro DVM, MSc | Marta L. P. Lisi DVM, PhD |
Andrea G. Asta DVM | Silvia Marches DVM, PhD, MSc | Paolo Sommaruga DVM |
Federico Massari DVM, Diplomate, ECVS

Clinica Veterinaria Nervianese, Nerviano, Milan, Italy

Correspondence

Lavinia Chiti, Clinica Veterinaria Nervianese, Nerviano, Milan, Italy
Email: lavinia.chiti@gmail.com

Abstract

Objective: To describe a local interpolation flap based on the upper lip for the reconstruction of the dorsal aspect of the nasal plane in dogs.

Study design: Ex vivo and in vivo case report.

Animals: A 2-year-old medium-sized spayed female mixed-breed dog and a canine cadaver of a medium-sized mixed-breed dog.

Methods: A dorsal defect of the nasal plane of uncertain origin in a dog was reconstructed with a unilateral interpolation flap obtained from the caudal upper lip skin and ~~attached to the labial~~ mucosa (lip-to-nose flap). The procedure was first performed in a cadaveric model to assess its feasibility.

Results: The lip-to-nose flap allowed the reconstruction of a dorsal nasal plane defect with a satisfactory cosmetic outcome. The incorporation of the labial mucosa guaranteed a complete and uncomplicated wound healing of the flap; a small area of necrosis occurred at the donor site, which resolved in a few days.

Conclusion: The use of a lip-to-nose flap was feasible in these 2 medium-sized dogs and led to satisfactory cosmetic outcome ~~in 1 dog~~.

Clinical significance: A lip-to-nose flap may be considered to reconstruct dorsal nasal plane defects in dogs and allows for a cosmetic outcome.

1 | INTRODUCTION

Defects of the external nose in dogs are often caused by traumatic injuries or surgical resections of neoplasms such as squamous cell carcinoma.^{1,2} Reconstruction of the nasal plane is particularly challenging in dogs because of the lack of free local tissue to reconstruct the original appearance of the nose.³ The canine nose is a highly specialized structure and contributes to the appearance of the dog.⁴ Thus, cosmetic and functional outcome are of great concern and may limit the extent of the surgical excision in dogs with nasal neoplasia. For the same reasons, surgeons are cautious when performing nasal plane reconstruction in patients with traumatic injuries of the external nose.^{3,4} In the veterinary literature, several techniques based on local flaps have been proposed for primary closure

of nasal defects. These techniques range from advancement flaps such as the dorsal muzzle and labial advancement to composed flaps such as the bilateral alar cartilage and the musculofascial island labial flap.⁴ Gallegos and colleagues⁵ reported a labial mucosal technique for reconstruction of defects involving the nasal plane and the rostral labial tissue. The labial mucosal inversion technique improves cosmetic outcome and provides a labial mucosal antechamber protecting the nasal mucosa from direct exposure. This technique involves excision of a large portion of the upper lip adjacent to the nasal plane; it is thus not ideal for the reconstruction of dorsal defects without involvement of the rostral aspect of the nasal plane and upper lip. Recently, a modified nasal rotational flap was described for the reconstruction of a rostral nasal defect involving the lateral nasal plane in a dog.³

Despite the good cosmetic and functional outcome obtained, this technique is less well suited to dorsal nasal planum defects because healthy tissue at the dorsocaudal part of the nasal plane is required to harvest the rotational flap correctly.

The lip-to-lid flap is a local interpolation flap that is based on the caudal upper lip skin and the attached labial mucosa that has been successfully applied to replace the lower eyelid in dogs and cats.⁶⁻⁸ The objective of this study is to report a modification of the lip-to-lid flap for the reconstruction of dorsal nasal defects in dogs and to assess its feasibility in a cadaveric and in vivo model. The technique described consists of a lip-to-nose interpolation flap based on the subdermal plexus, which allows direct mucosal apposition in order to obtain a good functional and cosmetic outcome.

2 | CLINICAL REPORT

A 2-year-old mixed-breed spayed female dog weighing 21 kg was referred to our practice in February 2017 for a defect of the dorsal aspect of the nasal plane. The dog had been found by a rescue association 10 days previously with no known clinical history. At clinical examination, the dog presented with a large defect of the nasal plane involving the rostral third of the nasal bones, the philtrum, and the dorsal lateral cartilages bilaterally. The lateral part of the alar cartilages and the ventral lateral cartilages were preserved (Figure 1). The rostral portion of the dorsal meatus was exposed, causing a chronic inflammation of the nasal mucosa and associated nasal discharge. No other abnormalities were identified during physical examination. Hematological analysis revealed regenerative anemia, neutrophilic lymphocytosis, and hypoalbuminemia. Serum proteins electrophoresis pattern was consistent with a protozoarian infection and *Hepatozoon canis* was detected in the blood smear. Imidocarb dipropionate was administered at a dose of 6 mg/kg subcutaneously twice 14 days apart. A computed tomography (CT) scan of the nose was performed, and no other abnormalities were detected. Although no biopsy of the nasal tissue was performed in this case, we hypothesized that the nasal plane defect in this dog was of traumatic origin.

FIGURE 1 View of the defect. Note the exposure of the rostral portion of the dorsal meatus and the subsequent inflammatory hyperemia of the nasal mucosa; rostral (A) and dorsal (B) views

Biopsies for histopathological examination and bacterial and fungal culture should have been performed to rule out other possible causes of nasal defects. However, even though such procedures were strongly recommended, it was eventually refused by the rescue association because of financial issues. Because the lesion was causing discomfort to the dog and gross appearance and CT findings were highly suggestive of a traumatic origin, it was decided to perform a reconstructive surgery despite the absence of a histopathological diagnosis.

A decision was made to close the nasal defect with a lip-to-nose flap. The procedure was first performed on a canine cadaver of similar size to assess its feasibility (Figure 2). The cadaver was placed in sternal recumbency and the nose and lip were clipped. The flap was obtained from the upper lip as described by Pavletic⁶ for the lip-to-lid technique. Two full-thickness incisions of the left upper lip were made starting from the medial canthus of the ipsilateral eye with a 45° angle from a line drawn between the medial canthus of each eye (Figure 2A-C). The 2 incisions were convergent and outlined so that the distance between the rostral and caudal edge of the flap corresponded to the minor diameter of the nasal plane lesion, with the caudal edge of the flap 3 cm rostral from the commissure (Figure 2A,B). The buccal mucosa was divided from the gingival mucosa, and the whole buccal mucosa was preserved to reconstruct the dorsal nasal mucosa. The skin was then undermined dorsally under the platysma muscle, and the skin flap with the attached lip margin was transposed to cover the nasal plane defect (Figure 2D-F). A bridging incision was created just caudal to the nasal plane lesion (Figure 2D). The mucosal edge of the flap was sutured to the remaining nasal mucosa with a simple interrupted pattern with monofilament absorbable sutures (poliglecaprone 25 USP 3-0; Ethicon, Somerville, New Jersey; Figure 2G). The skin edge of the flap was then sutured to the nasal plane with a 2-layer simple continuous suture pattern, and the defect at the donor site was routinely closed in 3 layers with monofilament absorbable sutures (Figure 2G,H).

The cosmetic outcome was considered satisfactory, and the same procedure was thus performed in-vivo. We assumed that the lip-to-nose flap could protect the nasal mucosa from the environment, thus limiting inflammation and resolving the associated nasal discharge.

The dog was premedicated with butorphanol (0.2 mg/kg IM) and medetomidine (10 mcg/kg IM); general anesthesia was induced with propofol (6 mg/kg IV) and maintained with isoflurane in oxygen after intubation. Saline solution 0.9% was administered IV at the rate of 5 mL/kg/h during surgery. The dog was positioned in sternal recumbency. The nose and upper lip were clipped and aseptically prepared for surgery (Figure 3A). Wound debridement was performed: the epithelizing edges of the dorsal aspect of the defect and the medial margins of the alar cartilages were sharply excised to obtain fresh wound edges. A composite flap was

FIGURE 2 Cadaveric model of the lip-to-nose flap for the reconstruction of dorsal nasal plane defects. **A**, The dorsal aspect of the nasal plane has been excised, and the rostral part of the resulting defect has been measured. **B**, The same distance has been measured on the lower lip to obtain the correct width of the mucosal edge of the flap. **C**, Two full-thickness incisions of the upper lip have been made to create the flap. The 2 incisions extend from the medial canthus of the eye to the lower lip margin and form a 45° angle from a line drawn between the medial and lateral canthi. The whole buccal mucosa has been preserved to reconstruct the dorsal nasal mucosa (green arrow). **D**, A bridging incision has been made just proximal to the nasal plane defect (red arrow). **E**, The flap has been transposed to cover the nasal plane defect. **F**, The skin edge of the flap has been sutured in a simple continuous pattern to the recipient site, and the lower lip has been reconstructed with an advancement flap. **G**, Completed reconstruction, right lateral view. **H**, Completed reconstruction, frontal view

then obtained by following the technique described in the cadaveric model, and the nasal plane defect was reconstructed with the lip-to-nose flap (Figure 3B-D). Great care was taken to preserve the subdermal plexus and to perfectly appose the skin edges of the flap. A vacuum drain (Emodren; Medicalplastic, Milan, Italy) was placed to prevent seroma formation (Figure 3D). The defect at the donor site was closed with a labial advancement flap. Pain was managed by fentanyl (5 mcg/kg/h continuous rate infusion IV) and local nerve block (bupivacaine 0.5% 1 mL)

FIGURE 3 Lip-to-nose flap for the reconstruction of a dorsal nasal plane defect in a 2-year-old mixed-breed, spayed, female dog. **A**, The defect involved the rostral third of the nasal bones, the philtrum, and the dorsal lateral cartilages bilaterally; part of the alar cartilages and the ventral lateral cartilages were preserved. **B**, The flap has been obtained from the upper lip skin and the attached oral mucosa (blue arrow). **C**, The flap has been transposed to cover the dorsal defect. **D**, Oblique view of the muzzle after reconstruction; an active-suction drain has been placed

intraoperatively and by methadone 0.1 mg/kg IV twice daily postoperatively until discharge. Cefazolin (25 mg/kg IV twice daily) and meloxicam (0.2-0.1 mg/kg IV once daily) were administered for 7 and 5 days, respectively, after surgery. The drain was removed 2 days later because the production of serosanguinolent fluid had steadily decreased; final production was 0.04 mL/kg/24 h.⁷ The dog was discharged 7 days postoperatively, and drug therapy for treatment of *H. canis* infection was discontinued 14 days later.

Although the lip-to-nose flap did not allow for the reconstruction of the philtrum, the cosmetic outcome was improved, and the motility of the external nose was preserved. At discharge, a small area of necrosis was noted at the donor site, probably due to the use of the monopolar electrosurgery; the lip-to-nose flap appeared to be healed completely (Figure 4A,B). Necrosis at the donor site was treated with topic silver sulfadiazine and healed completely in 7 days. The dog had minimal nasal discharge, which resolved spontaneously 48 hours postoperatively. A small dog ear resulted from the transposition of the flap, which underwent a partial regression during the initial phase of the healing process. Fifteen days after surgery, there were no signs of flap failure, and both the nasal plane and the donor site were perfectly healed without evidence of nasal orifice stenosis. At 45 days, the dog had recovered uneventfully, and the cosmetic and functional outcomes were satisfactory (Figure 5A-C).

FIGURE 4 Frontal (A) and left lateral (B) views at discharge. Note the small area of necrosis at the donor site (blue arrow)

3 | DISCUSSION

In this case, a dorsal defect of the nasal plane in a dog was successfully reconstructed with a lip-to-nose interpolation flap based on the subdermal plexus. Primary closure techniques for a cosmetic reconstruction of the canine nasal plane are a great concern in veterinary reconstructive surgery, and several techniques that are based on local flaps have been previously described.^{3,5} However, none of these techniques are ideal for the closure of dorsal defects. To the best of the authors' knowledge, no published report has yet described a surgical technique that is based on a subdermal plexus flap specifically designed to cover dorsal nasal plane defects. The lip-to-lid flap has been largely employed for the reconstruction of the lower eyelid in dogs and cats and has had a high rate of success.^{6,8,9} The lip-to-nose flap described here similarly relies on the subdermal plexus and permits first intention healing with minimal tension. The upper lip represents a highly vascularized structure, with contribution from the infraorbital and lateral nasal artery; this may result in an improved success rate of the flap.¹⁰

The incorporation of the buccal mucosa permits a mucosal apposition that promotes primary wound healing with minimal scar tissue formation. This may prevent stenosis of the nostril as well as changes in the appearance of the external nose. In the case reported here, the lip-to-nose flap healed completely without complications, and no stenosis of the nostril was observed; moreover, a total remission of the nasal discharge was achieved. In fact, nasal discharge in this case was probably due to chronic inflammation of the nasal mucosa caused by exposure to the environment. The lip-to-nose flap permitted us to create an interface between the

nasal mucosa and the environment, thus limiting the inflammation and solving nasal discharge.

The pigmented nonhaired lip margin resembles the original appearance of the nose, and its incorporation allowed for a better cosmetic outcome. The dog ear resulting from the flap transposition partially regressed in the first days after surgery and did not require a surgical correction because it did not compromise the overall cosmetic result.

The creation of the full-thickness flap from the upper lip is relatively simple to achieve, and the lip-to-nose technique can be performed in a limited amount of time. The lip-to-nose flap is designed to cover dorsal defects of the nasal plane. In the case that we report here, the defect also involved the philtrum, which was not reconstructed. Nonetheless, the cosmetic appearance was not compromised, and a functional reconstruction was achieved because the diameter of the nasal passage was preserved and no persistent nasal discharge occurred.

The defect that we treated was probably of traumatic origin. It is the authors' opinion that the lip-to-nose flap might be an effective option for the cosmetic closure of defects caused by surgical resection of neoplasm of the nasal plane. Additional studies are warranted to confirm the feasibility of the lip-to-nose flap for nasal plane reconstruction after tumor excision. Moreover, because the lip offers a discrete amount of free tissue, the flap obtained from the upper lip and the attached mucosa can be widened to cover larger defects of the dorsal nasal plane or lesions which extend laterally.

The results of this study provide evidence that the lip-to-nose flap is feasible for a single-stage reconstruction of nasal plane defects in dogs and allows for a good cosmetic and functional outcome. However, additional studies with greater numbers of cases are required to investigate additional applications of the lip-to-nose flap and to assess its success rate.

CONFLICT OF INTEREST

The authors declare no conflicts of interest related to this report.

REFERENCES

1. Lascelles BDX, Parry AT, Stidworthy MF, Dobson JM, White RA. Squamous cell carcinoma of the nasal planum in 17 dogs. *Vet Rec*. 2000; 147:473-476.
2. Thomson M. Squamous cell carcinoma of the nasal planum in cats and dogs. *Clin Tech Small Anim Pract*. 2007;22:42-45.
3. Haar G, Buiks SC, Kirpensteijn J. Cosmetic reconstruction of a nasal plane and rostral nasal skin defect using a modified nasal rotation flap in a dog. *Vet Surg*. 2013;42:176-179.
4. Pavletic MM. Nasal reconstruction technique. In: Pavletic MM, ed. *Atlas of Small Animal Wound Management and Reconstructive Surgery*. 3rd ed. Ames, IA: Wiley-Blackwell; 2010:574-600.
5. Gallegos J, Schmiedt CW, McNulty JF. Cosmetic rostral nasal reconstruction after nasal planum and premaxilla resection: technique and results in two dogs. *Vet Surg*. 2007;36:669-674.
6. Pavletic MM. Major eyelid reconstruction technique. In: Pavletic MM, ed. *Atlas of Small Animal Wound Management and Reconstructive Surgery*. 3rd ed. Ames, IA: Wiley-Blackwell; 2010:562-570.

FIGURE 5 Appearance of the dog 45 days after surgery; rostral (A), lateral (B), and dorsal (C) views

7. Bristow PC, Halfacree ZJ, Baines SJ. A retrospective study of the use of active suction drains in dogs. *J Small Anim Pract.* 2015;56:325-330.
8. Hunt GB. Use of the lip-to-lid flap for replacement of the lower eyelid in five cats. *Vet Surg.* 2006;35:284-286.
9. Pavletic MM, Nafe LA, Confer AW. Mucocutaneous subdermal plexus flap from the lip for lower eyelid restoration in the dog. *J Am Vet Med Assoc.* 1982;15:921-926.
10. Bezuidenbout A. The heart and arteries. In: Evans HE, de Lahunta A, eds. *Miller's Anatomy of the Dog.* 4th ed. Philadelphia, PA: Elsevier Saunders; 2013:455-456.

How to cite this article: Chiti LE, Montinaro V, Lisi MLP, et al. Lip-to-nose flap for nasal plane reconstruction in dogs: A cadaveric and in vivo feasibility study. *Veterinary Surgery.* 2018;1-5. <https://doi.org/10.1111/vsu.12965>

Uncorrected Proofs