

ORIGINAL ARTICLE

Impact of Fractional Flow Reserve Derived From Coronary Computed Tomography Angiography on Heart Team Treatment Decision-Making in Patients With Multivessel Coronary Artery Disease

Insights From the SYNTAX III Revolution Trial

AQ1

Daniele Andreini, MD; et al

BACKGROUND: Fractional flow reserve (FFR) is a reliable tool for the functional assessment of coronary stenoses. FFR computed tomography (CT) derived (FFR_{CT}) has shown to be accurate, but its clinical usefulness in patients with complex coronary artery disease remains to be investigated. The present study sought to determine the impact of FFR_{CT} on heart team's treatment decision-making and selection of vessels for revascularization in patients with 3-vessel coronary artery disease.

METHODS: The trial was an international, multicenter study randomizing 2 heart teams to make a treatment decision between percutaneous coronary interventions and coronary artery bypass grafting using either coronary computed tomography angiography or conventional angiography. The heart teams received the FFR_{CT} and had to make a treatment decision and planning integrating the functional component of the stenoses. Each heart team calculated the anatomic SYNTAX score, the noninvasive functional SYNTAX score and subsequently integrated the clinical information to compute the SYNTAX score III providing a treatment recommendation, that is, coronary artery bypass grafting, percutaneous coronary intervention, or equipoise coronary artery bypass grafting-percutaneous coronary intervention. The primary objective was to determine the proportion of patients in whom FFR_{CT} changed the treatment decision and planning.

RESULTS: Overall, 223 patients were included. Coronary computed tomography angiography assessment was feasible in 99% of the patients and FFR_{CT} analysis in 88%. FFR_{CT} was available for 1030 lesions (mean FFR_{CT} value 0.64±13). A treatment recommendation of coronary artery bypass grafting was made in 24% of the patients with coronary computed tomography angiography with FFR_{CT}. The addition of FFR_{CT} changed the treatment decision in 7% of the patients and modified selection of vessels for revascularization in 12%. With conventional angiography as reference, FFR_{CT} assessment resulted in reclassification of 14% of patients from intermediate and high to low SYNTAX score tertile.

CONCLUSIONS: In patients with 3-vessel coronary artery disease, a noninvasive physiology assessment using FFR_{CT} changed heart team's treatment decision-making and procedural planning in one-fifth of the patients.



VISUAL OVERVIEW: A visual overview is available for this article.

CLINICAL TRIAL REGISTRATION: URL: <https://www.clinicaltrials.gov>. Unique identifier: NCT02813473.

AQ3

Key Words: angiography ■ coronary artery disease ■ coronary computed tomography angiography ■ decision-making ■ percutaneous coronary intervention

AQ4

See Editorial by Author

Correspondence to: Antonio L. Bartorelli, MD, Centro Cardiologico Monzino, IRCCS, Milan, Italy. Email antonio.bartorelli@ccfm.it

The full author list is available on page XXX.

The Data Supplement is available at <https://www.ahajournals.org/doi/suppl/10.1161/CIRCINTERVENTIONS.118.007607>.

For Sources of Funding and Disclosures, see page XXX.

© 2019 American Heart Association, Inc.

Circulation: Cardiovascular Interventions is available at www.ahajournals.org/journal/circinterventions

WHAT IS KNOWN

- The SYNTAX III Revolution trial showed that in patients with 3-vessel coronary artery disease treatment decision-making based on coronary computed tomography angiography is in high agreement with the decision derived from conventional angiography.
- Physiology-guided (invasive fractional flow reserve [FFR]) percutaneous coronary revascularization has been associated with lower rate of MACE compared with invasive angiographic guidance and is recommended by the latest guidelines.

WHAT THE STUDY ADDS

- The heart team changed the treatment recommendation in 7% of the cases and modified the selection of vessels to be revascularized in 12% when functional evaluation with FFR_{CT} was added to an anatomic assessment with coronary computed tomography angiography alone.
- In patients assessed by coronary computed tomography angiography, FFR_{CT} reduced the proportion of patients with hemodynamically significant 3-vessel coronary artery disease from 92.3% to 78.8% and reclassified to a lower SYNTAX Score 15.5% of patients.

Nonstandard Abbreviations and Acronyms

CABG	coronary artery bypass grafting
CAD	coronary artery disease
CTA	coronary computed tomography angiography
FAME	Fractional Flow Reserve Versus Angiography for Multivessel Evaluation
FFR	fractional flow reserve
FFR_{CT}	fractional flow reserve derived from computed tomography
PCI	percutaneous coronary interventions

Revascularization by either percutaneous coronary intervention (PCI) or coronary artery bypass graft surgery (CABG) is indicated in flow-limiting coronary stenoses to reduce myocardial ischemia and its adverse clinical manifestations.¹ In patients with multivessel coronary artery disease (CAD), physiology-guided coronary revascularization has shown to improve clinical outcomes compared with an angiographic assessment alone.² Determination of pressure-wire indexes such as fractional flow reserve (FFR) or instantaneous wave-free ratio has demonstrated that almost half of the lesions with a diameter stenosis >50% are not hemodynamically significant.³

FFR derived from computed tomography angiography (FFR_{CT}) is a noninvasive method able to identify

lesion-specific ischemia. In patients with multivessel disease, FFR_{CT} has shown to have good diagnostic performance with invasive pressure-wire assessment as reference.⁴ Moreover, the extent, severity, and functional component of CAD can be objectively quantified using the functional SYNTAX score.⁴ The functional SYNTAX score has higher discrimination for clinical events compared with the anatomic SYNTAX score, while reducing inter-observer variability. The calculation of the SYNTAX score III, combining in a noninvasive setting anatomy, physiology and patient's clinical information provides the heart team with individualized risk stratification and treatment recommendation based on the predicted 4-year mortality in patients undergoing PCI or CABG.⁵

The SYNTAX III Revolution trial showed that in patients with left main or 3-vessel CAD treatment decision-making based on coronary computed tomography angiography (CTA) is in high agreement with the decision derived from conventional angiography.⁶ However, the influence of FFR_{CT} on treatment decision-making and selection of vessels for revascularization remains to be investigated. Thus, the present study sought to determine the impact of FFR_{CT} on heart team's treatment decision and procedural planning in patients with left main or 3-vessel CAD.

METHODS

The data that support the findings of this study are available from the corresponding author on reasonable request.

Study Design

The present study reports the secondary end point of the SYNTAX III REVOLUTION trial. The design of the SYNTAX III REVOLUTION Trial, a randomized study investigating the use of CT scan and angiography of the heart to help the doctors decide which method is the best to improve blood supply to the heart in patients with complex coronary artery disease, has been reported previously.⁵ The trial was an international, multicenter study randomizing 2 heart teams to make a treatment decision between PCI and CABG using either coronary CTA or conventional angiography, while blinded to the other imaging modality. The results of the primary end point based on coronary anatomy alone have been recently published.⁶ The present analysis focuses on a second level of the 2 heart teams decision-making after the incorporation of the physiology component (FFR_{CT}) in the coronary CTA diagnostic strategy arm, which represents the secondary end point of the SYNTAX III REVOLUTION trial.⁵ The study was approved by an institutional review committee, and the subjects gave informed consent. The European Cardiovascular Research Institute (Rotterdam, the Netherlands) with unrestricted grants from GE Healthcare (Chicago, IL) and Heart Flow Inc (Redwood City, CA) sponsored the study.

Enrolment and Randomization

Patients with left main or 3-vessel CAD diagnosed with either coronary CTA or conventional angiography and candidates

for either PCI or CABG were assessed for eligibility. Patients were consented to undergo coronary CTA using a whole-heart coverage, high-definition CT scanner (Revolution CT; GE Healthcare, Chicago, IL), and to participate in a randomized trial of decision-making between PCI and CABG performed by the local heart team and relying on alternative imaging techniques. Two heart teams composed by an interventional cardiologist, a cardiac surgeon and a radiologist specialized in cardiac imaging, were randomized to either assess the coronary anatomy with coronary CTA or conventional angiography in addition to the patient's clinical information. The heart team allocated to coronary CTA had to make a second treatment decision and procedural planning taking into consideration the functional component of the coronary stenoses provided by FFR_{CT} . Similarly, the heart team randomized to conventional angiography received coronary CTA with FFR_{CT} and had to make a second treatment decision based on the 3 diagnostic methods. Each heart team calculated the anatomic SYNTAX score based only on their allocated imaging modality, the noninvasive functional SYNTAX score, and subsequently integrated the clinical information to compute the SYNTAX score III providing a treatment recommendation, that is, CABG, PCI, or equipoise between CABG and PCI. Any anatomic SYNTAX score was eligible for screening, and patients with anatomic SYNTAX score >33 were not excluded. Patients with prior revascularization were excluded. Complete details of the inclusion and exclusion criteria have been previously described.^{5,6}

Image Acquisition and Analysis

Coronary CTA was performed with the GE Revolution CT scanner.⁷ A proprietary post-processing algorithm (Snap Shot Freeze) allowed for the correction of motion artifacts.⁸ The imaging acquisition guidelines are detailed in the Table I in the [Data Supplement](#). Image quality was assessed using the 5-point Likert scale at the patient level. The 2 local heart teams signed off their decision on the choice of revascularization mode based on the anatomic assessment alone. Subsequently, the FFR_{CT} was used to calculate the noninvasive functional SYNTAX score, computed by subtracting nonflow limiting stenoses ($\text{FFR}_{\text{CT}} > 0.80$) from the anatomic SYNTAX score. FFR_{CT} has a lower limit of detection of 0.50. Finally, the noninvasive functional SYNTAX score was used to calculate the SYNTAX Score III, which is conceptually a combination of coronary anatomy complexity with its functional reperfusion and patient's clinical characteristics and comorbidities. The anatomic and functional SYNTAX scores were also calculated by an independent core laboratory (Cardialysis BV, Rotterdam, the Netherlands) and were made available to each heart team for consultation. For the present article, the functional SYNTAX score was calculated from the heart team assessment, whereas lesion level data were analyzed by the core laboratory (Cardialysis BV, Rotterdam, the Netherlands).

Objectives

The primary objective of the present study was to determine the proportion of patients in whom FFR_{CT} changed the treatment decision and the selection of vessels for revascularization with respect to the management based on anatomic assessment with either coronary CTA or conventional angiography. The secondary objective was to assess the impact of FFR_{CT} on

patients risk reclassification compared to the anatomic assessment alone using the SYNTAX score tertiles (SYNTAX score: 0–22, 23–32, and >32).

Statistical Analysis

The heart team's treatment recommendation led to one of 3 decisions according to the SYNTAX Score III: (1) CABG only, patients should be treated by CABG due to a higher 4-year mortality with PCI; (2) PCI only, patients should be treated by PCI due to a higher 4-year mortality with CABG; and (3) equipoise between CABG and PCI, patients could be treated by either approach, considering that the 4-year mortality prediction is similar between them. The power calculation of the sample size of the SYNTAX III REVOLUTION Trial has been previously described.⁵ The risk reclassification is presented as the proportion of patients reclassified from the anatomic to the functional SYNTAX score tertiles (ie, low <22 , intermediate 23–32, and high >32). Comparison of the continuous anatomic SYNTAX scores was performed with the paired t test. Differences in categorical variables were assessed with the use of McNemar test, since they were from a matched population. Agreement between (1) SYNTAX II recommendation strategies derived from angiography only versus that derived from angiography with CTA and FFR_{CT} and between (2) tertiles of anatomic SYNTAX score derived from different imaging strategies were assessed with the concordance coefficient of κ . A 2-sided P value of 0.05 or less was considered to indicate statistical significance. All statistical analyses were performed with the use of SAS software, version 9.4 (SAS Institute).

RESULTS

From June 29, 2016 to February 8, 2018, 223 patients with left main or 3-vessel CAD were enrolled in 6 centers from 5 European countries. Baseline clinical characteristics and CT acquisition information are shown in Table II in the [Data Supplement](#). Coronary CTA assessment was feasible in 99% of the patients, and the FFR_{CT} analysis was available in 196 patients (88% of the entire study population; Figure 1). FFR_{CT} was available for 1030 lesions and was positive (≤ 0.80) in 89% of them (mean FFR_{CT} value 0.64 ± 13 ; Figure 2).

Impact of FFR_{CT} on the Treatment Decision Based on Coronary CTA

Using coronary CTA alone, 1108 stenoses (5.0 ± 1.7 per patient) were detected. The mean anatomic SYNTAX score derived from coronary CTA was 33.9 ± 13.0 ($n=233$ patients). The heart teams subtracted 205 lesions, assessed as nonfunctionally significant, for the functional SYNTAX score calculation. This led to a reduction of the noninvasive functional SYNTAX score ($n=196$) to 30.5 ± 13.0 (P value < 0.001 versus anatomic SYNTAX score). The inclusion of the physiology component led to a mean reduction of the SYNTAX score of 3.7 points. Table 1 shows the impact of the physiology information on the SYNTAX score components.




Figure 1. Revision SYNTAX III Study flow chart.

FFR_{CT} indicates fractional flow reserve derived from computed tomography angiography.

AQ15

Based on the functional SYNTAX score according to FFR_{CT}, SYNTAX score III recommended CABG only in 24% (47/194) of the patients and the heart team treatment decision was CABG in 69% (161/194) of the cases. FFR_{CT} changed the treatment decision between CABG and PCI in 7% (14/194) of the patients and modified the revascularization strategy in 12.1% of the vessels. Compared with anatomic coronary CTA assessment, addition of FFR_{CT} reduced the number of patients with significant 3-vessel CAD from 92.3% to 78.8%.

Impact of Coronary CTA With FFR_{CT} on Treatment Decision Based on Conventional Angiography

Conventional coronary angiography identified 1073 stenoses (4.8±1.7 per patient) with a mean SYNTAX score of 30.3±12.2. Further evaluation with coronary CTA and FFR_{CT} removed 129 nonhemodynamically significant

lesions from the functional SYNTAX score calculation; however, the mean SYNTAX score remained unchanged (31.2±13.4; Table 2). The SYNTAX score III recommended CABG only in 26% (50/196) of the patients, and CABG was selected as the treatment by the heart team in 69% (135/196) of the patients. The addition of FFR_{CT} changed the heart team's treatment selection in 6.6% of patients and modified treatment planning in 18.3%. The number of patients with significant 3-vessel CAD remained unchanged (86.1% versus 86.2%).

Risk Reclassification

Using FFR_{CT}, 15.5% (30/194) of the patients were reclassified to a lower risk SYNTAX score tertile based on coronary CTA alone (Table III in the [Data Supplement](#), Figure 3A). Similarly, the addition of noninvasive functional assessment to conventional angiography reclassified 14% (31/221) of the patients to a lower risk tertile

F3




Figure 2. Flow reserve derived from computed tomography angiography (FFR_{CT}) values distribution.

The diagram shows the number of lesions divided into incremental FFR_{CT} values.

*Because the lower limit of detection of FFR_{CT} is 0.5, values presented at this interval 0.50–0.525 also represent all values below 0.5.

(Table IV in the [Data Supplement](#), Figure 3B). The recommendation based on the SYNTAX II score changed in 6.7% of the patients (ie, in 3.1% of the patients, the recommendation changed from CABG only to equipoise by adding FFR_{CT} to conventional angiography, and in 3.6% the recommendation changed from equipoise to CABG only). The agreement was high with a κ of 0.82 (0.73–0.92; Table 3).

DISCUSSION

The main findings of the present study can be summarized as follows: (1) by including the noninvasive functional evaluation with FFR_{CT}, the heart team changed the treatment recommendation in 7% of the cases and modified the selection of vessels for revascularization in 12% as compared with a coronary CTA assessment alone. Moreover, inclusion of FFR_{CT} information on top of conventional angiography changed the treatment recommendation in 6.6% of the cases and modified the planning in 18.3%; (2) the noninvasive functional SYNTAX score reclassified 15.5% of the patients to a lower SYNTAX score tertile based on coronary CTA and 14% of the patients to a lower SYNTAX score tertile based on conventional angiography; and (3) in patients assessed by coronary CTA, FFR_{CT} reduced the proportion of patients with hemodynamically significant 3-vessel CAD from 92.3% to 78.8%.

It is noteworthy that in this study anatomically nonsignificant lesions were not scored in the calculation of the coronary CTA SYNTAX score, even if the FFR_{CT} in the distal vessel was positive. On the contrary, when FFR_{CT} was ≤ 0.8 in the proximal coronary segments, all distal lesions were considered significant. This approach might underestimate the clinical impact of FFR_{CT}, and technical improvements are ongoing to overcome the limitation.

Use of physiology-guided percutaneous revascularization has shown to reduce the rate of major adverse

cardiovascular events compared with angiographic guidance alone and has been recommended by the latest guidelines to identify hemodynamically relevant lesions.⁹ In the SYNTAX II study, which included patients with 3-vessel disease without left main involvement and moderate anatomic complexity (mean SYNTAX score 20.3 ± 6.4), routine physiological assessment (performed in 82.8% of patients) reduced the percentage of patients with 3-vessel disease to 37.2%.¹⁰ At variance with SYNTAX II, the SYNTAX III REVOLUTION trial included patients with more severe CAD, as underlined by a mean anatomic SYNTAX score of 30.0 ± 12.0 .⁶ Moreover, the proportion of lesions with a positive FFR was 89% in SYNTAX III and 74.6% in SYNTAX II. In SYNTAX II, inclusion of FFR_{CT} reclassified 30% of the patients to a lower SYNTAX score tertile, which is a higher proportion than that observed in the present study. The limited exclusion criteria of SYNTAX III, particularly the inclusion of patients with left main coronary artery stenosis, are likely responsible for this finding and underline the functional compromise of unselected patients with multivessel CAD. Of note, 11% of the coronary lesions in this complex population were found to have a value higher than 0.80 at FFR_{CT} and this finding influenced the treatment decision in 6% of the patients and modified treatment planning in 16%.

In a large, retrospective study, FFR-guided CABG is associated with a reduction in death and myocardial infarction compared with angiography-guided CABG at 6-year follow-up.¹¹ We may expect an increase in functional-guided CABG in the upcoming years. In particular, coronary CTA with FFR_{CT} may provide to interventional cardiologists and cardiac surgeons a combined anatomic and functional noninvasive assessment of multivessel disease for the type and modality of revascularization. Indeed, one of the potential advantages of FFR_{CT} is the possibility to interrogate the physiology of any segment in the epicardial coronary circulation.^{12–14} From the heart

Table 1. Anatomy and Functional SYNTAX Score Assessment in the Heart Team Randomized to Coronary CTA

Characteristics	Coronary CTA Only 223 Patients	Coronary CTA With FFR _{CT} 194 Patients	P Value
Number of lesions	1108	903	
Diseased vessels			
RCA	95.5% (212/222)	86.6% (168/194)	<0.001
LAD	99.5% (221/222)	96.9% (188/194)	0.063
LCX	95.0% (211/222)	90.7% (176/194)	0.039
Left main	36.0% (80/222)	30.4% (59/194)	0.006
Anatomy and functional SYNTAX score components			
Total occlusion	12.0% (133/1108)	12.8% (116/903)	0.5
Trifurcation	2.2% (24/1108)	1.6% (14/903)	1.0
Bifurcation	23.4% (259/1108)	22.5% (203/903)	1.0
Type of bifurcation			
Medina 1,0,0	1.1% (12/1108)	1.1% (10/903)	1.0
Medina 0,1,0	0.8% (9/1108)	0.9% (8/903)	1.0
Medina 1,1,0	6.9% (77/1108)	6.9% (62/903)	1.0
Medina 1,1,1	8.6% (95/1108)	8.4% (76/903)	1.0
Medina 0,0,1	1.8% (20/1108)	1.6% (14/903)	1.0
Medina 1,0,1	1.9% (21/1108)	1.8% (16/903)	1.0
Medina 0,1,1	2.3% (25/1108)	1.9% (17/903)	1.0
Aorto-ostial lesion	3.6% (40/1108)	3.0% (27/903)	0.13
Severe tortuosity	0.9% (10/1108)	1.1% (10/903)	1.0
Length >20 mm	30.4% (296/975)	30.0% (236/787)	0.25
Heavy calcification	28.9% (320/1108)	27.1% (245/903)	1.0
Thrombus	0.3% (3/1108)	0.3% (3/903)	1.0
Anatomy and functional SYNTAX score	33.9±13.0	30.5±13.0	
1-vessel disease	0.9% (2/222)	3.1% (6/194)	0.22
2-vessel disease	6.8% (15/222)	18.0% (35/194)	<0.001
3-vessel disease	91.9% (204/222)	78.4% (152/194)	<0.001
Left main only	0.5% (1/222)	0.5% (1/194)	1.0
3-vessel disease (or 3-VD equivalent)	92.3% (204/221)	78.8% (152/193)	<0.001

McNemar test used to compare matched categorical variables. CTA indicates computed tomography angiography; FFR_{CT}, fractional flow reserve derived from computed tomography angiography; LAD, left anterior artery; LCX, left circumflex; and RCA, right coronary artery.

team's perspective, a nonhemodynamically significant lesion in the left anterior descending artery may prompt a change in the treatment strategy, favoring a percutaneous interventional approach in a multivessel disease patient. Moreover, in patients undergoing CABG, the recognition of nonhemodynamically significant lesions may reduce unnecessary grafts, surgical time, and might result in higher graft patency. The ongoing FAME 3 Trial (Fractional Flow Reserve Versus Angiography for Multivessel Evaluation), a comparison of FFR-guided PCI and CABG surgery in patients with multivessel CAD, will provide further clinical evidence of the potential benefit of FFR-guided surgical revascularization.¹⁵

Table 2. Anatomy and Functional SYNTAX Score Assessment in The Heart Team Randomized to Conventional Angiography

Characteristics	Conventional Angiography First Strategy		P Value
	Conventional Angiography Only 223 Patients	Conventional Angiography and Coronary CTA With FFR _{CT} 196 Patients	
Number of lesions	1073	944	
Diseased vessels			
RCA	92.4% (206/223)	93.9% (184/196)	1.0
LAD	96.4% (215/223)	97.4% (191/196)	0.63
LCX	95.1% (212/223)	92.3% (181/196)	0.15
Left main diseased	28.7% (64/223)	28.1% (55/196)	1.0
Anatomy and functional SYNTAX score components:			
Total occlusion	14.0% (150/1073)	13.6% (128/944)	0.42
Trifurcation	2.4% (26/1073)	2.2% (21/944)	1.0
Bifurcation	22.0% (236/1073)	21.3% (201/944)	0.88
Type of bifurcation			
Medina 1,0,0	1.7% (18/1073)	1.2% (11/944)	0.07
Medina 0,1,0	2.6% (28/1073)	1.8% (17/944)	0.33
Medina 1,1,0	4.4% (47/1073)	5.0% (47/944)	0.28
Medina 1,1,1	6.8% (73/1073)	7.2% (68/944)	0.66
Medina 0,0,1	2.4% (26/1073)	1.5% (14/944)	0.21
Medina 1,0,1	2.2% (24/1073)	2.8% (26/944)	0.23
Medina 0,1,1	1.9% (20/1073)	1.9% (18/944)	0.84
Aorto-ostial lesion	3.3% (35/1073)	3.7% (35/944)	0.86
Severe tortuosity	2.2% (24/1073)	0.8% (8/944)	0.002
Length >20 mm	29.1% (269/923)	30.1% (246/816)	0.15
Heavy calcification	13.0% (140/1073)	20.9% (197/944)	<0.001
Thrombus	0.3% (3/1073)	0.6% (6/944)	0.125
Anatomy and functional SYNTAX score	30.3±12.2	31.2±13.4	
1-vessel disease	1.3% (3/223)	1.5% (3/196)	1.0
2-vessel disease	12.1% (27/223)	11.7% (23/196)	1.0
3-vessel disease	86.1% (192/223)	86.2% (169/196)	0.82
Left main only	0.4% (1/223)	0.5% (1/196)	1.0
3-vessel disease (or 3-VD equivalent)	86.5% (192/222)	86.7% (169/195)	0.82

McNemar test used to compare matched categorical variables. CTA indicates computed tomography angiography; FFR_{CT}, fractional flow reserve derived from computed tomography angiography; LAD, left anterior artery; LCX, left circumflex; and RCA, right coronary artery.

Practical implementation of pressure-wire assessment of coronary lesions is still low worldwide. Furthermore, multivessel FFR or instantaneous wave-free ratio evaluation is seldom performed before a revascularization procedure, particularly before CABG. FFR_{CT} has the potential to refine the heart team's treatment decision and planning by incorporating the functional component of coronary epicardial lesions. The results of the present study support the feasibility and utility of preprocedural physiology assessment to better select and guide the




Figure 3. Flow reserve derived from computed tomography angiography (FFR_{CT}) and risk reclassification.

AQ7 The figure shows the impact of the noninvasive functional SYNTAX score on risk reclassification of patients assessed by coronary computed tomography angiography (CTA; **A**) and **B**. The red, green, and blue squares show the proportion of patients in whom the 2 imaging techniques are concordant in SYNTAX risk classification, respectively.

revascularization strategy in patients with multivessel CAD. Moreover, the high accuracy of FFR_{CT} in patients with multivessel disease and the safety regarding radiation dose achievable with the latest generation scanners¹⁶ make the noninvasive physiology assessment a very promising tool for the evaluation of patients with multivessel disease before a revascularization procedure. Progress in the field of coronary CTA, FFR_{CT} , and the application of machine learning on fluid dynamic analysis have proven to be clinically relevant.¹⁷ Of note, the acceptance rate for FFR_{CT} analysis in the present study was very high (88%), particularly if compared with that reported in other prospective multicenter trials. For example, it was 33% in the PRO_{CTA} trial¹⁸ and 69% in the SYNTAX II.⁴ However, the high rate of coronary CTA suitability for FFR_{CT} appears not surprising if we consider the scanner used in the SYNTAX III trial that combines fast gantry rotation time and intracycle motion-correction

algorithm aimed at reducing the impact of motion artifacts on image quality.¹⁹

Future Perspectives

The present study opens new perspectives on the use of coronary CTA as a tool to provide the interventionalist and cardiac surgeon with an anatomy and functional noninvasive road-map for myocardial revascularization strategy. Moreover, full automation of the SYNTAX III score has the potential to further enhance the decision-making process in patients with multivessel disease. The interactive planner, which is a new application of FFR_{CT} , could improve treatment selection while tailoring procedural planning based on assessing functional outcomes after virtual treatment.²⁰ In this novel and growing clinical field, stress myocardial CT perfusion has been introduced as a new tool for evaluating the functional relevance of coronary stenoses.^{21–25} However, unlike FFR_{CT} , CT perfusion requires an additional scan, use of a stressor agent, and is associated with higher radiation exposure.²⁶ Moreover, use of an adenosine-stress protocol may raise some safety concerns particularly in patients with left main or 3-vessel CAD.

Limitations

Some limitations of this study should be acknowledged. First, due to the study design, no data on the diagnostic accuracy of FFR_{CT} versus invasive FFR were assessed. However, physiological evaluation with FFR_{CT} has been shown to be accurate in multivessel CAD patients.⁴ Second, SYNTAX III Revolution was a decision-making trial;

Table 3. Level of Agreement Between Conventional Angiography Only vs Coronary CTA and Conventional Angiography With Functional Assessment by FFR_{CT}

Recommendation Based on Angiography Only	Recommendation Based on Angiography and Coronary CTA With FFR_{CT}		
	CABG Only		PCI Only/Equipoise
CABG only	21.9% (43/196)		3.1% (6/196)
PCI only/equipoise	3.6% (7/196)		71.4% (140/196)
	Number	%, 95% CI	κ , 95% CI
Concordance	183	93.4 (88.9–96.4)	0.82 (0.73–0.92)

Recommendation are based on SYNTAX II Score. CABG indicates coronary artery bypass grafting; CTA, computed tomography angiography; FFR_{CT} , fractional flow reserve derived from computed tomography angiography; and PCI, percutaneous coronary intervention.

thus, the clinical application of FFR_{CT} derived SYNTAX III score requires further investigation in a clinical outcome trial. Third, we recognize that this study was not designed to indicate to what degree FFR_{CT} may influence patients management.

Conclusions

In patients with left main or 3-vessel CAD, a noninvasive physiology assessment using FFR_{CT} changed heart team's treatment decision-making and selection of vessels for revascularization in one-fifth of the patients. This may improve the appropriateness and clinical outcome of myocardial revascularization treatment.

ARTICLE INFORMATION

Received January 18, 2019; accepted September 11, 2019.

Authors

Daniele Andreini, MD, PhD; Rodrigo Modolo, MD; Yuki Katagiri, MD; Saima Mushtaq, MD; Jeroen Sonck, MD; Carlos Collet, MD; Stefano et al (Maurizio Roberto, MD; Kaoru Tanaka, MD; Yosuke Miyazaki, MD; Jens Czaplá, MD; Danny Schoors, MD; Andre Plass, MD; Francesco Maisano, MD, PhD; Philipp Kaufmann, MD, PhD; Xavier Orry, MD; Pierre-Adrien Metzendorf, MD; Thierry Folliguet, MD; Gloria Färber, MD; Ioannis Diamantis, MD; Marc Schönweiß, MD; Giorgia Bonalumi, MD; Marco Guglielmo, MD; Cristina Ferrari, MD; Paolo Olivares, MD; Laura Cavallotti, MD; Ingrid Leal, MSc; Wietze Lindeboom, MSc; Yoshinobu Onuma, MD, PhD; Patrick W. Serruys, MD, PhD; Antonio L. Bartorelli, MD; for the SYNTAX III REVOLUTION investigators

Affiliations

Centro Cardiologico Monzino, IRCCS, Milan, Italy (D.A., S.M., S.d.M., M.R., G.B., M.G., C.F., P.O., L.C., A.L.B.). Department of Clinical Sciences and Community Health, Cardiovascular Section (D.A.) and Department of Biomedical and Clinical Sciences "Luigi Sacco" (A.L.B.), University of Milan, Italy. Department of Cardiology, Amsterdam University Medical Center, the Netherlands (R.M., Y.K.). Department of Internal Medicine, Cardiology Division, Hospital de Clinicas, University of Campinas, SP, Brazil (R.M.). Cardiovascular Center Aalst, OLV Hospital, Belgium (J.S., C.C.). Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Belgium (K.T., J.C., D.S.). Thoraxcenter, Erasmus MC, the Netherlands (Y.M., Y.O.). University of Zurich, Switzerland (A.P., F.M., P.K.). CHRU Nancy and Université de Lorraine, Nancy, France (X.O., P.-A.M., T.F.). Jena University Hospital, Friedrich Schiller University of Jena, Germany (G.F., I.D.). Heinrich Braun Klinikum, Zwickau, Germany (M.S.). Cardialysis BV, Rotterdam, the Netherlands (I.L., W.L.). Department of Cardiology, Royal Brompton and Harefield Hospitals, Imperial College London, United Kingdom (P.W.S.).

Sources of Funding

The European Cardiovascular Research Institute sponsored this study with unrestricted research grants from General Electric Health Care and Heart Flow Inc.

Disclosures

Dr Collet reports receiving grants from Heartflow Inc and Biosensors. Dr Serruys reports consultancy fees from Abbott, Biosensors, Medtronic, Micell, Qualimed, Sinomedical Sciences, St Jude Medical, Stentys, Svelte Medical Systems, Philips/Volcano, Xeltis, Stentit, and HeartFlow. The other authors report no conflicts.

REFERENCES

- Windecker S, Kolh P, Alfonso F, Collet JP, Cremer J, Falk V, Filippatos G, Hamm C, Head SJ, Juni P, et al. 2014 ESC/EACTS guidelines on myocardial revascularization. *EuroIntervention*. 2015; 10: 1024–1094. doi: 10.1093/eurheartj/ehu278
- Tonino PA, De Bruyne B, Pijls NH, Siebert U, Ikeno F, van't Veer M, Klauss V, Manoharan G, Engström T, Oldroyd KG, et al; FAME Study Investigators. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med*. 2009;360:213–224. doi: 10.1056/NEJMoa0807611
- Götberg M, Cook CM, Sen S, Nijjer S, Escaned J, Davies JE. The evolving future of instantaneous wave-free ratio and fractional flow reserve. *J Am Coll Cardiol*. 2017;70:1379–1402. doi: 10.1016/j.jacc.2017.07.770
- Collet C, Miyazaki Y, Ryan N, Asano T, Tenekecioglu E, Sonck J, Andreini D, Sabate M, Brugaletta S, Stables RH, et al. Fractional flow reserve derived from computed tomographic angiography in patients with multivessel CAD. *J Am Coll Cardiol*. 2018;71:2756–2769. doi: 10.1016/j.jacc.2018.02.053
- Cavalcante R, Onuma Y, Sotomi Y, Collet C, Thomsen B, Rogers C, Zeng Y, Tenekecioglu E, Asano T, Miyazaki Y, et al. Non-invasive heart team assessment of multivessel coronary disease with coronary computed tomography angiography based on SYNTAX score II treatment recommendations: design and rationale of the randomised SYNTAX III Revolution trial. *EuroIntervention*. 2017;12:2001–2008. doi: 10.4244/EIJ-D-16-00612
- Collet C, Onuma Y, Andreini D, Sonck J, Pompilio G, Mushtaq S, La Meir M, Miyazaki Y, de Mey J, Gaemperli O, et al. Coronary computed tomography angiography for heart team decision-making in multivessel coronary artery disease. *Eur Heart Journal*. 2018;39:3689–3698. doi: 10.1093/eurheartj/ehy581
- Andreini D, Pontone G, Mushtaq S, Conte E, Perchinunno M, Guglielmo M, Volpato V, Annoni A, Baggiano A, Formenti A, et al. Atrial fibrillation: diagnostic accuracy of coronary CT angiography performed with a whole-heart 230-µm spatial resolution CT scanner. *Radiology*. 2017;284:676–684. doi: 10.1148/radiol.2017161779
- Andreini D, Lin FY, Rizvi A, Cho I, Heo R, Pontone G, Bartorelli AL, Mushtaq S, Villines TC, Carrascosa P, et al. Diagnostic performance of a novel coronary CT angiography algorithm: prospective multicenter validation of an intracycle cT motion correction algorithm for diagnostic accuracy. *AJR Am J Roentgenol*. 2018;18:1–8. doi: 10.2214/AJR.17.18670
- Patel MR, Calhoon JH, Dehmer GJ, Grantham JA, Maddox TM, Maron DJ, Smith PK. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 appropriate use criteria for coronary revascularization in patients with stable ischemic heart disease: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2017;69:2212–2241. doi: 10.1016/j.jacc.2017.02.001
- Escaned J, Collet C, Ryan N, De Maria GL, Walsh S, Sabate M, Davies J, Lesiak M, Moreno R, Cruz-Gonzalez I, et al. Clinical outcomes of state-of-the-art percutaneous coronary revascularization in patients with de novo three vessel disease: 1-year results of the SYNTAX II study. *Eur Heart J*. 2017;38:3124–3134. doi: 10.1093/eurheartj/ehx512
- Fournier S, Toth GG, De Bruyne B, Johnson NP, Ciccirelli G, Xaplanteris P, Milkas A, Strisciuglio T, Bartunek J, Vanderheyden M, et al. Six-year follow-up of fractional flow reserve-guided versus angiography-guided coronary artery bypass graft surgery. *Circ Cardiovasc Interv*. 2018;11:e006368. doi: 10.1161/CIRCINTERVENTIONS.117.006368
- Koo BK, Erglis A, Doh JH, Daniels DV, Jegere S, Kim HS, Dunning A, DeFrance T, Lansky A, Leipsic J, et al. Diagnosis of ischemia-causing coronary stenoses by noninvasive fractional flow reserve computed from coronary computed tomographic angiograms. Results from the prospective multicenter DISCOVER-FLOW (Diagnosis of Ischemia-Causing Stenoses Obtained Via Noninvasive Fractional Flow Reserve) study. *J Am Coll Cardiol*. 2011;58:1989–1997. doi: 10.1016/j.jacc.2011.06.066
- Min JK, Leipsic J, Pencina MJ, Berman DS, Koo BK, van Mieghem C, Erglis A, Lin FY, Dunning AM, Apruzzese P, et al. Diagnostic accuracy of fractional flow reserve from anatomic CT angiography. *JAMA*. 2012;308:1237–1245. doi: 10.1001/2012.jama.11274
- Nørgaard BL, Leipsic J, Gaur S, Seneviratne S, Ko BS, Ito H, Jensen JM, Mauri L, De Bruyne B, Bezerra H, et al; NXT Trial Study Group. Diagnostic performance of noninvasive fractional flow reserve derived from coronary computed tomography angiography in suspected coronary artery disease: the NXT trial (analysis of coronary blood flow using CT angiography: next steps). *J Am Coll Cardiol*. 2014;63:1145–1155. doi: 10.1016/j.jacc.2013.11.043
- Zimmermann FM, De Bruyne B, Pijls NH, Desai M, Oldroyd KG, Park SJ, Reardon MJ, Wendler O, Woo J, Yeung AC, et al. Rationale and design of the Fractional Flow Reserve versus Angiography for Multivessel Evaluation (FAME) 3 Trial: a comparison of fractional flow reserve-guided percutaneous coronary intervention and coronary artery bypass graft surgery in patients with multivessel coronary artery disease. *Am Heart J*. 2015;170:619–626. e2. doi: 10.1016/j.ahj.2015.06.024

16. Andreini D, Mushtaq S, Pontone G, Conte E, Guglielmo M, Annoni A, Baggiano A, Formenti A, Ditali V, Mancini ME, et al. Diagnostic performance of coronary CT angiography carried out with a novel whole-heart coverage high-definition CT scanner in patients with high heart rate. *Int J Cardiol*. 2018;257:325–331. doi: 10.1016/j.ijcard.2017.10.084
17. Coenen A, Kim YH, Kruk M, Tesche C, De Geer J, Kurata A, Lubbers ML, Daemen J, Itu L, Rapaka S, et al. Diagnostic accuracy of a machine-learning approach to coronary computed tomographic angiography-based fractional flow reserve: result from the MACHINE consortium. *Circ Cardiovasc Imaging*. 2018;11:e007217. doi: 10.1161/CIRCIMAGING.117.007217
18. Lu MT, Ferencik M, Roberts RS, Lee KL, Ivanov A, Adami E, Mark DB, Jaffer FA, Leipsic JA, Douglas PS, Hoffmann U. Noninvasive FFR derived from coronary CT angiography: management and outcomes in the PROMISE Trial. *JACC Cardiovasc Imaging*. 2017;10:1350–1358. doi: 10.1016/j.jcmg.2016.11.024
19. Andreini D, Pontone G, Mushtaq S, Mancini ME, Conte E, Guglielmo M, Volpato V, Annoni A, Baggiano A, Formenti A, et al. Image quality and radiation dose of coronary CT angiography performed with whole-heart coverage CT scanner with intra-cycle motion correction algorithm in patients with atrial fibrillation. *Eur Radiol*. 2018;28:1383–1392. doi: 10.1007/s00330-017-5131-2
20. Sonck J, Miyazaki Y, Mandry D, Andreini D. Non-invasive treatment planning of tandem coronary artery lesions using an interactive planner for PCI. *EuroIntervention*. 2018;14:924–925. doi: 10.4244/EIJ-D-17-00815
21. Rochitte CE, George RT, Chen MY, Arbab-Zadeh A, Dewey M, Miller JM, Niinuma H, Yoshioka K, Kitagawa K, Nakamori S, et al. Computed tomography angiography and perfusion to assess coronary artery stenosis causing perfusion defects by single photon emission computed tomography: the CORE320 study. *Eur Heart J*. 2014;35:1120–1130. doi: 10.1093/eurheartj/eh488
22. George RT, Arbab-Zadeh A, Miller JM, Vavere AL, Bengel FM, Lardo AC, Lima JA. Computed tomography myocardial perfusion imaging with 320-row detector computed tomography accurately detects myocardial ischemia in patients with obstructive coronary artery disease. *Circ Cardiovasc Imaging*. 2012;5:333–340. doi: 10.1161/CIRCIMAGING.111.969303
23. Osawa K, Miyoshi T, Koyama Y, Hashimoto K, Sato S, Nakamura K, Nishii N, Kohno K, Morita H, Kanazawa S, et al. Additional diagnostic value of first-pass myocardial perfusion imaging without stress when combined with 64-row detector coronary CT angiography in patients with coronary artery disease. *Heart*. 2014;100:1008–1015. doi: 10.1136/heartjnl-2013-305468
24. Greif M, von Ziegler F, Bamberg F, Tittus J, Schwarz F, D'Anastasi M, Marcus RP, Schenzle J, Becker C, Nikolaou K, et al. CT stress perfusion imaging for detection of haemodynamically relevant coronary stenosis as defined by FFR. *Heart*. 2013;99:1004–1011. doi: 10.1136/heartjnl-2013-303794
25. Pontone G, Andreini D, Guaricci AI, Baggiano A, Fazzari F, Guglielmo M, Muscogiuri G, Berzovini CM, Pasquini A, Mushtaq S, et al. Incremental diagnostic value of stress computed tomography myocardial perfusion with whole-heart coverage CT scanner in intermediate- to high-risk symptomatic patients suspected of coronary artery disease. *JACC Cardiovasc Imaging*. 2019;12:338–349. doi: 10.1016/j.jcmg.2017.10.025
26. Schuijff JD, Ko BS, Di Carli MF, Hislop-Jambrich J, Ithdayhid AR, Seneviratne SK, Lima JAC. Fractional flow reserve and myocardial perfusion by computed tomography: a guide to clinical application. *Eur Heart J Cardiovasc Imaging*. 2018;19:127–135. doi: 10.1093/ehjci/jex240

AQ14

AUTHOR QUERIES

Authors please note: Authors are responsible for any page charges as outlined in the acceptance letter or as indicated on the Instructions for Authors (<https://www.ahajournals.org/CIRc/revised-accepted-manuscripts>). Unless you have selected open access for your article, or it is otherwise noted on the acceptance letter, page charges are as follows: \$70 per black and white page (print articles only) or \$35 per page (online-only articles only). For each color page (print only), please add \$653/page. If there are any concerns regarding these charges, these should be addressed within 48 hours of receiving the s-proof. Author(s) will be invoiced for all color and page charges post publication. If you have selected open access for your article, please refer to details in the queries below.

AUTHOR PLEASE ANSWER ALL QUERIES

- AQ1–Please note only those terms that are used 5 times or more can be abbreviated, except trial names and proteins, which should be expanded at first use but then can be abbreviated throughout regardless of how many times they appear.
- AQ2–Please turn to page 3 of your proof and review the running head, which will appear in the upper right-hand margins of odd-numbered pages. Running heads must be 50 or fewer characters in length, including spaces and punctuation. If your original short title was longer than 50 characters, we may have shortened it. Please modify if necessary (but observe our length guidelines).
- AQ3–Please check and confirm that the url and trial number for the clinical trial information is correct.
- AQ4–Key words may have been edited to match the US National Library of Medicine's Medical Subject Headings (<http://www.nlm.nih.gov/mesh/MBrowser.html>). If they need modification, please limit the total number of key words to 7.
- AQ5–Please define "MACE" in box text.
- AQ6–Please define "REVOLUTION and PROMISE" in text.
- AQ7–Please define "SYNTAX" in text, figures, and tables.
- AQ8–Please define "ICA" in Figure 3.
- AQ9–Please check author's name "Stefano de Martini, Pierre-Adrien Metzdorf, and Gloria Färber".
- AQ10–Please confirm that all authors are included in the correct order in the byline and that all names are spelled correctly, including special characters, accents, middle initials, and degrees, if applicable. For indexing purposes, confirm author names have been correctly identified as given names (blue), surnames (red), and suffixes (black). Color in the byline will not appear on the final published version. Note that journal style discourages listing honorary degrees (FAHA, FRCP, etc.) in the byline; please delete such degrees from the author byline.

- AQ11–Please confirm that all authors' institutional affiliations (including city/state/country locations) are correct as shown in the affiliations footnote.
- AQ12–Please provide department or division details, if any, for the affiliations, if not listed already.
- AQ13–Please carefully review any Acknowledgments, Sources of Funding, and/or Disclosures listed at the end of the manuscript (before the References), and confirm that they are accurate and complete for all authors.
- AQ14–Please review the typeset tables carefully against copies of the originals to verify accuracy of editing and typesetting.
- AQ15–We've identified that all figures are low resolution and could result in poor-quality graphics in the final publication. Please send higher-resolution versions of these figures, or approve to proceed with the figures as shown in your proof.