

Title: LEPTIN, RESISTIN, AND PCSK9 - THE ROLE OF STAT3

Short title: STAT3 regulates PCSK9

Chiara Macchi¹, Maria Francesca Greco¹, Margherita Botta¹, Paola Sperandeo¹, Paola Dongiovanni², Luca Valenti^{3,4}, Arrigo F.G. Cicero⁵, Claudio Borghi⁵, Maria Giovanna Lupo⁶, Stefano Romeo^{7,8,9}, Alberto Corsini¹, Paolo Magni^{1,10}, Nicola Ferri⁶ and Massimiliano Ruscica¹

¹Department of Pharmacological and Biomolecular Sciences, Università degli Studi di Milano, Milan, Italy. ²General Medicine and Metabolic Diseases, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ³ Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Milan, Italy; ⁴Translational Medicine, Transfusion Medicine and Hematology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ⁵Department of Medicine and Surgery Sciences, Alma Mater Studiorum University of Bologna, Bologna, Italy; ⁶Department of Pharmaceutical and Pharmacological Sciences, Università degli Studi di Padova, Padova, Italy; ⁷Department of Molecular and Clinical Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; ⁸ Clinical Nutrition Unit, Department of Medical and Surgical Science, Magna Graecia University, Catanzaro, Italy; ⁹Department of Cardiology, Sahlgrenska University Hospital, Gothenburg, Sweden; ¹⁰IRCCS MultiMedica, Sesto S. Giovanni, Milan, Italy.

Corresponding author:

Massimiliano Ruscica

Dipartimento di Scienze farmacologiche e Biomolecolari

Università degli Studi di Milano

e-mail: massimiliano.ruscica@unimi.it

Chiara Macchi

Dipartimento di Scienze farmacologiche e Biomolecolari

Università degli Studi di Milano

e-mail: chiara.macchi@unimi.it

Funding statement: Cariplo Foundation (2015-0552 and 2018-0511 to MR); Grants from MIUR Progetto Eccellenza. Ricerca Finalizzata Ministero della Salute RF-2016-02364358

and Ricerca corrente Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and the European Union (EU) Programme Horizon 2020 (under grant agreement No. 777377) for the project LITMUS-“ Liver Investigation: Testing Marker Utility in Steatohepatitis” to LV.

Conflict of interest: CM, MFG, MB, PS, PD, MGL, PM, AFGC, CB and MR have nothing to declare. SR has been consulting, received lecture fees and research grants from AMGEN, Sanofi-aventis and Astra Zeneca. AC received honoraria from AstraZeneca, AMGEN, Sanofi, Recordati, Novartis, MSD, Mediolanum, DOC, Mylan and Pfizer. LV consulting for Gilead, Pfizer, Astra Zeneca, Novo Nordisk, Intercept pharmaceuticals, Diatech Pharmacogenetics, IONIS.

Abstract

In a condition of dysfunctional visceral fat depots, as in the case of obesity, alterations in adipokines levels may be detrimental for the cardiovascular system. The proinflammatory leptin and resistin adipokines have been described as possible links between obesity and atherosclerosis. The present study was aimed at evaluating whether proprotein convertase subtilisin/kexin type 9 (PCSK9), a key regulator of low-density lipoprotein metabolism, is induced by leptin and resistin through the involvement of the inflammatory pathway of signal transducer and activator of transcript 3 (STAT3). In HepG2 cells, leptin and resistin upregulated PCSK9 gene and protein expression as well as the phosphorylation of STAT3. Upon STAT3 silencing, leptin and resistin lost their ability to activate PCSK9. The knock-down of STAT3 did not affect the expression of leptin and resistin receptors as well as that of PCSK9. The analysis of human *PCSK9* promoter region showed that the two adipokines raise PCSK9 promoter activity via the involvement of sterol regulatory element motif. In healthy male, a positive association between circulating leptin and PCSK9 levels was found only when BMI was < 25 kg/m². In conclusion, our study identified STAT3 as one of the molecular regulators of leptin- and resistin-mediated transcriptional induction of PCSK9.

Keywords: PCSK9, leptin, resistin, STAT3

Introduction

Adipose tissue is an endocrine organ able to secrete active molecules, namely, adipokines which contribute to regulate appetite and satiety, fat distribution, and energy expenditure ¹. When the metabolic milieu is deranged, as in the case of obesity and adipose tissue metaflammation, adipokines may play a role in cardiovascular disease (CVD), by promoting atherogenesis, plaque progression and thrombosis ². Roughly 80% of patients with coronary heart disease are overweight or obese ³. Among the huge variety of adipokines produced by adipose tissue, the proinflammatory leptin and resistin have been described as possible links between obesity and atherosclerosis ^{4, 5}.

Leptin, a 16 kDa cytokine produced predominantly by adipose tissue, regulates feeding and promotes energy expenditure. This hormone is implicated also in the regulation of the immune system, autonomic and cardiovascular regulation ⁶. Leptin exerts its biological actions by binding to a class I cytokine receptor encoded by *LEPR* in humans. Among the six LepR isoforms (LepRa to LepRf), the LepRb is the only one with a full-length intracellular domain which allows the activation of a cell-signalling pathway. In response to leptin, Janus tyrosine kinase 2 (JAK2) phosphorylates LepR triggering the recruitment of the Signal Transducer and Activator of Transcript 3 (STAT3) through its SH2 domain. STAT3 is subsequently phosphorylated by JAK2, resulting in its dimerization and nuclear translocation in order to regulate the transcription of STAT3-target genes ⁷. Among STAT proteins (STAT 1, 2, 3, 4, 5a, 5b, and 6), STAT3 contributes to various metabolic processes like hyperphagia and obesity ⁸, being leptin one of the major players in these events ⁹.

Resistin is a 12.5 kDa sized C-terminal cysteine-rich signaling peptide secreted predominantly by adipose tissue and macrophages in humans and rodents. In humans, resistin adipose tissue expression and circulating levels are increased in overweight and obese subjects, strongly supporting the association with a raised CV risk in obese individuals ¹⁰. Among the designated receptors mediating resistin effects, *e.g.* the inflammatory cascade, the adenylyl cyclase associated protein 1 (CAP1) has been recently described as a *bona fide* resistin receptor ¹¹.

Proprotein convertase subtilisin/kexin type 9 (PCSK9), mainly secreted by the liver and to a lower extent by the intestine, post-translationally regulates the number of cell-surface LDL receptors (LDLR). At transcriptional level, PCSK9 synthesis is largely controlled by the involvement of transcription factor families sterol regulatory element (SRE)-binding proteins (SREBPs) and by the acute-phase response transcriptional

controller hepatocyte nuclear factor-1 (HNF-1) α ¹². This last evidence along with the knowledge that PCSK9 expression is also regulated by the pro-inflammatory cytokine tumor necrosis factors (TNF)- α , in a suppressor of cytokine signaling (SOCS)3-dependent manner, suggests a direct link between inflammation and the regulation of lipid metabolism via PCSK9 ¹³.

Considering that PCSK9 circulating levels have been associated with atherosclerosis and to support the hypothesis that the derangement in adipokine secretion is involved in obesity-associated CV risk ¹⁴, the aim of the present study was to evaluate whether leptin and resistin mediate PCSK9 activation through the inflammatory STAT3 pathway. Indeed, the JAK/STAT signaling pathway, a pillar downstream mediator of a variety of cytokines, is dysregulated in metabolic diseases including obesity ¹⁵.

Materials and Methods

Cell cultures - The human hepatocellular carcinoma cell line, HepG2, was routinely cultured in 10% Fetal Bovine Serum (FBS)/DMEM supplemented with penicillin, streptomycin, nonessential amino acids and sodium pyruvate (all from Sigma Aldrich, Milan, Italy). For the experiments, cells were starved overnight and then incubated with DMEM containing 10% of human lipoprotein plasma deprived serum (LPDS).

Reagents and antibodies - DMEM, trypsin EDTA, penicillin, streptomycin, nonessential amino acid solution, FBS, disposable culture flasks and petri dishes were from Merck (Milan, Italy). Molecular weight protein standard, precast polyacrylamide gels (4-12%), BCA assay for determination of protein concentrations, maxima 1strand cDNA and maxima SYBR Green/Fluorescein qPCR were all from Life Technologies (Monza, Italy). Bovine Serum Albumin was from SIGMA-Aldrich (Milan, Italy). Human recombinant leptin (Sigma Aldrich) and resistin (Bio Visoon, CA) were used at 100 ng/mL. Human recombinant PCSK9 was used at 2.5 μ g/mL (Cayman Chemicals, item 20631, US) ¹⁶.

Transfection of siRNA - ON-TARGET plus SMART pool siRNA directed to STAT3 and PCSK9 or scramble control were from DharmaconTM (Carlo Erba Reagents, Milan, Italy). Transfections were performed by using SilentFectTM Lipid Reagent (BIO-RAD laboratories,

Hercules, CA) according to the manufacturer's protocol. The day before the transfection, HepG2 cells were seeded in MEM/10%FBS at a density of 6×10^5 /well (6-well tray). Cells were then transfected with 40 nmol/L of siRNA. Forty-eight hours post transfection, the medium was replaced with DMEM/10%LPDS \pm leptin (100 ng/mL) or resistin (100 ng/mL) for an additional 48 h before performing the quantitative (q)PCR analysis.

In silico analyses for STAT3 binding motifs on human PCSK9 promoter - Human, gorilla, mouse and rat PCSK9 promoter sequences (up to 2000 bp upstream the start codon) were retrieved from Ensembl Genome Browser ¹⁷ by using the latest released genome assembly (GRCh38.p13, Kamilah_GGO_v0, GRCh38.p6 and Rnor_6.0 for human, gorilla, mouse and rat, respectively). JASPAR, an open-access database of curated, non-redundant transcription factors binding (TFB) profiles modelled as position-specific weight matrices (PSSMs), was used to scan the selected PCSK9 promoter region for any STAT3 binding site ¹⁸. The relative profile score threshold - defined as the minimum relative score required for reporting a match between the TFB model and the imputed sequence - was set at 80%. FIMO tool, from the open-access MEME-Suit 5.1.1, was used to double-check the results obtained from JASPAR ^{19, 20}. STAT3 PSSMs profile was retrieved from JASPAR and imputed in FIMO and the selected PCSK9 promoter region scanned. Matches with a p -value ≤ 0.0001 were taken into account. The same analyses were carried out for SRE and HNF-1.

Generation of PCSK9 promoter luciferase reporters with a STAT3 binding site deletion. D1-STAT3-mut plasmid was generated by deleting the STAT3 binding sequence predicted by JASPER in the PCSK9 promoter sequence in plasmid D1. Plasmid D1 is a derivative of the commercial promoter-reporter vector pGL3-Basic, generated by cloning the human PCSK9 promoter fragment spanning - 1,711 to -94 upstream of the *luc* reporter gene. Deletion of the STAT3 sequence (CTTCTGGAAAG), spanning - 916 to - 906 in the PCSK9 promoter, was obtained by two-step PCR according to the following protocol. Two fragments flanking the putative STAT3 sequence were PCR amplified using D1 plasmid as template and the following primer pairs: AP679 (5'-CGACGAGGTACCGAGCTCGGATCCACTAGTAAC-3') - AP680 (5'-ATTCAATTTGCAAAGATTTC-3'),

to generate Fragment UP (upstream), and AP681 (5'-CAAATTGAATCTGAGCTTGTGCCTACCATAG-3') - AP682 (5'-CCCAAGCTTACTGTGCAGGAGCTGAAGTTC-3'), to generate Fragment DOWN (downstream). Primer AP679 and AP682 sequences contain the KpnI and HindIII restriction sites, respectively. Fragments UP and DOWN were used as template for a second round of PCR amplification using AP679 and AP682 primers. The final PCR product was cloned in the D1 plasmid between KpnI and HindIII restriction sites in place of the PCSK9 promoter region, generating D1-STAT3 plasmid. The deletion of the STAT3 region from the PCSK9 promoter in D1-STAT3 plasmid was confirmed by DNA sequencing.

Transfections of Reporter Constructs - As previously described, HepG2 cells were transfected with the plasmid PCSK9 pGL3-PCSK9-D4 containing the 5' flanking region of the PCSK9 gene from -440 to -94, relative to the ATG start codon in front of the luciferase coding sequence. The promoter constructs contain wild-type, SRE mutated (SRE-mu) and HNF-1 alpha mutated (HNF1-mu) sequences ²¹. To measure the human PCSK9 promoter activity, HepG2 cells were seeded in 48-well plates at a density of 4×10^5 cells per well. The day after, cells were transiently transfected with pGL3-PCSK9-D1 (wildtype and STAT3 mutated) and with pGL3-PCSK9-D4 plasmids (wild-type, SRE-mu and HNF-1-mu), with TurboFectTM transfection reagent (Thermo Fisher). Forty-eight hours post transfection, cells were incubated with DMEM/10%LPDS \pm leptin (100 ng/mL), resistin (100 ng/mL) and simvastatin (20 μ M) for an additional 24 hours. Luciferase activities were measured by using Neolite reagent (Perkin Elmer, Milan, Italy) according to manufacturer's instructions.

RNA preparation and quantitative real time PCR - Total RNA was extracted with the iScript Sample Preparation Buffer cDNA synthesis preparation reagents (BIO-RAD laboratories) according to manufacturer's instructions or by spin column (Qiagen, Milan, Italy). Reverse transcription-polymerase first-strand cDNA synthesis was performed by using maxima 1strand cDNA synthesis kit (Thermo, Life Technologies). qPCR was then performed by using the Kit Thermo SYBR Green/ROX qPCR Master Mix (Thermo, Life Technologies) and specific primers for selected genes. The analyses were performed with the 9600 Biorad Real-Time PCR Detection Systems (Biorad). The primer sequences

have been listed in Table 1. PCR cycling conditions were as follows: 94°C for 5 min, 40 cycles at 94°C for 15 s, and 60°C for 30 s. Data were expressed as Ct values and used for the relative quantification of targets with the $2^{-\Delta\Delta Ct}$ calculation.

Western Blot Analysis - Total cytosolic protein extracts of HepG2 were obtained by collecting cells in 70 µl of Mammalian Protein Extraction Reagents (Thermo Fisher Scientific) containing a cocktail of protease and phosphatase inhibitors (Roche Diagnostics). Ten µg of proteins and a molecular mass marker (Novex Sharp Protein Standard, Invitrogen™; Life Technologies) were separated on 4-12% sodium dodecylsulfate-polyacrylamide gel (SDS-PAGE; Novex NuPAGE 4-12% Bis-Tris Mini Gels, Invitrogen; Life Technologies) under denaturing and reducing conditions. Proteins were then transferred to a nitrocellulose membrane by using the iBlot™ Gel Transfer Device (Invitrogen; Life Technologies). The membranes were washed with Tris-Buffered Saline-Tween 20 (TBS-T) and non-specific binding sites were blocked in TBS-T containing 5% BSA (SIGMA-Aldrich) for 90 min at room temperature. The blots were incubated overnight at 4°C with a diluted solution (5% BSA or non-fat dried milk) of the human primary antibodies (listed below). Membranes were washed with TBS-T and then exposed for 90 min at RT to a diluted solution (5% non-fat dried milk) of the secondary antibodies (anti-mouse and anti-rabbit peroxidase-conjugated secondary antibodies (New England Biolabs, MA). Immunoreactive bands were detected by exposing the membranes to Clarity Western ECL chemiluminescent substrates (Bio-Rad Laboratories) for 5 min and images were acquired with a ChemiDoc XRS System (Bio-Rad Laboratories). Densitometric readings were evaluated using the ImageLab software as previously described. The dilution of the human primary antibodies were: PCSK9 (1:1,000 - Genetex, CA); STAT3 (1:1,000 - Cell Signaling); pSTAT3 (1:10,000 - Abcam, UK); actin (1:1,000 - Santa Cruz, CA), tubulin (1:2,000 - Sigma Aldrich) and vinculin (1:1,000 - Genetex).

Study population - The association between PCSK9 and leptin and resistin plasma levels was evaluated in 149 healthy male subjects from the Brisighella Heart Study. This study represents a longitudinal population study on a randomized sample representative of the entire population of Brisighella, a rural Northern Italian village. The study has been

active since 1972 and is carried out in agreement with the Declaration of Helsinki. The protocol was approved by the institutional ethics board of the University Hospital of Bologna ²².

ELISA assays - Plasma PCSK9 concentrations were measured by a commercial enzyme-linked immunosorbent assay (ELISA) kit (R&D Systems, Minneapolis, MN). The minimum detectable PCSK9 concentration was 0.219 ng/mL ²³. Concerning the evaluation of PCSK9 in the conditioned medium, HepG2 cells were cultured in 12-well plates (6×10^5 cells/well). Medium (200 μ L) was collected after 24-h treatment with leptin, resistin and simvastatin. Samples were not diluted and results normalized for the total amount of proteins.

Plasma leptin concentrations were measured by a commercial ELISA kit (R&D Systems). The minimum detectable dose of human leptin is typically less than 7.8 pg/mL ²⁴.

Analysis of the data - Statistical analysis was performed using the Prism statistical analysis package version 6.0 (GraphPad Software, San Diego, CA). Data are given as mean \pm SD of three independent experiments. When possible, p-values were determined by Student's t-test. Otherwise, differences between treatment groups were evaluated by 1-way ANOVA. A probability value of $p < 0.05$ was considered statistically significant.

Results

Leptin and resistin induce PCSK9. First of all, a time course experiment (4-, 8-, 16- and 24 hours) was conducted to evaluate the gene expression of *PCSK9* upon leptin (100 ng/mL) and resistin (100 ng/mL) treatment (Supplemental Figure 1). Incubation of HepG2 cells with human recombinant leptin and resistin resulted in a significant rise in PCSK9 gene expression of roughly +50% (Fig. 1A) after 24 hours. Compared to medium alone, protein expression, evaluated by Western blot analysis, was increased roughly by 2- and 1.7-fold, upon treatment with leptin and resistin, respectively (Fig. 1B). Simvastatin (20 μ M) - used as a positive control - raised both PCSK9 gene and protein expression (Fig. 1D and E). To corroborate Western blot analyses, the release of PCSK9 was measured. In the HepG2 conditioned-medium, the levels of PCSK9 were increased by 42.6% ($p < 0.01$) and

36.3% ($p < 0.05$) following leptin and resistin treatments, respectively (Fig. 1C). Simvastatin (20 μ M) was used as a positive control (Fig. 1F).

STAT3 mediates leptin- and resistin-driven PCSK9 activation. In our cell-based model, 24-h treatment with leptin and resistin phosphorylated STAT3 (pSTAT3^{Y705}) without affecting that of total STAT3 (Fig. 2A). To follow-up the hypothesis that STAT3 was involved in leptin- and resistin-driven PCSK9 activation, HepG2 cells were knock-down for STAT3. Transfection with specific siRNA anti-STAT3 fully abolished the gene and protein expression of STAT3 (Fig. 2B-C). Under these experimental conditions, leptin and resistin did not upregulate PCSK9 expression (gene and protein), while basal PCSK9 gene and protein expression was not affected (Fig. 2D-F). These data suggest that PCSK9 upregulation by leptin and resistin in hepatocytes requires STAT3. To exclude any effect of PCSK9 on STAT3 expression, we showed that silencing PCSK9 did not result in any change in STAT3 levels (Fig. 2C). Moreover, application of siSTAT3 did not change the gene expression of leptin (*LEPR*) and resistin (*CAP1*) receptors (Supplemental Figure 2).

STAT3 and PCSK9 transcriptional activity. The involvement of STAT3 in the PCSK9 pathway was further investigated at the transcriptional level. First of all, an *in silico* search of STAT3 binding sites on proximal PCSK9 promoter region was performed by using JASPAR database and FIMO tool from MEME-Suit 5.1.1. The analyses revealed a strong STAT3 binding site (JASPAR score: 96%; FIMO threshold: $p = 2.07E-05$) spanning the region -916 to -906 in the human PCSK9 promoter (Table 2). This region is upstream the Sp1/HNF1- α /SRE triad already published by Li *et al.*²¹ (Fig. 3A). To corroborate this finding, a similar analysis was performed on gorilla, mouse, and rat PCSK9 promoters: a multiple alignment of the four sequences was generated thanks to Nucleotide BLAST Alignment tool by NCBI (Fig. 3A). The mild degeneration in the STAT3 consensus sequence has been already reported in literature²⁵. After transfection with D1 and D1-STAT3-mut plasmids (Fig. 3B), both containing the human PCSK9 promoter region spanning -1,711 to -94 nucleotides upstream of the gene coding for the luciferase enzyme, HepG2 cells were treated with leptin, resistin and simvastatin. Twenty-four-hour-treatment enhanced transcriptional activity, an effect which was not counteracted when STAT3 mutation was inserted (Fig. 3C). Simvastatin was used as a positive control (Fig. 3D). These data suggest that PCSK9 upregulation by leptin and resistin in hepatocytes is not directly mediated by binding of STAT3 to the PCSK9 promoter.

Leptin and resistin affect the transcriptional activity of PCSK9. To further explore the mechanism whereby leptin and resistin upregulate PCSK9 transcript levels, HepG2 cells were transiently transfected with constructs containing the proximal human PCSK9 promoter region (-440 to -94). As shown in Fig. 4A, leptin and resistin significantly incremented the luciferase activity by 54% and 15%, respectively. Similar results were obtained upon treatment with simvastatin (+141%). Being SREBPs and HNF-1 the major transcription factors regulating PCSK9²⁶, we investigated luciferase activity upon the insertion of SRE and HNF-1 mutations in PCSK9 promoter sequence. The presence of mutation in the SRE sequence completely abolished the PCSK9 transcriptional activity driven by leptin and resistin as well as that of simvastatin (Fig. 4B). Conversely, mutation in HNF-1 did not alter leptin- and resistin-driven luciferase activities, showing an increment of 43% and 58%, respectively (Fig 4C).

Association studies. The circulating levels of leptin and PCSK9 were measured by ELISA in a clinical setting of 149 males (56 ± 4 years) free of cardiovascular disease at enrolment and belonging to the cohort of Brisighella Heart Study²². A positive association between circulating leptin and PCSK9 levels ($\beta = 0.352$, $p = 0.014$) was found only in subjects with BMI < 25 kg/m² (Fig 5A). When BMI rose, *i.e.* between 25 and 30 kg/m² ($\beta = 0.147$, $p = 0.295$; Fig 5B) or > 30 kg/m² ($\beta = -0.030$, $p = 0.840$; Fig 5C), the association was lost. Concerning resistin, the circulating levels did not associate with PCSK9 (data not shown). The main characteristics of the selected participants have been reported in the supplemental table 1.

Discussion

Obesity and associated metabolic disorders are becoming major health care concerns worldwide. Obesity is highly associated with chronic low-grade inflammation and it is believed that this obesity-linked inflammatory state is due to changes in the expression of cytokines released by adipose tissue. Adipose tissue is a source of secreted adipokines that act as modulators of metabolic and cardiovascular processes²⁷. Among adipokines, the expression of LDLR has been described to be modulated by leptin and resistin, the last being able to stabilize the cellular expression of PCSK9^{5, 28}. Adiponectin has been also described to affect PCSK9 expression/production, although via the activation of PPAR γ and AMPK pathways²⁹.

In the tight liaison among adipokines, inflammation and atherosclerosis, the main finding of this study relies on the evidence that leptin and resistin, two adipokines associated to regulation of atherogenesis and inflammation, induce the expression of PCSK9 in hepatocytes, which are the main source of circulating protein, through the Janus kinase (JAK)/STAT3 inflammatory pathway. A growing body of evidence suggests that this pathway is dysregulated in the context of obesity and metabolic disease ¹⁵. The interplay between inflammation and PCSK9 has been investigated in the last years leading to the hypothesis that it represents a fed-forward loop in which inflammation activates PCSK9 and *vice versa* PCSK9 impacts on the inflammatory burden associated to atheroma formation ³⁰.

The cytokine responsive JAK/ STAT pathway is involved in adipokine-mediated crosstalk between adipocytes and liver or skeletal muscle ^{31, 32}. The involvement of STAT3 signaling upon leptin receptor activation represents the primary mechanism by which leptin regulates energy balance ³³ and resistin has also been reported to activate this pathway ³⁴. In particular, STAT3, a transcription factor expressed in multiple metabolic tissues, is activated in response to cytokines, growth factors, and nutrients ³⁵. STAT3 may account for the constitutive activation of NF- κ B during chronic inflammation. This activation contributes to the transcriptional regulation of inflammatory cytokines, such as ICAM-1, VCAM-1, MCP-1 and IL-6, as well as matrix metalloproteinases ³⁶. Moreover, it has been showed that the impairment of oncostatin M signaling, a secreted cytokine mainly involved in chronic inflammatory and cardiovascular diseases, has a protective effect on atherosclerosis partially due to the inhibition of the JAK2/STAT3 activation in macrophages ³⁷. Thus, overall, by silencing STAT3, we have demonstrated that this inflammatory pathway may be at the crossroad between adipose tissue and PCSK9 activation. Furthermore, STAT3 phosphorylation is markedly increased in atherosclerotic lesions, and interfering with STAT3 pathway prevents atherosclerotic lesion formation *in vivo* ³⁸. In line with this evidence, the direct role of PCSK9 in atheroma formation should be considered. The Atherosclerosis Risk in Communities (ARIC) epidemiological study demonstrated that genetically reduced levels of PCSK9 were associated with a reduction in myocardial infarction, fatal coronary heart diseases or coronary revascularization, despite a significant prevalence of other nonlipid-related CV-risk factors ³⁹. Higher serum PCSK9 levels were linearly associated with a higher necrotic core fraction in coronary atherosclerosis, regardless of serum LDL-C, confirming data in PCSK9 knock-out mice that were partially protected from neointimal formation

⁴⁰. In humans, PCSK9 has been also described to be associated with other components of atheroma formation, *i.e.* platelet aggregation ¹⁶, hypertension and inflammation ³⁰.

We have found that leptin and resistin activate the transcription of PCSK9 through the involvement of the SRE motif, apparently, without the involvement of HNF-1 ⁴¹. The involvement of STAT3 pathway at the transcriptional level was negligible. However, this latter analysis requires further investigation since the promoter region of human *PCSK9* contains other STAT3 responsible elements (Table 2). Since PCSK9 transcription is controlled through cis regulatory elements imbedded in the proximal promoter region of *PCSK9* gene where the Sp1 sites, SRE and HNF-1 are located, we utilized a PCSK9 promoter luciferase reporter plasmid containing mutations for both SRE and HNF responsive elements. We found that exposure of HepG2 hepatocytes to leptin and resistin increases the transcriptional activity of the PCSK9 promoter. Furthermore, while a construct variant containing a mutation in HNF-1 motif did not affect the ability of adipokines to upregulate luciferase activity, no induction was detected after introducing a mutation in the SRE motif. Relative to this latter evidence, our data are in line with findings demonstrating that, at transcriptional level PCSK9 synthesis is largely controlled by SREBPs ⁴². Our findings are in agreement with the observation that leptin induces SREBP1 activation in porcine granulosa cells ⁴³.

When looking at the correlation between circulating levels of PCSK9 and leptin, we found that this exists only in the group with BMI < 25 Kg/m², an evidence in line with that previous published by Kwakernaak et al. ⁴⁴ in 30 women. This becomes relevant considering that both PCSK9 and leptin levels follow a gender difference, being higher in female subjects than in male⁴⁵⁻⁴⁷. Overall, leptin may be a determinant of PCSK9 levels predominantly before a state of leptin resistance occurs and, most of all, in obese patients other factors may impact on the regulation of PCSK9 levels. Leptin resistance, proposed to take place during the onset and progression of obesity ⁴⁸, could be the result of an impaired leptin signaling pathway, including reduced access of the hormone to its receptor or due to epigenetic modifications that could contribute to leptin expression and signaling disturbances ⁴⁹. This complex scenario has been also highlighted in ob/ob mice lacking leptin in which the effects of leptin replacement on PCSK9 mRNA expression and protein levels were sex-dependent ⁵⁰. In male mice, leptin treatment reduced plasma levels of PCSK9, whereas no changes were observed in females ⁵⁰. On the other hand, although it was suggested that resistin may be involved in the interplay between inflammation and atherosclerosis, whether circulating resistin is a determinant of higher PCSK9 levels needs to be further explored ⁴⁴. Indeed, our data obtained in

healthy male subjects did not fit with a previous report by Li et al.⁵¹, who found that in patients with cardiovascular artery disease resistin positively correlate with PCSK9 levels. Some of these discrepancies may also rely on the tight liaison surrounding resistin and PCSK9, *i.e.* the C-terminal domain of PCSK9 has a structural homology with the human resistin C-terminal domain, and CAP1 represents a binding partner of PCSK9 in mediating LDLR degradation⁵². On this matter, we have previously demonstrated that PCSK9 induces a pro-inflammatory response in macrophages⁵³, an effect common to resistin, *i.e.* a rise in IL6, TNF- α , MCP-1 (data not published). Conversely, a possible *vicious circle* in the liver has been excluded based on the observation that treatment with human recombinant PCSK9 did not increase the gene expression of *IL6* in HepG2 cells (Supplemental Figure 3). This is in line with recent findings showing that PCSK9 is not induced in artificial human inflammation and is not correlated with inflammatory response⁵⁴.

Overall, we provided evidence that PCSK9 may be one of the players at crossroad of the interplay between obesity, adipose tissue dysfunction and CVD risk. Such effect appears to be dependent on the inflammatory JAK/STAT pathway.

References

- [1] Perez-Tilve D: 'Et Tu, Leptin?'. *Trends Endocrinol Metab* 2019, 30:232-3.
- [2] Barale C, Russo I: Influence of Cardiometabolic Risk Factors on Platelet Function. *Int J Mol Sci* 2020, 21.
- [3] Ades PA, Savage PD: Obesity in coronary heart disease: An unaddressed behavioral risk factor. *Prev Med* 2017, 104:117-9.
- [4] Rashid S, Kastelein JJ: PCSK9 and resistin at the crossroads of the atherogenic dyslipidemia. *Expert Rev Cardiovasc Ther* 2013, 11:1567-77.
- [5] Du Y, Li S, Cui CJ, Zhang Y, Yang SH, Li JJ: Leptin decreases the expression of low-density lipoprotein receptor via PCSK9 pathway: linking dyslipidemia with obesity. *J Transl Med* 2016, 14:276.
- [6] Cui H, Lopez M, Rahmouni K: The cellular and molecular bases of leptin and ghrelin resistance in obesity. *Nat Rev Endocrinol* 2017, 13:338-51.
- [7] Banks AS, Davis SM, Bates SH, Myers MG, Jr.: Activation of downstream signals by the long form of the leptin receptor. *J Biol Chem* 2000, 275:14563-72.
- [8] Gao Q, Wolfgang MJ, Neschen S, Morino K, Horvath TL, Shulman GI, Fu XY: Disruption of neural signal transducer and activator of transcription 3 causes obesity, diabetes, infertility, and thermal dysregulation. *Proc Natl Acad Sci U S A* 2004, 101:4661-6.
- [9] Bates SH, Stearns WH, Dundon TA, Schubert M, Tso AW, Wang Y, Banks AS, Lavery HJ, Haq AK, Maratos-Flier E, Neel BG, Schwartz MW, Myers MG, Jr.: STAT3 signalling is required for leptin regulation of energy balance but not reproduction. *Nature* 2003, 421:856-9.
- [10] Reilly MP, Lehrke M, Wolfe ML, Rohatgi A, Lazar MA, Rader DJ: Resistin is an inflammatory marker of atherosclerosis in humans. *Circulation* 2005, 111:932-9.
- [11] Lee S, Lee HC, Kwon YW, Lee SE, Cho Y, Kim J, Lee S, Kim JY, Lee J, Yang HM, Mook-Jung I, Nam KY, Chung J, Lazar MA, Kim HS: Adenylyl cyclase-associated protein 1 is a receptor for human resistin and mediates inflammatory actions of human monocytes. *Cell Metab* 2014, 19:484-97.
- [12] Macchi C, Banach M, Corsini A, Sirtori CR, Ferri N, Ruscica M: Changes in circulating pro-protein convertase subtilisin/kexin type 9 levels - experimental and clinical approaches with lipid-lowering agents. *Eur J Prev Cardiol* 2019, 26:930-49.
- [13] Momtazi-Borojeni AA, Sabouri-Rad S, Gotto AM, Pirro M, Banach M, Awan Z, Barreto GE, Sahebkar A: PCSK9 and inflammation: a review of experimental and clinical evidence. *Eur Heart J Cardiovasc Pharmacother* 2019, 5:237-45.
- [14] Ruscica M, Baragetti A, Catapano AL, Norata GD: Translating the biology of adipokines in atherosclerosis and cardiovascular diseases: Gaps and open questions. *Nutr Metab Cardiovasc Dis* 2017, 27:379-95.
- [15] Dodington DW, Desai HR, Woo M: JAK/STAT - Emerging Players in Metabolism. *Trends Endocrinol Metab* 2018, 29:55-65.
- [16] Camera M, Rossetti L, Barbieri SS, Zanolli I, Canciani B, Trabattini D, Ruscica M, Tremoli E, Ferri N: PCSK9 as a Positive Modulator of Platelet Activation. *J Am Coll Cardiol* 2018, 71:952-4.
- [17] Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov AG, Bennett R, Bhai J, Billis K, Boddu S, Marugan JC, Cummins C, Davidson C, Dodiya K, Fatima R, Gall A, Giron CG, Gil L, Grego T, Haggerty L, Haskell E, Hourlier T, Izuogu OG, Janacek SH, Juettemann T, Kay M, Lavidas I, Le T, Lemos D, Martinez JG, Maurel T, McDowall M, McMahon A, Mohanan S, Moore B, Nuhn M, Oheh DN, Parker A, Parton A, Patricio M, Sakthivel MP, Abdul Salam AI, Schmitt BM, Schuilenburg H, Sheppard D, Sycheva M, Szuba M, Taylor K, Thormann A, Threadgold

G, Vullo A, Walts B, Winterbottom A, Zadissa A, Chakiachvili M, Flint B, Frankish A, Hunt SE, G II, Kostadima M, Langridge N, Loveland JE, Martin FJ, Morales J, Mudge JM, Muffato M, Perry E, Ruffier M, Trevanion SJ, Cunningham F, Howe KL, Zerbino DR, Flicek P: Ensembl 2020. *Nucleic Acids Res* 2020, 48:D682-D8.

[18] Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, Modi BP, Correard S, Gheorghe M, Baranasic D, Santana-Garcia W, Tan G, Cheneby J, Ballester B, Parcy F, Sandelin A, Lenhard B, Wasserman WW, Mathelier A: JASPAR 2020: update of the open-access database of transcription factor binding profiles. *Nucleic Acids Res* 2020, 48:D87-D92.

[19] Grant CE, Bailey TL, Noble WS: FIMO: scanning for occurrences of a given motif. *Bioinformatics* 2011, 27:1017-8.

[20] Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS: MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* 2009, 37:W202-8.

[21] Li H, Dong B, Park SW, Lee HS, Chen W, Liu J: Hepatocyte nuclear factor 1alpha plays a critical role in PCSK9 gene transcription and regulation by the natural hypocholesterolemic compound berberine. *J Biol Chem* 2009, 284:28885-95.

[22] Ruscica M, Ferri N, Fogacci F, Rosticci M, Botta M, Marchiano S, Magni P, D'Addato S, Giovannini M, Borghi C, Cicero AFG, Brisighella Heart Study G: Circulating Levels of Proprotein Convertase Subtilisin/Kexin Type 9 and Arterial Stiffness in a Large Population Sample: Data From the Brisighella Heart Study. *J Am Heart Assoc* 2017, 6.

[23] Ruscica M, Simonelli S, Botta M, Ossoli A, Lupo MG, Magni P, Corsini A, Arca M, Pisciotta L, Veglia F, Franceschini G, Ferri N, Calabresi L: Plasma PCSK9 levels and lipoprotein distribution are preserved in carriers of genetic HDL disorders. *Biochim Biophys Acta Mol Cell Biol Lipids* 2018, 1863:991-7.

[24] Ruscica M, Macchi C, Gandini S, Morlotti B, Erzegovesi S, Bellodi L, Magni P: Free and bound plasma leptin in anorexia nervosa patients during a refeeding program. *Endocrine* 2016, 51:380-3.

[25] Tripathi SK, Chen Z, Larjo A, Kanduri K, Nousiainen K, Aijo T, Ricano-Ponce I, Hrdlickova B, Tuomela S, Laajala E, Salo V, Kumar V, Wijmenga C, Lahdesmaki H, Lahesmaa R: Genome-wide Analysis of STAT3-Mediated Transcription during Early Human Th17 Cell Differentiation. *Cell Rep* 2017, 19:1888-901.

[26] Ruscica M, Ricci C, Macchi C, Magni P, Cristofani R, Liu J, Corsini A, Ferri N: Suppressor of Cytokine Signaling-3 (SOCS-3) Induces Proprotein Convertase Subtilisin Kexin Type 9 (PCSK9) Expression in Hepatic HepG2 Cell Line. *J Biol Chem* 2016, 291:3508-19.

[27] Neeland IJ, Ross R, Despres JP, Matsuzawa Y, Yamashita S, Shai I, Seidell J, Magni P, Santos RD, Arsenault B, Cuevas A, Hu FB, Griffin B, Zambon A, Barter P, Fruchart JC, Eckel RH, International Atherosclerosis S, International Chair on Cardiometabolic Risk Working Group on Visceral O: Visceral and ectopic fat, atherosclerosis, and cardiometabolic disease: a position statement. *Lancet Diabetes Endocrinol* 2019, 7:715-25.

[28] Melone M, Wilsie L, Palyha O, Strack A, Rashid S: Discovery of a new role of human resistin in hepatocyte low-density lipoprotein receptor suppression mediated in part by proprotein convertase subtilisin/kexin type 9. *J Am Coll Cardiol* 2012, 59:1697-705.

[29] Sun L, Yang X, Li Q, Zeng P, Liu Y, Liu L, Chen Y, Yu M, Ma C, Li X, Li Y, Zhang R, Zhu Y, Miao QR, Han J, Duan Y: Activation of Adiponectin Receptor Regulates Proprotein Convertase Subtilisin/Kexin Type 9 Expression and Inhibits Lesions in ApoE-Deficient Mice. *Arterioscler Thromb Vasc Biol* 2017, 37:1290-300.

- [30] Ruscica M, Tokgozoglu L, Corsini A, Sirtori CR: PCSK9 inhibition and inflammation: A narrative review. *Atherosclerosis* 2019, 288:146-55.
- [31] Emanuelli B, Peraldi P, Filloux C, Chavey C, Freidinger K, Hilton DJ, Hotamisligil GS, Van Obberghen E: SOCS-3 inhibits insulin signaling and is up-regulated in response to tumor necrosis factor-alpha in the adipose tissue of obese mice. *J Biol Chem* 2001, 276:47944-9.
- [32] Ueki K, Kondo T, Kahn CR: Suppressor of cytokine signaling 1 (SOCS-1) and SOCS-3 cause insulin resistance through inhibition of tyrosine phosphorylation of insulin receptor substrate proteins by discrete mechanisms. *Mol Cell Biol* 2004, 24:5434-46.
- [33] Allison MB, Myers MG, Jr.: 20 years of leptin: connecting leptin signaling to biological function. *J Endocrinol* 2014, 223:T25-35.
- [34] Pirvulescu M, Manduteanu I, Gan AM, Stan D, Simion V, Butoi E, Calin M, Simionescu M: A novel pro-inflammatory mechanism of action of resistin in human endothelial cells: up-regulation of SOCS3 expression through STAT3 activation. *Biochem Biophys Res Commun* 2012, 422:321-6.
- [35] Mashili F, Chibalin AV, Krook A, Zierath JR: Constitutive STAT3 phosphorylation contributes to skeletal muscle insulin resistance in type 2 diabetes. *Diabetes* 2013, 62:457-65.
- [36] Bode JG, Albrecht U, Haussinger D, Heinrich PC, Schaper F: Hepatic acute phase proteins--regulation by IL-6- and IL-1-type cytokines involving STAT3 and its crosstalk with NF-kappaB-dependent signaling. *Eur J Cell Biol* 2012, 91:496-505.
- [37] Zhang X, Li J, Qin JJ, Cheng WL, Zhu X, Gong FH, She Z, Huang Z, Xia H, Li H: Oncostatin M receptor beta deficiency attenuates atherogenesis by inhibiting JAK2/STAT3 signaling in macrophages. *J Lipid Res* 2017, 58:895-906.
- [38] Dutzmann J, Daniel JM, Bauersachs J, Hilfiker-Kleiner D, Sedding DG: Emerging translational approaches to target STAT3 signalling and its impact on vascular disease. *Cardiovasc Res* 2015, 106:365-74.
- [39] Cohen JC, Boerwinkle E, Mosley TH, Jr., Hobbs HH: Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 2006, 354:1264-72.
- [40] Ferri N, Marchiano S, Tibolla G, Baetta R, Dhyani A, Ruscica M, Uboldi P, Catapano AL, Corsini A: PCSK9 knock-out mice are protected from neointimal formation in response to perivascular carotid collar placement. *Atherosclerosis* 2016, 253:214-24.
- [41] Dong B, Singh AB, Shende VR, Liu J: Hepatic HNF1 transcription factors control the induction of PCSK9 mediated by rosuvastatin in normolipidemic hamsters. *Int J Mol Med* 2017, 39:749-56.
- [42] Jeong HJ, Lee HS, Kim KS, Kim YK, Yoon D, Park SW: Sterol-dependent regulation of proprotein convertase subtilisin/kexin type 9 expression by sterol-regulatory element binding protein-2. *J Lipid Res* 2008, 49:399-409.
- [43] Ruiz-Cortes ZT, Martel-Kennes Y, Gevry NY, Downey BR, Palin MF, Murphy BD: Biphasic effects of leptin in porcine granulosa cells. *Biol Reprod* 2003, 68:789-96.
- [44] Kwakernaak AJ, Lambert G, Dullaart RP: Plasma proprotein convertase subtilisin-kexin type 9 is predominantly related to intermediate density lipoproteins. *Clin Biochem* 2014, 47:679-82.
- [45] Lakoski SG, Lagace TA, Cohen JC, Horton JD, Hobbs HH: Genetic and metabolic determinants of plasma PCSK9 levels. *J Clin Endocrinol Metab* 2009, 94:2537-43.
- [46] Ooi TC, Raymond A, Cousins M, Favreau C, Taljaard M, Gavin C, Jolly EE, Malone S, Eapen L, Chretien M, Mbikay M, Mayne J: Relationship between testosterone, estradiol and circulating PCSK9: Cross-sectional and interventional studies in humans. *Clin Chim Acta* 2015, 446:97-104.

- [47] Macchi C, Ferri N, Favero C, Cantone L, Vigna L, Pesatori AC, Lupo MG, Sirtori CR, Corsini A, Bollati V, Ruscica M: Long-term exposure to air pollution raises circulating levels of proprotein convertase subtilisin/kexin type 9 in obese individuals. *Eur J Prev Cardiol* 2019, 26:578-88.
- [48] Munzberg H, Myers MG, Jr.: Molecular and anatomical determinants of central leptin resistance. *Nat Neurosci* 2005, 8:566-70.
- [49] Crujeiras AB, Carreira MC, Cabia B, Andrade S, Amil M, Casanueva FF: Leptin resistance in obesity: An epigenetic landscape. *Life Sci* 2015, 140:57-63.
- [50] Levenson AE, Haas ME, Miao J, Brown RJ, de Ferranti SD, Muniyappa R, Biddinger SB: Effect of Leptin Replacement on PCSK9 in ob/ob Mice and Female Lipodystrophic Patients. *Endocrinology* 2016, 157:1421-9.
- [51] Li S, Xu RX, Zhang Y, Guo YL, Zhu CG, Liu G, Dong Q, Li JJ: Relation of resistin to proprotein convertase subtilisin-kexin type 9 levels in coronary artery disease patients with different nutritional status. *J Endocrinol Invest* 2015, 38:1291-9.
- [52] Jang HD, Lee SE, Yang J, Lee HC, Shin D, Lee H, Lee J, Jin S, Kim S, Lee SJ, You J, Park HW, Nam KY, Lee SH, Park SW, Kim JS, Kim SY, Kwon YW, Kwak SH, Yang HM, Kim HS: Cyclase-associated protein 1 is a binding partner of proprotein convertase subtilisin/kexin type-9 and is required for the degradation of low-density lipoprotein receptors by proprotein convertase subtilisin/kexin type-9. *Eur Heart J* 2020, 41:239-52.
- [53] Ricci C, Ruscica M, Camera M, Rossetti L, Macchi C, Colciago A, Zanotti I, Lupo MG, Adorni MP, Cicero AFG, Fogacci F, Corsini A, Ferri N: PCSK9 induces a pro-inflammatory response in macrophages. *Sci Rep* 2018, 8:2267.
- [54] Heinzl MW, Resl M, Klammer C, Egger M, Dieplinger B, Clodi M: Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Is Not Induced in Artificial Human Inflammation and Is Not Correlated with Inflammatory Response. *Infect Immun* 2020, 88.

Figure legend

Figure 1. HepG2 cells were seeded in DMEM/10% FBS. Cells were incubated with leptin (100 ng/mL) or resistin (100 ng/mL) in DMEM supplemented with 10%LPDS for 24 hours. Simvastatin (20 μ M) has been used as a positive control. At the end of the incubation, total RNA and proteins were extracted. β -actin was used as a housekeeping for both gene (*ACTB*) and protein expression. Panels A and D represent PCSK9 gene expression (qPCR); panels B and E represent PCSK9 protein expression (WB); panels C and F represent PCSK9 levels assayed by ELISA (data were normalized for the total amount of proteins). The values above the blot represent the densitometric readings. Differences between groups have been assessed by Student's t-test or one-way ANOVA; * $p < 0.05$, ** $p < 0.01$. CTR, medium alone. At least three independent experiments were conducted. qPCR, quantitative real-time PCR; WB, Western blot; *ACTB*, β -actin gene; ELISA, enzyme-linked immunosorbent assay; LPDS, human lipoprotein plasma deprived serum.

Figure 2. Panel A) HepG2 cells were seeded in DMEM/10% FBS. After 48h, the medium was replaced with DMEM with 10%LPDS supplemented with leptin (100 ng/mL) and resistin (100 ng/mL). After 24 h, the expression of pSTAT3^{Y705} and STAT3 was evaluated. β -actin and vinculin were used as housekeeping proteins. Panel B-C) HepG2 cells were seeded in DMEM/10% FBS and the day after transfected with siRNA scramble and siRNA anti-STAT3 or siRNA anti-PCSK9. After 48h, the expression of STAT3 was evaluated. β -actin and tubulin were used as housekeeping to normalize the data. Panel D-E) HepG2 transfected with siRNA scramble and siRNA anti-STAT3 were or were not treated with leptin (100 ng/mL) and resistin (100 ng/mL) and the gene expression of PCSK9 was evaluated by qPCR. Actin was used as a housekeeping gene. Panel F) HepG2 transfected with siRNA scramble and siRNA anti-STAT3 were or were not treated with leptin (100 ng/mL) and resistin (100 ng/mL) and the protein expression of PCSK9 was evaluated by Western blot. Vinculin was used as housekeeping to normalize the data. In panels A, C and F (WB), the values above the blots represent the densitometric readings evaluated by using the ImageLab software. Differences between groups were assessed by Student's t-test and one-way ANOVA; * $p < 0.05$, ** $p < 0.01$, † $p < 0.05$ refers to Leptin treatment vs Leptin treatment plus siSTAT3. CTR, medium alone. At least three independent experiments were conducted. qPCR, quantitative real-time PCR; WB, Western blot.

Figure 3. Panel A) Sequence comparison of proximal regions of PCSK9 promoter of human, gorilla, mouse, and rat species. Panel B) Schematic representation of the human PCSK9 promoter-luciferase reporter plasmid D1. Positions are referred to the nucleotide preceding the ATG start codon of human PCSK9 gene, arbitrarily designated as - 1. Positions - 94 and - 1,711 indicate the 3' and the 5' ends of PCSK9 promoter insert, respectively. Positions of the 5' and 3' ends of the STAT3 binding sequence are indicated. Panel C-D) HepG2 cells were transfected with pGL3-PCSK9-D1 and pGL3-PCSK9-D1-STAT3-mutated. The day after the transfection, the medium was replaced with DMEM containing 10%LPDS supplemented with leptin (100 ng/mL), resistin (100 ng/mL) or simvastatin (20 μ M). After an additional 24 hours, luciferase activities have been determined by Neolite reagent. Differences between groups were assessed by Student's t-test or one-way ANOVA; * $p < 0.05$, ** $p < 0.01$. CTR, medium alone. At least three independent experiments were conducted. Numbers inside the bars represent the percentage of increment. Nucleotides in red refer to STAT3 sequence in PCSK9 promoter region; Nucleotides in blue refer to Sp1 sequence in PCSK9 promoter region; Nucleotides in green refer to HNF-1 sequence in PCSK9 promoter region; Nucleotides in purple refer to SRE sequence in PCSK9 promoter region; ** refer to bp spanning between the inter-region sequences of PCSK9 promoter region. STAT3, signal transducer and activator of transcript 3; Sp1, Proximal specificity protein 1; HNF1, hepatocyte nuclear factor-1; SRE, sterol regulatory element.

Figure 4. HepG2 cells were transfected with pGL3-PCSK9-D4 (*panel A*), pGL3-PCSK9-SREmut (*panel B*) and pGL3-PCSK9-HNFmut (*panels C*). The day after the transfection, the medium was replaced with DMEM containing 10%LPDS supplemented with leptin (100 ng/mL), resistin (100 ng/mL) or simvastatin (20 μ M). After an additional 24 hours, luciferase activities have been determined by Neolite reagent. Differences between groups were assessed by Student's t-test; * $p < 0.05$. CTR, medium alone. At least three independent experiments were conducted.

Figure 5. Plasma levels of leptin and PCSK9 were measured by commercial ELISA assays. Subjects have been stratified according to BMI, i.e. $< 25 \text{ Kg/m}^2$ ($n=48$; *panel A*), between 25 and 30 Kg/m^2 ($n= 53$; *panel B*) and $> 30 \text{ Kg/m}^2$ ($n= 48$; *panel C*).

Table 1. List and sequence of primers used in this study

Primers	Forward	Reverse
<i>PCSK9</i>	5'-CCTGCGCGTGTCAACT-3'	5'-GCTGGCTTTTCCGAAACTC-3'
<i>STAT3</i>	5'-CAGCAGCTTGACACACGGTA -3'	5'-AAACACCAAAGTGGCATGT-3'
<i>ACTB</i>	5' TTCTACAATGAGCTGCGTGTG-3'	5' GGGGTGTTGAAGGTCTCAAA-3'
<i>LEPR</i>	5'-TACTTTGGAAGCCCCTGATG-3'	5' AAGCACTGAGTGACTGCACG-3'
<i>CAP1</i>	5'-ACTGGCCTGGAGCAAAACG-3'	5'-CGGCAGAGGGTCCAGATG-3'
<i>IL6</i>	5'-ACCCCCAGGAGAAGATTCCA-3'	5' GGTTGTTTTCTGCCAGTGCC-3'

PCSK9, proprotein convertase subtilisin/kexin type 9; STAT3, Signal Transducer and Activator of Transcript 3; ACTB, β -actin; LEPR, leptin receptor; CAP1, adenylyl cyclase associated protein 1; IL6, interleukin 6

Table 2. *In silico* search of STAT3 binding sites on proximal PCSK9 promoter region.

<i>Predicted Site Sequence</i>	<i>Start</i>	<i>End</i>	<i>Strand</i>	<i>JASPAR Score (%)</i>	<i>FIMO Threshold (p-value)</i>
CTTCTGGAAAG	-916	-906	+	95.92	2.07×10^{-5}
CTTCCAGAAAG	-1082	-1072	+	93.06	7.44×10^{-4}
CTTTCTGGAAG	-1072	-1082	-	92.42	1.01×10^{-4}
ATACTGGGAAG	-1091	-1101	-	89.88	2.93×10^{-4}
TTGCCTGTAAT	-1385	-1395	-	88.45	4.46×10^{-4}
CAGCAGGGAAA	-1485	-1475	+	87.79	4.61×10^{-4}
GTTCAAGAAAT	-950	-960	-	87.27	7.13×10^{-4}
GTGAAGGGAAA	-2184	-2174	+	87.08	6.22×10^{-4}

PCSK9, proprotein convertase subtilisin kexin type 9; STAT3, Signal Transducer and Activator of Transcript 3.