

Article

Investigation of Tick-Borne Pathogens in *Ixodes ricinus* in a Peri-Urban Park in Lombardy (Italy) Reveals the Presence of Emerging Pathogens

Alessandra Cafiso ^{1,*}, Emanuela Olivieri ², Anna Maria Floriano ², Giulia Chiappa ¹, Valentina Serra ¹, Davide Sassera ² and Chiara Bazzocchi ^{1,3}

¹ Department of Veterinary Medicine, University of Milan, Via dell'Università 6, 26900 Lodi, Italy; giulia.chiappa@unimi.it (G.C.); valentina.serra@unimi.it (V.S.); chiara.bazzocchi@unimi.it (C.B.)

² Department of Biology and Biotechnology, University of Pavia, Via Ferrata 9, 27100 Pavia, Italy; emanuelaolivieri84@gmail.com (E.O.); annamaria.floriano01@universitadipavia.it (A.M.F.); davide.sassera@unipv.it (D.S.)

³ Coordinated Research Center "EpiSoMI", University of Milan, 20133 Milan, Italy

* Correspondence: alessandra.cafiso@unimi.it; Tel.: +39-0250334533



Citation: Cafiso, A.; Olivieri, E.; Floriano, A.M.; Chiappa, G.; Serra, V.; Sassera, D.; Bazzocchi, C. Investigation of Tick-Borne Pathogens in *Ixodes ricinus* in a Peri-Urban Park in Lombardy (Italy) Reveals the Presence of Emerging Pathogens. *Pathogens* **2021**, *10*, 732. <https://doi.org/10.3390/pathogens10060732>

Academic Editors: Sergio E. Bermúdez, Victor Montenegro H., José Reck and Abdul Ghafar

Received: 14 May 2021

Accepted: 7 June 2021

Published: 10 June 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Ticks are important vectors of a great range of pathogens of medical and veterinary importance. Lately, the spread of known tick-borne pathogens has been expanding, and novel ones have been identified as (re)emerging health threats. Updating the current knowledge on tick-borne pathogens in areas where humans and animals can be easily exposed to ticks represents a starting point for epidemiological studies and public awareness. A PCR screening for tick-borne pathogens was carried out in *Ixodes ricinus* ticks collected in a peri-urban recreational park in Ticino Valley, Italy. The presence of *Rickettsia* spp., *Borrelia burgdorferi* sensu lato complex, *Anaplasma* spp. and *Babesia* spp. was evaluated in a total of 415 *I. ricinus* specimens. *Rickettsia* spp. (*R. monacensis* and *R. helvetica*) were detected in 22.96% of the samples, while *B. burgdorferi* s.l. complex (*B. afzelii* and *B. lusitaniae*) were present in 10.94%. *Neohrlichia mikurensis* (1.99%) and *Babesia venatorum* (0.73%) were reported in the area of study for the first time. This study confirmed the presence of endemic tick-borne pathogens and highlighted the presence of emerging pathogens that should be monitored especially in relation to fragile patients, the difficult diagnosis of tick-borne associated diseases and possible interactions with other tick-borne pathogens.

Keywords: tick-borne diseases; *Neohrlichia mikurensis*; Ehrlichia-like; Schotti variant; *Babesia venatorum*; EU1; *Babesia capreoli*; *Borrelia burgdorferi* s.l.; tick bite; Northern Italy

1. Introduction

Ticks are hematophagous arthropods considered, along with mosquitoes, the main vectors of important infectious diseases in humans, livestock and domestic animals worldwide [1]. In recent years, the spread of tick-borne-associated microorganisms and pathogens have been expanding [2,3]. The use of molecular biology and genomic analyses have indeed allowed the discovery of new microbial species, strains or genetic variants, increasing the number of potential health-threatening microorganisms associated with ticks (e.g., *Candidatus Neohrlichia mikurensis*—hereafter, *N. mikurensis*—and *Babesia venatorum* [3–10]). In the context of increasing numbers of immunocompromised patients, due to the extended use of novel therapeutic approaches involving extensive immunosuppression [11], the risk of tick-borne diseases (TBDs) needs even more awareness. Symptomatic infections from *N. mikurensis* can be triggered in immunosuppressed patients, with symptoms resembling diseases such as Lyme disease (LD) [12]. Additionally, tick-borne infections through blood transfusions and organ transplantations represent a challenging issue, one that warrants the better implementation of tick-borne pathogens (TBPs) surveillance [13].

Italy is endemic for a considerable number of TBPs, and *Ixodes ricinus*, one of the most common tick species in this country, is responsible for the transmission of over 90% of the TBDs occurring in the area and in Europe [14]. Due to its triphasic behavior and low host specificity, *I. ricinus* is considered one of the primary vectors of multiple pathogens that affect human and animal health [15,16], including *Borrelia burgdorferi* sensu lato complex genospecies, spotted fever group (SFG) rickettsiae, *Anaplasma phagocytophilum*, *N. mikurensis* and *Babesia* spp. [17–22]. The distribution of *I. ricinus* in Europe has expanded during the last decades as a result of multipartite interactions between global warming, anthropogenically induced factors and changes in forest and wildlife management that, in turn, affect the available habitats for ticks [23,24]. Peri-urban recreational areas, where populations of large mammals (e.g., roe deer and wild boar) are increasing, have also become a crucial meeting point between humans, pets and ticks and currently represent a risk for the shifting of the natural transmission cycles of some TBPs [25]. Furthermore, the expansion of recreational outdoor activities has greatly increased the incidence of tick bites, leading to a higher risk of infection with TBPs [26]. Thus, the evaluation of TBPs distribution needs a constant update to maintain the awareness of tick-related diseases. The first step for TBDs surveillance should consist of assessing the diversity of pathogens occurring in a given area and their relative epidemiological importance [27]. In addition, information on coinfections in ticks is pivotal, since different combinations of pathogens in humans and animals are likely to lead to different symptoms, the varying severity of the outgoing disease and to have a negative impact on the diagnosis [28]. In the coming years, TBDs are expected to become one of the major concerns for public health in Europe [27], and the surveillance of TBPs in certain areas has become essential for epidemiological studies and risk assessment for both humans and animals.

The aim of this study was to update and evaluate the presence of both endemic and emerging TBPs of significant public health importance occurring in a Northern Italy peri-urban park and nature reserve characterized by a growing number of human outdoor activities.

2. Results

A total of 415 ticks were collected and all identified as *I. ricinus*. Two hundred and ninety-seven out of 415 collected specimens were nymphs (grouped in 50 pools and 47 single specimens), while the remaining 118 were adults (58 females and 60 males), for a total of 215 DNA samples. Overall, 46.98% (101/215) of the screened DNA samples were positive for at least one pathogen.

The most prevalent TBP was *Rickettsia* spp., with 22.96% positive samples (78/215; CI: 18.61–27.76). Thirty out of 78 *Rickettsia* spp. *gltA* gene amplicons were sequenced to determine the bacterial species. Of these, 29 amplicons (accession MZ068233) showed 100% identity with *Rickettsia monacensis*, while one sample (accession MZ068234) showed 100% identity with *Rickettsia helvetica* (Supplementary Figure S1a).

The second-most common pathogen was *B. burgdorferi* s.l. complex, which showed an overall prevalence of 10.94% (42/215; CI: 8.07–14.34). Twenty out of 42 *groEL* amplicons were sequenced, resulting in a total of five *Borrelia afzelii* (accession MZ068235 and MZ068236, 98.35–100% identity with the sequences deposited in GenBank) and 15 *Borrelia lusitaniae* samples (accession MZ068237, 100% identity with the *B. lusitaniae* sequences deposited in GenBank; Supplementary Figure S1b).

Anaplasma spp. were detected in 1.99% of the samples screened by performing the related PCR protocol (8/215; CI: 0.91–3.67). After sequencing, all PCR products (accession MZ049694) revealed 100% identity with *N. mikurensis* (Supplementary Figure S1c).

Babesia spp. were detected in 0.98% of the samples (4/215; CI: 0.30–2.26), and all the obtained amplicons were subjected to sequencing. Three samples (accession MZ049960) showed 100% identity with *B. venatorum* (0.73% prevalence; CI: 0.18–1.88), and one sample (accession MZ050063) showed 100% identity with *Babesia capreoli* (0.24% prevalence, CI: 0.01–1.06; Supplementary Figure S1d).

In 118 adults and 47 single nymphs, the total observed rate of coinfection was 7.27% (12/165). The coinfection prevalence was 5.93% (7/118) in adult specimens and 10.64% (5/47) in single nymph samples. In detail, 11 of the 12 samples were coinfecting with SFG *Rickettsia* spp. and *B. burgdorferi* s.l. complex genospecies, while, in a nymph, a triple infection of *B. lusitaniae*, *R. monacensis* and *N. mikurensis* was observed.

3. Discussion

The increase of recreational outdoor activities and land usage in the last decades, as well as conservational and restocking programs for wild ungulates in nature reserves [4], have led to an increased risk for people and domestic animals to get tick bites. In this situation, a continuous monitoring on TBPs is pivotal in the context of a One Health approach. Despite Northern Italy encompassing the areas classified as endemic for important TBPs, including the *B. burgdorferi* s.l. complex [29], only fragmented investigations have been performed throughout the years [29–32].

The results of this study confirm the presence of endemic TBPs vectored by *I. ricinus* and also highlight the presence of emerging, previously undetected pathogens of public health concern in a Northern Italy peri-urban park close to the great urbanized area of the city of Milan and nature reserve characterized by human outdoor activities.

The study area, close to the Ticino River, represents the optimal environment for *I. ricinus*, with a well-established biocenosis of small and large mammals, birds and reptiles. Populations of large animals such as roe deer have recently become more abundant, thanks to repopulation programs in the 1990s [33,34]. All these features represent favorable factors for the maintenance of the *I. ricinus* complete life cycle and, consequently, of the vectored pathogens.

We detected *Rickettsia* spp. at a higher prevalence (23%) compared to the other studies performed in Northern Italy on field-collected *I. ricinus*. The prevalence in previous reports of *Rickettsia* spp. in host-seeking ticks in Italy ranged from 1.6% and 19.23% [35]. The *Rickettsia* prevalence could be influenced by several factors, including the seasonality or year of sampling, environment, tick hosts and differences in the sensitivity of the PCR protocols [36]. The sequencing indicated *R. monacensis* as the most common species, while only one sequenced amplicon belonged to *R. helvetica*. This is consistent with the geographical distribution of the two species throughout Europe, with *R. monacensis* being more common in *I. ricinus* populations in Southern Europe [37]. *Rickettsia monacensis* is an emerging pathogen that has been shown to cause a Mediterranean spotted fever-like illness in humans in different European countries, including Italy [38]. Although the bacterium has also been detected in domestic animals, such as dogs and cats [39], no cases of clinical illness in such hosts have been reported [40]; thus, systematic approaches to address the pathogenicity of this infectious bacterial species in nonhuman patients should also be undertaken [41]. The other detected *Rickettsia* species, *R. helvetica*, was previously considered to be nonpathogenic, but it was subsequently included among the SFG rickettsiae after being associated with human illness, with infections suspected to have caused perimyocarditis, unexplained febrile illness and sarcoidosis [42]. Additionally, three cases of a mild form of human rickettsiosis were attributed to *R. helvetica* in Northern Italy through serological analyses [43]. To the best of our knowledge, no information was previously available on the presence of *R. monacensis* and *R. helvetica* in questing ticks in this study area. The closest report of these species was a screening performed on the ticks collected from migratory birds, which highlighted the presence of both *R. monacensis* and *R. helvetica* in the province of Como, Lombardy region [32]. The obtained results suggested that the bacteria belonging to this species are not occasional findings brought by migratory animals but are more likely well-established actors in the *I. ricinus*–vertebrate host interaction.

The *Borrelia burgdorferi* s.l. complex showed an overall prevalence of 10.94%, in line with the values previously observed in Northern Italy [29,44]. Despite the presence of *B. afzelii* and *B. lusitaniae* was previously observed in a study performed in the same area,

the obtained results showed *B. lusitaniae* as being the more represented, in contrast to what reported by Pistone and colleagues in 2010 [29]. It must be noted that *B. lusitaniae* is considered the most prevalent genospecies of the *B. burgdorferi* s.l. complex in ticks in Southern Europe and the Mediterranean area [45]. Moreover, lizards are the most likely reservoirs for *B. lusitaniae* [46–48] and are known to be present in the area with high *I. ricinus* infestation rates [33]. Conversely, *B. afzelii* is maintained by rodents [49], which are reservoirs for this *B. burgdorferi* s.l. complex genospecies and are widespread in the study area [50,51]. *Borrelia afzelii* is one the most common *B. burgdorferi* s.l. complex genospecies, together with *B. garinii*, and infects ticks with the highest prevalence rates in the Central European countries [44].

No samples were found positive for *Anaplasma* spp. This result is in contrast with the reports on this bacterial genus performed in Italy [9,20,27,52], although scarce information about this pathogen in the questing *I. ricinus* is available for the Lombardy region. Future studies should be focused on the investigation of *Anaplasma phagocytophilum* in ticks and blood samples recovered from roe deer or wild boar (which have been demonstrated to be natural hosts for this pathogen [20]). *Neohrlichia mikurensis*, another member of the *Anaplasmataceae* family, was found in the area of study. Although *N. mikurensis* has been reported with a prevalence between 1% and over 20% throughout Europe [53], and previous surveys have highlighted its presence in Northeastern Italy [54], to the best of our knowledge, the present work represents the first detection of this emerging TBP in Northwestern Italy. *Neohrlichia mikurensis* has been raising attention for causing symptoms resembling those of LD, including erythema migrans (EM)-like rashes [12,55], leading to *N. mikurensis* infection misdiagnoses [56]. Since LD is relatively common in Northern Italy [29,57], the cooccurrence of *N. mikurensis* warrants deeper investigations and population's awareness. Furthermore, a *N. mikurensis* infection may show similarities to *Anaplasma* or *Ehrlichia* infections [58], as neohrlichiosis displays clinical signs that may vary in severity but are usually nonspecific, such as a fever, lethargy, myalgia, arthralgia and anorexia [53]. This should also be taken into consideration in light of coinfections in tick vectors, as observed in one of the screened samples. Since no serological test for *N. mikurensis* is currently available [54], molecular screenings of reservoir hosts, particularly small mammals such as rodents of the genera *Apodemus* spp. or *Microtus* spp. [59], could provide important information on the circulation of the pathogen.

Babesia venatorum (formerly known as *Babesia* sp. EU1), whose primary host is considered to be roe deer, was previously identified in ticks from the region of Northern Italy with similar low infection rates as those observed in this study [27,35,52]. Notably, *B. venatorum* can cause clinical manifestations of different severity in immunocompromised or splenectomized humans [5,60]. Nevertheless, it is estimated that *B. venatorum* infections may have been overlooked or misdiagnosed, possibly due to serologic cross-reactivity in laboratory diagnostic tests [61]. *Babesia capreoli*, a species closely related to *Babesia divergens*, is typically found to infect roe deer but seems to be apathogenic for sheep, cattle and humans [62,63]. *Babesia capreoli* was previously reported in the blood and ticks collected from roe deer in Italy [64,65].

Additionally, attention should be paid to coinfections, since hosts coinfecting with different pathogens may show more severe symptoms of diseases [66]. A positive association between *B. burgdorferi* s.l. complex genospecies and *Rickettsia* spp. has been observed in several studies [66,67], possibly resulting in higher replication rates of the two species both in the vector and the vertebrate host [68].

Considering the increasing role of peri-urban areas in recreational activities, the presence of TBPs represents a red flag for animal and human health that should be constantly monitored. Emerging pathogens such as *N. mikurensis* and *B. venatorum* should be kept under surveillance, especially in light of infections in fragile patients [69], and should be considered in cases where the etiological agents of commonly known TBDs are not detected.

4. Materials and Methods

4.1. Collection Site

Ticks were collected in the “La Fagiana” Nature Reserve (Pontevicchio, 45°26′07.1″ N 8°49′46.7″ E), within the central area of Ticino Valley Nature Park (Lombardy region, Northern Italy), 40 km from the city of Milan. The 500-ha peri-urban area, around 134 m.a.s.l., is characterized by forest areas, meadows, oxbow lakes and wetlands and represents the most important public nature reserve in Ticino Valley. The forest is normally used for leisure, educational activities and dog-walking, and represents an important site for wildlife. The fauna consists of small- and medium-sized mammals (e.g., roe deer and wild boar); birds and small reptiles, such as lizards.

4.2. Tick Collection

Host-seeking ticks were collected in 2019 by the unsystematic flagging and dragging of low vegetation during the period of high seasonal activity of *I. ricinus* in Central Europe (between April and October) using a 1 m² woolen blanket. Samplings were performed close to pathways and picnic areas, where humans and domestic animals are more likely to be exposed to ticks during outdoor activities.

Once collected, ticks were grouped according to their sex and developmental stage and preserved in 70% ethanol. Subsequently, ticks were identified on the basis of their morphological features [70], and nymphs were grouped in pools of 1–5 specimens for a total of 97 nymph samples, while the 118 collected adults were treated individually; the specimens were stored at +4 °C until further analyses.

All the specimens (118 adults and 97 nymph samples, for a total of 215 samples) were subsequently processed for DNA extraction and the molecular detection of TBPs.

4.3. Molecular Analyses

DNA was extracted from single adult ticks and pools of nymphs using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) following the manufacturer’s instructions. Extracted DNA samples were quantified and stored at –80 °C for subsequent analyses. Quality of the extracted DNA was assessed by amplifying a fragment of ~360 bp of the 12S rDNA of Ixodidae [71]. The DNA samples were then tested with specific qualitative PCR protocols designed for the amplification of SFG rickettsiae, *Anaplasma* spp., *B. burgdorferi* s.l. and *Babesia* spp. DNA. For *Babesia divergens*/*B. capreoli* species differentiation, a de novo reverse primer was designed to amplify a ~1200-bp portion of the 18S rDNA gene (all target genes, primers and references are reported in Table 1). According to Malandrin et al. [62], the three base differences in the 18S rDNA amplified fragment (positions 631, 663 and 1637, with AAC for *B. divergens* and GTT for *B. capreoli*) can discriminate the two species.

Table 1. List of tick-borne pathogens (TBPs), target genes, PCR primer names and nucleotide sequences and references of the PCR primers used in the TBP screening.

TBP	Target Gene	Primer Name	Nucleotide Sequence (5′-3′)	Reference
SFG rickettsiae	<i>gltA</i>	Rp877p Rp1258n	GGGGACCTGCTCACGGCGG ATTGCAAAAAGTACAGTGAACA	[72]
<i>Anaplasma</i> spp.	16S rRNA	16S8FE B-GA1B_mod	GGAATTCAGAGTTGGATCATGGCTCAG CGGGATCCCCGAGTTTGCCGGGACTT ¹	[73]
<i>B. burgdorferi</i> s.l.	<i>groEL</i>	groEL-F groEL-R	ACGATTCTTATGTTGAGGG TCTCAAGAACTGGTAAAAG	[74]
<i>Babesia</i> spp.	18S rDNA	PIRO-A PIRO-B	AATACCCAATCCTGACACAGGG TAAATACGAATGCCCCCAAC	[75] ²
<i>B. divergens</i> / <i>B. capreoli</i> ³	18S rDNA	PIRO-A Piro-900b	AATACCCAATCCTGACACAGGG AACCTTGTTACGACTTCTCC	[75] This work

¹ Modified by the authors. ² PCR annealing step performed at 62 °C. ³ PCR protocol and conditions: 95 °C for 3 min; 5 cycles at 95 °C for 30 s, 64 °C for 30 s and 72 °C for 60 s; 15 cycles at 95 °C for 30 s, 60 °C for 30 s and 72 °C for 60 s; 20 cycles at 95 °C for 30 s, 54 °C for 30 s and 72 °C for 60 s and final elongation 72 °C for 5 min. Amplicon size ~1200 bp.

The obtained PCR products were excised from agarose gel and purified using the Wizard[®] SV Gel and PCR Clean-Up System Kit (Promega, Madison, WI, USA) according to the manufacturer's protocols and bidirectionally Sanger-sequenced. Sequences were then assembled, manually curated with SeaView 4.7 [76] and compared with the representative sequences available in NCBI GenBank using BLAST. The obtained sequences were deposited in GenBank.

4.4. Phylogenetic and Statistical Analyses

The prevalence of each pathogen was calculated with the estimated pooled prevalence (EPP) with a 95% confidence interval (95% CI) using the online pool prevalence calculator Epitools [77]. The method estimates the prevalence and confidence limits for variable pool sizes and assumes 100% test sensitivity and specificity.

All the phylogenetic inferences were performed as follows: sequences were aligned with MUSCLE v3.8.31 [78], the evolutionary model to be used for phylogenetic inference was chosen according to the AIC (using modeltest-ng [79]) and the phylogeny was inferred using RAxML 8.2.4 (100 bootstraps, -p 123, -x 1234 [80]). The evolutionary models applied for phylogenetic inference were: GTR+I+G for *Babesia* spp. and *Anaplasma* spp./*Neoehrlichia* spp., HKY+I+G for the *B. burgdorferi* s.l. complex genospecies and *Rickettsia* spp.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/pathogens10060732/s1>: Figure S1: Phylogenetic inferences of the obtained gene sequences.

Author Contributions: Conceptualization, A.C. and E.O.; methodology, A.C., C.B. and G.C.; investigation, A.C., E.O., G.C. and V.S.; resources, C.B. and D.S.; data curation, A.C., A.M.F. and E.O.; formal analysis: A.C., A.M.F. and E.O.; writing—original draft preparation, A.C., E.O., C.B., D.S. and A.M.F.; writing—review and editing, A.C., C.B., E.O., D.S., A.M.F., G.C. and V.S.; supervision, C.B., D.S. and A.C. and project administration, A.C. and E.O. All authors have read and agreed to the published version of the manuscript.

Funding: The authors acknowledge support from the University of Milan through the APC initiative.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The obtained gene sequences were deposited in the NCBI GenBank database. The accession numbers of the obtained sequences are contained within the article and in the Supplementary Materials.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Parola, P.; Raoult, D. Ticks and tickborne bacterial diseases in humans: An emerging infectious threat. *Clin. Infect. Dis.* **2001**, *32*, 897–928. [CrossRef]
2. Ogden, N. Changing geographic ranges of ticks and tick-borne pathogens: Drivers, mechanisms and consequences for pathogen diversity. *Front. Cell. Infect. Microbiol.* **2013**, *3*. [CrossRef]
3. Rochlin, I.; Toledo, A. Emerging tick-borne pathogens of public health importance: A mini-review. *J. Med. Microbiol.* **2020**, *69*, 781–791. [CrossRef] [PubMed]
4. Randolph, S.E. Evidence that climate change has caused 'emergence' of tick-borne diseases in Europe? *Int. J. Med. Microbiol. Suppl.* **2004**, *293*, 5–15. [CrossRef]
5. Herwaldt, B.L.; Cacciò, S.; Gherlinzoni, F.; Aspöck, H.; Slemenda, S.B.; Piccaluga, P.; Martinelli, G.; Edelhofer, R.; Hollenstein, U.; Poletti, G.; et al. Molecular characterization of a non-*Babesia divergens* organism causing zoonotic babesiosis in Europe. *Emerg. Infect. Dis.* **2003**, *9*, 943–948. [CrossRef]
6. Häselbarth, K.; Tenter, A.M.; Brade, V.; Krieger, G.; Hunfeld, K.-P. First case of human babesiosis in Germany—Clinical presentation and molecular characterisation of the pathogen. *Int. J. Med. Microbiol.* **2007**, *297*, 197–204. [CrossRef] [PubMed]
7. Sanogo, Y.O.; Parola, P.; Shpynov, S.; Camicas, J.L.; Brouqui, P.; Caruso, G.; Raoult, D. Genetic diversity of bacterial agents detected in ticks removed from asymptomatic patients in northeastern Italy. *Ann. N. Y. Acad. Sci.* **2003**, *990*, 182–190. [CrossRef]
8. von Loewenich, F.D.; Geißdörfer, W.; Disqué, C.; Matten, J.; Schett, G.; Sakka, S.G.; Bogdan, C. Detection of "*Candidatus Neoehrlichia mikurensis*" in two patients with severe febrile illnesses: Evidence for a European sequence variant. *J. Clin. Microbiol.* **2010**, *48*, 2630–2635. [CrossRef] [PubMed]

9. Otranto, D.; Dantas-Torres, F.; Giannelli, A.; Latrofa, M.S.; Cascio, A.; Cazzin, S.; Ravagnan, S.; Montarsi, F.; Zanzani, S.A.; Manfredi, M.T.; et al. Ticks infesting humans in Italy and associated pathogens. *Parasites Vectors* **2014**, *7*, 328. [[CrossRef](#)] [[PubMed](#)]
10. Quarsten, H.; Grankvist, A.; Høyvoll, L.; Myre, I.B.; Skarpaas, T.; Kjelland, V.; Wenneras, C.; Noraas, S. *Candidatus* Neoehrlichia mikurensis and *Borrelia burgdorferi* sensu lato detected in the blood of Norwegian patients with erythema migrans. *Ticks Tick Borne Dis.* **2017**, *8*, 715–720. [[CrossRef](#)]
11. Bender, D.A.; Heilbroner, S.P.; Wang, T.J.C.; Shu, C.A.; Hyde, B.; Spina, C.; Cheng, S.K. Increased rates of immunosuppressive treatment and hospitalization after checkpoint inhibitor therapy in cancer patients with autoimmune disease. *J. Immunother. Cancer* **2020**, *8*, e001627. [[CrossRef](#)]
12. Ondruš, J.; Balážová, A.; Baláž, V.; Zechmeisterová, K.; Novobilský, A.; Široký, P. *Candidatus* Neoehrlichia mikurensis is widespread in questing *Ixodes ricinus* ticks in the Czech Republic. *Ticks Tick Borne Dis.* **2020**, *11*, 101371. [[CrossRef](#)]
13. Mohan, K.V.K.; Leiby, D.A. Emerging tick-borne diseases and blood safety: Summary of a public workshop. *Transfusion* **2020**, *60*, 1624–1632. [[CrossRef](#)]
14. Süss, J.; Schrader, C.; Falk, U.; Wohanka, N. Tick-borne encephalitis (TBE) in Germany—Epidemiological data, development of risk areas and virus prevalence in field-collected ticks and in ticks removed from humans. *Int. J. Med. Microbiol. Suppl.* **2004**, *293*, 69–79. [[CrossRef](#)]
15. Anderson, J.F. Epizootiology of Lyme borreliosis. *Scand. J. Infect. Dis. Suppl.* **1991**, *77*, 23–34. [[PubMed](#)]
16. Mihalca, A.D.; Gherman, C.M.; Magdaş, C.; Dumitrache, M.O.; Györke, A.; Sándor, A.D.; Domşa, C.; Oltean, M.; Mircean, V.; Mărcuţan, D.I.; et al. *Ixodes ricinus* is the dominant questing tick in forest habitats in Romania: The results from a countrywide dragging campaign. *Exp. Appl. Acarol.* **2012**, *58*, 175–182. [[CrossRef](#)] [[PubMed](#)]
17. Nilsson, K.; Jaenson, T.G.; Uhnöo, I.; Lindquist, O.; Pettersson, B.; Uhlén, M.; Friman, G.; Pålsson, C. Characterization of a spotted fever group rickettsia from *Ixodes ricinus* ticks in Sweden. *J. Clin. Microbiol.* **1997**, *35*, 243–247. [[CrossRef](#)]
18. Simser, J.A.; Palmer, A.T.; Fingerle, V.; Wilske, B.; Kurtti, T.J.; Munderloh, U.G. *Rickettsia monacensis* sp. nov., a spotted fever group rickettsia, from ticks (*Ixodes ricinus*) collected in a European city park. *Appl. Environ. Microbiol.* **2002**, *68*, 4559–4566. [[CrossRef](#)]
19. Stuenkel, S. *Anaplasma phagocytophilum*—the most widespread tick-borne infection in animals in Europe. *Vet. Res. Commun.* **2007**, *31*, 79–84. [[CrossRef](#)]
20. Carpi, G.; Bertolotti, L.; Pecchioli, E.; Cagnacci, F.; Rizzoli, A. *Anaplasma phagocytophilum* groEL gene heterogeneity in *Ixodes ricinus* larvae feeding on roe deer in northeastern Italy. *Vector Borne Zoonotic Dis.* **2008**, *9*, 179–184. [[CrossRef](#)] [[PubMed](#)]
21. Duh, D.; Saksida, A.; Petrovec, M.; Dedushaj, I.; Avšič-Županc, T. Novel one-step real-time RT-PCR assay for rapid and specific diagnosis of Crimean-Congo hemorrhagic fever encountered in the Balkans. *J. Virol. Methods* **2006**, *133*, 175–179. [[CrossRef](#)]
22. Vayssier-Taussat, M.; Moutailler, S.; Michelet, L.; Devillers, E.; Bonnet, S.; Cheval, J.; Hébert, C.; Eloit, M. Next generation sequencing uncovers unexpected bacterial pathogens in ticks in western Europe. *PLoS ONE* **2013**, *8*, e81439. [[CrossRef](#)]
23. Medlock, J.M.; Hansford, K.M.; Bormane, A.; Derdakova, M.; Estrada-Peña, A.; George, J.-C.; Golovljova, I.; Jaenson, T.G.; Jensen, J.-K.; Jensen, P.M.; et al. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasites Vectors* **2013**, *6*, 1. [[CrossRef](#)]
24. Sréter-Lancz, Z.; Sréter, T.; Széll, Z.; Egyed, L. Molecular evidence of *Rickettsia helvetica* and *R. monacensis* infections in *Ixodes ricinus* from Hungary. *Ann. Trop. Med. Parasitol.* **2005**, *99*, 325–330. [[CrossRef](#)]
25. Rizzoli, A.; Silaghi, C.; Obiegala, A.; Rudolf, I.; Hubálek, Z.; Földvári, G.; Plantard, O.; Vayssier-Taussat, M.; Bonnet, S.; Spitalská, E.; et al. *Ixodes ricinus* and its transmitted pathogens in urban and peri-urban areas in Europe: New hazards and relevance for public health. *Front. Public Health* **2014**, *2*, 251. [[CrossRef](#)]
26. Maioli, G.; Pistone, D.; Bonilauri, P.; Pajoro, M.; Barbieri, I.; Patrizia, M.; Vicari, N.; Dottori, M. Ethiological agents of rickettsiosis and anaplasmosis in ticks collected in Emilia-Romagna region (Italy) during 2008 and 2009. *Exp. Appl. Acarol.* **2012**, *57*, 199–208. [[CrossRef](#)]
27. Capelli, G.; Ravagnan, S.; Montarsi, F.; Ciocchetta, S.; Cazzin, S.; Porcellato, E.; Babiker, A.M.; Cassini, R.; Salviato, A.; Cattoli, G.; et al. Occurrence and identification of risk areas of *Ixodes ricinus*-borne pathogens: A cost-effectiveness analysis in north-eastern Italy. *Parasites Vectors* **2012**, *5*, 61. [[CrossRef](#)]
28. Lejal, E.; Moutailler, S.; Šimo, L.; Vayssier-Taussat, M.; Pollet, T. Tick-borne pathogen detection in midgut and salivary glands of adult *Ixodes ricinus*. *Parasites Vectors* **2019**, *12*, 152. [[CrossRef](#)]
29. Pistone, D.; Pajoro, M.; Fabbi, M.; Vicari, N.; Marone, P.; Genchi, C.; Novati, S.; Sasser, D.; Epis, S.; Bandi, C. Lyme borreliosis, Po river valley, Italy. *Emerg. Infect. Dis.* **2010**, *16*, 1289. [[CrossRef](#)]
30. De Meneghi, D. Wildlife, Environment and (re)-emerging zoonoses, with special reference to sylvatic tick-borne zoonoses in north-western Italy. *Ann. Dell'istituto Super. Sanita* **2006**, *42*, 405–409.
31. Pintore, M.D.; Ceballos, L.; Iulini, B.; Tomassone, L.; Pautasso, A.; Corbellini, D.; Rizzo, F.; Mandola, M.L.; Bardelli, M.; Peletto, S.; et al. Detection of invasive *Borrelia burgdorferi* strains in north-eastern Piedmont, Italy. *Zoonoses Public Health* **2015**, *62*, 365–374. [[CrossRef](#)]
32. Pajoro, M.; Pistone, D.; Varotto Boccazzi, I.; Mereghetti, V.; Bandi, C.; Fabbi, M.; Scattorin, F.; Sasser, D.; Montagna, M. Molecular screening for bacterial pathogens in ticks (*Ixodes ricinus*) collected on migratory birds captured in northern Italy. *Folia Parasitol.* **2018**. [[CrossRef](#)]
33. Scali, S.; Manfredi, M.T.; Guidali, E. *Lacerta bilineata* (Reptilia, Lacertidae) as a host of *Ixodes ricinus* (Acari, Ixodidae) in a protected area of northern Italy. *Parassitologia* **2001**, *43*, 165–168.

34. De Pasquale, D.; Dondina, O.; Scancarello, E.; Meriggi, A. Long-term viability of a reintroduced population of roe deer *Capreolus capreolus*, in a lowland area of northern Italy. *Folia Zool.* **2019**, *68*, 9–20. [[CrossRef](#)]
35. Castro, L.R.; Gabrielli, S.; Iori, A.; Cancrini, G. Molecular detection of *Rickettsia*, *Borrelia*, and *Babesia* species in *Ixodes ricinus* sampled in northeastern, central, and insular areas of Italy. *Exp. Appl. Acarol.* **2015**, *66*, 443–452. [[CrossRef](#)]
36. Estrada-Peña, A.; de la Fuente, J. The Ecology of ticks and epidemiology of tick-borne viral diseases. *Antivir. Res.* **2014**, *108*, 104–128. [[CrossRef](#)]
37. Silaghi, C.; Gilles, J.; Höhle, M.; Pradel, I.; Just, F.T.; Fingerle, V.; Küchenhoff, H.; Pfister, K. Prevalence of spotted fever group rickettsiae in *Ixodes ricinus* (Acari: Ixodidae) in southern Germany. *J. Med. Entomol.* **2008**, *45*, 948–955. [[CrossRef](#)]
38. Madeddu, G.; Mancini, F.; Caddeo, A.; Ciervo, A.; Babudieri, S.; Maida, I.; Fiori, M.L.; Rezza, G.; Mura, M.S. *Rickettsia monacensis* as cause of Mediterranean Spotted Fever-like illness, Italy. *Emerg. Infect. Dis.* **2012**, *18*, 702–704. [[CrossRef](#)]
39. Lauzi, S.; Maia, J.P.; Epis, S.; Marcos, R.; Pereira, C.; Luzzago, C.; Santos, M.; Puente-Payo, P.; Giordano, A.; Pajoro, M.; et al. Molecular detection of *Anaplasma platys*, *Ehrlichia canis*, *Hepatozoon canis* and *Rickettsia monacensis* in dogs from Maio Island of Cape Verde archipelago. *Ticks Tick Borne Dis.* **2016**, *7*, 964–969. [[CrossRef](#)]
40. Morganti, G.; Gavaudan, S.; Canonico, C.; Ravagnan, S.; Olivieri, E.; Diaferia, M.; Marenzoni, M.L.; Antognoni, M.T.; Capelli, G.; Silaghi, C.; et al. Molecular survey on *Rickettsia* spp., *Anaplasma phagocytophilum*, *Borrelia burgdorferi* sensu sato, and *Babesia* spp. in *Ixodes ricinus* ticks infesting dogs in central Italy. *Vector Borne Zoonotic Dis.* **2017**, *17*, 743–748. [[CrossRef](#)]
41. Schreiber, C.; Krücken, J.; Beck, S.; Maaz, D.; Pachnick, S.; Krieger, K.; Gross, M.; Kohn, B.; von Samson-Himmelstjerna, G. Pathogens in ticks collected from dogs in Berlin/Brandenburg, Germany. *Parasites Vectors* **2014**, *7*, 535. [[CrossRef](#)]
42. Nilsson, K.; Liu, A.; Pålsson, C.; Lindquist, O. Demonstration of intracellular microorganisms (*Rickettsia* spp., *Chlamydia pneumoniae*, *Bartonella* spp.) in pathological human aortic valves by PCR. *J. Infect.* **2005**, *50*, 46–52. [[CrossRef](#)]
43. Fournier, P.-E.; Allombert, C.; Supputamongkol, Y.; Caruso, G.; Brouqui, P.; Raoult, D. Aneruptive fever associated with antibodies to *Rickettsia helvetica* in Europe and Thailand. *J. Clin. Microbiol.* **2004**, *42*, 816–818. [[CrossRef](#)]
44. Rauter, C.; Hartung, T. Prevalence of *Borrelia burgdorferi* sensu lato genospecies in *Ixodes ricinus* ticks in Europe: A metaanalysis. *Appl. Environ. Microbiol.* **2005**, *71*, 7203–7216. [[CrossRef](#)]
45. Grego, E.; Bertolotti, L.; Peletto, S.; Amore, G.; Tomassone, L.; Mannelli, A. *Borrelia lusitaniae* *OspA* gene heterogeneity in Mediterranean basin area. *J. Mol. Evol.* **2007**, *65*, 512–518. [[CrossRef](#)] [[PubMed](#)]
46. Younsi, H.; Sarih, M.; Jouda, F.; Godfroid, E.; Gern, L.; Bouattour, A.; Baranton, G.; Postic, D. Characterization of *Borrelia lusitaniae* isolates collected in Tunisia and Morocco. *J. Clin. Microbiol.* **2005**, *43*, 1587–1593. [[CrossRef](#)]
47. Amore, G.; Tomassone, L.; Grego, E.; Ragagli, C.; Bertolotti, L.; Nebbia, P.; Rosati, S.; Mannelli, A. *Borrelia lusitaniae* in immature *Ixodes ricinus* (Acari: Ixodidae) feeding on common wall lizards in Tuscany, central Italy. *J. Med. Entomol.* **2007**, *44*, 303–307. [[CrossRef](#)] [[PubMed](#)]
48. Majláthová, V.; Majláth, I.; Hromada, M.; Tryjanowski, P.; Bona, M.; Antczak, M.; Víchová, B.; Dzimko, Š.; Mihalca, A.; Peťko, B. The role of the sand lizard (*Lacerta agilis*) in the transmission cycle of *Borrelia burgdorferi* sensu lato. *Int. J. Med. Microbiol.* **2008**, *298*, 161–167. [[CrossRef](#)]
49. Ferreri, L.; Perazzo, S.; Venturino, E.; Giacobini, M.; Bertolotti, L.; Mannelli, A. Modeling the effects of variable feeding patterns of larval ticks on the transmission of *Borrelia lusitaniae* and *Borrelia afzelii*. *Theo. Popul. Biol.* **2017**, *116*, 27–32. [[CrossRef](#)]
50. Ragagli, C.; Bertolotti, L.; Giacobini, M.; Mannelli, A.; Bisanzio, D.; Amore, G.; Tomassone, L. Transmission dynamics of *Borrelia lusitaniae* and *Borrelia afzelii* among *Ixodes ricinus*, lizards, and mice in Tuscany, central Italy. *Vector Borne Zoonotic Dis.* **2010**, *11*, 21–28. [[CrossRef](#)]
51. Castioni, C.; Debernardi, P.; Patriarca, E. L'alimentazione invernale nel gufo comune (*Asio otus*) nel Parco del Ticino (Italia nord-occidentale). *Riv. Piemont. Stor. Nat.* **1998**, *19*, 299–312.
52. Aureli, S.; Galuppi, R.; Ostanello, F.; Foley, J.E.; Bonoli, C.; Rejmanek, D.; Rocchi, G.; Orlandi, E.; Tampieri, M.P. Abundance of questing ticks and molecular evidence for pathogens in ticks in three parks of Emilia-Romagna region of northern Italy. *Ann. Agric. Environ. Med.* **2015**, *22*, 459–466. [[CrossRef](#)] [[PubMed](#)]
53. Silaghi, C.; Beck, R.; Oteo, J.A.; Pfeffer, M.; Sprong, H. Neoehrlichiosis: An emerging tick-borne zoonosis caused by *Candidatus Neoehrlichia mikurensis*. *Exp. Appl. Acarol.* **2016**, *68*, 279–297. [[CrossRef](#)] [[PubMed](#)]
54. Portillo, A.; Santibáñez, P.; Palomar, A.M.; Santibáñez, S.; Oteo, J.A. 'Candidatus Neoehrlichia mikurensis' in Europe. *New Microbes New Infect.* **2018**, *22*, 30–36. [[CrossRef](#)]
55. Grankvist, A.; Andersson, P.-O.; Mattsson, M.; Sender, M.; Vaht, K.; Höper, L.; Sakiniene, E.; Trysberg, E.; Stenson, M.; Fehr, J.; et al. Infections with the tick-borne bacterium "Candidatus Neoehrlichia mikurensis" mimic noninfectious conditions in patients with B cell malignancies or autoimmune diseases. *Clin. Infect. Dis.* **2014**, *58*, 1716–1722. [[CrossRef](#)]
56. Fehr, J.S.; Bloemberg, G.V.; Ritter, C.; Hombach, M.; Lüscher, T.F.; Weber, R.; Keller, P.M. Septicemia caused by tick-borne bacterial pathogen *Candidatus Neoehrlichia mikurensis*. *Emerg. Infect. Dis.* **2010**, *16*, 1127–1129. [[CrossRef](#)]
57. Zanzani, S.A.; Rimoldi, S.G.; Manfredi, M.; Grande, R.; Gazzonis, A.L.; Merli, S.; Olivieri, E.; Giacomet, V.; Antinori, S.; Cislighi, G.; et al. Lyme borreliosis incidence in Lombardy, Italy (2000–2015): Spatiotemporal analysis and environmental risk factors. *Ticks Tick Borne Dis.* **2019**, *10*, 101257. [[CrossRef](#)]
58. Wennerås, C. Infections with the tick-borne bacterium *Candidatus Neoehrlichia mikurensis*. *Clin. Microbiol. Infect.* **2015**, *21*, 621–630. [[CrossRef](#)]

59. Svitálková, Z.H.; Haruštiaková, D.; Mahríková, L.; Mojšová, M.; Berthová, L.; Slovák, M.; Kocianová, E.; Vayssier-Taussat, M.; Kazimírová, M. *Candidatus* Neoehrlichia mikurensis in ticks and rodents from urban and natural habitats of south-western Slovakia. *Parasites Vectors* **2016**, *9*, 2. [[CrossRef](#)]
60. Bläckberg, J.; Lazarevic, V.L.; Hunfeld, K.-P.; Persson, K.E.M. Low-Virulent *Babesia venatorum* infection masquerading as hemophagocytic syndrome. *Ann. Hematol.* **2018**, *97*, 731–733. [[CrossRef](#)]
61. Gray, A.; Capewell, P.; Loney, C.; Katzer, F.; Shiels, B.R.; Weir, W. Sheep as host species for zoonotic *Babesia venatorum*, United Kingdom. *Emerg. Infect. Dis.* **2019**, *25*, 2257–2260. [[CrossRef](#)]
62. Malandrín, L.; Jouglín, M.; Sun, Y.; Brisseau, N.; Chauvin, A. Redescription of *Babesia capreoli* (Enigk and Friedhoff, 1962) from roe deer (*Capreolus capreolus*): Isolation, cultivation, host specificity, molecular characterisation and differentiation from *Babesia divergens*. *Int. J. Parasitol.* **2010**, *40*, 277–284. [[CrossRef](#)] [[PubMed](#)]
63. Kogler, S.; Gotthalseder, E.; Shahi-Barogh, B.; Harl, J.; Fuehrer, H.-P. *Babesia* spp. and *Anaplasma phagocytophilum* in free-ranging wild ungulates in central Austria. *Ticks Tick Borne Dis.* **2021**, *12*, 101719. [[CrossRef](#)]
64. Tampieri, M.P.; Galuppi, R.; Bonoli, C.; Cancrini, G.; Moretti, A.; Pietrobelli, M. Wild ungulates as *Babesia* hosts in northern and central Italy. *Vector Borne Zoonotic Dis.* **2008**, *8*, 667–674. [[CrossRef](#)] [[PubMed](#)]
65. Zanet, S.; Trisciuglio, A.; Bottero, E.; de Mera, I.G.F.; Gortazar, C.; Carpignano, M.G.; Ferroglio, E. Piroplasmosis in wildlife: *Babesia* and *Theileria* affecting free-ranging ungulates and carnivores in the Italian Alps. *Parasites Vectors* **2014**, *7*, 70. [[CrossRef](#)] [[PubMed](#)]
66. Václav, R.; Ficová, M.; Prokop, P.; Betáková, T. Associations between coinfection prevalence of *Borrelia lusitaniae*, *Anaplasma* sp., and *Rickettsia* sp. in hard ticks feeding on reptile hosts. *Microb. Ecol.* **2011**, *61*, 245–253. [[CrossRef](#)]
67. Raileanu, C.; Moutailler, S.; Pavel, I.; Porea, D.; Mihalca, A.D.; Savuta, G.; Vayssier-Taussat, M. *Borrelia* diversity and co-infection with other tick borne pathogens in ticks. *Front. Cell. Infect. Microbiol.* **2017**, *7*. [[CrossRef](#)] [[PubMed](#)]
68. Raulf, M.-K.; Jordan, D.; Fingerle, V.; Strube, C. Association of *Borrelia* and *Rickettsia* spp. and bacterial loads in *Ixodes ricinus* ticks. *Ticks Tick Borne Dis.* **2018**, *9*, 18–24. [[CrossRef](#)]
69. Quarsten, H.; Salte, T.; Lorentzen, Å.R.; Hansen, I.J.W.; Hamre, R.; Forselv, K.J.N.; Øines, Ø.; Wennerås, C.; Noraas, S. Tick-borne pathogens detected in the blood of immunosuppressed Norwegian patients living in a tick-endemic area. *Clin. Infect. Dis.* **2020**. [[CrossRef](#)]
70. Manilla, G. *Acari: Ixodida*; Fauna d'Italia; Calderini: Bologna, Italy, 1998; Volume 36.
71. Beati, L.; Keirans, J.E. Analysis of the systematic relationships among ticks of the genera *Rhipicephalus* and *Boophilus* (Acari: Ixodidae) based on mitochondrial 12S ribosomal DNA gene sequences and morphological characters. *J. Parasitol.* **2001**, *87*, 32–48. [[CrossRef](#)]
72. Regnery, R.L.; Spruill, C.L.; Plikaytis, B.D. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J. Bacteriol.* **1991**, *173*, 1576–1589. [[CrossRef](#)]
73. Aktas, M.; Altay, K.; Dumanli, N.; Kalkan, A. Molecular detection and identification of *Ehrlichia* and *Anaplasma* species in ixodid ticks. *Parasitol. Res.* **2009**, *104*, 1243. [[CrossRef](#)]
74. Chiappa, G.; Cafiso, A.; Monza, E.; Serra, V.; Olivieri, E.; Romeo, C.; Bazzocchi, C. Development of a PCR for *Borrelia burgdorferi* sensu lato, targeted on the *groEL* gene. *Folia Parasitol.* **2020**. [[CrossRef](#)] [[PubMed](#)]
75. Olmeda, A.S.; Armstrong, P.M.; Rosenthal, B.M.; Valladares, B.; del Castillo, A.; de Armas, F.; Miguelez, M.; González, A.; Rodríguez Rodríguez, J.A.; Spielman, A.; et al. A subtropical case of human babesiosis. *Acta Trop.* **1997**, *67*, 229–234. [[CrossRef](#)]
76. Gouy, M.; Guindon, S.; Gascuel, O. SeaView Version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **2010**, *27*, 221–224. [[CrossRef](#)] [[PubMed](#)]
77. Sergeant, E.S.G. Epitools Epidemiological Calculators. *Ausvet Pty Ltd.* 2018. Available online: <http://epitools.ausvet.com.au> (accessed on 20 June 2019).
78. Edgar, R.C. MUSCLE: A multiple sequence alignment method with reduced time and space complexity. *BMC Bioinform.* **2004**, *5*, 113. [[CrossRef](#)]
79. Darriba, D.; Posada, D.; Kozlov, A.M.; Stamatakis, A.; Morel, B.; Flouri, T. ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Mol. Biol. Evol.* **2020**, *37*, 291–294. [[CrossRef](#)]
80. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **2014**, *30*, 1312–1313. [[CrossRef](#)]