


The COVID-19 pandemic and its global effects on dental practice. An international survey



G Campus , M Diaz Betancourt , MG Cagetti , RA Giacaman ,
DJ Manton , GVA Douglas , TS Carvalho , JC Carvalho ,
A Vukovic , FJ Cortés-Martinicorena , D Bourgeois ,
V Machiulskiene , R Sava-Rosianu , J Krithikadatta ,
N Sergeevna Morozova , AM Acevedo , AA Agudelo-Suarez ,
G Aguirre , K Aguirre , I Alfonso , HA Ghaffar , RA El Fadl ,
SA Al Maghlouth , T Almerich-Torres , F Amadori , K Antia ,
E Bajric , DM Barbosa-Liz , S Ben-Tanfous , H Bieber , S Bhusari ,
S Birch , G Bontà , P Bottenberg , JJ Bruers , L Bustillos ,
P Bührens , J Cai , JL Cairolì , FCA Carrer , MB Correa ,
B Cortés-Acha , F Carrouel , R de Carvalho Oliveira , F Cocco ,
F Crombie , JI Csikar , D Declerck , M Denkovski , J Deschner ,
J Dopico-San Martín , O Viktorovna Dudnik , W Y Escobar ,
A Elwishahy , CE Fernandez , M Fontana , A Frattaroli Pericchi ,
M Ghorbe , E. Gigineishvili , A Garcia Quintana , J Gray ,
N Gugnani , K Gambetta-Tessini , A Haider , M Hopcraft ,
J Hüttmann , N Hysenaj , A Jalal , M Jikia , J John ,
G Kaps-Richter , T Kerber Tedesco , S Leon , KA Levin ,
H Pau Lew , M Aperecida Moreira Machado ,
A Beneictovna MacLennan , J Onome Mafeni , M Minatel Braga ,
JM Montiel-Company , A Malerba , S Mandić-Rajčević ,
A Askerovich Mamedo , S A Mani , O Marouane , D Markovic ,
E Paredes Martinez , N Maroufidis , F Medeiros Mendes ,
S Musa , A Necibi , N Azlida Mohd Nor , B Tochukwu Ojukwu ,
N Opdam , L Ottolenghi , J Owen , A Pássaro , IF Persoon ,
T Peric , E Pesaressi-Torres , V Philippides , SP Plaza-Ruiz ,
D Procida Raggio , FJ Rivas Cartagena , F Ramos-Gomez ,
M Sabashvili , G Solis Sanchez , R Villena Sarmiento ,
H Schrader , S Serban , R Bairstow , A Senna , B Shi ,
CPC Sim , S Slabsinskiene , G Spagnulo , O Olufemi Taiwo ,
A Thodhorjani , P Tietler , CMC Volgenant , MH van der Veen ,
Z Vlahovic , A Visaria , Y Romero Uzcategui , E Xhajanka ,
Q Yan , O Zen , O Zeyer , A Zukanovic , TG Wolf

PII: S0300-5712(21)00170-6
DOI: <https://doi.org/10.1016/j.jdent.2021.103749>
Reference: JJOD 103749

To appear in: *Journal of Dentistry*

Received date: 14 February 2021
Revised date: 7 July 2021
Accepted date: 9 July 2021

Please cite this article as: G Campus , M Diaz Betancourt , MG Cagetti , RA Giacaman , DJ Manton , GVA Douglas , TS Carvalho , JC Carvalho , A Vukovic , FJ Cortés-Martínicorena , D Bourgeois , V Machiulskiene , R Sava-Rosianu , J Krithikadatta , N Sergeevna Morozova , AM Acevedo , AA Agudelo-Suarez , G Aguirre , K Aguirre , I Alfonso , HA Ghaffar , RA El Fadl , SA Al Maghlouth , T Almerich-Torres , F Amadori , K Antia , E Bajric , DM Barbosa-Liz , S Ben-Tanfous , H Bieber , S Bhusari , S Birch , G Bontà , P Bottenberg , JJ Bruers , L Bustillos , P Bührens , J Cai , JL Cairoli , FCA Carrer , MB Correa , B Cortés-Acha , F Carrouel , R de Carvalho Oliveira , F Cocco , F Crombie , JI Csikar , D Declerck , M Denkovski , J Deschner , J Dopico-San Martín , O Viktorovna Dudnik , W Y Escobar , A Elwishahy , CE Fernandez , M Fontana , A Frattaroli Pericchi , M Ghorbe , E. Gigineishvili , A Garcia Quintana , J Gray , N Gugni , K Gambetta-Tessini , A Haider , M Hopcraft , J Hüttmann , N Hysenaj , A Jalal , M Jikia , J John , G Kaps-Richter , T Kerber Tedesco , S Leon , KA Levin , H Pau Lew , M Aperecida Moreira Machado , A Beneictovna MacLennan , J Onome Mafeni , M Minatel Braga , JM Montiel-Company , A Malerba , S Mandić-Rajčević , A Askerovich Mamedo , S A Mani , O Marouane , D Markovic , E Paredes Martinez , N Maroufidis , F Medeiros Mendes , S Musa , A Necibi , N Azlida Mohd Nor , B Tochukwu Ojukwu , N Opdam , L Ottolenghi , J Owen , A Pássaro , IF Persoon , T Peric , E Pesaressi-Torres , V Philippides , SP Plaza-Ruiz , D Procida Raggio , FJ Rivas Cartagena , F Ramos-Gomez , M Sabashvili , G Solis Sanchez , R Villena Sarmiento , H Schrader , S Serban , R Bairstow , A Senna , B Shi , CPC Sim , S Slabsinskiene , G Spagnulo , O Olufemi Taiwo , A Thodhorjani , P Tietler , CMC Volgenant , MH van der Veen , Z Vlahovic , A Visaria , Y Romero Uzcategui , E Xhajanka , Q Yan , O Zen , O Zeyer , A Zukanovic , TG Wolf , The COVID-19 pandemic and its global effects on dental practice. An international survey, *Journal of Dentistry* (2021), doi: <https://doi.org/10.1016/j.jdent.2021.103749>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The COVID-19 pandemic and its global effects on dental practice. An international survey.

The COVID-19 pandemic and its global effects on dental practice. An international survey.

COVIDental Collaboration Group

Contributing Authors

G Campus (Department of Restorative, Preventive and Paediatric Dentistry, University of Bern, Bern, Switzerland; Department of Surgery, Microsurgery and Medicine Sciences, School of Dentistry, University of Sassari, Sassari, Italy; School of Dentistry Sechenov University, Moscow, Russia); M Diaz Betancourt (Department of Restorative, Preventive and Paediatric Dentistry, University of Bern, Bern, Switzerland); MG Cagetti (Department of Biomedical, Surgical and Dental Science, University of Milan, Milan, Italy); RA Giacaman (Cariology and Gerodontology Units, Department of Oral Rehabilitation, Faculty of Health Sciences, University of Talca, Talca, Chile); DJ Manton (Cariology and Paediatric Dentistry, Centrum voor Tandheelkunde en Mondzorgkunde, University Medical Centre Groningen, Groningen, the Netherlands); GVA Douglas (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); T S Carvalho (Department of Restorative, Preventive and Paediatric Dentistry, University of Bern, Bern, Switzerland); JC Carvalho (Department of Oral Health Sciences, Faculty of Medicine and Dentistry, UCLouvain, Louvain, Belgium); A Vukovic (Department of Pediatric and Preventive Dentistry School of Dental Medicine University of Belgrade, Belgrade, Serbia); F J Cortés-Martinicorena (Private Practitioner, Valencia, Spain); D Bourgeois (University Claude Bernard Lyon 1, Laboratory “Systemic Health Care,” University of Lyon, Lyon, France); V Machiulskiene (Clinic of Dental and Oral Pathology, Faculty of Odontology, Department of Pediatric and Preventive Dentistry, Lithuanian University of Health Sciences, Kaunas, Lithuania); R Sava-Rosianu (Department of Preventive, Community Dentistry and Oral Health, Faculty of Dentistry, University of Medicine and Pharmacy “Victor Babes”, Timisoara, Romania); J Krithikadatta (Department of Conservative Dentistry & Endodontics, Saveetha Dental College and Hospitals, Chennai, India); N Sergeevna Morozova (School of Dentistry Sechenov University, Moscow, Russia); AM Acevedo (Universidad Central de Venezuela, Facultad de Odontología, Caracas, Venezuela); AA Agudelo-Suarez (Faculty of Dentistry, University of Antioquia, Colombia, Medellín, Antioquia); G Aguirre, K Aguirre (Facultad de Odontología, Universidad de El Salvador, El Salvador); I Alfonso (Universidad Central de Venezuela, Facultad de Odontología, Caracas, Venezuela); H A Ghaffar, R A El Fadl (Faculty of Dentistry, Ain Shams University, Cairo, Egypt); S A Al Maghlouth (King Fahad Medical City, Riyadh, Saudi Arabia); T Almerich-Torres (Facultad de Odontología - Universidad San Martín de Porres, Lima, Peru); F Amadori (Dental Clinic, School of Pediatric Dentistry, University of Brescia, Brescia, Italy); K Antia, (School of Health Sciences, University of Georgia, Tbilisi, Georgia); E Bajric (Department of Preventive and Pediatric Dentistry, Faculty of Dentistry with Clinics of Sarajevo University, Sarajevo, Bosnia Hercegovina); D M Barbosa-Liz (Faculty of Dentistry, University of Antioquia, Colombia, Medellín, Antioquia); S Ben-Tanfous (Faculty of Dental Medicine of Monastir, Monastir, Tunisia); H Bieber (Bundeswehr Medical Service Headquarters, Koblenz, Germany); S Bhusari (Heidelberg Institute of Global Health (HIGH), Heidelberg, Germany); S Birch (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); G Bontà (Department of Biomedical,

Surgical and Dental Science, University of Milan, Milan, Italy); P Bottenberg (Oral Health Research Cluster, VUB, Bruxelles, Belgium); J J Bruers (University of Amsterdam and Vrije Universiteit Amsterdam, Amsterdam, the Netherlands); L Bustillos (Universidad de Los Andes, Facultad de Odontología, Merida, Venezuela); P Bührens (Freier Verband Deutscher Zahnärzte, Bonn, Germany); J Cai (The State Key Laboratory Breeding Base of Basic Science of Stomatology & Key Laboratory of Oral Biomedicine Ministry of Education, School & Hospital of Stomatology, Wuhan University, Wuhan, People's Republic of China); J L Cairoli (Department of Biotechnology and Life Sciences, University of Insubria, Varese, Italy); F CA Carrer (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); M B Correa Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); B Cortés-Acha, F Carrouel (University Claude Bernard Lyon 1, Laboratory "Systemic Health Care," University of Lyon, Lyon, France); R de Carvalho Oliveira (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); F Cocco (Department of Surgery, Microsurgery and Medicine Sciences, School of Dentistry, University of Sassari, Sassari, Italy); F Crombie (Melbourne Dental School, Faculty of Medicine, Dentistry & Health Sciences, The University of Melbourne, Melbourne, Australia); J I Csikar (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); D Declerck, M Denkovski (Private practitioner, Skopje, Macedonia); J Deschner (Department of Periodontology and Operative Dentistry, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany); J Dopico-San Martin (Private Practitioner, Pamplona, Spain); O Viktorovna Dudnik (School of Dentistry Sechenov University, Moscow, Russia); W Y Escobar (Facultad de Odontología, Universidad de El Salvador, El Salvador); A Elwishahy (Heidelberg Institute of Global Health at Heidelberg University Hospital, Heidelberg, Germany); C E Fernandez (Cariology and Gerodontology Units, Department of Oral Rehabilitation, Faculty of Health Sciences, University of Talca, Talca, Chile); M Fontana (University of Michigan School of Dentistry, Ann Arbor, USA); A Frattaroli Pericchi (Universidad Central de Venezuela, Facultad de Odontología, Caracas, Venezuela); M Ghorbe (Faculty of Dental Medicine of Monastir, Monastir, Tunisia); Gigineishvili E (School of Health Sciences, University of Georgia, Tbilisi, Georgia); A García Quintana (Universidad Central de Venezuela, Facultad de Odontología, Caracas, Venezuela); J Gray (King Fahad Medical City, Ryad, Saudi Arabia); N Gugni (Department of Pediatric and Preventive Dentistry, Dental college, Haryana, India); K Gambetta-Tessini (Cariology and Gerodontology Units, Department of Oral Rehabilitation, Faculty of Health Sciences, University of Talca, Talca, Chile); A Haider (Department of Oral and Maxillofacial Surgery, Fatima Memorial Hospital, Lahore, Pakistan); M Hopcraft (Melbourne Dental School, Faculty of Medicine, Dentistry & Health Sciences, The University of Melbourne, Melbourne, Australia); J Hüttmann (Freier Verband Deutscher Zahnärzte, Bonn, Germany); N Hysenaj (Department of Prosthodontic, Faculty of Dental Medicine, University of Medicine, Tirana, Albania); A Jalal (Department of Oral and Maxillofacial Surgery, Fatima Memorial Hospital, Lahore, Pakistan); M Jikia (School of Health Sciences, University of Georgia, Tbilisi, Georgia); J John (Faculty of Dentistry, University of Malaysia, Kuala Lumpur, Malaysia); G Kaps-Richter (Freier Verband Deutscher Zahnärzte, Bonn, Germany); T Kerber Tedesco (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); S Leon (Cariology and Gerodontology Units, Department of Oral Rehabilitation, Faculty of Health Sciences, University of Talca, Talca, Chile); K A Levin (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); H Pau Lew (Department of Restorative Dentistry National Dental Centre Singapore, Singapore); M Aperecida Moreira Machado (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); A Beneictovna MacLennan (School of Dentistry Sechenov University, Moscow, Russia); J Onome Mafeni (Intercountry Center for Oral Health for Africa, Jos, Nigeria); M Minatel Braga Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); J M Montiel-Company (School of Dentistry University of Valencia, Valencia, Spain); A Malerba (Department of Biomedical, Surgical and Dental Science, University of Milan, Milan, Italy); S Mandić-Rajčević (School of

Public Health and Health Management, Institute of Social Medicine, Faculty of Medicine, University of Belgrade, Belgrade, Serbia; Departmeny of Health Sciences; University of Milan, Milan, Italy); A Askerovich Mamedo (School of Dentistry Sechenov University, Moscow, Russia); S A Mani (Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia); O Marouane (Faculty of Dental Medicine of Monastir, Monastir, Tunisia); D Markovic (Department of Pediatric and Preventive Dentistry School of Dental Medicine University of Belgrade, Belgrade, Serbia); E Paredes Martinez (Facultad de Odontología - Universidad San Martin de Porres, Lima, Peru); N Maroufidis (Private Practioner, Athens, Greece); F Medeiros Mendes (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); S Musa (Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia); A Necibi (Faculty of Dental Medicine of Monastir, Monastir, Tunisia); N Azlida Mohd Nor (Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia); B Tochukwu Ojukwu (Intercountry Center for Oral Health for Africa, Jos, Nigeria); N Opdam (School of Dentistry, Radboud University Medical Centre, Nijmegen, the Netherlands); L Ottolenghi (Department of Dental and Maxillo-Facial Sciences, Sapienza Rome University, Rome, Italy); J Owen (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); A Pássaro (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); I F Persoon (Department of Preventive Dentistry, Academic Centre for Dentistry Amsterdam, Amsterdam , the Netherlands); T Peric (Department of Pediatric and Preventive Dentistry School of Dental Medicine University of Belgrade, Belgrade, Serbia); E Pesaressi-Torres (Facultad de Odontología - Universidad San Martin de Porres, Lima, Peru); V Philippides (President Cyprus Dental Association, Nicosia, Cyprus); S P Plaza-Ruiz (Orthodontic Department, Fundación Universitaria CIEO-UniCIEO Bogotá, Colombia); D Procida Raggio (Universidade de Sao Paulo- School of Dentistry - Department of Community Dentistry, Sao Paulo, Brazil); F J Rivas Cartagena (Facultad de Odontologia, Universidad de El Salvador, El Salvador); F Ramos-Gomez (UCLA Center for Children's Oral Health (UCCOH) UCLA School of Dentistry, Los Angeles, USA); M Sabashvili (School of Health Sciences, University of Georgia, Tbilisi, Georgia); G Solis Sanchez, R Villena Sarmiento (Facultad de Odontología - Universidad San Martin de Porres, Lima, Peru); H Schrader (Freier Verband Deutscher Zahnärzte, Bonn, Germany); S Serban (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); R Bairstow (Department of Dental Public Health, School of Dentistry, University of Leeds, Leeds, UK); A Senna (Department of Biomedical, Surgical and Dental Science, University of Milan, Milan, Italy); B Shi (The State Key Laboratory Breeding Base of Basic Science of Stomatology &Key Laboratory of Oral Biomedicine Ministry of Education, School & Hospital of Stomatology, Wuhan University, Wuhan, People's Republic of China); C PC Sim (Department of Restorative Dentistry National Dental Centre Singapore, Singapore); S Slabsinskiene (Clinic of Dental and Oral Pathology, Faculty of Odontology, Department of Pediatric and Preventive Dentistry, Lithuanian University of Health Sciences, Kaunas, Lithuania); G Spagnulo (Department of Neurosciences, Reproductive and Odontostomatological Sciences, University of Naples "Federico II", Naples, Italy; School of Dentistry Sechenov University, Moscow, Russia); O Olufemi Taiwo (Intercountry Center for Oral Health for Africa, Jos, Nigeria); A Thodhorjani (Department of Prosthodontic, Faculty of Dental Medicine, University of Medicine, Tirana, Albania); P Tietler (Freier Verband Deutscher Zahnärzte, Bonn, Germany); C M C Volgenant, M H van der Veen ((Department of Preventive Dentistry, Academic Centre for Dentistry Amsterdam, Amsterdam , the Netherlands); Z Vlahovic (Medical Faculty of Pristina - Kosovska Mitrovica, Pristina, Montenegro); A Visaria (Duke-NUS Medical School, Singapore); Y Romero Uzcategui (Universidad de Los Andes, Facultad de Odontología, Merida,Venezuela); E Xhajanka (Department of Prosthodontic, Faculty of Dental Medicine, University of Medicine, Tirana, Albania); Q Yan, O Zen (The State Key Laboratory Breeding Base of Basic Science of Stomatology &Key Laboratory of Oral Biomedicine Ministry of Education, School & Hospital of Stomatology, Wuhan University, Wuhan, People's Republic of China); O Zeyer (Schweizerische Zahnärzte-Gesellschaft SSO, Bern, Switzerland); A Zukanovic (Department of Preventive and Pediatric Dentistry, Faculty of Dentistry with Clinics of Sarajevo

University, Sarajevo; Bosnia Hergevozina); T G Wolf (Department of Periodontology and Operative Dentistry, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany).

Correspondence to:

Prof. Guglielmo Campus, Department of Restorative, Preventive and Paediatric Dentistry at the University of Bern School of Dental Medicine, Freiburgstrasse 7, CH-3012 Bern (Switzerland)

Tel +41 31 664 0987

guglielmo.campus@zmk.unibe.ch

Abstract

Objectives: A multicentre survey was designed to evaluate the impact of COVID-19 outbreak on dental practice worldwide, estimate the COVID-19 related symptoms/signs, work attitudes and behaviour and the routine use of protective measures and personal protective equipment (PPE).

Methods: A global survey using a standardized questionnaire with research groups from 36 countries was designed. The questionnaire was developed and pretested during April 2020 and contained three domains: 1) personal data; 2) COVID-19 positive rate and symptoms/signs presumably related to the coronavirus; 3) working conditions and PPE adopted after the outbreak. Countries' data were grouped by the country positive rate (CPR) during the survey period and by Gross-National-Income *per capita*. An ordinal multinomial logistic regression model was carried out with COVID-19 self-reported rate referred by dental professionals as dependent variable to assess the association with questionnaire items.

Results: A total of 52,491 questionnaires were returned with a male/female ratio of 0.63. Out of the total respondents, 7,859 dental professionals (15%) reported symptoms/signs compatible with COVID-19. More than half of the sample ($n=27,818$; 53%) stated to use FFP2/N95 masks, while 21,558 (41.07%) used eye protection. In the bivariate analysis, CPR and N95/FFP2 were significantly associated ($OR=1.80$ $_{95\%CI}=1.60/2.82$ and $OR=5.20$ $_{95\%CI}=1.44/18.80$, respectively), while Gross-National-Income was not statistically associated with CPR ($OR=1.09$ $_{95\%CI}=0.97/1.60$). The same significant associations were observed in the multivariate analysis.

Conclusions: Oral health service provision has not been significantly affected by COVID-19, although access to routine dental care was reduced due to country-specific temporary lockdown periods. While the dental profession has been identified at high-risk, the reported rates of COVID-19 for dental professionals were not significantly different to those reported for the general

population in each country. These findings may help to better plan oral health care for future pandemic events.

Keywords

COVID-19; SARS-CoV-2; Dentistry; Dental professionals; Personal Protective Equipment (PPE); Infection control; Oral Health; Global Health; Public Health Dentistry

Journal Pre-proof

Introduction

The COVID-19 outbreak worldwide has had severe repercussions on individual behaviours and social interaction. Viral transmission occurs mainly through inhalation/ingestion/direct mucous contact with saliva droplets. Additionally, the virus could stagnate on hands, objects or surfaces for a long time and remain infectious [1].

Among the many potential sources of transmission and spread of SARS-CoV-2, dental services have received high attention [2-4]. Dental professionals are at the front line of healthcare, working in the mouth, with a potentially increased viral exposure. Indeed, according to the USA's Occupational Safety and Health Administration agency, all dental healthcare personnel (dentist, dental hygienist and dental assistant) are included at the highest risk category for SARS-CoV-2 exposure [4]. The putative risk posed by the dental profession was widely conveyed on mainstream media [5]. Whether this assertion is justified and supported by worldwide data, is elusive. We, therefore, decided to explore this issue through a global multicentre research survey among dental professionals.

During the first months of the COVID-19 pandemic, with extremely high infection rates and relatively high mortality, many national authorities worldwide, including some dental organizations, imposed several restrictive measures on the profession. In most countries, dental care was restricted during the COVID-19 outbreak and limited only to urgent care. Implementation of effective infection control measures was recommended to prevent nosocomial coronavirus infection [6]. Thus, the dental profession was forced to change in March-April 2020 to accommodate to new regulations and guidelines. By the beginning of the pandemic and during lockdown periods, dental treatment has been generally suspended or postponed, except for urgent care. The response of the dental profession to COVID-19 has not been homogeneous. While some countries restricted dental care only to urgencies/emergencies, in others, where the pandemic had subsided or even during the phase of maximum infection rates, dental clinics and offices had returned/continued to also provide non-urgent dental care, but figures from global data are not available. Moreover, limited knowledge at that time, unavailability of tests, availability or incorrect use of personal protective equipment (PPE) and conflicting protocols may have lowered protection among the dental care workforce and patients, potentially increasing the infection rate and transmission [7-10]. This paper reports findings from the 2020 International Collaborative COVID-19 Disease Study giving, to the best of authors' knowledge, the only updated worldwide observational epidemiological study on COVID-19 and dental professionals. The study describes morbidity due to COVID-19 among dental personnel in 36 countries/areas from May to August 2020, as well as the approaches to protective measures used to reduce the risk of viral infection and transmission.

The aim of this global survey was to evaluate the impact of the COVID-19 outbreak among dental professionals in different countries worldwide, via an extensive multicentre study. Due to the numerous data collected of this global study, this is the first paper of a series of epidemiological publications of dental professionals regarding COVID-19. The main aims were to estimate the impact of COVID-19 on oral health service provisions, the disease rate among dental professionals globally, describe self-reported symptoms/signs presumably related to COVID-19, and to investigate the adoption of protective measures and the PPE used to reduce the risk of viral infection and transmission.

Materials and Methods

Study design

The structured closed cross-sectional survey was developed with a modified Delphi method using a standardized questionnaire developed and pretested in Italy during April 2020 [9]. Briefly, following the Stehr-Green scale, the questionnaire was built up and structured into three domains: 1) personal data including age, gender, area of living and working, and working status; 2) dental professionals infection rate and symptoms/signs presumably related to the COVID-19; 3) working conditions and PPE adopted after the outbreak of the infection. (Appendix 2). A pre-test before the Italian survey was performed with twelve dental professionals, reaching a satisfactory Intraclass Correlation Coefficient (ICC) value of 0.80 for each item.

The protocol of the study was previously described and registered in the World Pandemic Research Network (WPRN) WPRN-486352 [10]. The study follows the CHERRIES guidelines (the checklist is available as supplementary file). Thirty-six collaborating research groups around the world were contacted by the central management team and all agreed to participate. Each research team was instructed on the research protocol, the survey design and the participant search strategies by a central coordinator (GC). Research teams were formed individually in each country and tasks were assigned at their convenience.

Participants and study size

Dental professionals working in each national health system, either private or public, including general or specialists were enrolled. Each country performed a sample size calculation; the total number of working dental professionals was ascertained, and according to the previously published protocol, at least a high proportion of participants (5-20%) in each country was invited to participate, aiming to reach minimum 5% proportion [10]. In order to reach this purpose, each national team could use the strategies that best suited the context of their country. Study participants

were reached in each country via adequate platforms to avoid duplicate answers, and the invites were sent through the national Dental Board systems, official web pages of national Dental Boards and/or social media groups/pages. The number of invited dental professionals ranged from all personnel registered by the national Dental Board to members of the professional social media groups. A mixed strategy was used; direct email to individual dental professionals, invitations to participate through national dental associations and other related organizations and dissemination on social media (*i.e.* Facebook), so it was not possible to estimate how many invitations had been disseminated in many countries. The number of respondents and relevant demographic data for each country is shown in Table 1.

Bias

The platform used to collect data and run the survey in each country was set to avoid duplicate answers. The duration of the survey in each country was at least one week according to the previously published protocol [10] (Table 1). All national research teams sent accurately collected and organized data according to the protocol, with the written report on country level data.

Each country/region translated the questionnaire, performed a pilot test and semantically adjusted the questions, if necessary. The same core questionnaire was mandatorily applied to all participating countries, regardless of optional additional questions incorporated by each national team. The full questionnaire is presented as supplementary material (Appendix 2). A description of the survey's aim along with the link to the questionnaire and an online informed consent were provided to each participant in accordance with the applicable data protection laws of the respective countries. In some countries, the project protocol and the consent form were approved by the local Ethics Committee. Each participant was defined by a unique identification number based on PI address. Participants were allowed to review and correct their answers, but if they did not sign the consent, the questionnaire was automatically closed. Data collection timeframe was from May to August 2020. All data were stored in a repository using Research Electronic Data Capture (REDCap) account.

Independent variables and data sources

Data from all the participating countries were stratified in several ways:

- if the dental professionals in the different countries continued to work routinely during the lockdown or limited the clinical activity to emergencies only;
- by the country/region (*i.e.* California not U.S.A. etc.) community positive rate (CPR) of COVID-19 during the survey period for each country (Table 1) [11]. The CPR were

calculated using the Git Hub data [12] and, where internationally available data were not updated, by the national research teams obtaining data from the official national web page reporting COVID-19 statistical recourses in each local language.

- by Gross National Income (GNI) per capita 2019, Atlas method and Purchasing Power Parity (PPP, international dollar) [13].

Outcome variables

Ordinal multinomial linear regression analysis was run using the COVID self-reported rate as dependent variable to evaluate the association with questionnaire items. COVID self-reported rate was calculated using data derived from the questionnaire, reporting one or more specific or nonspecific COVID symptoms [14,15]. Clinical manifestations of COVID-19 appear after an incubation period of around 5–6 days and most frequently include fever, dry cough, tiredness with the possible onset of sputum production, headache, haemoptysis, diarrhoea, dyspnoea, and/or lymphopenia amongst others [16-20].

Statistical Analysis

Data were exported to an Excel (Microsoft Corp., WA, USA) spreadsheet and quality-checked to ensure accuracy, sent to the management team (GC, MDB, MGC, TGW) and transferred to STATA16™ (Statacorp, TX, USA) for statistical analysis. The raw data were added as a supplementary file (Appendix 3). Data analysis was performed centrally. Demographic data and the impact of COVID-19 on dental professionals and oral health provision were descriptively presented.

All demographic data were completed without missing data. Missing data were present on the outcome variables and complete case analysis was selected as the primary analysis with no imputation [21,22]. Absolute and relative frequencies were calculated for each item. Differences in proportions were evaluated with χ^2 test or Fisher exact test if one cell had a value of less than five. Multiple testing for *post hoc* estimation, such as the number of observed frequencies, expected frequencies, percentage, and contribution to the chi-square were run. Estimation of a nonparametric test for trend across the areas with different prevalence of COVID self-reported rate and questionnaire items were also calculated. The effect size was calculated using the Cramer's V, as a measure of the strength of association among the levels of the row and column variables. Ordinal multinomial linear regression analysis was run to evaluate the association with questionnaire items.

Results

The map of the countries participating in the survey, the country positive rate (CPR) to COVID-19 according to the survey period and COVID self-reported positive rate are displayed in Figure 1. A total of 52,491 dental professionals from the 36 countries completed the questionnaires. The total population, survey period (2020), total new cases during the survey period, the positive rate, the type of oral health care provision, the reported number of dental professionals, the hypothetical sample at the 5% level and the number of dental professionals reached in each country by continent, country/area is displayed in Table 1.

Descriptive data

In most countries, the dental professionals sex ratio favoured women (*i.e.* in Russia 100% were women), whilst Switzerland had the highest ratio for males (1.97). More than 50% of the dental professionals in countries such as Argentina, Peru, USA/California, Venezuela, Australia, Albania, Belgium, Cyprus, Chile, Germany, Greece, Italy, Lithuania, Macedonia, Montenegro, Netherlands, Romania, Spain and Switzerland, reported to work as owners of a private practice. Working within a National Health System or at Universities or for administrative staff was reported highly by Nigerian, Chinese (province of Hubei), Malaysian, Singaporean and UK dental professionals. The majority were general dental professionals (43.2% *versus* 23.61% specialists), one third of the participants did not enter this item.

Table 1 shows the impact of the COVID-19 pandemic on the provision of routine oral health services globally. Only 14 of the 36 countries reported to perform routine cares across both private and public sectors during the survey period. Thirteen countries reported provision of emergency care only. Almost 80% ($n=41,776$) of dental professionals continued providing some face-to-face clinical activities, many despite limited access to appropriate PPE.

Self-reported prevalence of COVID-19 in dental professionals

Overall, 15% of respondents ($n=7,859$) reported symptoms/signs pertaining to COVID-19. This was higher in the Americas, in particular, Perú and El Salvador where more than two thirds of the respondents reported experiencing at least one symptom assumed to be from COVID-19. Of those who reported signs of having COVID-19, more than 90% reported to have worked clinically during the outbreak, while the remainder had only community level exposure, many unable to work or working from home offering remote dental advice (Teledentistry) and triage services.

Participating dental professionals were grouped by the positive rates registered at the country level (CPR) and by the COVID self-reported rate (Table 2). The COVID self-reported rate ranged

between 0% in some Balkan Countries and Nigeria to over 10% in Russia (14.68%) and Pakistan (17.34%). At country level, the community rates of COVID-19 were usually higher than those reported by the dental participants, except for China, Malaysia, Netherlands, Pakistan, Russia, Saudi Arabia and Tunisia, where the results were higher for dental professionals than the CPR.

Protecting measures for patients and dental professionals

The precautionary measures adopted by dental professionals who continued to work after the COVID-19 outbreak are displayed in Table 3. More than half of the dental professionals (n=27,818; 53%) reported the use of FFP2/N95, while 41.07% (n=21,558) declared the use of eye protection equipment. The percentage of dental professionals who reported routine use of protective mask (FFP2/N95) was notably high in South American countries with low GNI (Brazil, Peru and Colombia with 100%). In most countries, more than half of the dental professionals reduced the number of patient appointments to reduce crowding in the waiting room, with the highest percentage in countries with the highest GNI, *i.e.* Germany (93.84%). The patients' health status and body temperature were checked by 57.6% of the respondents, with the highest prevalence in countries with the highest GNI (Australia 99.84% and Germany 91.05%). The use of mouthrinses intended to reduce SARS-CoV-2 salivary load was highly variable, but some trends between countries with different GNI were observed regarding the use of alcohol-containing mouthrinses and essential oils or cetylpyridinium chloride ($p=0.01$ and $p<0.01$, respectively) (*data not tabulated*).

When COVID self-reported rate among dental professionals was used as the dependent variable for an ordinal multinomial regression model (Table 4), CPR and N95/FFP2 were statistically significantly associated ($\beta=1.07$ and $\beta=-1.46$, respectively both $p<0.01$). Also, GNI was statistically associated to the test-positive rate among dental professionals ($\beta=0.72$ $p=0.03$). The face-to-face clinical activity and private practice rate (PPr) (<25%) were not statistically associated to the dependent variable.

Discussion

This is the first global population-based study, to the authors' knowledge, that describes the effect of COVID-19 on the dental profession, describing self-reported COVID-19 infection and presumably related symptoms/signs suffered, exploring work attitudes and behaviour, and investigating the routine use of protective measures and PPE. The questionnaire used also included other aspects such as COVID-related risk awareness and perception; however, the focus of this project was centred on the dissemination of the infection among dental professionals and the

implemented protection measures for dental professionals and patients. The survey included more than 50,000 participants across 36 countries. Findings showed that living in countries with high COVID-19 positive rates increased the probability that the dental professionals became infected, although these higher rates could be related to infection acquired within the community, as well as at work. In terms of protective measures, the use of N95/FFP2 masks significantly reduced the chances of the dental professionals reporting symptoms/signs related to the disease, while the other PPEs seem less relevant for SARS-CoV-2 protection. Overall, the probability of dental professionals becoming infected was associated to the Gross National Income (GNI) of the countries where they work.

The present study offers a global picture of the dental profession and COVID-19, even if the large number of respondents cannot be considered representative of the entire world population of dental professionals; nonetheless, these findings can provide a measurement framework and methods for countries and global stakeholders to better plan health service provision.

Although our study provided relevant information on the effect of COVID-19 on the dental profession, caution is necessary when interpreting the results due to some limitations. First of all, the obvious limitation could be choosing COVID self-reported positive rate as dependent variable as representative of the impact of the disease in the field. In most countries, testing was not available in the earlier stages of the pandemic. Moreover, any person who presented any of the symptoms related to COVID were asked to isolate. These people were, then, certainly impacted by the disease, even if they did not have an official positive testing. In this case, the number of professionals who had symptoms might be higher than the true prevalence of COVID. Also, there is need to highlight the possibility of false negative test results and the necessity to measure clinical sensitivity and specificity of each test according to real-life situations (symptoms, contacts etc) [23]. Further research would be needed to address other outcome data collected during this survey, such as tested "positive for COVID", and "hospitalized for COVID ". On the other hand, there is possibility that the lack of access to testing in low-income countries, may have led to an over-estimation of infection due to reliance on untested symptoms.

Secondly, the time periods for collection of the survey data and independent reports of dental professional infection rates do not coincide since survey was carried out from May to August 2020. The dental professionals reported infection rates from the beginning of the pandemic. However, this does not reduce the relevance of the results, since these data were only used to categorize countries and as a metric for comparing countries. In addition, these data provide useful information on how adequately countries were testing, also in relation to the size of the outbreak and thus facilitating the understanding of how widespread the virus was, in conjunction with confirmed case data [24].

According to World Health Organization (WHO) criteria, a positive testing rate of less than 5% is a reliable indicator that the epidemic is under control [24].

Although these data do not seem to support all the recommended enhanced protective measures, except for higher-rated masks, the results must be interpreted cautiously, nonetheless. The use of higher-rated masks, in itself, may be an indicator of higher levels of protection overall. This survey is based on self-reported information in a large sample of dental professionals from 36 countries. Importantly, the design of this research does not allow inferences to be drawn about cause and effect, therefore, conclusions must be carefully drawn. It is possible that at the individual level, the use of N95/FFP2 masks alone is not a sufficient protection against SARS-CoV-2 infection. It might be speculated that dental professionals wearing N95 masks at work are also more likely to wear them when they are in normal life settings, reducing the risk of community transmission. However, such appealing hypothesis cannot be verified from the present data.

The pandemic of COVID-19 has affected the entire world, with over 150 million cases and more than 3.2 million deaths by May 2021, worldwide [25], and an increasing infection and death rate in developing countries. The dental profession has not had universal guidelines on how to manage the virus, despite being perceived as having extremely high risk for exposure among all professions [2-4]. However, data from similar studies in Europe and USA showed that the prevalence and the positivity rates among dental professionals/dental hygienists was low, suggesting that current infection control recommendations against COVID-19 could be appropriate [8,9,25, 26]. Similarly, our findings follow the same trend for most countries, where infectivity rate during the performance of the survey could be retrieved (Tables 2 and 3). This trend could be at least partially explained by the preventive measures frequently implemented by dental professionals all over the world, that protect the dentist whilst treating symptomatic patients, reducing the risk of infection [27,28].

High variability could be observed among countries, but in general, dental professionals from countries with higher GNI reported lower infection rates, potentially because they were less well protected in lower GNI countries; however, it could also be a reflection on testing protocols in high income countries which may have necessitated that healthcare workers be tested disproportionately more than the general population. In a recent survey among other health professions in the USA, the majority of responders reported to have access to PPEs [27,29], but the situation is different in countries with lower GNI, which may lead to higher infection rates. In eight of the surveyed countries, the dental professionals' COVID self-reported rate was higher than that of the general population (Pakistan, Russia, Saudi Arabia, Tunisia, Netherlands, Lithuania, Malaysia and China).

When dental professionals continued practicing in countries with higher positive rates, our data showed that this also was associated with a higher COVID self-reported rate than those working in countries with lower rates (Table 4). The results also showed a trend for a higher rate of COVID self-reported infection among dental professionals working in public rather than private practice. It is reasonable to speculate that in hospitals and primary public care centres more people congregate at the same time, reducing time to ventilate rooms between patients, increasing the risk of virus transmission. It is also possible that, in such places, there are higher odds of COVID-19-positive people gathering, among patients, institutional staff and accompanying persons. It has been shown that other health professionals working in health centres are at a higher risk of acquiring the SARS-CoV-2 due to direct contact with positive patients, long working hours, lack of preparation or limited access to PPE [7,27,29]. This higher risk may be also true for community contacts when infection rates are high in the country.

Although droplets of contaminated secretions and direct contact were initially regarded as the main routes for SARS-CoV-2 infection [30], aerosol-generating procedures (AGPs) in contaminated environments is an indisputable route, in addition to exhalation, speaking, sneezing or coughing [3,4,31]. At the present time, nosocomial infection of SARS-CoV-2 through the airborne route has not been reported [32]. The ordinal multinomial logistic regression showed that only the use of N95/FFP2 masks significantly reduced the probability of reporting signs/symptoms of COVID-19. Information was not collected to explain, when N95/FFP2 masks were not used, the reason for this action; it is therefore impossible to know if N95/FFP2 masks were not available to dental professionals or if their non-use was due to cost or because they were not considered more effective than surgical masks. In other words, the use of high-quality masks appears to be the most effective measure to avoid infection by SARS-CoV-2. From the HIV/AIDS onset and the increased awareness of hepatitis B and C in the 1990's, the dental profession adopted strict infection control measures. Indeed, the Center of Disease Control (CDC) released guidelines for infection control starting to the assumption that every patient in a dental office could be infected with HIV or Hepatitis B viruses [33]. These guidelines were later updated, based on new evidence, but the philosophy was not altered [34]. This may explain the low infection rate among dental professionals, despite the aerosol generation and the close proximity to potentially infected patients. Dental professionals have a "culture" of infection control. When transmission rates of the pandemic are in the respective region high, dental professionals should act following a similar behaviour as they did in those days for HIV, considering every patient as COVID-19 positive. However, when transmission is low or the pandemic is alleviated, these measures will cause waste of resources, contradicts sustainability, and add burdens to countries where resources are limited. Using higher

levels of protection should be dependent on community transmission levels of COVID-19. The facilities and personnel must undergo an even stricter infection control protocol, reduce AGPs, space appointment times, establish correct ventilation and train dental and other associated health workers [35,36]. Thus, the extra recommendations raised after the COVID-19 outbreak were rapidly adopted for most dental professionals that continued treating patients, either restricting it to dental urgencies/emergencies or those having elective care [34,35]. Public health organizations rapidly released enhanced infection prevention and control recommendations with specific PPEs, based on the knowledge of SARS-CoV-2 at the time. Thus, dental AGPs must be performed wearing surgical masks, basic clinical PPE and eye protection. For those care providers with no access to high-standard N95/FFP2 masks and when performing AGPs, the guidelines strongly suggest using a high-level surgical mask along with a face shield over it simultaneously [35,37,38]. Handling of equipment is also important, as bacterial contamination of masks during manipulation has been recently reported in the AGP dental settings [38,39]. Changing the mask after treating a patient followed by hand disinfection has been recommended [40]. It is recommended that the current international and national guidelines are maintained until evidence dictates otherwise or when the pandemic is alleviated.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests

Authors Contributions

G Campus, contributed to conception, design, data acquisition, analysis, and interpretation, drafted and critically revised the manuscript; M D Betancourt contributed to design, data analysis, and interpretation and critically revised manuscript; M G Cagetti, T G Wolf, contributed to conception, design, interpretation, drafted and critically revised the manuscript; R A Giacaman D J. Manton, G V.A. Douglas, T Saads Carvalho, J C Carvalho, contributed to design, acquisition, analysis, and interpretation, drafted and critically revised the manuscript; A Vukovic, J F. Cortés-Martinicorena, R Sava-Rosianu, J Krithikadatta, contributed to design, acquisition and interpretation, critically revised the manuscript; D Bourgeois contributed to interpretation and critically revised the manuscript; N Sergeevna Morozova, AM Acevedo, A A Agudelo-Suarez, G Aguirre, K Aguirre, I Alfonso, H A Ghaffar, R A El Fadl, S A A AlMaghlouth, T Almerich-Torres, F Amadori, K Antia, E Bajric, D M Barbosa-Liz, S Ben-Tanfous, H Bieber, S Bhusari, S Birch, G Bontà, P Bottenberg, J J Bruers, L Bustillos, P

Bührens, J Cai, F CA Carrer, M B Correa, B Cortés-Acha, F Carrouel, R de Carvalho Oliveira, F Crombie, J I Csikar, D Declerck, M Denkovski, J Deschner, J Dopico-San Martin, O Viktorovna Dudnik, W Y Escobar, A Elwishahy, C E Fernandez, M Fontana, A Frattaroli Pericchi, M Ghorbe, Gigineishvili E, A Garcia Quintana, J Gray, N Gughani, K Gambetta-Tessini, A Haider, M Hopcraft, J Hüttmann, N Hysenaj, A Jalal, M Jikia, J John, G Kaps-Richter, T Kerber Tedesco, S Leon, K A Levin, H Pau Lew, M Aperecida Moreira Machado, A Beneictovna MacLennan, J Onome Mafeni, M Minatel Braga, J M Montiel-Company, A Malerba, S Mandić-Rajčević, A Askerovich Mamedo, S Ann Man, O Marouane, D Markovic, E Paredes Martinez, N Maroufidis, F Medeiros Mendes, S Musa, A Necibi, N Azlida Mohd Nor, B Tochukwu Ojukwu, N Opdam, L Ottolenghi, J Owen, A Pássaro, I F Persoon, T Peric, V Philippides, S P Plaza-Ruiz, D Procida Raggio, F J Rivas Cartagena, F Ramos-Gomez, M Sabashvili, G Solis Sanchez, H Schrader, S Serban, B Shi, C PC Sim, S Slabsinskiene, G Spagnulo, O Olufemi Taiwo, A Thodhorjani, P Tietler, E Pesaressi Torres, C M C Volgenant, M H van der Veen, Z Vlahovic, A Visaria, Y Romero Uzcategui, E Xhajanka, Q Yan, O Zen, O Zeyer, A Zukanovic, contributed to acquisition and critically revised the manuscript; J L Cairolí, A Senna, contributed to conception, design, acquisition, critically revised the manuscript; F Cocco contributed to analysis and interpretation, drafted and critically revised the manuscript. All authors gave final approval and agreed to be accountable for all aspects of the work.

Acknowledgments

The authors thank all participants of the study. Bernadette Rawyler and Ines Badertscher (University of Bern, Switzerland) This survey did not receive any specific grant from any funding agency in the public, commercial, or not-for-profit sectors. The survey was self-supported at country/group team level. The first three authors of the COVIDental Collaboration Group had full access to all the data. Individual participant data are available as supplementary material. The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

1. X. Peng, X. Xu, Y. Li, L. Cheng, X. Zhou, B. Ren, Transmission routes of 2019-nCoV and controls in dental practice. *Int. J. Oral Sci.* 3 (2020) (1) 9. <https://doi.org/10.1038/s41368-020-0075-9>.
2. M. Banakar, K. Bagheri Lankarani, D. Jafarpour, S. Moayedi, M. H. Banakar, A. Mohammad Sadeghi A, COVID-19 transmission risk and protective protocols in dentistry: a

- systematic review. *BMC Oral Health* 20 (2020) 275. <https://doi:10.1186/s12903-020-01270-9>.
3. J. E. Gallagher, K. C. Sukriti, I. G. Johnson, W. Al-Yaseen, R. Jones, S. McGregor, M. Robertson, R. Harris, N. Innes, W. G. Wade, A systematic review of contamination (aerosol, splatter and droplet generation) associated with oral surgery and its relevance to COVID-19. *BDJ Open* 6 (2020) 25. doi: 10.1038/s41405-020-00053-2.
 4. N. Innes, I. G. Johnson, W. Al-Yaseen, R. Harris, R. Jones, K. C. Sukriti, S. McGregor, M. Robertson, W. G. Wade, J. E. Gallagher. A systematic review of droplet and aerosol generation in dentistry. *J Dent.* 105 (2021) 103556. doi: 10.1016/j.jdent.2020.103556.
 5. D. Michaels, G.R. Wagner, Occupational Safety and Health Administration (OSHA) and Worker Safety During the COVID-19 Pandemic. *JAMA.* 324 (2020) 1389-1390. <https://doi:10.1001/jama.2020.16343>.
 6. The New York Times, The Workers Who Face the Greatest Coronavirus Risk. <https://www.nytimes.com/interactive/2020/03/15/business/economy/coronavirus-worker-risk.html?smid=em-share>, 2020 (accessed 8 December 2020).
 7. Y. Pan, H. Liu, C. Chu, X. Li, S. Liu, S. Lu, Transmission routes of SARS-CoV-2 and protective measures in dental clinics during the COVID-19 pandemic. *Am. J. Dent.* 33 (2020) 129-134.
 8. G. Bontà, G. Campus, M. G. Cagetti, COVID-19 pandemic and dental hygienists in Italy: a questionnaire survey. *BMC Health Serv Res.* 20 (2020) 994. <https://doi:10.1186/s12913-020-05842-x>.
 9. M.G. Cagetti, J. L. Cairolì, A. Senna, G. Campus, COVID-19 Outbreak in North Italy: An Overview on Dentistry. A Questionnaire Survey. *Int. J. Environ. Res. Public Health* 17 (2020) 3835. <https://doi:10.3390/ijerph17113835>.
 10. G. Campus, M. Diaz-Betancourt, M. G. Cagetti, J. C. Carvalho, T. S. Carvalho, J. F. Cortés-Martínez, J. Deschner, G.V.A. Douglas, R. S. Giacaman, V. Machiulskiene, et al., Study Protocol for an Online Questionnaire Survey on Symptoms/Signs, Protective Measures, Level of Awareness and Perception Regarding COVID-19 Outbreak among Dentists. A Global Survey. *Int. J. Environ. Res. Public Health* 17 (2020) 5598. <https://doi:10.3390/ijerph17155598>.

11. J.G. Hodge Jr, S. Wetter, E. Carey, E. Pendergrass, C.M. Reeves, H. Reinke, Legal "Tug-of-Wars" During the COVID-19 Pandemic: Public Health vs. Economic Prosperity. *J Law Med Ethics*. 48 (2020) 603-607. <https://doi:10.1177/1073110520958888>.
12. <https://github.com/datasets/covid-19>. 2021 (accessed 18 May 2021)
13. <https://datacatalog.worldbank.org/dataset/gni-capita-ranking-atlas-method-and-ppp-based>. 2021 (accessed 18 May 2021)
14. J. Baj, H. Karakuła-Juchnowicz, G. Teresiński, G. Buszewicz, M. Ciesielka, E. Sitarz, A. Forma, K. Karakuła, W. Flieger, P. Portincasa, et al., COVID-19: Specific and Non-Specific Clinical Manifestations and Symptoms: The Current State of Knowledge. *J Clin Med*. 9 (2020) 1753. <https://doi:10.3390/jcm9061753>.
15. Centers for Disease Control and Prevention. Symptoms of COVID-19 <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>, 2021(accessed 18 May 2021)
16. J.A. Belser, P.A. Rota, T. M. Tumpey, Ocular tropism of respiratory viruses. *Microbiol. Mol. Biol. Rev.* 77 (2013) 144-56. <https://doi:10.1128/MMBR.00058-12>.
17. M. Gandhi, D.S. Yokoe, D.V. Havlir, Asymptomatic Transmission, the Achilles' Heel of Current Strategies to Control Covid-19. *N. Engl. J. Med.* 382 (2020) 2158-2160. <https://doi:10.1056/NEJMe2009758>.
18. C.C. Lai, T.P. Shih, W.C. Ko, H.J. Tang, P.R. Hsueh, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int. J. Antimicrob. Agents*. 55 (2020) 105924. <https://doi:10.1016/j.ijantimicag.2020.105924>.
19. M.A. Lake, What we know so far: COVID-19 current clinical knowledge and research. *Clin. Med*. 20 (2020) 124-127. <https://doi:10.7861/clinmed.2019-coron>.
20. R. Holliday, J. R. Allison, C. C. Currie, D. C. Edwards, C. Bowes, K. Pickering, S. Reay, J. Durham, J. Lumb, N. Rostami, J. Coulter, C. Nile, N. Jakubovics, Evaluating contaminated dental aerosol and splatter in an open plan clinic environment: Implications for the COVID-19 pandemic. *J. Dent*. 105 (2021) 103565. doi: 10.1016/j.jdent.2020.103565.

21. J.A. Sterne, I.R. White, J.B. Carlin, M. Spratt, P. Royston, M.G. Kenward, A.M. Wood, J.R. Carpenter, Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ*. 338 (2009) b2393. <https://doi:10.1136/bmj.b2393>.
22. J.C. Jakobsen, C. Gluud, J. Wetterslev, P. Winkel, When and how should multiple imputation be used for handling missing data in randomised clinical trials - a practical guide with flowcharts. *BMC Med. Res. Methodol.* 17 (2017) 162. <https://doi:10.1186/s12874-017-0442-1>.
23. S. Woloshin, N. Patel, A. S. Kesselheim, False Negative Tests for SARS-CoV-2 Infection — Challenges and Implications. *N. Engl. J. Med.* 383 (2020), e38. <https://doi:10.1056/NEJMp2015897>
24. R.A. Middelburg, F.R. Rosendaal, COVID-19: How to make between-country comparisons. *Int. J. Infect. Dis.* 96 (2020) 477-481. <https://doi:10.1016/j.ijid.2020.05.066>.
25. Johns Hopkins University – Coronavirus Resource Center. COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). <https://coronavirus.jhu.edu/map.html>, 2021 (accessed 12 May 2021)
26. C.G. Estrich, M. Mikkelsen, R. Morrissey, M.L. Geisinger, E. Ioannidou, M. Vujicic, M.W.B. Araujo, Estimating COVID-19 prevalence and infection control practices among US dentists. *J. Am. Dent. Assoc.* 151 (2020) 815-824. <https://doi:10.1016/j.adaj.2020.09.005>.
27. M. Yanes-Lane, N. Winters, F. Fregonese, M. Bastos, S. Perlman-Arrow, J.R. Campbell, D. Menzies, Proportion of asymptomatic infection among COVID-19 positive persons and their transmission potential: A systematic review and meta-analysis. *PLoS One*. 15 (2020) e0241536. <https://doi:10.1371/journal.pone.0241536>.
28. F. Carrouel, L. S. Gonçalves, M. P. Conte, G. Campus, J. Fisher, L. Fraticelli, E. Gadea-Deschamps, L. Ottolenghi, D. Bourgeois, Antiviral Activity of Reagents in Mouth Rinses against SARS-CoV-2. *J Dent Res.* 100 (2021) 124-132. <https://doi:10.1177/0022034520967933>.
29. F. Calò, A. Russo, C. Camaioni, S. De Pascalis, N. Coppola, Burden, risk assessment, surveillance and management of SARS-CoV-2 infection in health workers: a scoping review. *Infect. Dis. Poverty* 9 (2020) 139. <https://doi:10.1186/s40249-020-00756-6>.

30. J. Coto, A. Restrepo, I. Cejas, S. Prentiss, The impact of COVID-19 on allied health professions. PLoS One 15 (2020) 0241328. <https://doi:10.1371/journal.pone.0241328>
31. T. Greenhalgh, J.L. Jimenez, K.A. Prather, Z. Tufekci, D. Fisman, R. Schooley, Ten scientific reasons in support of airborne transmission of SARS-CoV-2. Lancet. 397 (2021) 1603-1605. [https://doi.org/10.1016/S0140-6736\(21\)00869-2](https://doi.org/10.1016/S0140-6736(21)00869-2).
32. World Health Organization: Prevention, identification and management of health worker infection in the context of COVID-19, <https://www.who.int/publications/i/item/10665-336265>, 2020 (accessed 28 June 2021).
33. S.C.Y. Wong, R.T. Kwong, T.C. Wu, J.W.M. Chan, M.Y. Chu, S.Y. Lee, H.Y. Wong, D.C. Lung, Risk of nosocomial transmission of coronavirus disease 2019: an experience in a general ward setting in Hong Kong. J Hosp Infect. 105 (2020) 119-127. <https://doi:10.1016/j.jhin.2020.03.036>.
34. Centers for Disease Control and Prevention. Summary of infection prevention practices in dental settings: Basic expectations for safe care. <https://www.cdc.gov/oralhealth/infectioncontrol/summary-infection-prevention-practices/index.html>, 2020 (accessed 28th June 2021).
35. Centers for Disease Control and Prevention. Interim Infection Prevention and Control Recommendations for Healthcare Personnel During the Coronavirus Disease 2019 (COVID-19) Pandemic. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html>, 2021 (accessed 28th June 2021).
36. World Health Organization: Roadmap to improve and ensure good indoor ventilation in the context of COVID-19, <https://www.who.int/publications/i/item/9789240021280>, 2021 (accessed 28th June 2021).
37. S. Al Kawas, N. Al-Rawi, W. Talaat, Z. Hamdoon, B. Salman, S. Al Bayatti, W. Jerjes, A.B.R. Samsudin, Post COVID-19 lockdown: measures and practices for dental institutes. BMC Oral Health. 20(2020) 291. <https://doi:10.1186/s12903-020-01281-6>.
38. World Health Organization: Mask use in the context of COVID-19, [https://www.who.int/publications/i/item/advice-on-the-use-of-masks-in-the-community-during-home-care-and-in-healthcare-settings-in-the-context-of-the-novel-coronavirus-\(2019-ncov\)-outbreak](https://www.who.int/publications/i/item/advice-on-the-use-of-masks-in-the-community-during-home-care-and-in-healthcare-settings-in-the-context-of-the-novel-coronavirus-(2019-ncov)-outbreak), 2020 (accessed 28th June 2021).

39. European Centre for Disease Prevention and Control. Infection prevention and control and preparedness for COVID-19 in healthcare settings – Fifth update. <https://www.ecdc.europa.eu/en/publications-data/infection-prevention-and-control-and-preparedness-covid-19-healthcare-settings>. 2020 (accessed 28th June 2021).
40. M. Gund, J. Isack, M. Hannig, S. Thieme-Ruffing, B. Gärtner, G. Boros, S. Rupf, Contamination of surgical mask during aerosol-producing dental treatments. Clin Oral Investig. 7 (2020) 1–8. <https://doi:10.1007/s00784-020-03645-2>.

Journal Pre-proof

Table 1. Participating Countries survey Information.

Country	Population size (as of date)	Survey period (2020)	Total new cases in population (survey period)	N° dentists reported by the teams	Responders	% Positive/rate COVID-19 test in population	Oral health service provision/restrictions nationally (survey period)						
							Private	Public (Hospitals, etc.)					
		Start	End					Urgent	Routine	None	Urgent	Routine	None
Africa													
Egypt	102,334,404	16.07	31.07	9,827	19,111	925	24.41		Yes		Yes, only		
Nigeria	24,206,644	11.06	29.07	28,340	4,000	438	2.10		Yes		Yes, only		
Tunisia	11,818,619	08.07	25.07	237	4,500	871	1.51		Yes			Yes	
America													
Argentina	44,938,712	16.06	31.07	147,419	54,000	1,422	34.43	Yes, only			Yes, only		
Brazil	212,559,417	27.07	31.07	215,589	645,889	1,837	30.15	Yes, only			Yes, only		

Chile	19,116,201	20.07	01.08	26,821	21,541	2,307	10.98	Yes, only			Yes, only		
Colombia	50,882,891	19.06	24.07	212,382	36,000	5,375	26.60	Yes, only			Yes, only		
El Salvador	6,486,205	12.07	21.07	3,065	2,100	543	12.40	Yes, only					None
Peru	32,971,854	25.07	31.07	38,744	30,000	1,759	23.55		Yes				None
USA ^a	39,512,223	05.06	31.07	370,687	28,800	562	7.00	Yes, only			Yes, only		
Venezuela	28,435,940	18.07	28.07	5,134	12,000	638	5.46		Yes				None
Asia and Oceania													
China ^b	58,500,000	13.07	28.07	0	7,750	1,116	0		Yes			Yes	
India	1,380,004,385	11.07	25.07	543,059	277,303	3,265	10.92	Yes, only			Yes, only		
Malaysia	32,365,999	01.07	15.07	92	11,108	2,904	0.09		Yes		Yes, only		
Pakistan ^c	220,892,340	16.07	30.07	21,633	2,226	425	16.27		Yes		Yes, only		
Saudi Arabia	34,813,871	02.07	30.07	81,767	16,752	857	5.17	Yes, only			Yes, only		
Singapore	5,850,342	15.07	02.08	6,229	2,390	355	7.30		Yes			Yes	
Australia	25,667,000	13.07	31.07	6,750	16,861	815	0.69		Yes			Yes	
Europe													
Albania	2,877,797	16.07	31.07	2,850	2,776	206	39.1	Yes, only			Yes, only		

											y		
Belgium	115,89,623	30.07	04.08	3,273	10,080	782	2.68						
Bosnia	3,280,819	29.06	12.07	2,863	2,233	213	17.0		Yes			Yes	
Cyprus	1,207,359	16.07	31.07	62	1,017	176	0.49		Yes			Yes	
Georgia	3,989,167	10.07	31.07	192	3,050	317	1.48		Yes			Yes	
Greece	10,423,054	16.07	31.07	518	10,200	46	0.44						
Germany	83,783,942	04.06	16.06	4,469	70,740	3,542	0.70		Yes			Yes	
Italy	60461826	04.05	18.05	15,293	49,413	8,284	2.90	Yes, only			Yes, only		
Lithuania	2,722,289	12.07	24.07	99	3,800	204	0.46		Yes			Yes	
Macedonia	2,083,374	16.07	31.07	2,232	2,800	24	10.4	Yes, only			Yes, only		
Montenegro	628,066	16.07	31.07	1,184	591	110	14.84		Yes			Yes	
Netherlands	17,134,872	08.07	06.08	5,724	8,600	372	1.00		Yes			Yes	
Romania	19,237,691	16.07	31.07	16,006	16,442	1,051	4.72		Yes			Yes	
Russia	145,934,462	09.06	07.26	339,047	50,000	1,011	12.55	Yes, only			Yes, only		
Serbia	8,737,371	16.07	31.07	6,230	4,677	1,460	4.06		Yes			Yes	
Spain	46,754,778	29.05	11.06	4,374	38,000	2,318	0.80	Yes, only			Yes, only		
Switzerland	8,654,622	02.07	15.07	1,300	5,777	1,324	1.26		Yes			Yes	
UK	67,886,011	01.06	04.07	31,751	39,500	4,637	1.09	Until 08.06	Yes		Yes, only		
TOTAL						52,491							

^aUSA/California; ^bChina/Hubei; ^cPakistan/Lahore; ^d<https://www.worldometers.info/coronavirus>.

Table 2. Dentists positive rate to COVID-19 and work condition in the participating dentists.

Country	Dentists positive rate	Positive rate COVID-19 %	One or more symptoms	Dentist M/F ratio	Private	Public	Regular working load	Only emergencies
	% [□]		N (%)		N (%)	N (%)	N (%)	N (%)
Pakistan ^c	17.83	16.27	151 (36.39)	0.40	149 (36.43)	260 (63.57)	98 (24.62)	118 (29.65)
Russia	14.68	12.55	118 (11.67)	0.00	443 (43.81)	552 (34.59)	368 (36.44)	475 (47.03)
Saudi Arabia	7.84	5.17	257 (30.09)	0.86	210 (24.59)	628 (73.54)	47 (5.52)	383 (45.06)
Colombia	6.46	26.60	721 (13.41)	0.38	4327 (100.00)	0 (0.00)	126 (2.34)	1002 (18.64)
Brazil [^]	5.02	30.15	188 (10.22)	0.25	1161 (63.18)	664 (36.13)	87 (4.73)	1057 (57.54)
Tunisia	4.13	1.51	198 (22.73)	0.52	494 (64.98)	305 (35.02)	15 (1.73)	351 (40.53)
Spain	3.19	4.06	291 (12.55)	0.53	2073 (89.43)	245 (10.57)	18 (0.78)	916 (39.52)
Peru	3.19	23.55	1721 (97.84)	0.60	1358 (77.22) [°]	720 (40.93)	42 (2.39)	1187 (67.48)
Egypt [^]	3.09	24.41	191 (21.08)	1.31	361 (39.99)	545 (60.15)	146 (15.79)	416 (44.97)
Argentina	2.41	34.43	130 (9.20)	0.30	1166 (82.51)	247 (17.48)	20 (1.42)	557 (39.42)
Belgium	2.35	2.68	77 (9.51)	0.73	661 (93.95)	47 (6.05)	280 (35.90)	366 (46.92)
Chile	2.12	10.98	38 (1.65)	0.49	1592 (69.04)	747 (32.93)	96 (5.16)	1047 (56.26)
Netherlands	1.90	1.00	65 (17.91)	0.47	368 (100.00)	--	253 (69.70)	109 (30.03)
Bosnia	1.88	17.0	52 (24.41)	0.43	123 (57.75)	90 (42.25)	13 (6.10)	85 (39.91)
India	1.81	10.92	430 (13.97)	0.86	3302 (100.00)	0 (0.00)	24 (0.74)	3240 (99.26)
El Salvador	1.47	12.40	235 (43.26)	0.50	265 (48.36)	278 (51.19)	20 (3.68)	191 (35.17)
Georgia [^]	1.26	1.48	7 (2.20)	0.38	242 (76.48)	75 (23.66)	31 (9.80)	71 (22.40)
Italy	1.12	2.90	1346 (16.25)	0.66	7211 (87.05)	1073 (22.95)	171 (2.06)	4075 (49.19)
UK [□]	1.08	1.09	672 (14.49)	0.85	893 (19.26)	3738 (80.74)	2121 (45.74)	1102 (23.77)
Switzerland	0.91	1.26	35 (2.65)	1.97	1288 (97.32)	36 (2.72)	100 (7.58)	296 (22.44)
USA ^a	0.89	7.00	10 (1.78)	0.63	467 (83.10) [°]	95 (16.90)	226 (40.14)	222 (39.43)

Singapore	0.84	7.30	121 (34.03)	0.41	101 (28.44)	254 (71.56)	316 (89.01)	14 (3.94)
Romania	0.76	4.72	47 (4.47)	0.26	736 (70.03)	315 (29.97)	0 (0.00)	575 (54.71)
China ^b	0.72	0.00	12 (1.08)	0.55	476 (42.69)	639 (57.31)	40 (3.59)	228 (20.45)
Germany	0.69	0.70	77 (2.95)	1.43	2619 (99.28)	37 (0.72)	170 (6.84)	2140 (86.15)
Australia	0.61	0.69	115 (14.11)	0.63	674 (82.85)	140 (17.15)	78 (11.32)	391 (56.75)
Lithuania	0.49	0.46	13 (6.40)	0.13	138 (67.98)	25 (12.31)	4 (1.97)	42 (20.69)
Malaysia	0.39	0.09	223 (7.67)	0.13	134 (4.62)	2770 (95.38)	2161 (86.47)	226 (9.04)
Albania	0.00	39.14	5 (3.29)	0.52	128 (84.21)	24 (15.79)	6 (3.95)	23 (15.13)
Montenegro	0.00	14.84	1 (1.91)	1.11	104 (94.55)	6 (5.41)	18 (16.36)	42 (38.18)
Macedonia	0.00	10.4	0 (0.00)	0.60	22 (91.67)	2 (8.33)	1 (4.17)	6 (25.00)
Venezuela ^a	0.00	5.46	132 (20.69)	0.26	614 (96.23)	144 (3.77)	17 (2.65)	503 (78.60)
Serbia	0.00	4.06	175 (11.99)	0.35	1019 (69.80)	441 (30.20)	156 (10.68)	697 (47.74)
Nigeria	0.00	2.10	0 (0.00)	1.62	98 (23.62)	317 (76.39)	81 (18.54)	164 (37.53)
Cyprus ^a	0.00	0.49	4 (2.27)	0.71	165 (93.75)	170 (96.59)	7 (3.98)	78 (44.32)
Greece	0.00	0.44	1 (2.17)	0.80	40 (86.96)	6 (13.04)	8 (17.77)	19 (42.22)
p-value			<0.01	XXX	<0.01	<0.01	<0.01	<0.01

^aUSA/California; ^bChina/Hubei; ^cPakistan/Lahore; ^ddata calculated by the authors; ^eprivate, no differentiated;

^fIn these Countries other dental personnel participate to the survey; their replies are not reported in this table; ^gPositive/Hospitalized; ^hIn UK the survey involved only dentists working in primary care

Table 3. Precautionary measures taken by dentists who continued to provide face to face clinical care after the outbreak of COVID-19. Countries were ranked by the COVID-19 positive rate by country. In Figure 2 the COVID-19 positive rate by country and Dentists positive rates are reported.

		Before the arrival of the patients	In the operating room							
Country	Gross national income	Face-face activity	Phone triage	Appointment numbers reduced	Patients' health status	Surface disinfection	Eye protection equipment	N95/FP2 mask	TNT disposal gown	
						70% ethyl	0.5% sodium			

						alcohol	hypochlorite			
	range	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Egypt	<5,000	679 (73.40)	427 (62.88)	410 (60.38)	245 (36.08)	599 (88.22)	234 (34.46)	78 (11.49)	431 (63.48)	48 (7.07)
El Salvador	<5,000	351 (64.64)	169 (48.15)	231 (65.81)	283 (80.63)	230 (65.53)	229 (65.24)	329 (60.58)*	347 (98.86)	--
Georgia	<5,000	311 (98.11)	212 (68.17)	209 (67.20)	196 (63.02)	221 (71.06)	130 (13.53)	167 (53.70)	216 (69.45)	125 (40.19)
India	<5,000	3025 (92.65)	1213 (40.10)	1021 (33.75)	1457 (48.17)	2976 (98.38)	1851 (61.19)	1687 (55.77)	1368 (45.22)	2562 (84.69)
Nigeria	<5,000	399 (91.10)	50 (12.53)	135 (33.83)	105 (26.32)	89 (22.31)	83 (20.80)	155 (38.85)*	120 (30.08)	63 (15.79)
Pakistan ^c	<5,000	272 (64.00)	222 (81.62)	52 (19.12)	--	155 (56.99)	73 (26.84)	45 (16.54)	196 (72.06)	53 (19.49)
Tunisia	<5,000	745 (85.53)	466 (62.55)	588 (78.93)	595 (79.87)	277 (37.18)	448 (60.13)	720 (96.64)	570 (76.51)	--
Venezuela	<5,000	567 (88.87)	438 (77.25)	443 (78.13)	489 (86.24)	416 (73.37)	359 (63.32)	369 (65.08)	295 (52.03)	184 (32.45)
Albania	5,000/<10,000	206 (100.00)	21 (10.19)	22 (10.67)	22 (10.68)	22 (10.68)	11 (5.34)	27 (13.11)	22 (10.68)	10 (4.85)
Argentina	5,000/<10,000	836 (58.79)	434 (51.91)	502 (60.05)	472 (56.46)	405 (48.44)	399 (47.73)	557 (66.63)*	375 (44.86)	239 (28.59)§
Bosnia	5,000/<10,000	213 (100.00)	122 (57.28)	79 (37.09)	86 (40.38)	84 (39.44)	34 (15.96)	128 (60.09)	66 (30.99)	34 (15.96)
Brazil	5,000/<10,000	1270 (69.13)	684 (53.86)	1110 (87.14)	1059 (83.39)	1152 (90.71)	462 (36.38)	127 (100.00)	1270 (100.00)	696 (54.80)
Colombia	5,000/<10,000	5030 (95.58)	4137 (97.41)	4064 (95.69)	4064 (95.69)	2929 (68.97)	2329 (54.84)	4123 (97.08)	5030 (100.00)	1535 (36.14)
Macedonia	5,000/<10,000	22 (91.67)	6 (27.27)	2 (9.09)	2 (9.09)	2 (9.09)	--	7 (31.82)	5 (22.73)	3 (13.64)
Montenegro	5,000/<10,000	110 (100.00)	47 (42.73)	50 (45.45)	49 (44.55)	52(47.27)	25 (22.73)	36 (32.73)	11 (10.00)	6 (5.45)
Peru	5,000/<10,000	1229 (69.87)	1055 (59.97)	1337 (76.00)	1245 (70.77)	1041 (59.18)	1028 (58.44)	1229 (100.00)*	1229 (100.00)	1218 (69.24)§
Serbia	5,000/<10,000	742	553	652	665	725	363	742	284	76

	000	(50.8 2)	(74.53)	(87.87)	(89.6 2)	(97.71)	(48.92)	(100.0 0)	(38.2 7)	(10.24)
Chile	10,000 / <20,000	1589 (68.8 8)	921 (57.96)	1241 (78.10)	725 (45.6 3)	1086 (68.34)	573 (36.06)	1455 (91.57)	1291 (81.2 5)	280 (17.62)
China ^b	10,000 / <20,000	269 (14.6 4)	128 (47.58)	196 (72.86)	214 (79.5)	175 (65.06)	88 (32.71)	230 (85.50)	230 (85.5 0)	117 (43.49)
Cyprus	20,000 / 35,000	170 (96.5 9)	77 (45.29)	66 (38.82)	74 (43.5 3)	59 (34.71)	11 (6.47)	74 (43.53)	57 (33.5 3)	37 (21.76)
Greece	20,000 / 35,000	24 (100. 00)	19 (41.30)	25 (54.35)	19 (41.3 0)	16 (34.78)	3 (6.52)	22 (47.83)	14 (30.4 3)	11 (23.91)
Italy	20,000 / 35,000	7521 (90.7 9)	4226 (56.19)	3810 (50.66)	7452 (99.0 8)	5164 (68.66)	808 (10.74)	2552 (33.93)	6562 (87.2 5)	940 (12.50)
Lithuania	10,000 / <20,000	71 (34.8 0)	--	14 (19.72)	19 (26.7 6)	11 (15.49)	1 (1.41)	176 (86.70)	100 (49.2 6)	160 (78.82)
Malaysia	10,000 / <20,000	2183 (75.1 7)	483 (22.13)	1342 (61.48)	1383 (63.3 5)	1110 (50.85)	284 (13.01)	548 (25.10)	952 (43.6 1)	1280 (58.63)
Romania	10,000 / <20,000	1016 (96.6 7)	51 (5.02)	46 (4.53)	44 (4.33)	33 (3.25)	21 (2.07)	58 (5.71)	45 (4.43)	13 (1.28)
Russia	10,000/<2 0,000	946 (93.5 7)	118 (12.47)	561 (59.30)	620 (65.5 4)	335 (35.41)	116 (12.26)	793 (83.83)	312 (32.9 8)	91 (9.62)
Saudi Arabia	20,000 / 35,000	458 (53.4 4)	237 (51.75)	325 (70.96)	337 (73.5 8)	275 (60.04)	105 (22.93)	196 (42.79)	369 (80.5 7)	231 (50.44)
Spain	20,000 / 35,000	918 (28.5 3)	747 (81.37)	808 (88.02)	758 (82.5 7)	376 (40.96)	545 (59.37)	870 (94.77)	805 (87.6 9)	231 (25.16)
Australia	>35,000	718 (88.1 0)	506 (70.47)	448 (62.40)	631 (87.8 8)	525 (73.12)	152 (21.17)	243 (33.84)	191 (26.6 0)	86 (11.98)
Belgium	>35,000	732 (93.6 1)	596 (81.42)	567 (77.46)	573 (78.2 8)	523 (71.45)	116 (15.85)	429 (58.61)	570 (77.8 7)	106 (14.48)
Germany	>35,000	2510 (70.8 6)	2463 (98.13)	2485 (99.00)	2023 (80.6 0)	2414 (96.18)	2460 (98.01)	175 (6.97)	2319 (92.3 9)	841 (33.51)
Netherlands	>35,000	360 (82.2 6)	359 (99.72)	263 (73.06)	—	277 (76.94)	31 (8.61)	353 (94.89)	59 (16.3 9)	27 (7.50)
Singapore	>35,000	315 (88.7 3)	149 (47.30)	141 (44.76)	311 (98.7 3)	275 (87.30)	19 (6.05)	145 (46.03 **)	177 (56.1 9)	263 (83.49)**
Switzerland	>35,000	1243 (93.8)	1085 (87.29)	1083 (87.13)	1171 (94.2)	231 (18.58)	13 (1.05)	1154 (92.84)	696 (55.9)	57 (4.59)

		8))		1)				9)	
UK	>35,000	4264 (91.9 6)	2233 (52.37)	2146 (50.33)	2404 (56.3 8)	1337 (31.36)	1055 (24.74)	1553 (36.42)	262 (6.14)	120 (2.81)
USA ^a	>35,000	462 (82.2 1)	361 (78.14)	387 (83.77)	451 (97.6 2)	155 (33.55)	58 (12.55)	84 (18.18)	48 (10.3 9)	106 (22.94)
<i>p-value</i>	<0.01		<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01

^aUSA/California; ^bChina/Hubei; ^cPakistan/Lahore; ^{*}safety glasses or visor; ^{**}Reserve for aerosol generating procedures; [§] Water-repellent gown or disposable sterile microfiber gown; [°] Rotating instrument;

Table 4. Ordinal multinomial regression analysis using dentists' test-positive as dependent variable. For the multivariate analysis the forward stepwise method was used.

	Dentists positive rate		
Variables	β	Standard Error	p-value
GNI (>\$35,000 a year)	0.72	0.43	0.03
CPR (>10)	1.07	0.24	<0.01
ffp2/N95 (<25 %)	-1.46	0.43	<0.01
Face-to face	0.61	0.35	0.05
PPr (<25%)	0.32	0.21	0.59

GNI= Gross National Income per capita 2019; CPR=Country Positive Rate % to COVID-19;

ffp2= routine use of ffp2/N95 mask; Face-to-face= Face-to-face clinical activity;

PPr=Private Practice rate

Figure 1. COVID-19 positive rates by countries. In blue colour, is reported the COVID-19 positive rate of the population (%) at the time point when the survey was conducted, while in red the cumulative COVID-19 positive rate (%) among dental professionals since the beginning of the pandemic.