

The motivic Steenrod algebra in positive characteristic

Marc Hoyois, Shane Kelly, Paul Arne Østvær

August 1, 2013

Abstract

Let S be an essentially smooth scheme over a field and $\ell \neq \text{char } S$ a prime number. We show that the algebra of bistable operations in the mod ℓ motivic cohomology of smooth S -schemes is generated by the motivic Steenrod operations. This was previously proved by Voevodsky for S a field of characteristic zero. We follow Voevodsky's proof but remove its dependence on characteristic zero by using étale cohomology instead of topological realization and by replacing resolution of singularities with a theorem of Gabber on alterations.

Contents

1	Introduction	2
2	Background	4
2.1	Main categories and functors	4
2.2	Change of base scheme	8
2.3	Eilenberg–Mac Lane spaces and spectra	12
2.4	Operations in motivic cohomology	14
3	Main Results	15
3.1	The fundamental square	15
3.2	Motives of Eilenberg–Mac Lane spaces	16
3.3	Comparison with étale Steenrod operations	17
3.4	Proof of the main theorem	21
4	Commutativity of the fundamental square	23
4.1	The ℓ dh-topology	23
4.2	Descent for $\mathbf{MZ}_{(\ell)}$ -modules	26
4.3	End of the proof	31
5	Applications	32
5.1	The structure of the motivic Steenrod algebra and its dual	32
5.2	Modules over motivic cohomology	36

1 Introduction

This work is a continuation of [Voe10b]. The main results of *loc. cit.* are the computations, over fields of characteristic zero, of the motives of motivic Eilenberg–Mac Lane spaces and of the algebra of bistable operations in motivic cohomology. Here we generalize these computations to fields of positive characteristic, and further to essentially smooth schemes over fields.

Let k be a perfect field and $\ell \neq \text{char } k$ a prime number. We denote by \mathcal{M}^{**} the algebra of bistable operations in the motivic cohomology of smooth k -schemes with coefficients in \mathbf{Z}/ℓ introduced in [Voe03b, §2]. Examples of such operations are the reduced power operations $P^i \in \mathcal{M}^{2i(\ell-1), i(\ell-1)}$ for $i \geq 1$, the Bockstein operation $\beta \in \mathcal{M}^{1,0}$, and the operations given by multiplication by cohomology classes in $\mathrm{H}^{**}(k, \mathbf{Z}/\ell)$; we denote by \mathcal{A}^{**} the subalgebra of \mathcal{M}^{**} generated by these operations. The structure of the algebra \mathcal{A}^{**} was completely determined in [Voe03b]. On the other hand, motivic cohomology with coefficients in \mathbf{Z}/ℓ is represented, in the stable motivic homotopy category $\mathcal{SH}(\text{Sm}_k)$, by the motivic Eilenberg–Mac Lane spectrum \mathbf{MZ}/ℓ . It follows that any morphism $\mathbf{MZ}/\ell \rightarrow \Sigma^{p,q}\mathbf{MZ}/\ell$ in $\mathcal{SH}(\text{Sm}_k)$ induces a bistable operation of bidegree (p, q) , and in fact that every operation arises in this way. We therefore have the following situation:

$$\mathcal{A}^{**} \hookrightarrow \mathcal{M}^{**} \leftarrow \mathbf{MZ}/\ell^{**}\mathbf{MZ}/\ell.$$

Everything just described can be extended to essentially smooth schemes over fields (see Definition 2.9). Our main results are gathered in the following theorem.

Theorem 1.1: *Suppose S is a Noetherian scheme of finite Krull dimension that is essentially smooth over a field, and let $\ell \neq \text{char } S$ be a prime number.*

(1) *The operations*

$$\{\beta^{\epsilon_r} P^{i_r} \dots \beta^{\epsilon_1} P^{i_1} \beta^{\epsilon_0} \mid r \geq 0, i_j > 0, \epsilon_j \in \{0, 1\}, i_{j+1} \geq \ell i_j + \epsilon_j\}$$

*form a basis of \mathcal{M}^{**} as a left $\mathrm{H}^{**}(S, \mathbf{Z}/\ell)$ -module. In particular, $\mathcal{A}^{**} = \mathcal{M}^{**}$.*

(2) *The map $\mathbf{MZ}/\ell^{**}\mathbf{MZ}/\ell \rightarrow \mathcal{M}^{**}$ is an isomorphism.*

(3) *There is an equivalence of \mathbf{MZ}/ℓ -modules*

$$\mathbf{MZ}/\ell \wedge \mathbf{MZ}/\ell \simeq \bigvee_{\alpha} \Sigma^{p_{\alpha}, q_{\alpha}} \mathbf{MZ}/\ell$$

where (p_{α}, q_{α}) are the bidegrees of the operations in (1).

In characteristic zero Theorem 1.1 is an easy consequence of the results of [Voe03b] and [Voe10b]. We emphasize that the identification of the mod ℓ motivic cohomology of \mathbf{MZ}/ℓ with \mathcal{M}^{**} is not trivial and is a new result in positive characteristic. It implies that the mod

ℓ motivic cohomology of any motivic spectrum over S has a well-defined left \mathcal{A}^{**} -module structure.

One crucial step in the proof of Theorem 1.1 makes use of base change arguments. It reduces the proof to the case when S is a *perfect* field k .

Voevodsky’s proof of Theorem 1.1 (1) for fields of characteristic zero ([Voe10b, §3.4]) proceeds by realizing the inclusion $\mathcal{A}^{**} \hookrightarrow \mathcal{M}^{**}$ as the cohomology of a map of split proper Tate motives $\mathcal{M} \rightarrow \mathcal{A}$ [Voe10b, Lemma 3.50]. Since field extensions are conservative for split proper Tate motives [Voe10b, Corollary 2.70], this allows him to assume that k is the field of complex numbers. The calculations concluding the proof use the topological realization functor to compare \mathcal{M}^{**} with the classical Steenrod algebra. The main problems with attempting this proof in positive characteristic are:

- (a) There is no topological realization functor.
- (b) The only known technique for proving that motives of motivic Eilenberg-Mac Lane spaces are split proper Tate motives uses symmetric products, a construction which takes us out of the category Sm_k of smooth schemes. Resolution of singularities is then used to bridge the gap between the motivic homotopy category and its non-smooth analog, and likewise for the category of motives.

We solve these problems by replacing the topological realization functor with a cohomological étale realization functor (§3.3), and resolution of singularities with a theorem of Gabber on alterations [Ill, ILO12] (see Theorem 4.4). To apply the latter we use the ℓ dh-topology introduced in [Kel13]. The ℓ dh-topology is a refinement of the cdh-topology that allows finite flat surjective maps of degree prime to ℓ as coverings (see Definition 4.1). While the cdh-topology is designed to make schemes locally smooth in the presence of resolutions of singularities, Gabber’s theorem implies that every separated scheme of finite type over a perfect field admits an ℓ dh-covering by smooth schemes.

In §2 we first present what is relevant from the theory of presheaves with transfers. We proceed by explaining our base change arguments for motivic Eilenberg-Mac Lane spaces and operations in motivic cohomology. Theorem 1.1 is proved in §3 modulo the commutativity of the “fundamental square” formulated in Theorem 3.1, whose proof is deferred to §4. Our proof of the latter relies on the central result of [Kel13, §5] which states that $\mathbf{MZ}_{(\ell)}$ -modules satisfy ℓ dh-descent. For completeness, we repeat in §4.2, with minor modifications, the proof of this result from [Kel13, §5].

In §5 we gather some applications of our results and of Gabber’s theorem. As a direct consequence of Theorem 1.1, we show that for any motivic spectrum \mathbf{E} over S , $\mathbf{MZ}/\ell_{**}\mathbf{E}$ is a left comodule over the dual motivic Steenrod algebra \mathcal{A}_{**} . This fact is essential for adapting Adams spectral sequence techniques to motivic homotopy theory [DI10], [HKO11]. Secondly, we generalize the main result of [RØ08] by showing that Voevodsky’s category of motives

$\mathcal{DM}(\mathcal{S}m_k, \mathbf{Z}_{(\ell)})$ is equivalent to the homotopy category of $M\mathbf{Z}_{(\ell)}$ -modules when k is perfect and ℓ is invertible in k .

The results of this paper are further applied in [AF13] to study Euler classes and splittings of vector bundles, and in [Hoy13] to prove the Hopkins–Morel equivalence relating algebraic cobordism to motivic cohomology as well as the computation of the slices of Landweber exact motivic spectra and of the motivic sphere spectrum. Further applications are being developed in work on Morel’s π_1 -conjecture [OØ13] and on Milnor’s conjecture on quadratic forms [RØ13].

Acknowledgements: It is our pleasure to thank Andrei Suslin for useful suggestions during the preparation of this paper, and Chuck Weibel for valuable comments. The third author received partial support from the Leiv Eriksson mobility programme and RCN ES479962. He would also like to thank the MIT Mathematics Department for its kind hospitality.

2 Background

2.1 Main categories and functors

Let S be a Noetherian scheme of finite Krull dimension. We refer to such schemes as *base schemes* for short. In this section we set notation by reviewing the homotopy theories of motivic spaces, spaces with transfers, spectra, and spectra with transfers over S , and the various adjunctions between them. We refer to [RØ08, §2] and [Voe10b, §1.1] for more detailed expositions.

Let $\mathcal{S}ch_S$ be the category of separated schemes of finite type over S . A full subcategory \mathcal{C} of $\mathcal{S}ch_S$ is called an *admissible category* if the following hold [Voe10b, §0]:

- (1) $S \in \mathcal{C}$ and $\mathbf{A}_S^1 \in \mathcal{C}$.
- (2) If $X \in \mathcal{C}$ and $U \rightarrow X$ is étale of finite type, then $U \in \mathcal{C}$.
- (3) \mathcal{C} is closed under finite products and finite coproducts.

The most important example is the admissible category $\mathcal{S}m_S$ of separated smooth S -schemes of finite type, which we will simply call *smooth schemes*.

If \mathcal{C} is an admissible category, let $\mathcal{C}or(\mathcal{C})$ be the category with the same objects as \mathcal{C} but whose morphisms are the finite S -correspondences. In the notation of [CD12, §9.1.1], the set of morphisms from X to Y in $\mathcal{C}or(\mathcal{C})$ is the abelian group $c_0(X \times_S Y/X, \mathbf{Z})$.¹ This is a subgroup of the free abelian group generated by the closed integral subschemes Z of $X \times_S Y$ such that the induced morphism $Z \rightarrow X$ is finite and dominates an irreducible component of X . If X is regular, then $c_0(X \times_S Y/X, \mathbf{Z})$ is the entire free abelian group ([SV00b, Corollary 3.4.6]) but in general it is a proper subgroup ([SV00b, Example 3.4.7]). The category $\mathcal{C}or(\mathcal{C})$ is

¹In [SV00b] this group is denoted by $c_{equi}(X \times_S Y/X, 0)$; its definition is however incorrect for nonreduced schemes, see [CD12, §D.3].

additive, with direct sum given by disjoint union, and it has a symmetric monoidal structure such that the graph functor $\Gamma: \mathcal{C} \rightarrow \text{Cor}(\mathcal{C})$ is symmetric monoidal.

We denote by $s\text{Pre}^*(\mathcal{C})$ the category of pointed simplicial presheaves on \mathcal{C} . If R is any commutative ring, we denote by $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ the category of additive presheaves of simplicial R -modules on $\text{Cor}(\mathcal{C})$. By left Kan extension, we obtain an adjunction

$$R^{\text{tr}}: s\text{Pre}^*(\mathcal{C}) \rightleftarrows s\text{Pre}^{\text{tr}}(\mathcal{C}, R): u^{\text{tr}}, \quad (1)$$

where the right adjoint is the obvious forgetful functor. By the Dold–Kan correspondence, $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ is equivalent to the category of nonnegatively graded (homological) chain complexes of additive presheaves of R -modules on $\text{Cor}(\mathcal{C})$, but the simplicial description gives the functor R^{tr} a symmetric monoidal structure. In fact, the adjunction (1) is a symmetric monoidal Quillen adjunction for the usual projective model structures (i.e., the model structures where weak equivalences and fibrations are determined schemewise). The associated homotopy categories will be denoted by $\mathcal{H}^*(\mathcal{C})$ and $\mathcal{H}^{\text{tr}}(\mathcal{C}, R)$, respectively.

If τ is a Grothendieck topology in \mathcal{C} , consider the classes of maps

$$\begin{aligned} W_{\mathbf{A}^1} &= \{(\mathbf{A}^1 \times X)_+ \rightarrow X_+ \mid X \in \mathcal{C}\}, \\ W_\tau &= \{\mathcal{X}_+ \rightarrow X_+ \mid X \in \mathcal{C} \text{ and } \mathcal{X} \rightarrow X \text{ is a } \tau\text{-hypercover}\} \end{aligned}$$

in $s\text{Pre}^*(\mathcal{C})$. If E is any set of maps in $s\text{Pre}^*(\mathcal{C})$, we can consider the left Bousfield localization (see [Hir09]) of $s\text{Pre}^*(\mathcal{C})$ at E (resp. of $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ at $R^{\text{tr}}E$). Although the class W_τ is not essentially small, it is well-known that there exists a set $S \subset W_\tau$ such that an object in $\mathcal{H}^*(\mathcal{C})$ is W_τ -local if and only if it is S -local. It follows that the left Bousfield localization of $s\text{Pre}^*(\mathcal{C})$ at W_τ (resp. of $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ at $R^{\text{tr}}W_\tau$) also exists.

The left Bousfield localization of the projective structure on $s\text{Pre}^*(\mathcal{C})$ at $W_{\mathbf{A}^1} \cup W_{\text{Nis}}$ (resp. of $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ at $R^{\text{tr}}(W_{\mathbf{A}^1} \cup W_{\text{Nis}})$) is called the *motivic model structure* and its weak equivalences are the *motivic weak equivalences*. The associated motivic homotopy categories will be denoted by $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\mathcal{C})$ and $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\mathcal{C}, R)$, respectively. Occasionally we will consider intermediate localizations or other topologies, in which case we will use self-explanatory notations for the associated homotopy categories, e.g., $\mathcal{H}_{\text{ét}}^*(\mathcal{C})$, $\mathcal{H}_{\text{cdh}, \mathbf{A}^1}^{\text{tr}}(\mathcal{C}, R)$, etc.

In general, $\mathcal{H}_\tau^{\text{tr}}(\mathcal{C}, R)$ is *not* equivalent to the unstable derived category of the additive category of τ -sheaves of R -modules with transfers on \mathcal{C} . The following lemma provides necessary and sufficient conditions for this to be true.

Lemma 2.1: *Let $\mathcal{C} \subset \text{Sch}_S$ be an admissible category, R a commutative ring, and τ a topology on \mathcal{C} . The following assertions are equivalent.*

- (1) $u^{\text{tr}}R^{\text{tr}}: s\text{Pre}^*(\mathcal{C}) \rightarrow s\text{Pre}^*(\mathcal{C})$ sends W_τ to W_τ -local equivalences.
- (2) A morphism f in $s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ is an $R^{\text{tr}}W_\tau$ -local equivalence if and only if $u^{\text{tr}}(f)$ is an W_τ -local equivalence.

(3) *The square*

$$\begin{array}{ccc}
\mathcal{H}^{\mathrm{tr}}(\mathcal{C}, R) & \xrightarrow{\mathbf{L}\mathrm{id}} & \mathcal{H}_{\tau}^{\mathrm{tr}}(\mathcal{C}, R) \\
\mathbf{R}u^{\mathrm{tr}} \downarrow & & \downarrow \mathbf{R}u^{\mathrm{tr}} \\
\mathcal{H}^*(\mathcal{C}) & \xrightarrow{\mathbf{L}\mathrm{id}} & \mathcal{H}_{\tau}^*(\mathcal{C})
\end{array}$$

is commutative.

Proof. The implication (2) \Rightarrow (1) is obvious, and the equivalence (2) \Leftrightarrow (3) is essentially formal. Let us prove (1) \Rightarrow (2). Assume (1) and let f be a morphism in $s\mathrm{Pre}^{\mathrm{tr}}(\mathcal{C}, R)$. Since the functor $u^{\mathrm{tr}}: s\mathrm{Pre}^{\mathrm{tr}}(\mathcal{C}, R) \rightarrow s\mathrm{Pre}^*(\mathcal{C})$ takes values in the subcategory of simplicial radditive functors on \mathcal{C}_+ , [Voe10c, Theorem 4.20] shows that, if f is an $R^{\mathrm{tr}}W_{\tau}$ -local equivalence, then $u^{\mathrm{tr}}(f)$ is an W_{τ} -local equivalence. Conversely, suppose that $u^{\mathrm{tr}}(f)$ is an W_{τ} -local equivalence. Choose a map \tilde{f} between $R^{\mathrm{tr}}W_{\tau}$ -local objects in $s\mathrm{Pre}^{\mathrm{tr}}(\mathcal{C}, R)$ which is $R^{\mathrm{tr}}W_{\tau}$ -locally equivalent to f . Since we already proved that u^{tr} preserve equivalences, $u^{\mathrm{tr}}(\tilde{f})$ is W_{τ} -locally equivalent to $u^{\mathrm{tr}}(f)$, and since the source and target of $u^{\mathrm{tr}}(\tilde{f})$ are W_{τ} -local, $u^{\mathrm{tr}}(\tilde{f})$ is a projective equivalence. Therefore, \tilde{f} is a projective equivalence, and hence f is an $R^{\mathrm{tr}}W_{\tau}$ -local equivalence as desired. \square

Definition 2.2: Let $\mathcal{C} \subset \mathrm{Sch}_S$ be an admissible category and let R be a commutative ring. A topology τ on \mathcal{C} is *compatible with R -transfers* if the equivalent conditions of Lemma 2.1 are satisfied.

This definition agrees with [CD12, Definition 10.3.2]. By [CD12, Proposition 10.3.3], the Nisnevich topology is compatible with transfers on any admissible category \mathcal{C} , and by [CD12, Proposition 10.4.8 and Proposition 10.3.17], the cdh-topology is compatible with transfers on Sch_S . Both facts are also proved in [Voe10b, §1].

An object of $\mathcal{H}^*(\mathcal{C})$ is called \mathbf{A}^1 -*local* (resp. τ -*local*) if it is $W_{\mathbf{A}^1}$ -local (resp. W_{τ} -local) in the sense of [Hir09, Definition 3.1.4 (1) (a)]. Since S is Noetherian and of finite Krull dimension, a simplicial presheaf F is Nisnevich-local if and only if $F(\emptyset)$ is contractible and $F(Q)$ is homotopy cartesian for any cartesian square

$$Q = \begin{array}{ccc}
W & \longrightarrow & V \\
\downarrow & & \downarrow p \\
U & \xrightarrow{i} & X
\end{array}$$

in \mathcal{C} , where i is an open immersion, p is étale, and p induces an isomorphism $Z \times_X V \cong Z$ for some closed complement Z of $i(U)$; we will call such a square a *Nisnevich square*. By the general principles of left Bousfield localization, $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathcal{C})$ can be identified with the full subcategory of $\mathcal{H}^*(\mathcal{C})$ spanned by the \mathbf{A}^1 - and Nisnevich-local objects. Similarly, $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathcal{C}, R)$ is the full subcategory of $\mathcal{H}^{\mathrm{tr}}(\mathcal{C}, R)$ spanned by the \mathbf{A}^1 - and Nisnevich-local objects (i.e., those objects F such that $u^{\mathrm{tr}}F$ is \mathbf{A}^1 - and Nisnevich-local).

We note that the forgetful functor $u^{\text{tr}}: s\text{Pre}^{\text{tr}}(\mathcal{C}, R) \rightarrow s\text{Pre}^*(\mathcal{C})$ detects motivic weak equivalences. Indeed, u^{tr} sends $R^{\text{tr}}W_{\text{Nis}}$ to equivalences since the Nisnevich topology is compatible with R -transfers, and it is an easy exercise to show that it also sends $R^{\text{tr}}W_{\mathbf{A}^1}$ to \mathbf{A}^1 -local equivalences (see [Voe10b, Theorem 1.7]). As a result, we will also denote by u^{tr} the induced functor $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\mathcal{C}, R) \rightarrow \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\mathcal{C})$, right adjoint to $\mathbf{L}R^{\text{tr}}$.

An inclusion of admissible categories $i: \mathcal{C} \hookrightarrow \mathcal{D}$ induces Quillen adjunctions $(i_!, i^*)$ on the model categories $s\text{Pre}^*(-)$ and $s\text{Pre}^{\text{tr}}(-, R)$ with any of the model structures considered, where i^* is the restriction functor. These adjunctions commute with the adjunctions (1) since the right adjoints obviously commute. Since the functor $i_!$ is fully faithful, its derived functor $\mathbf{L}i_!$ will also be fully faithful provided that i^* preserve weak equivalences. It is clear that i^* always preserves \mathbf{A}^1 -local equivalences. If τ is a topology on \mathcal{D} , then $i^*: s\text{Pre}^*(\mathcal{D}) \rightarrow s\text{Pre}^*(\mathcal{C})$ preserves τ -local equivalences if and only if i is cocontinuous for τ [SGA72, III, Definition 2.1]. For $i^*: s\text{Pre}^{\text{tr}}(\mathcal{D}, R) \rightarrow s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ to preserve $R^{\text{tr}}W_{\tau}$ -local equivalences, we have the following criterion.

Lemma 2.3: *Let $i: \mathcal{C} \hookrightarrow \mathcal{D}$ be an inclusion of admissible subcategories of Sch_S , τ a topology on \mathcal{D} , and R a commutative ring. Suppose that i is cocontinuous for τ and that τ is compatible with R -transfers on \mathcal{D} . Then τ is also compatible with R -transfers on \mathcal{C} and the restriction functor $i^*: s\text{Pre}^{\text{tr}}(\mathcal{D}, R) \rightarrow s\text{Pre}^{\text{tr}}(\mathcal{C}, R)$ preserves $R^{\text{tr}}W_{\tau}$ -local equivalences.*

Proof. For the first claim we must verify that $u^{\text{tr}}R^{\text{tr}}: s\text{Pre}^*(\mathcal{C}) \rightarrow s\text{Pre}^*(\mathcal{C})$ sends τ -hypercovers in \mathcal{C} to τ -local equivalences. Since the functors $i_!$ are fully faithful, we have

$$u^{\text{tr}}R^{\text{tr}} \cong u^{\text{tr}}i^*i_!R^{\text{tr}} \cong i^*u^{\text{tr}}R^{\text{tr}}i_!.$$

Now $u^{\text{tr}}R^{\text{tr}}i_!$ sends τ -hypercovers in \mathcal{C} to τ -local equivalences on \mathcal{D} by the assumption that τ is compatible with R -transfers on \mathcal{D} , and since i is cocontinuous, i^* preserves τ -local equivalences. The second claim follows from the isomorphism $i^*u^{\text{tr}} \cong u^{\text{tr}}i^*$ and Lemma 2.1 (2). \square

The Nisnevich topology satisfies the hypotheses of Lemma 2.3 for any $i: \mathcal{C} \hookrightarrow \mathcal{D}$, so that the functors i^* preserve motivic weak equivalences. In particular, the derived functors $\mathbf{L}i_!: \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\mathcal{C}) \rightarrow \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\mathcal{D})$ and $\mathbf{L}i_!: \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\mathcal{C}, R) \rightarrow \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\mathcal{D}, R)$ are fully faithful.

We now turn to the stable theory. As $(\mathbf{G}_m, 1)$ is not projectively cofibrant, we choose a projectively cofibrant replacement \mathbf{G} (cf. [RØ08, Section 2.2]). If $p \geq q \geq 0$, define as usual

$$S^{p,q} = (S^1)^{\wedge p-q} \wedge \mathbf{G}^{\wedge q} \in s\text{Pre}^*(\mathcal{C}).$$

According to the general principles of [Hov01], the motivic model structures induce symmetric monoidal stable model structures on the category

$$\text{Spt}(\mathcal{C}) = \text{Sp}^{\Sigma}(s\text{Pre}^*(\mathcal{C}), S^{2,1})$$

of symmetric $S^{2,1}$ -spectra and on the category

$$\text{Spt}^{\text{tr}}(\mathcal{C}, R) = \text{Sp}^{\Sigma}(s\text{Pre}^{\text{tr}}(\mathcal{C}, R), R^{\text{tr}}S^{2,1})$$

of symmetric $R^{\text{tr}}S^{2,1}$ -spectra. The fibrant objects are the levelwise motivically fibrant spectra (E_0, E_1, \dots) such that the maps $E_i \rightarrow \Omega^{2,1}E_{i+1}$ adjoint to the bonding maps are motivic weak equivalences ([RØ08, Definition 14]). The associated homotopy categories will be denoted $\mathcal{SH}(\mathcal{C})$ and $\mathcal{DM}(\mathcal{C}, R)$, respectively.

There is a commutative diagram of symmetric monoidal Quillen adjunctions

$$\begin{array}{ccc}
s\text{Pre}^*(\mathcal{C}) & \begin{array}{c} \xleftarrow{R^{\text{tr}}} \\ \xrightarrow{u^{\text{tr}}} \end{array} & s\text{Pre}^{\text{tr}}(\mathcal{C}, R) \\
\Sigma^\infty \updownarrow \Omega^\infty & & \Sigma^\infty \updownarrow \Omega^\infty \\
\text{Spt}(\mathcal{C}) & \begin{array}{c} \xleftarrow{R^{\text{tr}}} \\ \xrightarrow{u^{\text{tr}}} \end{array} & \text{Spt}^{\text{tr}}(\mathcal{C}, R)
\end{array} \tag{2}$$

where the functors in the lower row are defined levelwise. We will denote by $\mathbf{1}$ the motivic sphere spectrum $\Sigma^\infty S_+$ in $\text{Spt}(\mathcal{C})$. When there is no danger of confusion, we will sometimes omit Σ^∞ from the notations.

We make the following observation which is lacking from [RØ08, §2].

Lemma 2.4: *The functor $u^{\text{tr}}: \text{Spt}^{\text{tr}}(\mathcal{C}, R) \rightarrow \text{Spt}(\mathcal{C})$ detects stable motivic weak equivalences.*

Proof. It detects levelwise motivic equivalences since $u^{\text{tr}}: s\text{Pre}^{\text{tr}}(\mathcal{C}, R) \rightarrow s\text{Pre}^*(\mathcal{C})$ detects motivic equivalences. Define a functor $Q: \text{Spt}(\mathcal{C}) \rightarrow \text{Spt}(\mathcal{C})$ by $(QE)_n = \text{Hom}(S^{2,1}, E_{n+1})$ (with action of Σ_n induced by that of Σ_{n+1}), and let $Q^\infty E = \text{colim}_{n \rightarrow \infty} Q^n E$. Similarly, let $Q^{\text{tr}}: \text{Spt}^{\text{tr}}(\mathcal{C}, R) \rightarrow \text{Spt}^{\text{tr}}(\mathcal{C}, R)$ be given by $(Q^{\text{tr}}E)_n = \text{Hom}(R^{\text{tr}}S^{2,1}, E_{n+1})$. Then a morphism f in $\text{Spt}(\mathcal{C})$ (resp. in $\text{Spt}^{\text{tr}}(\mathcal{C}, R)$) is a stable motivic equivalence if and only if $Q^\infty(f)$ (resp. $Q_{\text{tr}}^\infty(f)$) is a levelwise motivic equivalence. The proof is completed by noting that $u^{\text{tr}}Q_{\text{tr}}^\infty \cong Q^\infty u^{\text{tr}}$. \square

As a result, we simply denote by $u^{\text{tr}}: \mathcal{DM}(\mathcal{C}, R) \rightarrow \mathcal{SH}(\mathcal{C})$ the induced functor on homotopy categories.

2.2 Change of base scheme

In what follows we suppose given a class \mathcal{C} of morphisms in the category of base schemes which is stable under base change and such that, for any base scheme S , the category \mathcal{C}_S of S -schemes whose structure map is in \mathcal{C} forms an admissible category. Denote by $\mathcal{E}(-)$ any of the model categories $s\text{Pre}^*(-)$, $s\text{Pre}^{\text{tr}}(-, R)$, $\text{Spt}(-)$, $\text{Spt}^{\text{tr}}(-, R)$. As S varies, $\mathcal{E}(\mathcal{C}_S)$ is then a monoidal \mathcal{C} -fibered model category over the category of base schemes, in the sense of [CD12, §1.3.d]. In particular, a morphism of base schemes $f: T \rightarrow S$ induces a symmetric monoidal Quillen adjunction

$$f^*: \mathcal{E}(\mathcal{C}_S) \rightleftarrows \mathcal{E}(\mathcal{C}_T) : f_*$$

where f^* is induced by the base change functors $f^*: \mathcal{C}_S \rightarrow \mathcal{C}_T$ and $f^*: \text{Cor}(\mathcal{C}_S) \rightarrow \text{Cor}(\mathcal{C}_T)$. If f belongs to \mathcal{C}_S , there is also a Quillen adjunction

$$f_\# : \mathcal{E}(\mathcal{C}_T) \rightleftarrows \mathcal{E}(\mathcal{C}_S) : f^*$$

where $f_{\#}$ is induced by the forgetful functors $f_{\#}: \mathcal{C}_T \rightarrow \mathcal{C}_S$ and $f_{\#}: \text{Cor}(\mathcal{C}_T) \rightarrow \text{Cor}(\mathcal{C}_S)$. The \mathcal{C} -fibred structure of $\mathcal{E}(\mathcal{C})$ is moreover compatible with all the standard adjunctions between the various choices of \mathcal{E} , in the sense that the functors f^* and $f_{\#}$ commute with all left adjoints. For the adjunction $(R^{\text{tr}}, u^{\text{tr}})$, this follows from the commutativity of the squares

$$\begin{array}{ccc} \mathcal{C}_S & \xrightarrow{\Gamma} & \text{Cor}(\mathcal{C}_S) \\ f^* \downarrow & & \downarrow f^* \\ \mathcal{C}_T & \xrightarrow{\Gamma} & \text{Cor}(\mathcal{C}_T) \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathcal{C}_T & \xrightarrow{\Gamma} & \text{Cor}(\mathcal{C}_T) \\ f_{\#} \downarrow & & \downarrow f_{\#} \\ \mathcal{C}_S & \xrightarrow{\Gamma} & \text{Cor}(\mathcal{C}_S) \end{array}$$

from [CD12, Lemmas 9.3.3 and 9.3.7]. For the other adjunctions it is obvious.

Define a new class of morphisms $\widehat{\mathcal{C}}$ as follows. A map of base schemes $T \rightarrow S$ belongs to $\widehat{\mathcal{C}}$ if it is the limit of a cofiltered diagram (T_{α}) in \mathcal{C}_S whose transition maps $T_{\beta} \rightarrow T_{\alpha}$ are affine and dominant (the dominance condition is needed in the proof of Lemma 2.5 (2) below). If U is any T -scheme of finite type, then by [Gro66, Théorème 8.8.2] it is the limit of a diagram of schemes of finite type (U_{α}) over the diagram (T_{α}) . Moreover, if the morphism $U \rightarrow T$ is either

- separated,
- smooth or étale,
- an open immersion or a closed immersion,

then we can choose each $U_{\alpha} \rightarrow T_{\alpha}$ to have the same property (this follows from [Gro66, Proposition 8.10.4], [Gro67, Proposition 17.7.8], and [Gro66, Proposition 8.6.3], respectively). We shall assume that this is also true for \mathcal{C} -morphisms: we require any $U \in \mathcal{C}_T$ to be the limit of schemes U_{α} with $U_{\alpha} \in \mathcal{C}_{T_{\alpha}}$ (by the above, this holds if $\mathcal{C} = \text{Sm}$). It is then clear that $\widehat{\mathcal{C}}$ is closed under composition.

From now on we fix a $\widehat{\mathcal{C}}$ -morphism of base schemes $f: T \rightarrow S$, the cofiltered limit of \mathcal{C} -morphisms $f_{\alpha}: T_{\alpha} \rightarrow S$.

Lemma 2.5: *Consider the categories $s\text{Pre}^*$ and $s\text{Pre}^{\text{tr}}$ with the schemewise model structures. Let $X \in \mathcal{C}_T$ be a cofiltered limit of schemes X_{α} in $\mathcal{C}_{T_{\alpha}}$ and let $p \geq q \geq 0$.*

- (1) *For any $F \in s\text{Pre}^*(\mathcal{C}_S)$, the canonical map*

$$\text{hocolim}_{\alpha} \mathbf{R} \text{Map}(\Sigma^{p,q}(X_{\alpha})_+, \mathbf{L}f_{\alpha}^* F) \rightarrow \mathbf{R} \text{Map}(\Sigma^{p,q} X_+, \mathbf{L}f^* F)$$

is an equivalence.

- (2) *For any $F \in s\text{Pre}^{\text{tr}}(\mathcal{C}_S, R)$, the canonical map*

$$\text{hocolim}_{\alpha} \mathbf{R} \text{Map}(\mathbf{L}R^{\text{tr}} \Sigma^{p,q}(X_{\alpha})_+, \mathbf{L}f_{\alpha}^* F) \rightarrow \mathbf{R} \text{Map}(\mathbf{L}R^{\text{tr}} \Sigma^{p,q} X_+, \mathbf{L}f^* F)$$

is an equivalence.

Proof. Note that $\Sigma^{p,q}X_+$ can be obtained from copies of $(\mathbf{G}_m^{\times n} \times X)_+$ using finite homotopy colimits in a way which is compatible with base change. Since filtered homotopy colimits commute with finite homotopy limits, we can assume that $p = q = 0$. Both sides then preserve homotopy colimits in F , so we may further assume that $F = Y_+$ (resp. that $F = \mathbf{L}R^{\mathrm{tr}}Y_+$) where $Y \in \mathcal{C}_S$. Then $\mathbf{L}f^*F$ is represented by $Y \times_S T$ and the claim follows from [Gro66, Théorème 8.8.2] (resp. from [CD12, Proposition 9.3.9]). \square

We now make the following observations.

- Any trivial line bundle in \mathcal{C}_T is the cofiltered limit of trivial line bundles in \mathcal{C}_{T_α} .
- Any Nisnevich square in \mathcal{C}_T is the cofiltered limit of Nisnevich squares in \mathcal{C}_{T_α} .

The former is obvious. Any Nisnevich square in \mathcal{C}_T is a cofiltered limit of cartesian squares

$$\begin{array}{ccc} W_\alpha & \longrightarrow & V_\alpha \\ \downarrow & & \downarrow p_\alpha \\ U_\alpha & \xrightarrow{i_\alpha} & X_\alpha \end{array}$$

in \mathcal{C}_{T_α} , where i_α is an open immersion and p_α is étale. Let Z_α be the reduced complement of $i_\alpha(U_\alpha)$ in X_α . It remains to show that $Z_\alpha \times_{X_\alpha} V_\alpha \rightarrow Z_\alpha$ is eventually an isomorphism. By [Gro66, Corollaire 8.8.2.5], it suffices to show that $Z = \lim_\alpha Z_\alpha$ as closed subschemes of X . Now $\lim_\alpha Z_\alpha \cong Z_\alpha \times_{X_\alpha} X$ for large α , and so $\lim_\alpha Z_\alpha$ is a closed subscheme of X with the same support as Z . Moreover, it is reduced by [Gro66, Proposition 8.7.1], so it coincides with Z .

Lemma 2.6: *The functors $\mathbf{L}f^*: \mathcal{H}^*(\mathcal{C}_S) \rightarrow \mathcal{H}^*(\mathcal{C}_T)$ and $\mathbf{L}f^*: \mathcal{H}^{\mathrm{tr}}(\mathcal{C}_S, R) \rightarrow \mathcal{H}^{\mathrm{tr}}(\mathcal{C}_T, R)$ preserve \mathbf{A}^1 -local objects and Nisnevich-local objects.*

Proof. If f is in \mathcal{C} this follows from the existence of the Quillen left adjoint f_{\sharp} to f^* and the observation that f_{\sharp} sends trivial line bundles to trivial line bundles and Nisnevich squares to Nisnevich squares. Thus, each $\mathbf{L}f_\alpha^*$ preserves \mathbf{A}^1 -local objects and Nisnevich-local objects. Since any trivial line bundle (resp. Nisnevich square) over T is a cofiltered limit of trivial line bundles (resp. Nisnevich squares) over T_α , Lemma 2.5 shows that $\mathbf{L}f^*$ preserves \mathbf{A}^1 -local objects and Nisnevich-local objects in general. \square

Lemma 2.7: *Consider the categories $s\mathrm{Pre}^*$ and $s\mathrm{Pre}^{\mathrm{tr}}$ with the motivic model structures. Let $X \in \mathcal{C}_T$ be a cofiltered limit of schemes X_α in \mathcal{C}_{T_α} and let $p \geq q \geq 0$.*

- (1) *For any $F \in s\mathrm{Pre}^*(\mathcal{C}_S)$, the canonical map*

$$\mathrm{hocolim}_\alpha \mathbf{R}\mathrm{Map}(\Sigma^{p,q}(X_\alpha)_+, \mathbf{L}f_\alpha^*F) \rightarrow \mathbf{R}\mathrm{Map}(\Sigma^{p,q}X_+, \mathbf{L}f^*F)$$

is an equivalence.

(2) For any $F \in \text{sPre}^{\text{tr}}(\mathcal{C}_S, R)$, the canonical map

$$\text{hocolim}_\alpha \mathbf{R} \text{Map}(\mathbf{L}R^{\text{tr}}\Sigma^{p,q}(X_\alpha)_+, \mathbf{L}f_\alpha^*F) \rightarrow \mathbf{R} \text{Map}(\mathbf{L}R^{\text{tr}}\Sigma^{p,q}X_+, \mathbf{L}f^*F)$$

is an equivalence.

Proof. Combine Lemmas 2.5 and 2.6. \square

It is now easy to deduce a stable version of Lemma 2.7. Recall that objects in $\mathcal{SH}(\mathcal{C}_S)$ and $\mathcal{DM}(\mathcal{C}_S, R)$ can be modeled by Ω -spectra, i.e., sequences (E_0, E_1, \dots) of \mathbf{A}^1 - and Nisnevich-local objects E_i with motivic weak equivalences $E_i \simeq \Omega^{2,1}E_{i+1}$. If $E \in \mathcal{SH}(\mathcal{C}_S)$ is represented by the Ω -spectrum (E_0, E_1, \dots) and $X \in \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\mathcal{C}_S)$, we have

$$[\Sigma^{p,q}\Sigma^\infty X, E] \cong [\Sigma^{p+2r, q+r} X, E_r] \quad (3)$$

for any $r \geq 0$ such that $p + 2r \geq q + r \geq 0$, and similarly if $E \in \mathcal{DM}(\mathcal{C}_S, R)$ and $X \in \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\mathcal{C}_S, R)$.

If f is a \mathcal{C} -morphism, then the existence of the Quillen left adjoint f_{\sharp} to f^* shows that $\mathbf{L}f^*$ commutes with the unstable bigraded loop functors $\mathbf{R} \text{Hom}(S^{p,q}, -)$. An easy application of Lemma 2.7 then shows that this is still true if f is a $\widehat{\mathcal{C}}$ -morphism. Thus, we can explicitly describe the base change functors

$$\mathbf{L}f^* : \mathcal{SH}(\mathcal{C}_S) \rightarrow \mathcal{SH}(\mathcal{C}_T) \quad \text{and} \quad \mathbf{L}f^* : \mathcal{DM}(\mathcal{C}_S, R) \rightarrow \mathcal{DM}(\mathcal{C}_T, R)$$

by sending an Ω -spectrum (E_0, E_1, \dots) to the Ω -spectrum $(\mathbf{L}f^*E_0, \mathbf{L}f^*E_1, \dots)$. From (3) and Lemma 2.7 we obtain the following result.

Lemma 2.8: *Let $X \in \mathcal{C}_T$ be a cofiltered limit of schemes X_α in \mathcal{C}_{T_α} and let $p, q \in \mathbf{Z}$.*

- (1) For any $E \in \mathcal{SH}(\mathcal{C}_S)$, $[\Sigma^{p,q}\Sigma^\infty X_+, \mathbf{L}f^*E] \cong \text{colim}_\alpha [\Sigma^{p,q}\Sigma^\infty(X_\alpha)_+, \mathbf{L}f_\alpha^*E]$.
- (2) For any $E \in \mathcal{DM}(\mathcal{C}_S, R)$, $[\mathbf{L}R^{\text{tr}}\Sigma^{p,q}\Sigma^\infty X_+, \mathbf{L}f^*E] \cong \text{colim}_\alpha [\mathbf{L}R^{\text{tr}}\Sigma^{p,q}\Sigma^\infty(X_\alpha)_+, \mathbf{L}f_\alpha^*E]$.

A morphism in $\widehat{\text{Sm}}$ will be called essentially smooth:

Definition 2.9: A morphism of Noetherian schemes of finite Krull dimension $T \rightarrow S$ is said to be *essentially smooth* if it is the limit of a cofiltered diagram (T_α) of smooth S -schemes whose transition maps $T_\beta \rightarrow T_\alpha$ are affine and dominant.

For example, if $X \in \widehat{\text{Sm}}_S$ and $x \in X$, the local schemes $\text{Spec } \mathcal{O}_{X,x}$, $\text{Spec } \mathcal{O}_{X,x}^h$, and $\text{Spec } \mathcal{O}_{X,x}^{sh}$ (corresponding to the Zariski, Nisnevich, and étale topologies, respectively) are essentially smooth over S . The following lemma shows that an essentially smooth scheme over a field is in fact essentially smooth over a finite field \mathbf{F}_p or over \mathbf{Q} , and in particular over a perfect field. Together with the previous continuity results, this will allow us to extend many results from perfect fields to essentially smooth schemes over arbitrary fields.

Lemma 2.10: *Let k be a perfect field and L a field extension of k . Then the map $\text{Spec } L \rightarrow \text{Spec } k$ is essentially smooth.*

Proof. Since $\mathrm{Spec} L = \lim_K \mathrm{Spec} K$ where K ranges over the finitely generated field extensions of k contained in L , we may assume that $L = k(x_1, \dots, x_n)$ for some $x_i \in L$. Since k is perfect, $\mathrm{Spec} k[x_1, \dots, x_n]$ has a smooth dense open subset U ([Gro67, Corollaire 17.15.13]). Then $\mathrm{Spec} L$ is the cofiltered limit of the nonempty affine open subschemes of U . \square

2.3 Eilenberg–Mac Lane spaces and spectra

Let $\mathcal{C} \subset \mathrm{Sch}_S$ be an admissible category and let R be a commutative ring. Given $p \geq q \geq 0$ and an R -module A , the motivic Eilenberg–Mac Lane space $K(A(q), p)_{\mathcal{C}} \in \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathcal{C})$ is defined by

$$K(A(q), p)_{\mathcal{C}} = u^{\mathrm{tr}}(\mathbf{L}R^{\mathrm{tr}} S^{p,q} \otimes_{\mathbf{L}R}^{\mathbf{L}} A),$$

cf. [Voe10b, §3.2]. Note that this space does not depend on the ring R since $K(A(q), p)_{\mathcal{C}} = u^{\mathrm{tr}}(\mathbf{L}\mathbf{Z}^{\mathrm{tr}} S^{p,q} \otimes_{\mathbf{L}\mathbf{Z}}^{\mathbf{L}} A)$. The motivic Eilenberg–Mac Lane spectrum $\mathbf{M}A_{\mathcal{C}} \in \mathcal{SH}(\mathcal{C})$ is defined by

$$\mathbf{M}A_{\mathcal{C}} = u^{\mathrm{tr}}(\mathbf{L}R^{\mathrm{tr}} \mathbf{1} \otimes_{\mathbf{L}R}^{\mathbf{L}} A).$$

More explicitly, $\mathbf{M}A_{\mathcal{C}}$ is given by the sequence of spaces $K(A(n), 2n)_{\mathcal{C}}$ with bonding maps $\Sigma^{2,1} K(A(n), 2n)_{\mathcal{C}} \rightarrow K(A(n+1), 2n+2)_{\mathcal{C}}$. Note that $\mathbf{1} \in \mathrm{Spt}(\mathcal{C})$ is cofibrant and therefore $\mathbf{M}R_{\mathcal{C}}$ is equivalent to the commutative monoid $u^{\mathrm{tr}} R^{\mathrm{tr}} \mathbf{1}$ in the symmetric monoidal model category $\mathrm{Spt}(\mathcal{C})$. According to [RØ08, Proposition 38], there is a symmetric monoidal model category $\mathbf{M}R_{\mathcal{C}}\text{-mod}$ of $\mathbf{M}R_{\mathcal{C}}$ -modules. In addition, since $\mathbf{M}A_{\mathcal{C}} \simeq u^{\mathrm{tr}}(R^{\mathrm{tr}} \mathbf{1} \otimes_{\mathbf{L}R} \tilde{A})$ where $\tilde{A} \rightarrow A$ is a cofibrant replacement of A in $\mathrm{Spt}^{\mathrm{tr}}(\mathcal{C}, R)$, the object $\mathbf{M}A_{\mathcal{C}}$ has a canonical structure of $\mathbf{M}R_{\mathcal{C}}$ -module. When $\mathcal{C} = \mathrm{Sm}_S$, we will write $K(A(q), p)$ for $K(A(q), p)_{\mathcal{C}}$ and $\mathbf{M}A$ for $\mathbf{M}A_{\mathcal{C}}$.

The symmetric monoidal adjunction $(R^{\mathrm{tr}}, u^{\mathrm{tr}})$ between $\mathrm{Spt}(\mathcal{C})$ and $\mathrm{Spt}^{\mathrm{tr}}(\mathcal{C}, R)$ lifts to a symmetric monoidal adjunction

$$\Phi: \mathbf{M}R_{\mathcal{C}}\text{-mod} \rightleftarrows \mathrm{Spt}^{\mathrm{tr}}(\mathcal{C}, R) : \Psi$$

such that $\Phi(\mathbf{M}R_{\mathcal{C}} \wedge -) = R^{\mathrm{tr}}$. This is in fact a symmetric monoidal Quillen adjunction (see [RØ08, §2]). In §5.2 we will show that it is a Quillen equivalence when $\mathcal{C} = \mathrm{Sm}_S$ and S is the spectrum of a perfect field whose characteristic exponent is invertible in R . Here we recall a weaker result which holds over any base. Call an $\mathbf{M}R$ -module *cellular* if it is an iterated homotopy colimit of $\mathbf{M}R$ -modules of the form $\Sigma^{p,q} \mathbf{M}R$ with $p, q \in \mathbf{Z}$. Similarly, an object in $\mathcal{DM}(\mathrm{Sm}_S, R)$ is cellular if it is an iterated homotopy colimit of objects of the form $R^{\mathrm{tr}} \Sigma^{p,q} \mathbf{1}$ with $p, q \in \mathbf{Z}$.

Lemma 2.11: *The derived adjunction $(\mathbf{L}\Phi, \mathbf{R}\Psi)$ between $\mathrm{Ho}(\mathbf{M}R\text{-mod})$ and $\mathcal{DM}(\mathrm{Sm}_S, R)$ restricts to an equivalence between the full subcategories of cellular objects.*

Proof. The proof of [RØ08, Corollary 62] works with any ring R instead of \mathbf{Z} . \square

Let \mathcal{C} be a class of morphisms of schemes as in §2.2, and let $f: T \rightarrow S$ be a morphism of base schemes. For any $p \geq q \geq 0$ and any R -module A there is a canonical map

$$\mathbf{L}f^* K(A(q), p)_{\mathcal{C}_S} \rightarrow K(A(q), p)_{\mathcal{C}_T}, \quad (4)$$

adjoint to the composition

$$\begin{aligned} \mathbf{L}R^{\mathrm{tr}}\mathbf{L}f^*u^{\mathrm{tr}}(\mathbf{L}R^{\mathrm{tr}}S_S^{p,q} \otimes_R^{\mathbf{L}} A) &\simeq \mathbf{L}f^*\mathbf{L}R^{\mathrm{tr}}u^{\mathrm{tr}}(\mathbf{L}R^{\mathrm{tr}}S_S^{p,q} \otimes_R^{\mathbf{L}} A) \\ &\rightarrow \mathbf{L}f^*(\mathbf{L}R^{\mathrm{tr}}S_S^{p,q} \otimes_R^{\mathbf{L}} A) \simeq \mathbf{L}R^{\mathrm{tr}}\mathbf{L}f^*S_S^{p,q} \otimes_R^{\mathbf{L}} \mathbf{L}f^*A \simeq \mathbf{L}R^{\mathrm{tr}}S_T^{p,q} \otimes_R^{\mathbf{L}} A. \end{aligned}$$

Similarly, by applying (4) levelwise we obtain a canonical map

$$\mathbf{L}f^*MA_{\mathcal{C}_S} \rightarrow MA_{\mathcal{C}_T}. \quad (5)$$

Theorem 2.12: *Let U be a base scheme and let $f: T \rightarrow S$ a morphism in $\widehat{\mathcal{C}}_U$. Then (4) and (5) are equivalences.*

Proof. It suffices to show that (4) is an equivalence. We may clearly assume $S = U$, so that f itself is a $\widehat{\mathcal{C}}$ -morphism. It suffices to show that the canonical map

$$\mathbf{L}f^*u^{\mathrm{tr}} \rightarrow u^{\mathrm{tr}}\mathbf{L}f^* \quad (6)$$

is an equivalence. If f is in \mathcal{C} , then the functors f^* have Quillen left adjoints f_{\sharp} such that $\mathbf{L}f_{\sharp}\mathbf{L}R^{\mathrm{tr}} \simeq \mathbf{L}R^{\mathrm{tr}}\mathbf{L}f_{\sharp}$, and so (6) is an equivalence by adjunction. In the general case, let f be the cofiltered limit of \mathcal{C} -morphisms $f_{\alpha}: T_{\alpha} \rightarrow S$. Let $F \in \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathcal{C}_S, R)$. To show that $\mathbf{L}f^*u^{\mathrm{tr}}F \rightarrow u^{\mathrm{tr}}\mathbf{L}f^*F$ is an equivalence in $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathcal{C}_T)$, it suffices to show that for any $X \in \mathcal{C}_T$, the induced map

$$\mathbf{R}\mathrm{Map}(X_+, \mathbf{L}f^*u^{\mathrm{tr}}F) \rightarrow \mathbf{R}\mathrm{Map}(X_+, u^{\mathrm{tr}}\mathbf{L}f^*F) \simeq \mathbf{R}\mathrm{Map}(\mathbf{L}R^{\mathrm{tr}}X_+, \mathbf{L}f^*F)$$

is an equivalence. Write X as a cofiltered limit of schemes X_{α} in $\mathcal{C}_{T_{\alpha}}$. Then by Lemma 2.7, the above map is the homotopy colimit of the maps

$$\mathbf{R}\mathrm{Map}((X_{\alpha})_+, \mathbf{L}f_{\alpha}^*u^{\mathrm{tr}}F) \rightarrow \mathbf{R}\mathrm{Map}(\mathbf{L}R^{\mathrm{tr}}(X_{\alpha})_+, \mathbf{L}f_{\alpha}^*F),$$

which are equivalences since f_{α} is in \mathcal{C} . □

Now we specialize to the case when $\mathcal{C} = \mathrm{Sm}_S$ and S is essentially smooth over a field. If k is a field and X is a smooth k -scheme, the motivic cohomology groups $\mathbf{H}^{p,q}(X, A)$ are defined in [MVW06, Definition 3.4] for any abelian group A . By [MVW06, Proposition 3.8], these groups do not depend on the choice of the field k on which X is smooth. More generally, if X is an essentially smooth scheme over a field k , cofiltered limit of smooth k -schemes X_{α} , we define

$$\mathbf{H}^{p,q}(X, A) = \mathrm{colim}_{\alpha} \mathbf{H}^{p,q}(X_{\alpha}, A).$$

This definition does not depend on the choice of the diagram (X_{α}) since it is unique as a pro-object in the category of smooth k -schemes [Gro66, Corollaire 8.13.2]. Moreover, by [MVW06, Lemma 3.9], it is also independant of the choice of k . Theorem 4.24 in [Hoy13] shows that the motivic Eilenberg-Mac Lane spaces and spectra represent motivic cohomology:

Theorem 2.13: *Assume that S is essentially smooth over a field. Let A be an R -module and $X \in \mathbf{Sm}_S$. For any $p \geq q \geq 0$ and $r \geq s \geq 0$, there is a natural isomorphism*

$$\mathbf{H}^{p-r, q-s}(X, A) \cong [\Sigma^{r,s} X_+, K(A(q), p)]$$

and the canonical maps

$$K(A(q), p) \rightarrow \mathbf{R}\mathrm{Hom}(S^{r,s}, K(A(q+s), p+r)) \quad (7)$$

are equivalences. For any $p, q \in \mathbf{Z}$, there is a natural isomorphism

$$\mathbf{H}^{p,q}(X, A) \cong [\Sigma^\infty X_+, \Sigma^{p,q} \mathbf{M}A].$$

The following consequence of Theorem 2.13 summarizes the standard vanishing results for motivic cohomology, cf. [Hoy13, Corollary 4.26].

Corollary 2.14: *Assume that S is essentially smooth over a field. Let $X \in \mathbf{Sm}_S$ and $p, q \in \mathbf{Z}$ satisfy any of the following conditions:*

- (1) $q < 0$,
- (2) $p > q + d$,
- (3) $p > 2q$,

where d is the least integer such that X can be written as a cofiltered limit of smooth d -dimensional schemes over a field. Then, for any abelian group A , $[\Sigma^\infty X_+, \Sigma^{p,q} \mathbf{M}A] = 0$.

2.4 Operations in motivic cohomology

Let S be essentially smooth over a field. We fix a prime number $\ell \neq \mathrm{char} S$ and we abbreviate $K(\mathbf{Z}/\ell(n), 2n)$ to K_n and $\mathbf{M}\mathbf{Z}/\ell$ to \mathbf{M} . Denote by \mathcal{M}^{**} the algebra of bistable operations in the motivic cohomology of motivic spaces over S : an element $\phi \in \mathcal{M}^{p,q}$ is a collection of natural transformations

$$\phi_{r,s}: \tilde{\mathbf{H}}^{r,s}(-, \mathbf{Z}/\ell) \rightarrow \tilde{\mathbf{H}}^{r+p, s+q}(-, \mathbf{Z}/\ell), \quad r, s \in \mathbf{Z},$$

on $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathbf{Sm}_S)$ such that, under the bistability isomorphisms, $\phi_{r,s} \Sigma^{2,1} = \phi_{r-2, s-1}$. If $\iota_n \in \tilde{\mathbf{H}}^{2n, n}(K_n, \mathbf{Z}/\ell)$ is the tautological class, it is clear that we have an isomorphism

$$\mathcal{M}^{**} \cong \lim_{n \geq 0} \tilde{\mathbf{H}}^{*+2n, *+n}(K_n, \mathbf{Z}/\ell), \quad \phi \mapsto (\phi_{2n, n}(\iota_n))_n$$

(see [Voe03b, Proposition 2.7]). As a result, by Theorem 2.12, if S and T are both essentially smooth over a field, a map $f: T \rightarrow S$ induces a map of algebras

$$f^*: \mathcal{M}_S^{**} \rightarrow \mathcal{M}_T^{**}. \quad (8)$$

Note that any morphism $M \rightarrow \Sigma^{p,q}M$ in $\mathcal{SH}(\mathrm{Sm}_S)$ induces a bistable operation of bidegree (p, q) , which defines a canonical map

$$M^{**}M \rightarrow M^{**}. \quad (9)$$

In particular, the Bockstein map $M \rightarrow \Sigma^{1,0}M$ induces an operation $\beta \in \mathcal{M}^{1,0}$, and any cohomology class $\alpha \in H^{p,q}(S, \mathbf{Z}/\ell)$ defines an element of $\mathcal{M}^{p,q}$. We can also define the reduced power operations

$$P^i \in \mathcal{M}^{2i(\ell-1), i(\ell-1)}$$

as follows. If S is the spectrum of a perfect field, these are defined in [Voe03b, §9]. It is easy to see by inspecting their definition that if $f: \mathrm{Spec} k' \rightarrow \mathrm{Spec} k$ is an extension of perfect fields, $f^*(P^i) = P^i$. Thus, if $f: S \rightarrow \mathrm{Spec} k$ is essentially smooth where k is a perfect field, the operation $f^*(P^i) \in \mathcal{M}_S^{**}$ is independent of the choice of f .

Let $\mathcal{A}^{**} \subset \mathcal{M}^{**}$ be the subalgebra generated by

- the reduced power operations P^i for $i \geq 0$,
- the Bockstein β ,
- the operations $u \mapsto \alpha u$ for $\alpha \in H^{**}(S, \mathbf{Z}/\ell)$.

Clearly, the map (8) sends \mathcal{A}_S^{**} to \mathcal{A}_T^{**} , so that we have a commutative square of algebras:

$$\begin{array}{ccc} \mathcal{A}_S^{**} & \xrightarrow{f^*} & \mathcal{A}_T^{**} \\ \downarrow & & \downarrow \\ \mathcal{M}_S^{**} & \xrightarrow{f^*} & \mathcal{M}_T^{**}. \end{array} \quad (10)$$

3 Main Results

3.1 The fundamental square

The following theorem is our key technical result. It will be proved in §4.

Theorem 3.1: *Let k be a perfect field, $\ell \neq \mathrm{char} k$ a prime number, R a $\mathbf{Z}/(\ell)$ -algebra, and $i: \mathcal{C} \hookrightarrow \mathcal{D}$ an inclusion of admissible subcategories of Sch_k with $\mathcal{C} \subset \mathrm{Sm}_k$. Then the square*

$$\begin{array}{ccc} \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathcal{D}) & \xrightarrow{i^*} & \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^*(\mathcal{C}) \\ \mathbf{L}R^{\mathrm{tr}} \downarrow & & \downarrow \mathbf{L}R^{\mathrm{tr}} \\ \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathcal{D}, R) & \xrightarrow{i^*} & \mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathcal{C}, R) \end{array}$$

is commutative.

Commutativity of the fundamental square means the canonical natural transformation

$$\mathbf{L}R^{\mathrm{tr}}i^* \rightarrow i^*\mathbf{L}R^{\mathrm{tr}}$$

is an isomorphism. If k admits resolution of singularities in the sense of [FV00, Definition 3.4], this holds for any commutative ring R by [Voe10b, Theorem 1.21]. However, resolution of singularities is only known to hold for fields of characteristic zero.

3.2 Motives of Eilenberg–Mac Lane spaces

Let F be a field. Recall that a split proper Tate motive of weight $\geq n$ in $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathcal{C}, F)$ is a direct sum of object of the form $\mathbf{L}F^{\mathrm{tr}}S^{p,q}$ with $p \geq 2q$ and $q \geq n$ [Voe10b, Definition 2.60].

Theorem 3.2: *Let S be essentially smooth over a field of characteristic exponent c . Let A be a finitely generated $\mathbf{Z}[1/c]$ -module, F a field of characteristic $\neq c$, and $p \geq 2q \geq 0$. Then $\mathbf{L}F^{\mathrm{tr}}K(A(q), p)_{\mathrm{Sm}_S}$ is a split proper Tate motive of weight $\geq q$ in $\mathcal{H}_{\mathrm{Nis}, \mathbf{A}^1}^{\mathrm{tr}}(\mathrm{Sm}_S, F)$.*

Proof. We abbreviate $K(A(q), p)_{\mathcal{C}}$ to $K_{\mathcal{C}}$. By Theorem 2.12 we can assume that S is the spectrum of a perfect field. The theorem is obvious if $p = 0$, so assume further that $p > 0$. Let $\mathcal{D} \subset \mathrm{Sch}_S$ be the admissible subcategory of normal schemes and let $i: \mathrm{Sm}_S \hookrightarrow \mathcal{D}$ be the inclusion. Note that $i^*K_{\mathcal{D}} \simeq K_{\mathrm{Sm}_S}$. By Theorem 3.1, we have

$$i^*\mathbf{L}F^{\mathrm{tr}}K_{\mathcal{D}} \simeq \mathbf{L}F^{\mathrm{tr}}i^*K_{\mathcal{D}} \simeq \mathbf{L}F^{\mathrm{tr}}K_{\mathrm{Sm}_S}. \quad (11)$$

By [Voe10b, Corollary 3.28], $\mathbf{L}F^{\mathrm{tr}}K_{\mathcal{D}}$ is split proper Tate of weight $\geq q$. We conclude by noting that the adjunction $(\mathbf{L}i_!, i^*)$ restricts to an equivalence between the subcategories of split proper Tate motives of weight $\geq q$ since $\mathbf{L}i_!$ is fully faithful. \square

We now fix a prime number $\ell \neq \mathrm{char} S$ and we abbreviate $\mathbf{M}\mathbf{Z}/\ell$ to \mathbf{M} . The following corollary is assertion (2) of Theorem 1.1:

Corollary 3.3: *The canonical map $\mathcal{M}^{**}\mathbf{M} \rightarrow \mathcal{M}^{**}$ is an isomorphism.*

Proof. This map fits in the exact sequence

$$0 \rightarrow \lim^1 \tilde{\mathbf{H}}^{p-1+2n, q+n}(K_n, \mathbf{Z}/\ell) \rightarrow \mathbf{M}^{p, q}\mathbf{M} \rightarrow \lim \tilde{\mathbf{H}}^{p+2n, q+n}(K_n, \mathbf{Z}/\ell) \rightarrow 0,$$

and we must show that the \lim^1 term vanishes. By Theorem 3.2, $\mathbf{L}\mathbf{Z}/\ell^{\mathrm{tr}}K_n \simeq \Sigma^{2n, n}M_n$ where M_n is split proper Tate of weight ≥ 0 . All functors should be derived in the following computations. Using the standard adjunctions, we get

$$\begin{aligned} \tilde{\mathbf{H}}^{p-1+2n, q+n}(K_n, \mathbf{Z}/\ell) &\cong [\Sigma^\infty K_n, \Sigma^{p-1+2n, q+n}\mathbf{M}] \cong [\Sigma^\infty \mathbf{Z}/\ell^{\mathrm{tr}}K_n, \Sigma^{p-1+2n, q+n}\mathbf{Z}/\ell^{\mathrm{tr}}\mathbf{1}] \\ &\cong [\Sigma^{2n, n}\Sigma^\infty M_n, \Sigma^{p-1+2n, q+n}\mathbf{Z}/\ell^{\mathrm{tr}}\mathbf{1}] \cong [\Sigma^\infty M_n, \Sigma^{p-1, q}\mathbf{Z}/\ell^{\mathrm{tr}}\mathbf{1}]. \end{aligned}$$

To show that $\lim^1[\Sigma^\infty M_n, \Sigma^{p-1, q} \mathbf{Z}/\ell^{\text{tr}} \mathbf{1}] = 0$, it remains to show that the cofiber sequence

$$\bigoplus_{n \geq 0} \Sigma^\infty M_n \rightarrow \bigoplus_{n \geq 0} \Sigma^\infty M_n \rightarrow \text{hocolim}_{n \rightarrow \infty} \Sigma^\infty M_n$$

splits in $\mathcal{DM}(\mathfrak{Sm}_S, \mathbf{Z}/\ell)$. If S is the spectrum of a perfect field, this follows from [Voe10b, Corollary 2.71]. In general, let $f: S \rightarrow \text{Spec } k$ be essentially smooth where k is a perfect field. Then by Theorem 2.12, the above cofiber sequence is the image by f^* of the corresponding cofiber sequence over k , and hence it splits. \square

Corollary 3.4: $M \wedge M$ is equivalent to an M -module of the form $\bigvee_\alpha \Sigma^{p_\alpha, q_\alpha} M$.

Proof. By Theorem 3.2, $\mathbf{LZ}/\ell^{\text{tr}} K_n \simeq \Sigma^{2n, n} M_n$ where M_n is split proper Tate of weight ≥ 0 . By [Voe10b, Corollary 2.71] and Theorem 2.12, $\text{hocolim}_{n \rightarrow \infty} M_n$ is again a split proper Tate object of weight ≥ 0 , i.e., can be written in the form

$$\text{hocolim } M_n \simeq \bigoplus_\alpha \mathbf{LZ}/\ell^{\text{tr}} S^{p_\alpha, q_\alpha}$$

with $p_\alpha \geq 2q_\alpha \geq 0$. In the following computations, all functors must be appropriately derived. We have the equivalences

$$\begin{aligned} \mathbf{Z}/\ell^{\text{tr}} M &\simeq \mathbf{Z}/\ell^{\text{tr}} \text{colim } \Sigma^{-2n, -n} \Sigma^\infty K_n \simeq \text{colim } \Sigma^{-2n, -n} \Sigma^\infty \mathbf{Z}/\ell^{\text{tr}} K_n \\ &\simeq \text{colim } \Sigma^{-2n, -n} \Sigma^\infty \Sigma^{2n, n} M_n \simeq \text{colim } \Sigma^\infty M_n \simeq \Sigma^\infty \text{colim } M_n \\ &\simeq \Sigma^\infty \bigoplus_\alpha \mathbf{Z}/\ell^{\text{tr}} S^{p_\alpha, q_\alpha} \simeq \mathbf{Z}/\ell^{\text{tr}} \bigvee_\alpha \Sigma^\infty S^{p_\alpha, q_\alpha}. \end{aligned}$$

In particular, $\mathbf{Z}/\ell^{\text{tr}} M = \Phi(M \wedge M)$ is cellular. By Lemma 2.11, we obtain the equivalences

$$M \wedge M \simeq M \wedge \bigvee_\alpha \Sigma^\infty S^{p_\alpha, q_\alpha} \simeq \bigvee_\alpha \Sigma^{p_\alpha, q_\alpha} M. \quad \square$$

3.3 Comparison with étale Steenrod operations

In this paragraph, k is an algebraically closed field. We will denote by $a_{\text{ét}}$ the localization functors

$$a_{\text{ét}}: \mathcal{H}^*(\mathfrak{Sm}_k) \rightarrow \mathcal{H}_{\text{ét}}^*(\mathfrak{Sm}_k) \quad \text{and} \quad a_{\text{ét}}: \mathcal{D}(\mathfrak{Sm}_k) \rightarrow \mathcal{D}_{\text{ét}}(\mathfrak{Sm}_k).$$

Here $\mathcal{D}(\mathfrak{Sm}_k)$ is the homotopy category of chain complexes of presheaves of abelian groups on \mathfrak{Sm}_k , and $\mathcal{D}_{\text{ét}}(\mathfrak{Sm}_k)$ is the full subcategory spanned by chain complexes satisfying étale hyperdescent or, equivalently, the category obtained from $\mathcal{D}(\mathfrak{Sm}_k)$ by inverting maps inducing isomorphisms on étale homology sheaves.

The shifted Suslin–Voevodsky motivic complex $\mathbf{Z}(1)[1] \in \mathcal{D}_{\text{Nis}}(\mathfrak{Sm}_k)$ of weight one is quasi-isomorphic, as a chain complex of Nisnevich sheaves, to the presheaf represented by the multiplicative group scheme \mathbf{G}_m [MVW06, Theorem 4.1]. By assuming $1/m \in k$, the

Kummer short exact sequence of étale sheaves $0 \rightarrow \mu_m \rightarrow \mathbf{G}_m \xrightarrow{m} \mathbf{G}_m \rightarrow 0$ produces a quasi-isomorphism $a_{\text{ét}}\mathbf{Z}/m(1) \simeq \mu_m$, whence

$$a_{\text{ét}}\mathbf{Z}/m(q)[p] \simeq \mu_m^{\otimes q}[p] \quad (1/m \in k, \quad p, q \in \mathbf{Z}). \quad (12)$$

This equivalence induces for every pointed simplicial presheaf X on Sm_k a canonical map

$$\tilde{H}^{p,q}(X, \mathbf{Z}/m) = \tilde{H}_{\text{Nis}}^p(X, \mathbf{Z}/m(q)) \rightarrow \tilde{H}_{\text{ét}}^p(X, \mu_m^{\otimes q}) \quad (13)$$

from motivic to étale cohomology. Moreover, it is easy to show that (13) is compatible with cup products and the bigraded suspension isomorphisms. In étale cohomology the latter is the canonical isomorphism

$$\tilde{H}_{\text{ét}}^{p+2}(\mathbf{G}_m \wedge \Sigma X, \mu_n^{\otimes q+1}) \cong \tilde{H}_{\text{ét}}^p(X, \mu_n^{\otimes q}).$$

Lemma 3.5: *Let X be a pointed simplicial presheaf on Sm_k . The canonical map (13)*

$$\tilde{H}^{p,q}(X, \mathbf{Z}/\ell) \rightarrow \tilde{H}_{\text{ét}}^p(X, \mu_\ell^{\otimes q})$$

is an isomorphism when $p \leq q$.

Proof. The lemma holds for pointed simplicial smooth schemes by [Voe11, Theorem 6.17]. A standard argument implies that it automatically holds for all simplicial presheaves. In more details, consider the map

$$a_{\text{ét}}: \mathbf{R} \text{Map}(\mathbf{Z}X, \mathbf{Z}/\ell(q)[p]) \rightarrow \mathbf{R} \text{Map}(\mathbf{Z}X, \mu_\ell^{\otimes q}[p]) \quad (14)$$

between derived mapping spaces, which on π_0 gives the map of the lemma. If $X = Z_+$ for some $Z \in \text{Sm}_k$ and K is any simplicial set, applying the functor $[K, -]$ to (14) yields the instance of (13) with the pointed simplicial smooth scheme $(Z \times K)_+$. Hence, (14) is a weak equivalence for such X . Since an arbitrary simplicial presheaf is a homotopy colimit of representable presheaves and both sides of (14) transform homotopy colimits into homotopy limits, the general case follows. \square

Since mod ℓ motivic cohomology vanishes in negative weights, we obtain:

Corollary 3.6: *Let X be a pointed simplicial presheaf on Sm_k . Étale sheafification induces an isomorphism*

$$\tilde{H}^{**}(X, \mathbf{Z}/\ell)[\tau^{-1}] \cong \tilde{H}_{\text{ét}}^*(X, \mu_\ell^{\otimes *}),$$

where $\tau \in \mathbf{H}^{0,1}(\text{Spec } k, \mathbf{Z}/\ell) \cong \mu_\ell(k)$ is a primitive ℓ th root of unity.

Remark 3.7: In [Lev00] Levine constructs for $X \in \text{Sm}_k$ an isomorphism $\mathbf{H}^{**}(X, \mathbf{Z}/\ell)[\tau^{-1}] \cong \mathbf{H}_{\text{ét}}^*(X, \mu_\ell^{\otimes *})$. It is likely that this isomorphism is the same as that of Corollary 3.6 for representable presheaves.

When $p \geq q \geq 0$, the chain complex $\mathbf{Z}/m(q)[p] \in \mathcal{D}_{\text{Nis}}(\mathcal{S}m_k)$ is concentrated in non-negative degrees and its underlying simplicial presheaf (via the Dold–Kan correspondence) is, by definition, the motivic Eilenberg–Mac Lane space $K(\mathbf{Z}/m(q), p) \in \mathcal{H}_{\text{Nis}}^*(\mathcal{S}m_k)$ which represents the functor $\tilde{H}^{p,q}(-, \mathbf{Z}/m)$. The underlying simplicial presheaf of $\mu_m^{\otimes q}[p]$ is the Eilenberg–Mac Lane object $K(\mu_m^{\otimes q}, p) \in \mathcal{H}_{\text{ét}}^*(\mathcal{S}m_k)$ which represents $\tilde{H}_{\text{ét}}^p(-, \mu_m^{\otimes q})$. In view of the commutativity of the square

$$\begin{array}{ccc} \mathcal{D}_{\text{Nis}}^{\leq 0}(\mathcal{S}m_k) & \xrightarrow{a_{\text{ét}}} & \mathcal{D}_{\text{ét}}^{\leq 0}(\mathcal{S}m_k) \\ \downarrow & & \downarrow \\ \mathcal{H}_{\text{Nis}}^*(\mathcal{S}m_k) & \xrightarrow{a_{\text{ét}}} & \mathcal{H}_{\text{ét}}^*(\mathcal{S}m_k), \end{array}$$

where the vertical arrows are the forgetful functors, we may restate (12) as an equivalence

$$a_{\text{ét}}K(\mathbf{Z}/m(q), p) \simeq K(\mu_m^{\otimes q}, p) \quad (1/m \in k, p \geq q \geq 0) \quad (15)$$

in $\mathcal{H}_{\text{ét}}^*(\mathcal{S}m_k)$. Using the shorthands $K_n = K(\mathbf{Z}/\ell(n), 2n)$ and $K_n^{\text{ét}} = K(\mu_\ell^{\otimes n}, 2n)$, we can write $a_{\text{ét}}K_n \simeq K_n^{\text{ét}}$. Combining (15) and Corollary 3.6, we get:

Corollary 3.8: *Étale sheafification induces an isomorphism*

$$\tilde{H}^{**}(K_n, \mathbf{Z}/\ell)[\tau^{-1}] \cong \tilde{H}_{\text{ét}}^*(K_n^{\text{ét}}, \mu_\ell^{\otimes *}).$$

Let $\mathcal{M}_{\text{ét}}^{**}$ be the algebra of bistable étale cohomology operations on $\mathcal{H}_{\text{ét}}^*(\mathcal{S}m_k)$ with twisted μ_ℓ -coefficients. We then have a canonical isomorphism

$$\mathcal{M}_{\text{ét}}^{**} \cong \lim_{n \geq 0} \tilde{H}_{\text{ét}}^{*+2n}(K_n^{\text{ét}}, \mu_\ell^{\otimes *+n})$$

where the transition map

$$\tilde{H}_{\text{ét}}^{*+2n+2}(K_{n+1}^{\text{ét}}, \mu_\ell^{\otimes *+n+1}) \rightarrow \tilde{H}_{\text{ét}}^{*+2n}(K_n^{\text{ét}}, \mu_\ell^{\otimes *+n})$$

is induced by the canonical map $\mathbf{G}_m \wedge \Sigma K_n^{\text{ét}} \rightarrow K_{n+1}^{\text{ét}}$ and the suspension isomorphism

$$\tilde{H}_{\text{ét}}^{*+2n+2}(\mathbf{G}_m \wedge \Sigma K_n^{\text{ét}}, \mu_\ell^{\otimes *+n+1}) \cong \tilde{H}_{\text{ét}}^{*+2n}(K_n^{\text{ét}}, \mu_\ell^{\otimes *+n}).$$

We now use for the first time our hypothesis on k . Since k is algebraically closed, μ_ℓ is a constant sheaf and so $K_n^{\text{ét}}$ is equivalent to the constant simplicial presheaf with value $K(\mu_\ell(k)^{\otimes n}, 2n) \cong K(\mathbf{Z}/\ell, 2n)$. Moreover, if $\mathbf{R}\Gamma: \mathcal{H}_{\text{ét}}(\mathcal{S}m_k) \rightarrow \text{Ho}(\text{sSet})$ is the derived global section functor, with left adjoint c , then $\mathbf{R}\Gamma(F) \simeq F(\text{Spec } k)$ since $\text{Spec } k$ has no nontrivial étale hypercovers. Thus, the unit $\text{id} \rightarrow \mathbf{R}\Gamma \circ c$ is an isomorphism, i.e., c is fully faithful. It therefore induces an isomorphism

$$\tilde{H}^*(K(\mu_\ell(k)^{\otimes n}, 2n), A) \cong \tilde{H}_{\text{ét}}^*(K_n^{\text{ét}}, A)$$

for any abelian group A . In particular, there is a canonical isomorphism

$$\chi: \mathcal{A}^* \xrightarrow{\cong} \mathcal{M}_{\text{ét}}^{*,0}$$

where \mathcal{A}^* is the topological Steenrod algebra, and multiplication by τ induces isomorphisms $\mathcal{M}_{\text{ét}}^{*,i} \cong \mathcal{M}_{\text{ét}}^{*,i+1}$. If we define

$$P_{\text{ét}}^i = \tau^{i(\ell-1)} \chi(P^i) \in \mathcal{M}_{\text{ét}}^{2i(\ell-1), i(\ell-1)}$$

(note that $\tau^{\ell-1}$ is independent of the choice of τ), we obtain the following presentation of the algebra $\mathcal{M}_{\text{ét}}^{**}$: it is generated by $\tau^{\pm 1} \in \mathcal{M}_{\text{ét}}^{0, \pm 1}$, the Bockstein $\beta_{\text{ét}} \in \mathcal{M}_{\text{ét}}^{1,0}$, and the operations $P_{\text{ét}}^i \in \mathcal{M}_{\text{ét}}^{2i(\ell-1), i(\ell-1)}$ for $i \geq 1$; if ℓ is odd, the relations are the usual topological Adem relations, while if $\ell = 2$ (in which case $P^i = \text{Sq}^{2^i}$) we get the topological Adem relations with additional τ -multiples dictated by the second grading.

Since (13) is compatible with the suspension isomorphisms and $a_{\text{ét}}$ is a functor, it induces an algebra map

$$\phi: \mathcal{M}^{**} \rightarrow \mathcal{M}_{\text{ét}}^{**}.$$

On the other hand, comparing the motivic and étale Adem relations and sending generators to generators yields a well-defined algebra map

$$\psi: \mathcal{A}^{**} \rightarrow \mathcal{M}_{\text{ét}}^{**}$$

identifying \mathcal{A}^{**} with the subalgebra $\mathcal{M}_{\text{ét}}^{*, \geq 0}$.

Lemma 3.9: *The following triangle commutes:*

$$\begin{array}{ccc} \mathcal{A}^{**} & \hookrightarrow & \mathcal{M}^{**} \\ & \searrow \psi & \downarrow \phi \\ & & \mathcal{M}_{\text{ét}}^{**} \end{array}$$

Proof. It suffices to show that ϕ maps P^i to $P_{\text{ét}}^i$ and β to $\beta_{\text{ét}}$. Since the identification of $a_{\text{ét}} K_n$ with $K_n^{\text{ét}}$ is compatible with cup products, the motivic Cartan formula [Voe03b, Proposition 9.7] shows that the operations $\chi^{-1}(\tau^{-i(\ell-1)} \phi(P^i)) \in \mathcal{A}^*$ satisfy the axioms (1)–(5) of [Ste62, VI, §1] (if $\ell = 2$, it shows that the operations $\chi^{-1}(\tau^{-\lfloor i/2 \rfloor} \phi(\text{Sq}^i))$ satisfy the axioms (1)–(5) of [Ste62, I, §1]). Therefore, by [Ste62, VIII, Theorems 3.9 and 3.10] and the definition of $P_{\text{ét}}^i \in \mathcal{M}_{\text{ét}}^{**}$, $\phi(P^i) = P_{\text{ét}}^i$.

It remains to show that $\phi(\beta) = \beta_{\text{ét}}$. Let α denote the identification (15). The right

column in the diagram

$$\begin{array}{ccc}
\Omega a_{\acute{e}t}K(\mathbf{Z}/\ell^2(n), 2n+1) & \xrightarrow[\simeq]{\Omega\alpha} & \Omega K(\mu_{\ell^2}^{\otimes n}, 2n+1) \\
\downarrow & & \downarrow \\
a_{\acute{e}t}K(\mathbf{Z}/\ell(n), 2n) & \xrightarrow[\simeq]{\gamma} & K(\mu_{\ell}^{\otimes n}, 2n) \\
\downarrow a_{\acute{e}t}\beta & & \downarrow \beta \\
a_{\acute{e}t}K(\mathbf{Z}/\ell(n), 2n+1) & \xrightarrow[\simeq]{\alpha} & K(\mu_{\ell}^{\otimes n}, 2n+1) \\
\downarrow & & \downarrow \\
a_{\acute{e}t}K(\mathbf{Z}/\ell^2(n), 2n+1) & \xrightarrow[\simeq]{\alpha} & K(\mu_{\ell^2}^{\otimes n}, 2n+1)
\end{array}$$

is a fiber sequence, and so is the left column because $a_{\acute{e}t}$ preserves finite homotopy limits. Since the bottom square commutes, there exists an equivalence γ rendering the diagram commutative [Hov99, Proposition 6.3.5]. To show that $\phi(\beta) = \beta_{\acute{e}t}$, it suffices to show that $\gamma = \alpha$. Taking into account that the canonical equivalences

$$\Omega a_{\acute{e}t}K(\mathbf{Z}/\ell^2(n), 2n+1) \simeq a_{\acute{e}t}K(\mathbf{Z}/\ell^2(n), 2n) \quad \text{and} \quad \Omega K(\mu_{\ell^2}^{\otimes n}, 2n+1) \simeq K(\mu_{\ell^2}^{\otimes n}, 2n)$$

are compatible with α , we can identify the top square with:

$$\begin{array}{ccc}
a_{\acute{e}t}K(\mathbf{Z}/\ell^2(n), 2n) & \xrightarrow{\alpha} & K(\mu_{\ell^2}^{\otimes n}, 2n) \\
\downarrow \text{mod } \ell & & \downarrow \ell \\
a_{\acute{e}t}K(\mathbf{Z}/\ell(n), 2n) & \xrightarrow{\gamma} & K(\mu_{\ell}^{\otimes n}, 2n).
\end{array}$$

Here all objects are Eilenberg–Mac Lane objects of degree $2n$. The full subcategory of $\mathcal{H}_{\acute{e}t}^*(\mathcal{S}m_k)$ spanned by these objects is equivalent, via π_{2n} , to the category of étale sheaves of abelian groups on $\mathcal{S}m_k$ [Lur09, Proposition 7.2.2.12]. Therefore there exists at most one bottom horizontal map making the diagram commutative. Since this square commutes with α instead of γ , we must have $\gamma = \alpha$. \square

3.4 Proof of the main theorem

In this section we prove the remaining parts of Theorem 1.1, namely (1) and (3). Let S be an essentially smooth scheme of a field and $\ell \neq \text{char } S$ a prime number.

We first prove (1) when $S = \text{Spec } k$ for a perfect field k . We already know from the computation of \mathcal{A}^{**} in [Voe03b] that the claimed basis of \mathcal{M}^{**} is a basis of \mathcal{A}^{**} , so it will suffice to prove that $\mathcal{A}^{**} = \mathcal{M}^{**}$. Theorem 3.2 implies, cf. [Voe10b, Lemma 3.50], that there exist split proper Tate motives \mathcal{A} and \mathcal{M} whose cohomology agrees with \mathcal{A}^{**} and \mathcal{M}^{**} , respectively, and a map $\mathcal{M} \rightarrow \mathcal{A}$ which is invariant under change of perfect base field and

induces the inclusion $\mathcal{A}^{**} \hookrightarrow \mathcal{M}^{**}$. Thus, $\mathcal{A}^{**} = \mathcal{M}^{**}$ if and only if $\mathcal{M} \rightarrow \mathcal{A}$ is an equivalence. By [Voe10b, Corollary 2.70 (2)] we may assume that k is algebraically closed. In this case, $\mathcal{M}^{**} \cong \mathbf{Z}/\ell[\tau]$ where $\tau \in H^{0,1}(\mathrm{Spec} k, \mathbf{Z}/\ell) \cong \mu_\ell(k)$ is a primitive ℓ th root of unity.

We claim that (1) follows from the following statements:

- (i) $\mathcal{A}^{**}/\tau\mathcal{A}^{**} \rightarrow \mathcal{M}^{**}/\tau\mathcal{M}^{**}$ is injective.
- (ii) $\mathcal{A}^{**}[\tau^{-1}] \rightarrow \mathcal{M}^{**}[\tau^{-1}]$ is surjective.

Indeed, let $x \in \mathcal{M}^{**}$. By (ii), $\tau^n x$ belongs to \mathcal{A}^{**} for some $n \geq 0$. If $n > 0$, $\tau^n x$ becomes zero in $\mathcal{M}^{**}/\tau\mathcal{M}^{**}$. By (i) it is zero in $\mathcal{A}^{**}/\tau\mathcal{A}^{**}$. Thus, there exists $y \in \mathcal{A}^{**}$ such that $\tau^n x = \tau y$. Since \mathcal{M}^{**} is the cohomology of a split Tate object, it has no τ -torsion. Hence, $\tau^{n-1}x = y$ and so $\tau^{n-1}x$ belongs to \mathcal{A}^{**} . Induction on n implies $x \in \mathcal{A}^{**}$.

The injectivity of $\mathcal{A}^{**}/\tau\mathcal{A}^{**} \rightarrow \mathcal{M}^{**}/\tau\mathcal{M}^{**}$ is proved in [Voe10b, Proposition 3.56]. To finish the proof we show that $\mathcal{A}^{**}[\tau^{-1}] \rightarrow \mathcal{M}^{**}[\tau^{-1}]$ is surjective. By Lemma 3.9, there is a commutative diagram

$$\begin{array}{ccccc}
\mathcal{A}^{**}[\tau^{-1}] & \longrightarrow & \mathcal{M}^{**}[\tau^{-1}] & \longrightarrow & \tilde{H}^{*+2n, *+n}(K_n, \mathbf{Z}/\ell)[\tau^{-1}] \\
\searrow \psi & & \downarrow \phi & & \downarrow \simeq \\
& & \mathcal{M}_{\acute{e}t}^{**} & \longrightarrow & \tilde{H}_{\acute{e}t}^{*+2n}(K_n^{\acute{e}t}, \mu_\ell(k)^{\otimes *+n})
\end{array}$$

where the rightmost map is an isomorphism by Corollary 3.8. We are thus reduced to showing that $\mathcal{M}^{**}[\tau^{-1}] \rightarrow \mathcal{M}_{\acute{e}t}^{**}$ is injective. If $x = (x_0, x_1, \dots) \in \mathcal{M}^{**}$ maps trivially to $\mathcal{M}_{\acute{e}t}^{**}$ then x_n maps trivially to $\tilde{H}^{*+2n, *+n}(K_n, \mathbf{Z}/\ell)[\tau^{-1}]$ for all n . By Theorem 3.2, $\mathbf{L}\mathbf{Z}/\ell^{\mathrm{tr}}K_n$ is a split proper Tate motive and in particular $\tilde{H}^{*+2n, *+n}(K_n, \mathbf{Z}/\ell)$ has no τ -torsion. It follows that $x_n = 0$ for all n , whence $x = 0$. This concludes the proof of assertion (1) of Theorem 1.1 when the base is a perfect field.

We now turn to the proof of assertion (3). By Corollaries 3.3 and 3.4, we have

$$\mathcal{M}^{**} \cong [\mathbf{M}, \Sigma^{**}\mathbf{M}] \cong [\mathbf{M} \wedge \mathbf{M}, \Sigma^{**}\mathbf{M}]_{\mathbf{M}} \cong \left[\bigvee_{\alpha} \Sigma^{p_{\alpha}, q_{\alpha}} \mathbf{M}, \Sigma^{**}\mathbf{M} \right]_{\mathbf{M}} \cong \prod_{\alpha} \mathbf{M}^{*-p_{\alpha}, *-q_{\alpha}}. \quad (16)$$

To determine this family of bidegrees (p_{α}, q_{α}) , we can again assume, by Theorem 2.12, that S is the spectrum of an algebraically closed field, so that $\mathcal{M}^{**} \cong \mathbf{Z}/\ell[\tau]$. In this case we also know by part (1) that $\mathcal{M}^{**} \cong \bigoplus_{\gamma} \mathbf{M}^{*-r_{\gamma}, *-s_{\gamma}}$ where the family of bidegrees (r_{γ}, s_{γ}) is the desired one. In particular, $\mathcal{M}^{p,q}$ is finite for every $(p, q) \in \mathbf{Z}$, which shows that the product (16) is a direct sum. Thus, we find a bigraded isomorphism of free $\mathbf{Z}/\ell[\tau]$ -modules

$$\bigoplus_{\alpha} \Sigma^{-p_{\alpha}, -q_{\alpha}} \mathbf{Z}/\ell[\tau] \cong \bigoplus_{\gamma} \Sigma^{-r_{\gamma}, -s_{\gamma}} \mathbf{Z}/\ell[\tau],$$

(where $\Sigma^{i,j}$ indicates shifting of the bidegree), and it follows that the indexing families (p_{α}, q_{α}) and (r_{γ}, s_{γ}) coincide.

Finally, we prove (1) of Theorem 1.1 in general. Choose an essentially smooth morphism $f: S \rightarrow \text{Spec } k$ where k is a perfect field. By Corollary 2.14, for any $(p, q) \in \mathbf{Z}$, there are at most finitely many α such that $M_S^{p+p\alpha, q+q\alpha} \neq 0$. It follows that the product in (16) is a direct sum, and hence that the map

$$f^*: \mathcal{M}_k^{**} \rightarrow \mathcal{M}_S^{**}$$

induces an isomorphism of left M_S^{**} -modules $M_S^{**} \otimes_{M_k^{**}} M_k^{**} \cong M_S^{**}$. From the commutative square (10), we obtain $\mathcal{A}_S^{**} = \mathcal{M}_S^{**}$. The more precise statement of assertion (1) is automatic since $f^*: \mathcal{M}_k^{**} \rightarrow \mathcal{M}_S^{**}$ is an algebra map.

4 Commutativity of the fundamental square

In this section we prove Theorem 3.1. Throughout, the base scheme is a perfect field k of characteristic exponent c , and $\ell \neq c$ is a fixed prime number.

4.1 The ℓ dh-topology

We start with the definition of the ℓ dh-topology introduced in [Kel13]. The basic idea is to enlarge the cdh-topology of Suslin–Voevodsky [SV96] by including finite flat surjective maps of degree prime to ℓ .

We say that a family of maps $\{V_j \rightarrow X\}_{j \in J}$ is a *refinement* of another family $\{U_i \rightarrow X\}_{i \in I}$ if for each $j \in J$ there exists an $i_j \in I$ and a factorization $V_j \rightarrow U_{i_j} \rightarrow X$.

Definition 4.1: Let $\ell \neq \text{char } k$ be a prime number.

- (1) An *fps ℓ' -cover* (fini-plat-surjectif-premier-à- ℓ) is a finite flat surjective map $f: U \rightarrow X$ such that $f_*\mathcal{O}_U$ is a free \mathcal{O}_X -module of rank prime to ℓ .
- (2) An *ℓ dh-cover* is a finite family of maps of finite type $\{U_i \rightarrow X\}$ which admits a refinement $\{V'_j \rightarrow V_j \rightarrow X\}$ where $\{V_j \rightarrow X\}$ is a cdh-cover and each $V'_j \rightarrow V_j$ is an fps ℓ' -cover.

Definition 4.2: The *ℓ dh-topology* on Sch_k is the topology generated by the ℓ dh-covers. If $\mathcal{C} \subset \text{Sch}_k$ is a subcategory, the ℓ dh-topology on \mathcal{C} is the topology induced by the ℓ dh-topology on Sch_k .

Remark 4.3: The ℓ dh-topology is a “global” version of the topology of ℓ' -alterations appearing in [Ill] and [ILO12].

Theorem 4.4 (Gabber [Ill, Theorem 1.3], [ILO12, IX, Theorem 1.1]): *Let $X \in \text{Sch}_k$ and let $Z \subset X$ be a nowhere dense closed subset. There exists a map $f: Y \rightarrow X$ in Sch_k such that:*

- (1) *Y is smooth and quasi-projective over k .*

- (2) f is proper, surjective, and sends generic points to generic points.
- (3) For each generic point ξ of X there is a unique point η of Y over it, and $[k(\eta) : k(\xi)]$ is finite of degree prime to ℓ .
- (4) $f^{-1}(Z) \subset Y$ is a divisor with strict normal crossings.

We shall make use of the following formulation of the Raynaud–Gruson flattening theorem [RG71].

Theorem 4.5 ([SV00b, Theorem 2.2.2]): *Let $p : X \rightarrow S$ be a map of Noetherian schemes and U an open subscheme in S such that p is flat over U . Then there exists a closed subscheme Z in S such that $U \cap Z = \emptyset$, and the proper transform of X with respect to the blow-up $\text{Bl}_Z S \rightarrow S$ with center in Z is flat over S .*

Corollary 4.6: *For every $X \in \text{Sch}_k$ there exists an ℓ dh-cover $\{U_i \rightarrow X\}$ where each U_i is smooth and quasi-projective over k .*

Proof. The proof uses Noetherian induction. Suppose the result holds for all proper closed subschemes of X . We may assume that X is integral since the set of inclusions of irreducible components is a cdh-cover. Let $Y \rightarrow X$ be the map provided by Theorem 4.4, so that Y is smooth and quasi-projective over k . By Theorem 4.5, there exists a blowup with nowhere dense center $X' \rightarrow X$ such that the proper transform $Y' \rightarrow X'$ of $Y \rightarrow X$ is Zariski-locally an fps^{ℓ} -cover. If $Z \subset X$ is a proper closed subscheme such that $X' \rightarrow X$ is an isomorphism outside of Z , then $\{Z \rightarrow X, X' \rightarrow X\}$ is a cdh-cover. By the inductive hypothesis, there exists an ℓ dh-cover $\{Z_j \rightarrow Z\}_{j \in J}$ with each Z_j quasi-projective and smooth over k . We claim that $\{Z_j \rightarrow X\}_{j \in J} \cup \{Y \rightarrow X\}$ is an ℓ dh-cover. Indeed, it is refined by $\{Z_j \rightarrow X\}_{j \in J} \cup \{Y' \rightarrow X\}$ which is the composition of the cdh-cover $\{Z \rightarrow X, X' \rightarrow X\}$ with the ℓ dh-covers $\{Z_j \rightarrow Z\}_{j \in J}$ and $\{Y' \rightarrow X'\}$. \square

It is easy to show that any presheaf of $\mathbf{Z}_{(\ell)}$ -modules with transfers on Sch_k is an fps^{ℓ} -sheaf. In particular, such a presheaf is a cdh-sheaf if and only if it is an ℓ dh-sheaf. More generally, we have:

Theorem 4.7 ([Kel13, Theorem 3.4.17]): *Suppose F is a presheaf of $\mathbf{Z}_{(\ell)}$ -modules with transfers on Sch_k . Then for every $X \in \text{Sch}_k$, the canonical map*

$$H_{\text{cdh}}^n(X, a_{\text{cdh}}F) \rightarrow H_{\ell\text{dh}}^n(X, a_{\ell\text{dh}}F)$$

is an isomorphism for all $n \geq 0$.

Lemma 4.8: *Let \mathcal{C} be a small category, $\sigma \leq \tau$ Grothendieck topologies on \mathcal{C} , and F a presheaf of spectra on \mathcal{C} . Assume that:*

- (1) F is σ -local.

(2) every $X \in \mathcal{C}$ has finite σ -cohomological dimension.

(3) for every $X \in \mathcal{C}$ and every $p, q \in \mathbf{Z}$, the canonical map $H_\sigma^p(X, a_\sigma \pi_q F) \rightarrow H_\tau^p(X, a_\tau \pi_q F)$ is an isomorphism.

Then F is τ -local.

Proof. Denote by $\mathrm{Sp}(\mathcal{C})$ the triangulated category of presheaves of spectra on \mathcal{C} . It is endowed with a t -structure for which the truncation functors $\tau_{\leq n}$ and $\tau_{\geq n}$ are defined objectwise. For a topology ρ on \mathcal{C} , let $i_\rho: \mathrm{Sp}_\rho(\mathcal{C}) \hookrightarrow \mathrm{Sp}(\mathcal{C})$ be the inclusion of the triangulated subcategory of ρ -local presheaves and $a_\rho: \mathrm{Sp}(\mathcal{C}) \rightarrow \mathrm{Sp}_\rho(\mathcal{C})$ its left adjoint. The subcategory $\mathrm{Sp}_\rho(\mathcal{C})$ inherits a t -structure whose truncation functors are $a_\rho \tau_{\leq n} i_\rho$ and $a_\rho \tau_{\geq n} i_\rho$. Moreover, the functors a_ρ and i_ρ are both left t -exact (see [Lur11, Remark 1.9]).

By (1), we have $F \in \mathrm{Sp}_\sigma(\mathcal{C})$. We first prove the lemma under the assumption that F is bounded below for the t -structure on $\mathrm{Sp}_\sigma(\mathcal{C})$, i.e., that $F \simeq a_\sigma \tau_{\geq k} F$ for some $k \in \mathbf{Z}$. By (2), the t -structure on $\mathrm{Sp}_\sigma(\mathcal{C})$ is left complete (see [Jar97, §6.1]), so that $F \simeq \mathrm{holim}_{n \rightarrow \infty} a_\sigma \tau_{\leq n} F$. Since the inclusion $\mathrm{Sp}_\tau(\mathcal{C}) \hookrightarrow \mathrm{Sp}(\mathcal{C})$ preserves homotopy limits, it suffices to show that $a_\sigma \tau_{\leq n} F$ is τ -local for all $n \in \mathbf{Z}$. Using the fiber sequences

$$(\pi_n F)[n] \rightarrow \tau_{\leq n} F \rightarrow \tau_{\leq n-1} F$$

and the fact that $a_\sigma \tau_{\leq n} F = 0$ for $n < k$, we are reduced to proving that $a_\sigma((\pi_n F)[n])$ is τ -local for all $n \geq k$, or equivalently that the canonical map $a_\sigma((\pi_n F)[n]) \rightarrow a_\tau((\pi_n F)[n])$ induces an isomorphism on presheaves of homotopy groups. By definition of cohomology, we have

$$\pi_* a_\sigma((\pi_n F)[n]) = H_\sigma^{n-*}(-, a_\sigma \pi_n F),$$

so the desired result holds by (3).

For a general F , the previous proof applies to the presheaves of spectra $a_\sigma \tau_{\geq k} F$ for all $k \in \mathbf{Z}$ and shows that

$$a_\sigma \tau_{\geq k} F \simeq a_\tau \tau_{\geq k} F.$$

To complete the proof we observe that, for *any* topology ρ on \mathcal{C} , the canonical map

$$\mathrm{colim}_{k \rightarrow -\infty} i_\rho a_\rho \tau_{\geq k} F \rightarrow i_\rho a_\rho F$$

is an equivalence of presheaves of spectra. Its cofiber is $\mathrm{colim}_{k \rightarrow -\infty} i_\rho a_\rho \tau_{< k} F$. Since both i_ρ and a_ρ are left t -exact, $i_\rho a_\rho \tau_{< k} F$ is a presheaf of $(k-1)$ -truncated spectra, and hence $\mathrm{colim}_{k \rightarrow -\infty} i_\rho a_\rho \tau_{< k} F$ is a presheaf of spectra which are $(k-1)$ -truncated for all $k \in \mathbf{Z}$, i.e., it is contractible. \square

Remark 4.9: We sketch a different proof of Lemma 4.8 assuming that the topologies σ and τ have enough points, which holds for all the topologies we consider. The cosimplicial Godement resolutions give rise for every $X \in \mathcal{C}$ to spectral sequences $\{\sigma E_r^{**}\}_{r \geq 1}$ and $\{\tau E_r^{**}\}_{r \geq 1}$ with

$$\sigma E_2^{p,q} \cong H_\sigma^p(X, a_\sigma \pi_q F) \quad \text{and} \quad \tau E_2^{p,q} \cong H_\tau^p(X, a_\tau \pi_q F),$$

and to a morphism of spectral sequences from the former to the latter, which by (3) is an isomorphism starting from the second page. By (2), ${}_{\sigma}E_2^{p,*} = 0$ for $p \gg 0$, so both spectral sequences stabilize after finitely many steps. It follows from the second part of [Tho85, Proposition 5.47] that ${}_{\rho}E_{\infty}^{**}$ is the associated graded of a complete, Hausdorff, and (trivially) exhaustive filtration on $\pi_*(a_{\rho}F)(X)$, for $\rho = \sigma, \tau$. This shows that the canonical map $\pi_*(a_{\sigma}F)(X) \rightarrow \pi_*(a_{\tau}F)(X)$ is an isomorphism for all $X \in \mathcal{C}$, and hence that $a_{\sigma}F \simeq a_{\tau}F$.

Proposition 4.10: *Let R be a $\mathbf{Z}_{(\ell)}$ -algebra.*

- (1) *The ℓ dh-topology on Sch_k is compatible with R -transfers.*
- (2) *The ℓ dh-topology on Sm_k is compatible with R -transfers.*
- (3) *The functor $i^*: s\mathrm{Pre}^{\mathrm{tr}}(\mathrm{Sch}_k, R) \rightarrow s\mathrm{Pre}^{\mathrm{tr}}(\mathrm{Sm}_k, R)$ preserves $R^{\mathrm{tr}}W_{\ell\mathrm{dh}}$ -local equivalences.*

Proof. To prove (1) we must show that, for any ℓ dh-hypercover $f \in W_{\ell\mathrm{dh}}$, $u^{\mathrm{tr}}R^{\mathrm{tr}}(f)$ is an ℓ dh-local equivalence; we will show that it is in fact a cdh-local equivalence. Since cdh is compatible with transfers on Sch_k [Voe10b, Lemma 1.24], we know from Lemma 2.1 that $u^{\mathrm{tr}}R^{\mathrm{tr}}(f)$ is a cdh-local equivalence if (and only if) $R^{\mathrm{tr}}(f)$ is an $R^{\mathrm{tr}}W_{\mathrm{cdh}}$ -local equivalence, and by assumption it is an $R^{\mathrm{tr}}W_{\ell\mathrm{dh}}$ -equivalence. By adjunction, it will suffice to show that if $F \in s\mathrm{Pre}^{\mathrm{tr}}(\mathrm{Sch}_k, R)$ is $R^{\mathrm{tr}}W_{\mathrm{cdh}}$ -local, then F is in fact $R^{\mathrm{tr}}W_{\ell\mathrm{dh}}$ -local or, equivalently, $u^{\mathrm{tr}}(F)$ is ℓ dh-local.

We have a canonical identification of presheaves $\pi_q u^{\mathrm{tr}}(F) \cong u^{\mathrm{tr}}\pi_q(F)$ and since the latter have a structure of presheaves with transfers, it follows from Theorem 4.7 that

$$H_{\mathrm{cdh}}^p(X, a_{\mathrm{cdh}}\pi_q u^{\mathrm{tr}}(F)) \cong H_{\ell\mathrm{dh}}^p(X, a_{\ell\mathrm{dh}}\pi_q u^{\mathrm{tr}}(F)).$$

Since the cdh-topology is cohomologically bounded [SV00a], Lemma 4.8 shows that $u^{\mathrm{tr}}(F)$ is ℓ dh-local (to apply the lemma, we view simplicial abelian groups as connective spectra). This proves (1).

By Corollary 4.6, $i: \mathrm{Sm}_k \hookrightarrow \mathrm{Sch}_k$ is cocontinuous for the ℓ dh-topology. Assertions (2) and (3) now follow from (1) by virtue of Lemma 2.3. \square

4.2 Descent for $\mathbf{MZ}_{(\ell)}$ -modules

In this section we review the proof from [Kel13, §5] that $\mathbf{MZ}_{(\ell)}$ -modules are ℓ dh-local (see Definition 4.23), which is a key ingredient in the proof of Theorem 3.1. The argument can be summarized as follows. We know from Ayoub’s proper base change theorem that motivic spectra are cdh-local. The notion of “structure of traces” (Definition 4.11) is designed to bridge the gap between cdh-descent and ℓ dh-descent. A first step is therefore to show that the motivic cohomology spectrum $\mathbf{MZ}_{(\ell)} \in \mathcal{SH}(\mathrm{Sm}_k)$ has a structure of traces. Roughly speaking, this follows from the facts that homotopy algebraic K -theory \mathbf{KGL} has a structure of traces and that $\mathbf{MZ}_{(\ell)}$ is the zeroth slice of $\mathbf{KGL}_{(\ell)}$.

We first recall the notion of a structure of traces introduced in [Kel13].

Definition 4.11: Let $\mathcal{E}: \text{Sch}_k \rightarrow \text{Cat}$ be a 2-functor (or equivalently, a cofibered category over Sch_k) with values in additive categories and F an additive section of \mathcal{E} (i.e., a section of the associated cofibered category such that $F_{X \amalg Y} \cong (i_1)_* F_X \oplus (i_2)_* F_Y$). A *structure of traces* on F is a family of maps $\text{Tr}_f: f_* F_Y \rightarrow F_X$, defined for every finite flat surjective map $f: Y \rightarrow X$ in Sch_k , subject to the following coherence conditions (where f and g are finite flat surjective maps).

- (Additivity) $\text{Tr}_{f \amalg g} = (i_1)_* \text{Tr}_f \oplus (i_2)_* \text{Tr}_g$.
- (Functoriality) $\text{Tr}_{\text{id}_X} = \text{id}_{F_X}$, and if $g: Z \rightarrow Y$ and $f: Y \rightarrow X$, $\text{Tr}_{f \circ g} = \text{Tr}_f \circ f_* \text{Tr}_g$.
- (Base change) For every cartesian square

$$\begin{array}{ccc} W & \xrightarrow{g} & Z \\ q \downarrow & & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

in Sch_k , we have $F_p \circ \text{Tr}_f = p_* \text{Tr}_g \circ f_* F_q$.

- (Degree) If f is globally free of degree d (i.e., $f_* \mathcal{O}_Y \cong \mathcal{O}_X^d$), then $\text{Tr}_f \circ F_f = d \cdot \text{id}_{F_X}$.

When \mathcal{E} is the constant functor with value Ab , this definition specializes to the notion of *presheaf with traces* on Sch_k . The other example that we will use is the 2-functor $X \mapsto \mathcal{SH}(\text{Sm}_X)$ on Sch_k , which takes values in triangulated categories. This is a stable homotopy 2-functor in the sense of [Ayo07]. In this context, we will simply write f_* , f^* , and f_{\sharp} for the *derived* functors $\mathbf{R}f_*$, $\mathbf{L}f^*$, $\mathbf{L}f_{\sharp}$ associated to a change of base scheme f (see §2.2).

Suppose that the cofibered category \mathcal{E} is bifibered, i.e., that for every $f: Y \rightarrow X$ in Sch_k the functor $f_*: \mathcal{E}(Y) \rightarrow \mathcal{E}(X)$ has a left adjoint f^* . Then, to any object $E \in \mathcal{E}(\text{Spec } k)$, we can associate a section $X \mapsto E_X$ of \mathcal{E} as follows: if $a: X \rightarrow \text{Spec } k$ is the structure map of $X \in \text{Sch}_k$, then $E_X = a^* E$, and given $f: Y \rightarrow X$, the map $E_X \rightarrow f_* E_Y$ is the unit of the adjunction (f^*, f_*) . Under some mild assumptions on \mathcal{E} (for example, axioms 1 to 3 in [Ayo07, §1.4.1]), this is an additive section of \mathcal{E} . A *structure of traces* on $E \in \mathcal{E}(\text{Spec } k)$ is a structure of traces on the corresponding section $X \mapsto E_X$.

Remark 4.12: When $E = \text{MA} \in \mathcal{SH}(\text{Sm}_k)$ there is a conflict of notations in that, if X is not smooth over k , the object E_X is not known to coincide with the motivic Eilenberg–Mac Lane spectrum MA_X defined in §2.3. When we speak of a structure of traces on MA , we mean a structure of traces on the cartesian section $(a: X \rightarrow \text{Spec } k) \mapsto a^* \text{MA}_k$, *not* on the section $X \mapsto \text{MA}_X$.

Remark 4.13: If $\phi: \mathcal{E}_1 \rightarrow \mathcal{E}_2$ is a 2-natural transformation and F is an additive section of \mathcal{E}_1 , it is clear that a structure of traces on F induces a structure of traces on ϕF . For example, if F is a section of $X \mapsto \mathcal{SH}(\text{Sm}_X)$ with a structure of traces, then for any $E \in \mathcal{SH}(\text{Spec } k)$, $X \mapsto [E_X, F_X]$ is a presheaf with traces on Sch_k .

We recall two important results on traces from [Kel13]. The first one relates traces to transfers:

Theorem 4.14 ([Kel13, Theorem 3.8.1 (3)]): *Suppose F is a presheaf of $\mathbf{Z}(\ell)$ -modules with traces on Sch_k such that:*

- (1) $F(X) \rightarrow F(X_{\mathrm{red}})$ is an isomorphism for every $X \in \mathrm{Sch}_k$.
- (2) $a_{\mathrm{Nis}}F|_{\mathrm{Sm}_k}$ is an unramified sheaf in the sense of [Mor12, Definition 2.1].

Then $a_{\mathrm{cdh}}F$ has a canonical structure of presheaf with transfers.

The second result on traces from [Kel13] concerns the compatibility of traces with the slice filtration. To state it we need a few preliminary definitions.

Definition 4.15: A *weak structure of smooth traces* on $\mathbf{E} \in \mathcal{SH}(\mathrm{Sm}_k)$ is a family of maps $\mathrm{Tr}_f: f_*\mathbf{E}_Y \rightarrow \mathbf{E}_X$, defined for every finite, flat, globally free, and surjective map $f: Y \rightarrow X$ in Sm_k , such that, if f is of degree d , $\mathrm{Tr}_f \circ \mathbf{E}_f = d \cdot \mathrm{id}_{\mathbf{E}_X}$.

Clearly, a structure of traces on \mathbf{E} induces a weak structure of smooth traces on \mathbf{E} .

Definition 4.16: Let $R \subset \mathbf{Q}$ be a subring of the rational numbers. We say that a motivic spectrum $\mathbf{E} \in \mathcal{SH}(\mathrm{Sm}_k)$ is *R -local* if the abelian group $[\mathbf{E}, \mathbf{E}]$ is an R -module.

Theorem 4.17 ([Kel13, Proposition 4.3.7]): *Suppose $\mathbf{E} \in \mathcal{SH}(\mathrm{Sm}_k)$ is $\mathbf{Z}[1/c]$ -local with a structure of traces, and suppose that $s_q\mathbf{E}$ has a weak structure of smooth traces for every $q \in \mathbf{Z}$. Then $s_q\mathbf{E}$ has a structure of traces for every $q \in \mathbf{Z}$.*

Proposition 4.18 ([Kel13, Proposition 5.2.3]): *The homotopy algebraic K -theory spectrum $\mathrm{KGL} \in \mathcal{SH}(\mathrm{Sm}_k)$ has a structure of traces.*

Proof. Let $f: Y \rightarrow X$ be a finite flat surjective map in Sch_k . It induces an exact functor f_* between the biWaldhausen categories of perfect complexes, whence a morphism of K -theory spectra $\mathrm{Tr}_f: K(Y) \rightarrow K(X)$ in the sense of Thomason–Trobaugh. It follows from standard properties of algebraic K -theory that this defines a structure of traces on the presheaf of spectra $X \mapsto K(X)$: for additivity use [TT90, 1.7.2], for functoriality [TT90, 1.5.4], for base-change [TT90, 3.18], and for degree [TT90, 1.7.3]. Given $X \in \mathrm{Sch}_k$, let $K_X \in \mathrm{Ho}(\mathrm{Spt}^{\mathrm{sm}, \mathrm{op}}_X)$ be the restriction of K to Sm_X . The trace maps $\mathrm{Tr}_{f \times_X U}: K(Y \times_X U) \rightarrow K(U)$ define morphisms of presheaves $\mathrm{Tr}_f: f_*K_Y \rightarrow K_X$ which endow $X \mapsto K_X$ with a structure of traces.

The Bass K -theory presheaf K_X^B can be defined by

$$K_X^B = K_X^\sharp = \mathrm{hocolim}_n \mathbf{R} \mathrm{Hom}(\Sigma^\infty(\mathbf{P}^1)^{\wedge n}, K_X),$$

(see [Cis13, end of paragraph 2.5, Proposition 2.7, Proposition 2.10]). For any morphism $f: Y \rightarrow X$ in Sch_k , the derived pushforward functor $f_*: \mathrm{Ho}(\mathrm{Spt}^{\mathrm{sm}, \mathrm{op}}_Y) \rightarrow \mathrm{Ho}(\mathrm{Spt}^{\mathrm{sm}, \mathrm{op}}_X)$ right

adjoint to f^* commutes with homotopy colimits by abstract nonsense, and so $f_*K_X^B \simeq (f_*K_X)^\sharp$. It follows that the structure of traces on the section $X \mapsto K_X$ induces a structure of traces on the section $X \mapsto K_X^B$.

The homotopy algebraic K -theory presheaf of spectra $KH_X: \mathcal{S}m_X \rightarrow \mathcal{S}pt$ is by definition the \mathbf{A}^1 -localization of the presheaf K_X^B , that is,

$$KH_X = L_{\mathbf{A}^1}K_X^B = \text{hocolim}_n \mathbf{R} \text{Hom}(\Sigma^\infty(\Delta_X^n)_+, K_X^B),$$

where Δ_X^\bullet is the usual cosimplicial diagram in $\mathcal{S}m_X$ with $\Delta_X^n = \mathbf{A}_X^n$. As before, we have $f_*KH_Y \simeq L_{\mathbf{A}^1}f_*K_Y^B$ and hence an induced structure of traces on the section $X \mapsto KH_X$.

Denote by $\mathcal{E}(X)$ the homotopy category of \mathbf{P}^1 -spectra in the category of presheaves of spectra on $\mathcal{S}m_X$; its objects are sequences (E_0, E_1, \dots) of presheaves of spectra on $\mathcal{S}m_X$ together with equivalences $E_i \simeq \mathbf{R} \text{Hom}(\Sigma^\infty \mathbf{P}^1, E_{i+1})$. The 2-functor $X \mapsto \mathcal{S}\mathcal{H}(\mathcal{S}m_X)$ can then be identified with the sub-2-functor of \mathcal{E} spanned by the \mathbf{A}^1 - and Nisnevich-local objects. Under this identification, the motivic spectrum $\mathbf{KGL}_X \in \mathcal{S}\mathcal{H}(\mathcal{S}m_X)$ is the \mathbf{P}^1 -spectrum (KH_X, KH_X, \dots) where the equivalences $KH_X \simeq \mathbf{R} \text{Hom}(\Sigma^\infty \mathbf{P}^1, KH_X)$ are given by Bott periodicity. Thus, it remains to observe that the trace maps $\text{Tr}_f: f_*KH_Y \rightarrow KH_X$ are compatible with these equivalences, which follows easily from the definitions. \square

Proposition 4.19: *For any $\mathbf{E} \in \mathcal{D}\mathcal{M}(\mathcal{S}m_k, R)$, $u^{\text{tr}}\mathbf{E}$ has a weak structure of smooth traces.*

Proof. If $f: Y \rightarrow X$ is a finite flap surjective of degree d map in $\mathcal{S}m_k$, its transpose ${}^t f$ is a finite correspondence $X \rightarrow Y$ such that $f \circ {}^t f = d \cdot \text{id}_X$ in $\text{Cor}(\mathcal{S}m_k)$. It therefore induces, for any $F \in s\text{Pre}^{\text{tr}}(\mathcal{S}m_k, R)$, a map $\text{Tr}_f: f_*F_Y \rightarrow F_X$ such that $\text{Tr}_f \circ F_f = d \cdot \text{id}_{F_X}$. Since $a: X \rightarrow \text{Spec } k$ is smooth the functor $\mathbf{L}a^*: \mathcal{H}^{\text{tr}}(\mathcal{S}m_k, R) \rightarrow \mathcal{H}^{\text{tr}}(\mathcal{S}m_X, R)$ preserves \mathbf{A}^1 - and Nisnevich-local objects and commutes with $\Omega^{2,1}$ (see §2.2), so we obtain trace maps for the section $(a: X \rightarrow \text{Spec } k) \mapsto \mathbf{L}a^*\mathbf{E}$ of the 2-functor $X \mapsto \mathcal{D}\mathcal{M}(\mathcal{S}m_X, R)$ on $\mathcal{S}m_k$. Since u^{tr} is a morphism of 2-functors which moreover commutes with $\mathbf{L}a^*$ for a smooth, there is an induced weak structure of smooth traces on $u^{\text{tr}}\mathbf{E}$. \square

Corollary 4.20 ([Kel13, Corollary 5.2.4]): *Let $1/c \in R \subset \mathbf{Q}$. Then the motivic cohomology spectrum MR has a structures of traces in $\mathcal{S}\mathcal{H}(\mathcal{S}m_k)$.*

Proof. By [Lev08, Theorem 6.4.2] (or [GP12, Theorem 7.10]), the q th slice of \mathbf{KGL} is $\Sigma^{2q,q}\mathbf{M}\mathbf{Z}$. As s_q preserves homotopy colimits, the q th slice of $\mathbf{KGL} \otimes R$ is $\Sigma^{2q,q}MR$, which has a weak structure of smooth traces by Proposition 4.19. Since $\mathbf{KGL} \otimes R$ itself has a structure of traces by Proposition 4.18, we may apply Theorem 4.17 to deduce that $s_0(\mathbf{KGL} \otimes R) \simeq MR$ has a structure of traces. \square

Proposition 4.21: *Let $\mathbf{E} \in \mathcal{S}\mathcal{H}(\mathcal{S}m_k)$ be a motivic spectrum with a structure of traces (resp. a weak structure of smooth traces). Then for any $\mathbf{F} \in \mathcal{S}\mathcal{H}(\mathcal{S}m_k)$, $\mathbf{E} \wedge \mathbf{F}$ has a structure of traces (resp. a weak structure of smooth traces).*

Proof. By [Ayo07, Theorems 2.3.40 and 1.7.17], if $f: Y \rightarrow X$ is a projective morphism in Sch_k , $\mathbf{E} \in \mathcal{SH}(\text{Sm}_Y)$, and $\mathbf{F} \in \mathcal{SH}(\text{Sm}_X)$, the canonical map

$$f_*\mathbf{E} \wedge \mathbf{F} \rightarrow f_*(\mathbf{E} \wedge f^*\mathbf{F}) \quad (17)$$

is an isomorphism. Thus, if $f: Y \rightarrow X$ is a finite flat surjective map, we can define Tr_f as the composition

$$f_*(\mathbf{E} \wedge \mathbf{F})_Y \cong f_*(\mathbf{E}_Y \wedge f^*\mathbf{F}_X) \cong f_*\mathbf{E}_Y \wedge \mathbf{F}_X \xrightarrow{\text{Tr}_f \wedge \mathbf{F}_X} \mathbf{E}_X \wedge \mathbf{F}_X \cong (\mathbf{E} \wedge \mathbf{F})_X,$$

where the second isomorphism is an instance of (17). It is then easy to verify that this definition satisfies the axioms for a structure of traces (resp. a weak structure of smooth traces). \square

Corollary 4.22: *Let $1/c \in R \subset \mathbf{Q}$ and $\mathbf{E} \in \mathcal{SH}(\text{Sm}_k)$. Then $\mathbf{M}R \wedge \mathbf{E}$ has a structure of traces.*

Proof. Combine Corollary 4.20 and Proposition 4.21. \square

We now turn to the proof that $\mathbf{M}\mathbf{Z}_{(\ell)}$ -modules are ℓ dh-local.

Definition 4.23: Let τ be a Grothendieck topology on Sm_k . A motivic spectrum $\mathbf{E} \in \mathcal{SH}(\text{Sm}_k)$ is τ -local if it is local with respect to the class of maps

$$\{\Sigma^{p,q}\Sigma^\infty \mathcal{X}_+ \rightarrow \Sigma^{p,q}\Sigma^\infty X_+ \mid \mathcal{X} \rightarrow X \text{ is a } \tau\text{-hypercover and } p, q \in \mathbf{Z}\}.$$

Since $\text{Spt}(\text{Sm}_k)$ is a stable model category, $\mathcal{SH}(\text{Sm}_k)$ is canonically enriched in the stable homotopy category $\text{Ho}(\text{Spt})$. That is, for every $\mathbf{E}, \mathbf{F} \in \mathcal{SH}(\text{Sm}_k)$, we have a derived mapping spectrum $\mathbf{R}\text{Hom}(\mathbf{E}, \mathbf{F})$ such that $\Omega^\infty \mathbf{R}\text{Hom}(\mathbf{E}, \mathbf{F}) \simeq \mathbf{R}\text{Map}(\mathbf{E}, \mathbf{F})$. By [CD12, Theorem 3.2.15], there exists a presheaf of (symmetric) spectra $\underline{\mathbf{E}}: \text{Sch}_k^{\text{op}} \rightarrow \text{Spt}$ such that the composition of $\underline{\mathbf{E}}$ with the canonical functor $\text{Spt} \rightarrow \text{Ho}(\text{Spt})$ coincides with the functor

$$(a: X \rightarrow \text{Spec } k) \mapsto \mathbf{R}\text{Hom}(\mathbf{1}, \mathbf{R}a_*\mathbf{L}a^*\mathbf{E}).$$

Note that the restriction of $\underline{\mathbf{E}}$ to Sm_k is equivalent to $X \mapsto \mathbf{R}\text{Hom}(\Sigma^\infty X_+, \mathbf{E})$. We can thus rephrase Definition 4.23 as follows: $\mathbf{E} \in \mathcal{SH}(\text{Sm}_k)$ is τ -local if and only if, for every $p, q \in \mathbf{Z}$, the presheaf of spectra $\underline{\Sigma}^{p,q}\underline{\mathbf{E}}|_{\text{Sm}_k}$ satisfies τ -descent.

Theorem 4.24 ([Kel13, Theorem 5.3.7]): *Every $\mathbf{Z}_{(\ell)}$ -local motivic spectrum $\mathbf{E} \in \mathcal{SH}(\text{Sm}_k)$ with a structure of traces is ℓ dh-local.*

Proof. We must show that the presheaf of spectra $\underline{\Sigma}^{p,q}\underline{\mathbf{E}}|_{\text{Sm}_k}$ satisfies ℓ dh-descent for every $p, q \in \mathbf{Z}$. We will even prove that $\underline{\Sigma}^{p,q}\underline{\mathbf{E}}$ satisfies ℓ dh-descent. We may clearly assume that $p = q = 0$ since the hypotheses on \mathbf{E} are bistable.

As the cdh topology has finite cohomological dimension [SV00a, Theorem 12.5] and \underline{E} is cdh-local [Cis13, Proposition 3.7], Lemma 4.8 will show that \underline{E} is ℓ dh-local provided that the canonical maps

$$H_{\text{cdh}}^s(X, a_{\text{cdh}}\pi_t\underline{E}) \rightarrow H_{\ell\text{dh}}^s(X, a_{\ell\text{dh}}\pi_t\underline{E})$$

are isomorphisms for all $s, t \in \mathbf{Z}$. By Theorems 4.7 and 4.14, this will be the case if

- (0) $\pi_t\underline{E}$ is a presheaf of $\mathbf{Z}_{(\ell)}$ -modules with traces,
- (1) $\pi_t\underline{E}(X) \rightarrow \pi_t\underline{E}(X_{\text{red}})$ is an isomorphism for every $X \in \text{Sch}_k$,
- (2) $a_{\text{Nis}}\pi_t\underline{E}|_{\text{Sm}_k}$ is unramified.

(0) holds by assumption on \underline{E} (see Remark 4.13) and (1) is clear because the restriction functor $\mathcal{SH}(\text{Sm}_X) \rightarrow \mathcal{SH}(\text{Sm}_{X_{\text{red}}})$ is an equivalence of categories. The sheaf $a_{\text{Nis}}\pi_t\underline{E}|_{\text{Sm}_k}$ is strictly \mathbf{A}^1 -invariant by [Mor03, Remark 5.1.13], and in particular is unramified (see [Mor12, Example 2.3]). \square

Corollary 4.25: *Every $\mathbf{MZ}_{(\ell)}$ -module spectrum in $\mathcal{SH}(\text{Sm}_k)$ is ℓ dh-local.*

Proof. Let \underline{E} be an $\mathbf{MZ}_{(\ell)}$ -module spectrum. Then \underline{E} is a retract of $\mathbf{MZ}_{(\ell)} \wedge \underline{E}$ in $\mathcal{SH}(\text{Sm}_k)$, and so it suffices to show that $\mathbf{MZ}_{(\ell)} \wedge \underline{E}$ is ℓ dh-local. This follows from Corollary 4.22 and Theorem 4.24. \square

Corollary 4.26 ([Kel13, Corollary 5.3.8]): *Let R be a $\mathbf{Z}_{(\ell)}$ -algebra. If $\mathcal{X} \rightarrow X$ is a smooth ℓ dh-hypercover of $X \in \text{Sm}_k$, then the induced map $\mathbf{L}R^{\text{tr}}\mathcal{X}_+ \rightarrow \mathbf{L}R^{\text{tr}}X_+$ is an isomorphism in $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R)$.*

Proof. By [Voe10b, Theorem 1.15] and Voevodsky's Cancellation Theorem [Voe10a], the stabilization functor $\Sigma^\infty: \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R) \rightarrow \mathcal{DM}(\text{Sm}_k, R)$ is fully faithful. Thus, it suffices to show that any smooth ℓ dh-hypercover gives rise to an isomorphism in $\mathcal{DM}(\text{Sm}_k, R)$. Recall from §2.3 that the functor $\mathbf{L}R^{\text{tr}}\Sigma^\infty(-)_+$ factors through the functor $\mathbf{M}R \wedge \Sigma^\infty(-): \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^*(\text{Sm}_k) \rightarrow \text{Ho}(\mathbf{M}R\text{-mod})$ to the highly structured category of modules over $\mathbf{M}R$, and this reduces the problem to showing that the map $\mathbf{M}R \wedge \Sigma^\infty\mathcal{X}_+ \rightarrow \mathbf{M}R \wedge \Sigma^\infty X_+$ is an isomorphism for every smooth ℓ dh-hypercover $\mathcal{X} \rightarrow X$. This follows from Corollary 4.25. \square

Corollary 4.27: *Let R be a $\mathbf{Z}_{(\ell)}$ -algebra. Then the localization functor $\mathcal{H}^{\text{tr}}(\text{Sm}_k, R) \rightarrow \mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R)$ factors through $\mathcal{H}_{\ell\text{dh}}^{\text{tr}}(\text{Sm}_k, R)$.*

Proof. This is just a rephrasing of Corollary 4.26. \square

4.3 End of the proof

We now give the proof of Theorem 3.1. As in [Voe10b, end of §1.3], we may assume that $\mathcal{C} = \text{Sm}_k$ and $\mathcal{D} = \text{Sch}_k$. Since the restriction functors $i^*: \mathcal{H}^*(\text{Sch}_k) \rightarrow \mathcal{H}^*(\text{Sm}_k)$ and

$i^* : \mathcal{H}^{\text{tr}}(\text{Sch}_k, R) \rightarrow \mathcal{H}^{\text{tr}}(\text{Sm}_k, R)$ preserve \mathbf{A}^1 -Nisnevich-local equivalences, it suffices to show that for any $F \in \mathcal{H}^*(\text{Sch}_k)$ the canonical map

$$\mathbf{L}R^{\text{tr}}i^*F \rightarrow i^*\mathbf{L}R^{\text{tr}}F$$

in $\mathcal{H}^{\text{tr}}(\text{Sm}_k, R)$ becomes an isomorphism in $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R)$. In the commutative square

$$\begin{array}{ccc} \mathbf{L}R^{\text{tr}}i^*\mathbf{L}i_!i^* & \longrightarrow & \mathbf{L}R^{\text{tr}}i^* \\ \downarrow & & \downarrow \\ i^*\mathbf{L}R^{\text{tr}}\mathbf{L}i_!i^* & \longrightarrow & i^*\mathbf{L}R^{\text{tr}} \end{array}$$

the upper horizontal arrow and the left vertical arrow are isomorphisms since $\mathbf{L}i_!$ is fully faithful. Thus, it suffices to show that the lower horizontal arrow becomes an isomorphism in $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R)$. Since $\mathbf{L}i_!i^*F \rightarrow F$ is an isomorphism when restricted to Sm_k , it is an ℓdh -local equivalence by Corollary 4.6. Thus, $\mathbf{L}R^{\text{tr}}\mathbf{L}i_!i^*F \rightarrow \mathbf{L}R^{\text{tr}}F$ is an $R^{\text{tr}}W_{\ell\text{dh}}$ -local equivalence. By Proposition 4.10 (3), $i^*\mathbf{L}R^{\text{tr}}\mathbf{L}i_!i^*F \rightarrow i^*\mathbf{L}R^{\text{tr}}F$ is therefore an $R^{\text{tr}}W_{\ell\text{dh}}$ -local equivalence in $\mathcal{H}^{\text{tr}}(\text{Sm}_k, R)$. Finally, by Corollary 4.27, $R^{\text{tr}}W_{\ell\text{dh}}$ -local equivalences in $\mathcal{H}^{\text{tr}}(\text{Sm}_k, R)$ become equivalences in $\mathcal{H}_{\text{Nis}, \mathbf{A}^1}^{\text{tr}}(\text{Sm}_k, R)$, as was to be shown.

5 Applications

In this section we record some applications of the results of this paper which were previously only known for fields of characteristic zero.

5.1 The structure of the motivic Steenrod algebra and its dual

The goal of this paragraph is to generalize the structure theorems of Voevodsky for the motivic Steenrod algebra over perfect fields to essentially smooth schemes over fields. Throughout, the base scheme S is essentially smooth over a field, and $\ell \neq \text{char } S$ is a fixed prime number. We abbreviate $\mathbf{M}\mathbf{Z}/\ell$ to \mathbf{M} , and we write \mathcal{A}^{**} for the motivic Steenrod algebra at ℓ , which by Theorem 1.1 is the algebra of all bistable operations in mod ℓ motivic cohomology of smooth S -schemes, and also the algebra of bigraded endomorphisms of the motivic Eilenberg–Mac Lane spectrum \mathbf{M} . Recall that \mathcal{A}^{**} is generated by the reduced power operations P^i , the Bockstein β , and the subalgebra $\mathbf{H}^{**}(S, \mathbf{Z}/\ell)$. As usual, we write

$$B^i = \beta P^i,$$

and if $\ell = 2$,

$$\text{Sq}^{2i} = P^i \quad \text{and} \quad \text{Sq}^{2i+1} = B^i.$$

If $\ell = 2$, let ρ be the image of $-1 \in \mathbf{G}_m(S)$ in

$$\mathbf{H}^{1,1}(S, \mathbf{Z}/2) = \mathbf{H}_{\text{ét}}^1(S, \mu_2)$$

and let τ be the nonvanishing element of

$$H^{0,1}(S, \mathbf{Z}/2) = \mu_2(S) \cong \text{Hom}(\pi_0(S), \mathbf{Z}/2)$$

(recall that $\text{char } S \neq 2$ if $\ell = 2$).

Theorem 5.1 (The Adem relations):

(1) Assume $\ell \neq 2$. If $0 < a < \ell b$, then

$$P^a P^b = \sum_{t=0}^{\lfloor a/\ell \rfloor} (-1)^{a+t} \binom{(\ell-1)(b-t)-1}{a-\ell t} P^{a+b-t} P^t.$$

If $0 < a \leq \ell b$, then

$$\begin{aligned} P^a B^b &= \sum_{t=0}^{\lfloor a/\ell \rfloor} (-1)^{a+t} \binom{(\ell-1)(b-t)}{a-\ell t} B^{a+b-t} P^t \\ &\quad + \sum_{t=0}^{\lfloor (a-1)/\ell \rfloor} (-1)^{a+t-1} \binom{(\ell-1)(b-t)-1}{a-\ell t} P^{a+b-t} B^t. \end{aligned}$$

(2) Assume $\ell = 2$ and $0 < a < 2b$. If a and b are even, then

$$\text{Sq}^a \text{Sq}^b = \sum_{t=0}^{\lfloor a/2 \rfloor} \tau^{t \bmod 2} \binom{b-t-1}{a-2t} \text{Sq}^{a+b-t} \text{Sq}^t.$$

If a is even and b is odd, then

$$\begin{aligned} \text{Sq}^a \text{Sq}^b &= \sum_{t=0}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t} \text{Sq}^{a+b-t} \text{Sq}^t \\ &\quad + \sum_{\substack{t=0 \\ t \text{ odd}}}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t} \rho \text{Sq}^{a+b-t-1} \text{Sq}^t. \end{aligned}$$

If a is odd and b is even, then

$$\begin{aligned} \text{Sq}^a \text{Sq}^b &= \sum_{\substack{t=0 \\ t \text{ even}}}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t} \text{Sq}^{a+b-t} \text{Sq}^t \\ &\quad + \sum_{\substack{t=0 \\ t \text{ odd}}}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t-1} \rho \text{Sq}^{a+b-t-1} \text{Sq}^t. \end{aligned}$$

If a and b are odd, then

$$\text{Sq}^a \text{Sq}^b = \sum_{\substack{t=0 \\ t \text{ odd}}}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t} \text{Sq}^{a+b-t} \text{Sq}^t.$$

Proof. If S is a perfect field, this was originally proved in [Voe03b, Theorems 10.2 and 10.3], but with some typos that were corrected in [Rio12, Théorèmes 4.5.1 et 4.5.2]. In general, choose an essentially smooth morphism $f: S \rightarrow \text{Spec } k$ where k is a perfect field. The theorem follows from the fact that the base change map $f^*: \mathcal{A}_k^{**} \rightarrow \mathcal{A}_S^{**}$ is an algebra homomorphism which preserves P^i and β , and, when $\ell = 2$, τ and ρ . \square

Note that in the case $\ell = 2$, the Adem relations for a odd are obtained from the ones for a even by application of β , using that β is a derivation, that $\beta(\tau) = \rho$, and that $\beta(\rho) = 0$. The Adem relations (together with the relation $\beta^2 = 0$) can be used to express any monomial in the operations P^i and β as a linear combination of the basis operations described in Theorem 1.1. The only additional piece of information needed to obtain a presentation of \mathcal{A}^{**} as an algebra is thus the action of \mathcal{A}^{**} on $\mathbf{H}^{**}(S, \mathbf{Z}/\ell)$.

Let now \mathcal{A}_{**} be the dual motivic Steenrod algebra over S , that is,

$$\mathcal{A}_{**} = \text{Hom}_{\mathbf{M}_{**}}(\mathcal{A}^{-*,-*}, \mathbf{M}_{**}).$$

By duality, \mathcal{A}_{**} has a structure of commutative Hopf algebroid which is described in details in [Voe03b, §12] in the case where S is the spectrum of a perfect field. Our goal is now to identify \mathcal{A}_{**} with the Hopf algebroid of co-operations in mod ℓ motivic cohomology, i.e., $\mathbf{M}_{**}\mathbf{M}$. This is achieved in Propositions 5.3 and 5.5 below. For the following lemma, recall that $\text{Ho}(\mathbf{M}\text{-mod})$ is a closed symmetric monoidal category.

Lemma 5.2: *Let (p_α, q_α) be a family of bidegrees with $p_\alpha \geq 2q_\alpha \geq 0$ and such that, for every $q \in \mathbf{Z}$, there are only finitely many α with $q_\alpha \leq q$. Let $\mathbf{E} = \bigvee_\alpha \Sigma^{p_\alpha, q_\alpha} \mathbf{M} \in \text{Ho}(\mathbf{M}\text{-mod})$. Then the canonical map $\mathbf{E} \rightarrow \text{Hom}_{\mathbf{M}}(\text{Hom}_{\mathbf{M}}(\mathbf{E}, \mathbf{M}), \mathbf{M})$ is an equivalence of \mathbf{M} -modules, and the pairing $\pi_{**} \text{Hom}_{\mathbf{M}}(\mathbf{E}, \mathbf{M}) \otimes_{\mathbf{M}_{**}} \pi_{**} \mathbf{E} \rightarrow \mathbf{M}_{**}$ is perfect.*

Proof. The hypothesis on (p_α, q_α) together with Corollary 2.14 implies that

$$\begin{aligned} \mathbf{E} &= \bigvee_\alpha \Sigma^{p_\alpha, q_\alpha} \mathbf{M} \simeq \prod_\alpha \Sigma^{p_\alpha, q_\alpha} \mathbf{M}, \\ \text{Hom}_{\mathbf{M}}(\mathbf{E}, \mathbf{M}) &\simeq \prod_\alpha \Sigma^{-p_\alpha, -q_\alpha} \mathbf{M} \simeq \bigvee_\alpha \Sigma^{-p_\alpha, -q_\alpha} \mathbf{M}. \end{aligned}$$

This immediately implies the first statement and, taking homotopy groups, the second. \square

Proposition 5.3: *There is a canonical isomorphism $\mathcal{A}_{**} \cong \mathbf{M}_{**}\mathbf{M}$ and the canonical map $\mathcal{A}^{-*,-*} \rightarrow \text{Hom}_{\mathbf{M}_{**}}(\mathcal{A}_{**}, \mathbf{M}_{**})$ is an isomorphism.*

Proof. By Theorem 1.1, $\mathcal{A}^{**} \cong \mathbf{M}^{**}\mathbf{M}$. Therefore it suffices to show there is a perfect pairing $\mathbf{M}^{-*,-*} \otimes_{\mathbf{M}_{**}} \mathbf{M}_{**}\mathbf{M} \rightarrow \mathbf{M}_{**}$. This follows from Theorem 1.1 (3) and Lemma 5.2. \square

Lemma 5.4: *Let \mathbf{R} be a motivic E_∞ -ring spectrum and let \mathbf{E} and \mathbf{F} be \mathbf{R} -modules. The canonical map*

$$\pi_{**} \mathbf{E} \otimes_{\mathbf{R}_{**}} \pi_{**} \mathbf{F} \rightarrow \pi_{**}(\mathbf{E} \wedge_{\mathbf{R}} \mathbf{F})$$

is an isomorphism under either of the following conditions:

(1) F is a cellular R -module and $\pi_{**}E$ is flat over R_{**} .

(2) $F \simeq \bigvee_{\alpha} \Sigma^{p_{\alpha}, q_{\alpha}} R$ as an R -module.

Proof. Assuming (1), this is a natural transformation between homological functors of F that preserve sums, so we may assume (2), in which case the result is obvious. \square

Proposition 5.5: *Let E be an M -module. Then there is a canonical isomorphism*

$$\pi_{**}E \otimes_{M_{**}} \mathcal{A}_{**} \cong \pi_{**}(E \wedge M).$$

*In particular, $\mathcal{A}_{**}^{\otimes i} \cong M_{**}(M^{\wedge i})$ for all $i \geq 0$ (where the tensor product is over M_{**}).*

Proof. Since $M \wedge M \simeq \bigvee_{\alpha} \Sigma^{p_{\alpha}, q_{\alpha}} M$ and $M_{**}M \cong \mathcal{A}_{**}$, this is a special case of Lemma 5.4. \square

Proposition 5.5 is of computational interest because it implies that for any $E \in S\mathcal{H}(\text{Sm}_S)$, $M_{**}E$ is a left comodule over \mathcal{A}_{**} . For instance, the E_2 -page of the homological motivic Adams spectral sequence is comprised of Ext-groups in the category of \mathcal{A}_{**} -comodules. See e.g., [DI10] and [HKO11] for precise statements concerning the construction and convergence of the motivic Adams spectral sequence, which can be generalized to essentially smooth schemes over fields by the above results.

We can now give a complete description of the Hopf algebroid $\mathcal{A}_{**} = M_{**}M$. First, define a Hopf algebroid (A, Γ) as follows. Let

$$\begin{aligned} A &= \mathbf{Z}/\ell[\rho, \tau], \\ \Gamma &= A[\tau_0, \tau_1, \dots, \xi_1, \xi_2, \dots] / (\tau_i^2 - \tau\xi_{i+1} - \rho\tau_{i+1} - \rho\tau_0\xi_{i+1}). \end{aligned}$$

The structure maps η_L , η_R , ϵ , and Δ are given by the formulas

$$\begin{aligned} \eta_L: A &\rightarrow \Gamma, & \eta_L(\rho) &= \rho, \eta_L(\tau) = \tau, \\ \eta_R: A &\rightarrow \Gamma, & \eta_R(\rho) &= \rho, \eta_R(\tau) = \tau + \rho\tau_0, \\ \epsilon: \Gamma &\rightarrow A, & \epsilon(\rho) &= \rho, \\ & & \epsilon(\tau) &= \tau, \\ & & \epsilon(\tau_r) &= 0, \\ & & \epsilon(\xi_r) &= 0, \\ \Delta: \Gamma &\rightarrow \Gamma \otimes_A \Gamma, & \Delta(\rho) &= \rho \otimes 1, \\ & & \Delta(\tau) &= \tau \otimes 1, \\ & & \Delta(\tau_r) &= \tau_r \otimes 1 + 1 \otimes \tau_r + \sum_{i=0}^{r-1} \xi_{r-i}^{l^i} \otimes \tau_i, \\ & & \Delta(\xi_r) &= \xi_r \otimes 1 + 1 \otimes \xi_r + \sum_{i=1}^{r-1} \xi_{r-i}^{l^i} \otimes \xi_i. \end{aligned}$$

The coinverse map $c: \Gamma \rightarrow \Gamma$ is determined by the identities it must satisfy. Namely, we have

$$\begin{aligned} c(\rho) &= \rho, \\ c(\tau) &= \tau + \rho\tau_0, \\ c(\tau_r) &= -\tau_r - \sum_{i=0}^{r-1} \xi_{r-i}^{l^i} c(\tau_i), \\ c(\xi_r) &= -\xi_r - \sum_{i=1}^{r-1} \xi_{r-i}^{l^i} c(\xi_i). \end{aligned}$$

We view M_{**} as an A -algebra via the map $A \rightarrow M_{**}$ defined as follows: if ℓ is odd it sends both ρ and τ to 0, while if $\ell = 2$ it sends ρ and τ to the elements of the same name in M_{**} .

Theorem 5.6: \mathcal{A}_{**} is isomorphic to $\Gamma \otimes_A M_{**}$ with

$$|\tau_r| = (2\ell^r - 1, \ell^r - 1) \quad \text{and} \quad |\xi_r| = (2\ell^r - 2, \ell^r - 1).$$

The map $M_{**} \rightarrow \mathcal{A}_{**}$ dual to the left action of \mathcal{A}^{**} on M^{**} is a left coaction of (A, Γ) on the ring M_{**} , and the Hopf algebroid $(M_{**}, \mathcal{A}_{**})$ is isomorphic to the twisted tensor product of (A, Γ) with M_{**} , i.e.,

- η_L and ϵ are extended from (A, Γ) ,
- $\eta_R: H_{**} \rightarrow \mathcal{A}_{**}$ is the coaction,
- $\Delta: \mathcal{A}_{**} \rightarrow \mathcal{A}_{**} \otimes_{M_{**}} \mathcal{A}_{**}$ is induced by the diagonal of Γ and η_R to the second factor,
- $c: \mathcal{A}_{**} \rightarrow \mathcal{A}_{**}$ is induced by the coinverse of Γ and η_R .

Proof. If S is the spectrum of a perfect field, this is proved in [Voe03b, §12]. In general, choose an essentially smooth morphism $f: S \rightarrow \text{Spec } k$ where k is a perfect field. Note that the induced map $(M_k)_{**} \rightarrow (M_S)_{**}$ is a map of A -algebras. It remains to observe that the Hopf algebroid \mathcal{A}_{**} is obtained from $(M_k)_{**} M_k$ by extending scalars from $(M_k)_{**}$ to $(M_S)_{**}$, which follows formally from the following facts: $\mathbf{L}f^*$ is a symmetric monoidal functor, Theorem 2.12, and Corollary 3.4. \square

5.2 Modules over motivic cohomology

In this paragraph we generalize the the main result in [RØ06], [RØ08]. Recall that there is a symmetric monoidal Quillen adjunction

$$\Phi: MR\text{-mod} \rightleftarrows \text{Spt}^{\text{tr}}(\text{Sm}_S, R) : \Psi$$

which is a Quillen equivalence when S is a field of characteristic zero [RØ08, Theorem 1] or when S is any perfect field and $\mathbf{Q} \subset R$ [RØ08, Theorem 68]. In the following, all functors are derived by default.

Lemma 5.7: *Let k be a perfect field, $g: V \rightarrow U$ an fpsl^ℓ -cover in Sm_k , and R a $\mathbf{Z}_{(\ell)}$ -algebra. Then $\text{MR} \wedge g_+$ has a section in the homotopy category of MR -modules.*

Proof. Since $\text{Ho}(\text{MR-mod})$ is a triangulated category, it suffices to show that $\text{MR} \wedge g_+$ is an epimorphism, i.e., that for every MR -module \mathbf{E} ,

$$[U_+, \mathbf{E}] \rightarrow [V_+, \mathbf{E}] \quad (18)$$

is injective. If $a: U \rightarrow \text{Spec } k$ is the structure map, we have $U_+ = a_{\sharp} a^* \mathbf{1}$ and $V_+ = (ag)_{\sharp} (ag)^* \mathbf{1}$. Now for any smooth map f , $f_{\sharp} f^*$ is left adjoint to $f_* f^*$. Using this adjunction, we can identify (18) with the map

$$[\mathbf{1}, a_* a^* \mathbf{E}] \rightarrow [\mathbf{1}, (ag)_* (ag)^* \mathbf{E}]$$

induced by the unit of the adjunction (g^*, g_*) . Since \mathbf{E} is a retract of $\text{MR} \wedge \mathbf{E}$ in $\mathcal{SH}(\text{Sm}_k)$, we can assume that \mathbf{E} is a free MR -module. By Propositions 4.19 and 4.21, \mathbf{E} then has a weak structure of smooth traces. Since R is a $\mathbf{Z}_{(\ell)}$ -algebra, it follows that $a_* a^* \mathbf{E} \rightarrow (ag)_* (ag)^* \mathbf{E}$ has a retraction, namely $\frac{1}{d} a_* (\text{Tr}_g)$ where d is the degree of g . \square

Theorem 5.8: *Assume that S is the spectrum of a perfect field of characteristic $p > 0$ and let R be a commutative ring in which p is invertible. Then (Φ, Ψ) is a Quillen equivalence.*

Proof. As in [RØ08], it suffices to prove that the unit $\eta_{X_+}: \text{MR} \wedge X_+ \rightarrow \Psi \Phi(\text{MR} \wedge X_+)$ is a weak equivalence for all quasi-projective and connected $X \in \text{Sm}_S$, knowing that this holds if X is projective (more generally, η_E is an equivalence if $E \in \mathcal{SH}(\text{Sm}_S)$ is dualizable). A map of \mathbf{MZ} -modules is an equivalence if and only if it induces equivalences of $\mathbf{MZ}_{(\ell)}$ -modules for every prime ℓ , and since both Φ and Ψ preserve filtered homotopy colimits we can assume that R is a $\mathbf{Z}_{(\ell)}$ -algebra for some prime $\ell \neq \text{char } k$. In this case we follow the proof of the rational result [RØ08, Theorem 68] but use Gabber's theorem instead of de Jong's theorem.

We proceed by induction on the dimension of X , the case $d = 0$ being taken care of since X is then necessarily projective. Let $\mathcal{SH}_d(k)$ be the localizing subcategory of $\mathcal{SH}(\text{Sm}_k)$ generated by shifted suspension spectra of smooth connected k -schemes of dimension $\leq d$. By induction hypothesis, η_E is an equivalence for every $E \in \mathcal{SH}_{d-1}(k)$. Choose an open embedding $j: X \hookrightarrow Y$ into an integral projective k -scheme. Let $f: Y' \rightarrow Y$ be the map given by Gabber's Theorem 4.4, so that $X' = f^{-1}(X)$ is the complement of a divisor with strict normal crossings. Let $U \subset X$ be an open subset on which f restricts to an fpsl^ℓ -cover $g: V = f^{-1}(U) \rightarrow U$. Since Y' is smooth and projective, Y'_+ is dualizable in $\mathcal{SH}(\text{Sm}_k)$. By homotopy purity and induction on the number of irreducible components of $Y' \setminus X'$ (cf. the proof of [RØ08, Theorem 52]), X'_+ is dualizable in $\mathcal{SH}(\text{Sm}_k)$ and hence $\eta_{X'_+}$ is an equivalence. Consider the cofiber sequences

$$\begin{aligned} V_+ &\hookrightarrow X'_+ \rightarrow X'/V, \\ U_+ &\hookrightarrow X_+ \rightarrow X/U. \end{aligned}$$

By [RØ08, Lemma 66], X'/V and X/U belong to $\mathcal{SH}_{d-1}(k)$. By induction hypothesis, $\eta_{X'/V}$ and $\eta_{X/U}$ are equivalences. Thus, η_{V_+} is also an equivalence, and it remains to prove that η_{U_+} is an equivalence. This follows from Lemma 5.7, since $MR \wedge U_+$ is a retract of $MR \wedge V_+$ in $\mathrm{Ho}(MR\text{-mod})$. \square

References

- [AF13] A. Asok and J. Fasel, *Motivic secondary characteristic classes and the Euler class*, 2013, arXiv:1307.6831v1 [math.AG]
- [Ayo07] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I*, Astérisque (2007), no. 314, pp. x+466 pp. (2008)
- [CD12] D.-C. Cisinski and F. Déglise, *Triangulated categories of mixed motives*, 2012, arXiv:0912.2110v3 [math.AG]
- [Cis13] D.-C. Cisinski, *Descente par éclatements en K-théorie invariante par homotopie*, Ann. of Math. (2) **177** (2013), no. 2, pp. 425–448
- [DI10] D. Dugger and D. C. Isaksen, *The motivic Adams spectral sequence*, Geom. Topol. **14** (2010), no. 2, pp. 967–1014, <http://dx.doi.org/10.2140/gt.2010.14.967>
- [FV00] E. M. Friedlander and V. Voevodsky, *Bivariant cycle cohomology*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 138–187
- [GP12] G. Garkusha and I. Panin, *On the motivic spectral sequence*, 2012, arXiv:1210.2242v1 [math.AG]
- [Gro66] A. Grothendieck, *Éléments de Géométrie Algébrique: IV. Étude locale des schémas et des morphismes de schémas, Troisième partie*, Publ. Math. I.H.É.S. **28** (1966)
- [Gro67] ———, *Éléments de Géométrie Algébrique: IV. Étude locale des schémas et des morphismes de schémas, Quatrième partie*, Publ. Math. I.H.É.S. **32** (1967)
- [HKO11] P. Hu, I. Kriz, and K. Ormsby, *Convergence of the motivic Adams spectral sequence*, J. K-Theory **7** (2011), no. 3, pp. 573–596, <http://dx.doi.org/10.1017/is011003012jkt150>
- [Hir09] P. S. Hirschhorn, *Model Categories and Their Localizations*, AMS, 2009

- [Hov99] M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999
- [Hov01] ———, *Spectra and symmetric spectra in general model categories*, J. Pure Appl. Alg. **165** (2001), no. 63-127, preprint arXiv:math/0004051v3 [math.AT]
- [Hoy13] M. Hoyois, *From algebraic cobordism to motivic cohomology*, to appear in J. Reine Angew. Math., 2013, arXiv:1210.7182 [math.AG]
- [ILO12] L. Illusie, Y. Laszlo, and F. Orgogozo, *Travaux de Gabber sur l'uniformisation locale et la cohomologie étale des schémas quasi-excellents. Séminaire à l'École polytechnique 2006–2008*, 2012, http://www.math.polytechnique.fr/~orgogozo/travaux_de_Gabber/
- [Ill] L. Illusie, *On Gabber's refined uniformization*, Talks at the Univ. Tokyo, Jan. 17, 22, 31, Feb. 7, 2008, http://www.math.u-psud.fr/~illusie/refined_uniformization3.pdf
- [Jar97] J. F. Jardine, *Generalized étale cohomology theories*, Progress in Mathematics, vol. 146, Birkhäuser Verlag, Basel, 1997, <http://dx.doi.org/10.1007/978-3-0348-0066-2>
- [Kel13] S. Kelly, *Triangulated categories of motives in positive characteristic*, Ph.D. Thesis, Université Paris 13, Australian National University, 2013, arXiv:1305.5349 [math.AG]
- [Lev00] M. Levine, *Inverting the motivic Bott element*, K-Theory **19** (2000), no. 1, pp. 1–28, <http://dx.doi.org/10.1023/A:1007874218371>
- [Lev08] ———, *The homotopy coniveau tower*, J. Topol. **1** (2008), no. 1, pp. 217–267, <http://dx.doi.org/10.1112/jtopol/jtm004>
- [Lur09] J. Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009
- [Lur11] ———, *Derived Algebraic Geometry VII: Spectral Schemes*, 2011, <http://www.math.harvard.edu/~lurie/papers/DAG-VII.pdf>
- [MVW06] C. Mazza, V. Voevodsky, and C. Weibel, *Lecture notes on motivic cohomology*, Clay Mathematics Monographs, vol. 2, American Mathematical Society, Providence, RI, 2006
- [Mor03] F. Morel, *An introduction to \mathbf{A}^1 -homotopy theory*, Contemporary Developments in Algebraic K-theory, ICTP, 2003, pp. 357–441

- [Mor12] ———, *\mathbf{A}^1 -Algebraic Topology over a Field*, Lecture Notes in Mathematics, vol. 2052, Springer, 2012
- [OØ13] K. Ormsby and P. A. Østvær, *Stable motivic π_1 for low-dimensional fields*, Preprint, 2013
- [Rio12] J. Riou, *Opérations de Steenrod motiviques*, 2012, arXiv:1207.3121v1 [math.AG]
- [RG71] M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), pp. 1–89
- [RØ06] O. Röndigs and P. A. Østvær, *Motives and modules over motivic cohomology*, C. R. Math. Acad. Sci. Paris **342** (2006), no. 10, pp. 751–754, <http://dx.doi.org/10.1016/j.crma.2006.03.013>
- [RØ08] ———, *Modules over motivic cohomology*, Adv. Math. **219** (2008), no. 2, pp. 689–727, <http://dx.doi.org/10.1016/j.aim.2008.05.013>
- [RØ13] ———, *Slices of Hermitian K-theory and Milnor’s conjecture on quadratic forms*, Preprint, 2013
- [SGA72] *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*, Lecture Notes in Mathematics, Vol. 269, Springer-Verlag, Berlin, 1972, Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J.-L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat
- [SV96] A. Suslin and V. Voevodsky, *Singular homology of abstract algebraic varieties*, Invent. Math. **123** (1996), no. 1, pp. 61–94, <http://dx.doi.org/10.1007/BF01232367>
- [SV00a] ———, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), NATO Sci. Ser. C Math. Phys. Sci., vol. 548, Kluwer Acad. Publ., Dordrecht, 2000, pp. 117–189
- [SV00b] ———, *Relative cycles and Chow sheaves*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 10–86
- [Ste62] N. E. Steenrod, *Cohomology operations*, Lectures by N. E. Steenrod written and revised by D. B. A. Epstein. Annals of Mathematics Studies, No. 50, Princeton University Press, Princeton, N.J., 1962
- [TT90] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift III, Progress in Mathematics, vol. 88, Birkhäuser, 1990, pp. 247–435

- [Tho85] R. W. Thomason, *Algebraic K-theory and étale cohomology*, Ann. Sci. École Norm. Sup. (4) **18** (1985), no. 3, pp. 437–552,
http://www.numdam.org.virtual.anu.edu.au/item?id=ASENS_1985_4_18_3_437_0
- [Voe00] V. Voevodsky, *Triangulated categories of motives over a field*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 188–238
- [Voe03b] ———, *Reduced power operations in motivic cohomology*, Publ. Math. Inst. Hautes Études Sci. (2003), no. 98, pp. 1–57,
<http://dx.doi.org/10.1007/s10240-003-0009-z>
- [Voe10a] ———, *Cancellation theorem*, Doc. Math. (2010), no. Extra volume: Andrei A. Suslin sixtieth birthday, pp. 671–685
- [Voe10b] ———, *Motivic Eilenberg-MacLane spaces*, Publ. Math. Inst. Hautes Études Sci. (2010), no. 112, pp. 1–99,
<http://dx.doi.org/10.1007/s10240-010-0024-9>
- [Voe10c] ———, *Simplicial additive functors*, J. K-Theory **5** (2010), no. 2, pp. 201–244,
<http://dx.doi.org/10.1017/is010003026jkt097>
- [Voe11] ———, *On motivic cohomology with \mathbf{Z}/l -coefficients*, Ann. of Math. (2) **174** (2011), no. 1, pp. 401–438,
<http://dx.doi.org/10.4007/annals.2011.174.1.11>

Department of Mathematics, Northwestern University, USA.
 e-mail: hoyois@math.northwestern.edu

Fakultät Mathematik, Universität Duisburg-Essen, Germany.
 e-mail: shanekelly64@gmail.com

Department of Mathematics, University of Oslo, Norway.
 e-mail: paularne@math.uio.no