

Quantum Dynamics with Electronic Friction

Rocco Martinazzo^{1,2,*}, Irene Burghardt³

¹*Department of Chemistry, Università degli Studi di Milano, Via Golgi 19, 20133 Milano, Italy**

²*Istituto di Scienze e Tecnologie Molecolari, CNR, via Golgi 19, 20133 Milano, Italy and*

³*Institute of Physical and Theoretical Chemistry, Goethe University Frankfurt, Max-von-Laue-Str. 7, D-60438 Frankfurt/Main, Germany*

A theory of electronic friction is developed using the exact factorization of the electron-nuclear wavefunction. No assumption is made regarding the electronic bath, which can be made of independent or interacting electrons, and the nuclei are treated quantumly. The ensuing equation of motion for the nuclear wavefunction is a non-linear Schrödinger equation including a friction term. The resulting friction kernel agrees with a previously derived mixed quantum-classical result by Dou, Miao & Subotnik (*Phys. Rev. Lett.* **119**, 046001 (2017)), except for a *pseudo*-magnetic contribution in the latter that is here removed. More specifically, it is shown that the electron dynamics generally washes out the *gauge* fields appearing in the adiabatic dynamics. However, at T=0 K, the *pseudo*-magnetic force is fully re-established in the typical situation where the electrons respond rapidly on the slow time-scale of the nuclear dynamics (Markov limit). Hence, we predict Berry's phase effects to be observable also in the presence of electronic friction, and non-trivial geometric phases should be attainable for molecules on metallic magnetic surfaces.

Introduction. When molecules scatter off, react or simply vibrate at metal surfaces they can induce electronic excitations in the substrate, in addition to the usual phonon excitations. This energy loss mechanism is a clear violation of the Born-Oppenheimer (BO) approximation and can give rise to intriguing phenomena, including electron transfer processes and generation of chemically-induced currents [1]. Often, though, electron excitation is of limited extent and reduces to a frictional force of electronic origin that acts on the molecular degrees of freedom, in addition to the usual Born-Oppenheimer forces, a situation where the so-called “BO dynamics with electronic friction” description is appropriate.

Electronic friction has a long history [2]. One early derivation of the frictional forces that electrons exert on a set of (classically) moving nuclei is due to Head-Gordon and Tully [3], who first derived an expression for the relevant friction kernel γ . The result of Ref. [3],

$$\gamma_{kj}^{\text{HGT}} = \pi \hbar \sum_{ab} \langle a | \partial_k h | b \rangle \langle b | \partial_j h | a \rangle \delta(\epsilon_a - \epsilon_F) \delta(\epsilon_b - \epsilon_F) \quad (1)$$

where a, b label single-particle states, k, j label nuclear degrees of freedom, h is the one-particle Hamiltonian and ϵ_F the Fermi energy, was obtained at zero temperature in the independent electron approximation, and found to be consistent with earlier results on vibrational relaxation at metal surfaces [4–6]. It was later re-derived using different methodologies, including influence functionals [7] and nonequilibrium Green's functions [8] (see Ref. [2] for a comprehensive account). This form of electronic friction, combined with *first principles* electronic structure theory [9–11], has been applied to a variety of problems [11–17], and Langevin dynamics with electronic friction and density-functional theory potentials is nowadays a standard tool to investigate the dynamics of molecules at

metal surfaces. Other works addressed the issue of non-thermal, yet steady-state, electronic baths (*e.g.*, current-carrying metals) [18, 19] and of the electron-electron interactions [20, 21], and showed the importance of going beyond a mean-field treatment of the electronic dynamics [21]. In particular, Dou, Miao & Subotnik (DMS), using a mixed quantum-classical approach, derived a completely general friction kernel that applies to interacting electrons, is valid out of equilibrium, and reduces to previously published expressions for independent electrons [21]. DMS wrote the electronic friction tensor, in the Markov limit, as

$$\gamma_{kj}^{\text{DMS}} = - \int_0^\infty \text{tr}_e \left((\partial_k H_{\text{el}}) e^{-\frac{i}{\hbar} H_{\text{el}} \tau} (\partial_j \rho) e^{+\frac{i}{\hbar} H_{\text{el}} \tau} \right) d\tau \quad (2)$$

where ρ is the steady-state electron density-operator, H_{el} is the electronic Hamiltonian, possibly including interactions between electrons, and tr_e denotes the trace over the electron degrees of freedom.

The electronic friction limit can be considered a first order “realization” of the adiabatic approximation in a situation where a continuum of electronic states and the ensuing fast relaxation guarantee that, on the time scale of nuclear motion, the electrons follow *adiabatically* the nuclei. However, the adiabatic approximation, in addition to the usual BO forces, introduces *gauge* fields that reflect the geometric properties of the electronic eigenspaces when viewed as functions of the slow parameters (the nuclear degrees of freedom), and it is not clear whether and how these disappear when the electron dynamics is taken into account. The answer to this question lies into the nuclear wavefunction, since the electronic degrees of freedom are traced out in the above electronic-friction description and the phase of the electronic wavefunction cannot be tracked in any realistic experiment. Hence, a full quantum description of the

dynamics is made necessary, and this is the purpose of the present Letter.

The Letter is organized as follows. After summarizing some basic properties of the adiabatic approximation we analyze the $T = 0$ K exact quantum dynamics of the combined electron-nuclear system using a representation that closely resembles the adiabatic one. We shall show that the introduction of the electron dynamics generally washes out the above mentioned *gauge* fields, and thus removes any Berry's phase effect from the dynamics. Later, we analyze the case of an electronic bath that relaxes quickly on the time-scale relevant for the nuclear motion, and derive an electronic-friction kernel that describes the corresponding electronic-friction dynamical regime. In the Markov limit of a memoryless friction, we shall show that the *pseudo*-magnetic forces are fully restored, thereby making geometric phase effects potentially observable. Importantly, it is further shown how the adiabatic Hamiltonian has to be modified to include friction in the nuclear dynamics, and how the equation of motion for the nuclear wavefunction is turned into a non-linear equation of the Schrödinger-Langevin type.

Adiabatic approximation. The adiabatic approximation [22] and the related adiabatic theorem [23, 24] has under-pinned research into quantum systems with slowly evolving parameters, and form the basis of the theory of energy level crossings in molecules, of the Gell-Mann–Low theorem in quantum field theory and of Berry's geometrical phase. When the slow parameters \mathbf{x} (here, the nuclear coordinates) are considered as dynamical variables the approximation can be recast as a variational approximation with the wavefunction *ansatz*

$$|\Psi_t\rangle = \int_X d\mathbf{x} \psi_t(\mathbf{x}) |u_n(\mathbf{x})\rangle |\mathbf{x}\rangle$$

where $|u_n(\mathbf{x})\rangle$ is a chosen time-independent “frame” of the n^{th} electronic eigenstate, $|\mathbf{x}\rangle$ are position eigenstates of the slow variables, and $\psi(\mathbf{x})$ is the nuclear wavefunction. The latter satisfies the variational equation of motion $H_n^{\text{eff}}\psi = i\hbar\partial_t\psi$ with the effective Hamiltonian

$$H_n^{\text{eff}} = \frac{1}{2} \sum_{ij} \xi^{ij} \hat{\pi}_i \hat{\pi}_j + (E_n - \hbar A_0 + \phi) \quad (3)$$

Here, $E_n = \langle u_n | H_{\text{el}} | u_n \rangle$ is the n^{th} potential energy surface, $H_{\text{el}} \equiv H_{\text{el}}(\mathbf{x})$ the electronic Hamiltonian, ξ^{ij} is a coordinate-independent inverse-mass tensor of the slow variables and the mechanical momentum for the j^{th} degree of freedom $\hat{\pi}_j = \hat{p}_j - \hbar A_j$ is defined in terms of the Berry's connection [25] $A_j = i \langle u_n | \partial_j u_n \rangle$ and the coordinate representation of the momentum operator \hat{p}_j .

The geometric properties of the adiabatic approximation are subsumed in the quantum geometric tensor [26, 27]

$$q_{ij} = \langle \partial_i u_n | Q_n | \partial_j u_n \rangle \quad Q_n \equiv 1 - |u_n(\mathbf{x})\rangle \langle u_n(\mathbf{x})|$$

which determines both the scalar potential $\phi(\mathbf{x})$ and the commutation properties of the $\hat{\pi}_j$'s, through its real ($\Re q_{ij} = g_{ij}$) and imaginary ($\Im q_{ij} = -B_{ij}/2$) parts, respectively, *i.e.*,

$$\phi = \frac{\hbar^2}{2} \sum \xi^{ij} g_{ij} \quad [\hat{\pi}_i, \hat{\pi}_j] = i\hbar^2 B_{ij}$$

Here, g_{ij} is a quantum metric and B_{ij} is the ij^{th} component of the Berry's curvature, *i.e.*, the exterior derivative of the linear differential form $\omega = \sum_j A_j dx^j$. Furthermore, for later convenience, we have introduced a term $A_0 = i \langle u_n | \partial_t u_n \rangle$ that allows more general, time-dependent *gauge* transformations ($A_0 \equiv 0$ is the usual choice) [28].

The ensuing nuclear dynamics is governed by the Born-Oppenheimer force, $F_k^{\text{BO}} = -\partial_k E_n$, in conjunction with a *pseudo*-Lorentz force comprising an electric

$$F_k^{\text{el}} = -\partial_k \phi = -\frac{\hbar^2}{2} \sum_{ij} \xi^{ij} \frac{\partial g_{ij}}{\partial x^k}$$

and a magnetic contribution

$$F_k^{\text{mag}} = \frac{\hbar}{2} \sum_j (\hat{v}^j B_{kj} + B_{kj} \hat{v}^j)$$

($\hat{v}^j = \sum_i \xi^{ji} \hat{\pi}_i$ being the velocity operator) that have a purely geometrical origin and represent the legacy of the adiabatic constraint to the wavefunction. The above forces are separately *gauge* invariant. An additional *gauge*-invariant component related to the electron dynamics (ED) formally exists

$$F_k^{\text{ED}} = \hbar (\partial_k A_0 - \partial_t A_k) = -2\hbar \Im \langle \partial_k u | Q_n | \partial_t u \rangle \quad (4)$$

but vanishes identically in the adiabatic approximation.

As shown in the Supplemental Material (SM), the quantum geometric tensor also determines the local-in-time error [29, 30] of the adiabatic approximation. The latter takes approximately the form of an expectation value of the quantized quantum geometric tensor

$$\varepsilon^2 \approx \langle \psi | \sum_{ij} \hat{v}^i q_{ij} \hat{v}^j | \psi \rangle_X$$

(where X denotes integration w.r.t. the nuclear DOFs only) and measures the tendency of the system to undergo a non-adiabatic transition at short-time. In fact, $P_{\text{nad}} \approx \varepsilon^2 (t - t_s)^2$ is the total transition probability, if the adiabatic approximation were suddenly lifted at $t = t_s$. For comparison, we notice that if the slow variables were simply some parameters, and not dynamical variables, we would have exactly

$$\varepsilon^2 = \sum_{ij} V^i q_{ij} V^j \equiv \sum_{ij} V^i g_{ij} V^j$$

where V^i is the classical velocity of the i^{th} parameter (see SM for details).

Exact dynamics. The *exact* wavefunction can be yet represented in a form similar to above,

$$|\Psi_t\rangle = \int_{\mathbf{X}} d\mathbf{x} \psi_t(\mathbf{x}) |u_t(\mathbf{x})\rangle |\mathbf{x}\rangle$$

but now with a time-dependent electronic state $|u_t(\mathbf{x})\rangle$ for each nuclear configuration \mathbf{x} . This is the so-called exact-factorization of the wavefunction [31, 32], which is an ‘‘intermediate’’ representation that can be obtained by introducing a local basis of nuclear states $\{|\mathbf{x}\rangle\}$, and imposing a normalization condition on the ensuing local electronic states,

$$\langle \mathbf{x} | \Psi_t \rangle_X = \psi_t(\mathbf{x}) |u_t(\mathbf{x})\rangle \langle u_t(\mathbf{x}) | u_t(\mathbf{x}) \rangle = 1$$

The corresponding equations of motion [31, 32] are derived in SM using a projection-operator technique that emphasizes their *gauge* transformation properties. They can be summarized as follows. The nuclear wavefunction satisfies $H^{\text{eff}}\psi = i\hbar\partial_t\psi$, where the effective Hamiltonian takes precisely the form of Eq. 3, but now $|u(\mathbf{x}, t)\rangle$ replaces $|u_n(\mathbf{x})\rangle$ everywhere and $A_0 = i\langle u | \partial_t u \rangle$ is an arbitrary real *gauge* constraint that guarantees normalization of the local electronic state. The ensuing equation of motion for the *exact* nuclear wavefunction are thus formally *very* similar to the adiabatic ones. The additional force, due solely to the electron dynamics, is the *gauge*-invariant term of Eq. 4 that, differently from the adiabatic approximation, is generally non zero. That is, the total force reads *exactly* as $F_k^{\text{tot}} = F_k^{\text{BO}} + F_k^{\text{el}} + F_k^{\text{mag}} + F_k^{\text{ED}}$, where $F_k^{\text{BO}} \equiv -\partial_k \langle u | H_{\text{el}} | u \rangle = -\partial_k E_{\text{el}}$ is a time-dependent BO force, the geometric forces F_k^{el} and F_k^{mag} stem from the time-dependent geometric tensor q_{ij} and F_k^{ED} describes the electron dynamics.

The electronic equation of motion takes the form

$$i\hbar Q \partial_t |u\rangle = Q H_{\text{el}} |u\rangle + K[\psi_t] |u\rangle \quad (5)$$

where the second term on the r.h.s. describes the ‘electron drag’ with the nuclear motion and reads as

$$K[\psi_t] |u(t)\rangle = -i\hbar \sum_j V^j Q |\partial_j u(t)\rangle - \hbar R |u(t)\rangle \quad (6)$$

Here $V^j(\mathbf{x}) = (\hat{v}^j \psi_t) / \psi_t$ is the (*gauge*-invariant) time-dependent, complex-valued nuclear velocity field, $Q = 1 - |u\rangle \langle u|$, $R |u\rangle = \frac{\hbar}{2} \sum_{ij} \xi^{ij} D_{ij} |u\rangle$, and

$$D_{ij} |u\rangle = iA_i Q |\partial_j u\rangle + iA_j Q |\partial_i u\rangle + Q |\partial_i \partial_j u\rangle$$

is a second derivative of the electronic states. In this form the effective electronic Hamiltonian is the sum of *gauge* tensorial terms G , *i.e.*, of terms that behave simply as $G |u\rangle \rightarrow e^{-i\varphi} G |u\rangle$ under the *gauge* transformation $|u\rangle \rightarrow e^{-i\varphi} |u\rangle$. This Hamiltonian can be used to include

explicitly the electron reaction into the nuclear equation of motion. Specifically, plugging $Q\partial_t |u\rangle$ into F_k^{ED} above one obtains a correction to the previous *pseudo*-electric and *pseudo*-magnetic forces, that are turned into

$$F_k^{\text{el,c}} = 2\hbar \sum_j g_{kj} \Im V^j - \hbar^2 \sum_{ij} \xi^{ij} (\Re \langle \partial_i u | D_{kj} u \rangle + \Re \langle \partial_k u | D_{ij} u \rangle) \quad (7)$$

$$F_k^{\text{mag,c}} = \frac{\hbar}{2} \sum_j (\hat{v}^j B_{kj} + B_{kj} \hat{v}^j) - \hbar \sum_j B_{kj} \Re V^j \quad (8)$$

plus a genuine non-Born-Oppenheimer term

$$F_k^{\text{NBO}} = 2\Re \langle \partial_k u | Q H'_{\text{el}} | u \rangle$$

where $H'_{\text{el}} = H_{\text{el}} - E_{\text{el}}$ for later convenience. The latter force vanishes identically when $|u\rangle$ is an eigenstate of H_{el} and, more generally, is bound by the size of the (local) energy fluctuations in the electronic subsystem, $\Delta E_{\text{el}}^2 = \langle (H_{\text{el}} - E_{\text{el}})^2 \rangle$, through the k^{th} diagonal component of the quantum geometric tensor, namely $|F_k^{\text{NBO}}| \leq 2\Delta E_{\text{el}} \sqrt{g_{kk}}$. Thus, the total force acting on the k^{th} nuclear degree of freedom can be written as

$$F_k^{\text{tot}} = F_k^{\text{BO}} + F_k^{\text{el,c}} + F_k^{\text{mag,c}} + F_k^{\text{NBO}}$$

and represents the same total force given above but now explicitly including the electron reaction. The key point about this expression is that, as shown in SM, the dynamically corrected *pseudo*-Lorentz force vanishes *identically* when averaged over an arbitrary nuclear state since

$$\langle \psi | F_k^{\text{el,c}} | \psi \rangle_X = \langle \psi | F_k^{\text{mag,c}} | \psi \rangle_X = 0$$

and thus it disappears in the quantum-classical limit where the electronic friction approximation is often invoked. Hence, electronic friction must come from the term F_k^{NBO} .

We stress that the above results are *exact* and tell us that, in general, the effect of the electron dynamics is to wash out the *pseudo*-Lorentz force appearing in the adiabatic limit. As shown in SM, the vanishing of the average *pseudo*-electric force expresses conservation of the quantum metric, since such force takes the form of a(n expectation value of the) Ricci-Levi Civita covariant derivative of the metric tensor. This appears reasonable since the quantum metric and the related *pseudo*-electric force measure the error in the adiabatic *approximation* while the dynamics considered here is exact. Likewise, the vanishing of the *pseudo*-magnetic force signals the quenching of geometric phase effects that, in fact, should disappear when the dynamics is far from the adiabatic regime[33].

Electronic friction. Let us now focus on the electronic equation Eq. 5 in the situation where the electronic system relaxes quickly to the ground-state $|u_0\rangle$

and the deviation $|\Delta u\rangle = |u(t)\rangle - |u_0(t)\rangle$ remains small throughout the nuclear dynamical evolution (here, $|u_0(t)\rangle$ is the time-evolving ground electronic state). This is the condition where linear response theory (LRT) applies and also the situation where the electronic-friction picture is appropriate (see SM for details). Under these circumstances, $\langle \Delta u | u_0 \rangle \equiv 0$, the BO forces F_k^{BO} acting on the molecular degrees of freedom are the same as the ground-state ones, $E_{\text{el}} \approx E_0 + 2E_0 \Re \langle \Delta u | u_0 \rangle \equiv E_0$, and $F_k^{\text{NBO}} \approx 2 \Re \langle \partial_k u_0 | Q_0 H'_{\text{el}} | \Delta u \rangle$ represents the main effect of the electron reaction. Hence, upon plugging the LRT result for $|\Delta u\rangle$ into the above expression for F_k^{NBO} one obtains a friction-like term $F_k^\gamma = -2 \sum_j \Re \left(\int_0^\infty \Gamma_{kj}(\tau) V^j(t-\tau) d\tau \right)$, where V^j becomes the real particle velocity in the classical limit, and Γ is the kernel

$$\Gamma_{kj}(\tau) = \langle \partial_k u_0 | Q_0 H'_{\text{el}} e^{-\frac{i}{\hbar} H'_{\text{el}} \tau} | \partial_j u_0 \rangle$$

In the Markov limit we can set $V^j(t-\tau) \approx V^j(t)$ and the frictional force takes the form $F_k^\gamma = -\sum_j \bar{\gamma}_{kj} V^j$ where $\bar{\gamma}_{kj}$ is the zero-frequency limit of the frequency-dependent kernel $\bar{\gamma}_{kj}(\omega) = 2 \lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon \tau} e^{i\omega \tau} \Gamma_{kj}(\tau) d\tau$. As shown in SM, the real part of the tensor $\bar{\gamma}_{kj}$ is precisely the $T \rightarrow 0$ limit of the mixed quantum-classical DMS expression [21], Eq. 2, and contains a *pseudo*-magnetic contribution $2\hbar \Im \langle \partial_k u_0 | Q_0 | \partial_j u_0 \rangle = 2\hbar \Im q_{kj} = -\hbar B_{kj}$. The latter, when used to evaluate the forces acting on the nuclei, is seen to give a term $+\hbar \sum_j B_{kj} \Re V^j$ that precisely cancels the magnetic dynamical correction introduced above (see Eq. 8). This corrective effect is physically sound: electronic friction cools the nuclear motion and enforces the adiabatic limit, with its *gauge* fields. Upon removing the spurious *pseudo*-magnetic contribution, the ordinary electronic friction tensor is the real part of the expression

$$\gamma_{kj} = 2\pi \hbar \langle \partial_k u_0 | Q_0 (H_{\text{el}} - E_0) \delta(E_0^+ - H_{\text{el}}) | \partial_j u_0 \rangle \quad (9)$$

where $E_0^+ = E_0 + \hbar\omega$ in the limit $\omega \rightarrow 0^+$ is understood. Notice the symmetries $\gamma'_{kj} = \gamma'_{jk}$ and $\gamma''_{kj} = -\gamma''_{jk}$ for the real and imaginary parts of γ_{kj} , respectively, and the fact that γ'_{kj} satisfies the second fluctuation-dissipation relation as described in Ref. [21] (in the $T \rightarrow 0$ K limit considered here), since the argument used by Dou *et al.* remain valid in this context. Furthermore, upon using $\langle \partial_k u_0 | Q_0 H'_{\text{el}} = -\langle u_0 | (\partial_k H_{\text{el}}) Q_0$ we can equivalently rewrite γ'_{kj} as

$$\gamma'_{kj} = -2\pi \hbar \Re \langle u_0 | (\partial_k H_{\text{el}}) \delta(E_0^+ - H_{\text{el}}) | \partial_j u_0 \rangle \quad (10)$$

It is this expression that reduces to Eq. 1 in the independent electron approximation (see SM), and that shows more explicitly why γ'_{kj} needs a manifold of states lying infinitesimally close to the ground state to be non-vanishing. [The required excitation energy $\hbar\omega \rightarrow 0$ can also be viewed as the “running” correction to E_0 in the dynamical phase factor of the evolving electronic state.]

In the context of the exact factorization of the wavefunction, the electronic-friction regime considered here is best handled by using local electronic states in the “standard” *gauge*, *i.e.*, $\langle u^+ | \partial_t u^+ \rangle = 0$ where $^+$ denotes this *gauge*. In the LRT limit this amounts to setting $E_{\text{el}} = E_0 = 0$ in the electronic problem and writing $|u^+\rangle = |u_0\rangle + |\Delta u^+\rangle$, where now only the second term is time-dependent. This implies that the nuclear Hamiltonian in the chosen *gauge*, H^+ , very closely resembles the $n = 0$ adiabatic Hamiltonian of Eq. 3. In fact, as shown in SM, the main difference is a modified vector potential $A_k \rightarrow A_k + \delta A_k$ where

$$\begin{aligned} \delta A_k &= 2 \Im \langle \partial_k u_0 | \Delta u_0 \rangle \\ &\approx -2 \sum_j \Im \int_0^\infty \langle \partial_k u_0 | e^{-\frac{i}{\hbar} H'_{\text{el}} t'} Q_0 | \partial_j u_0 \rangle V_{t-t'}^j dt' \end{aligned}$$

This term is of second order in the spatial derivatives of the electronic states and generates a correction of the same order to the adiabatic forces through its time dependence, $F_k \rightarrow F_k - \hbar \partial_t (\delta A_k)$. An integration by parts transforms δA_k into

$$\delta A_k = -2 \sum_j \Im (q_{kj} X^j) + \frac{2}{\hbar} \sum_j \Re \left(\int_0^\infty \Gamma_{kj}(t') X_{t-t'}^j dt' \right) \quad (11)$$

where X^j 's are the integrated velocity fields, $X^j(\mathbf{x}, t) = \int_{-\infty}^t V^j(\mathbf{x}, t') dt'$ and the first term is seen to be the source of the dynamical correction to the *pseudo*-Lorentz force (Eq.s 7,8). In the Markov limit, as seen above, only the integrated kernel $\bar{\gamma}_{kj} = 2 \lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon \tau} \Gamma_{kj}(\tau) d\tau$ matters, for which we have $\bar{\gamma}_{kj} = -2i\hbar q_{kj} + \gamma_{kj}$. Hence, in this limit Eq. 11 reduces to

$$\delta A_k = \hbar^{-1} \sum_j \Re (\gamma_{kj} X_t^j) \quad (12)$$

where the same cancellation effect occurs as discussed above.

Eq. (12) is a key result of this Letter. It represents the simple amendment to the adiabatic Hamiltonian that is necessary in order to include the effect of electronic friction into the quantum dynamics of the nuclei, in the most relevant case where the Markov limit applies. This term turns the Schrödinger equation into a non-linear equation of motion, which conserves the wavefunction norm and describes energy dissipation. In the simplest case where $\xi^{ij} = \delta^{ij} M^{-1}$, if γ_{kj} can be taken *diagonal* and *uniform* in the configuration space of the system where the dynamics occurs, one finds

$$\delta A_k = \Re (\gamma X_t^k) \approx \partial_k \Re \left(-i\hbar M^{-1} \gamma \int_{-\infty}^t \ln \psi_{t'}(\mathbf{x}) dt' \right)$$

upon neglecting the contribution of the vector potential to the velocity field. Hence, δA_k becomes longitudinal and can be replaced by an appropriate *scalar* field

$\delta\phi \approx \hbar M^{-1} \gamma \Im(\ln \psi_i(\mathbf{x}))$ for negligible γ'' . This scalar potential is precisely the “phase potential” appearing in the effective Hamiltonian of the (non-linear) Kostin equation [34], and represents a very simple way to introduce dissipation into a Schrödinger-like equation.

We remark that the resulting equation for the nuclear wavefunction has no fluctuating term since it describes the evolution of electronically averaged quantities, *i.e.*, the nuclear observables dressed by the electronic state.

Conclusions. We have developed a theory of electronic friction that describes the nuclear dynamics in a quantum setting at $T = 0$ K. Friction is seen to turn the equation of motion for the nuclear wavefunction into a non-linear equation, where the vector potential depends on the past wavefunction behaviour. This low-temperature, frictional limit seems to be ideal to explore the nuclear dynamics in a situation where energy dissipation through excitation of electron-hole pairs of a metallic substrate *enforces* the adiabatic dynamics. We have shown that in this limit the *gauge* fields appropriate for an adiabatic dynamics are fully restored, hence we expect Berry’s phase effects to be operative. In molecular problems, the latter typically reduce to a sign change, since the Berry’s connection is flat in the presence of time-reversal symmetry. However, in the presence of a magnetic field (that here can come from the metallic surface itself) a non-zero curvature is present, thereby allowing the geometric phase to attain arbitrary values. A magnetic field also modifies the topology of conical intersections by increasing the dimensionality of the branching space, and turns the intersection points into *pseudo*-magnetic monopoles [27]. More generally, the *pseudo*-magnetic field is known to shield the physical magnetic field and allow the nuclei to behave essentially as neutral particles (as far as magnetic effects are of concern) [35]. However, the degree of shielding depends on the nuclear geometry and may change along the dynamical evolution [36]. The effects of these phenomena on elementary gas-surface processes remain yet to be explored.

* rocco.martinazzo@unimi.it

- [1] H. Nienhaus, Surface Science Reports **45**, 1 (2002).
 [2] W. Dou and J. E. Subotnik, Journal of Chemical Physics **148** (2018), 10.1063/1.5035412.
 [3] M. Head-Gordon and J. C. Tully, The Journal of Chemical Physics **103**, 10137 (1995).
 [4] B. N. Persson and M. Persson, Solid State Communications **36**, 175 (1980).
 [5] M. Persson and B. Hellsing, Physical Review Letters **49**, 662 (1982).
 [6] M. Head-Gordon and J. C. Tully, The Journal of Chemical Physics **96**, 3939 (1992).
 [7] M. Brandbyge, P. Hedegård, T. F. Heinz, J. A. Misewich, and D. M. Newns, Physical Review B **52**, 6042 (1995).
 [8] J. T. Lü, M. Brandbyge, P. Hedegård, T. N. Todorov, and D. Dundas, Physical Review B - Condensed Matter and Materials Physics **85**, 201401 (2012), arXiv:1205.0745.
 [9] R. J. Maurer, M. Askerka, V. S. Batista, and J. C. Tully, Physical Review B **94**, 115432 (2016).
 [10] Z. Jin and J. E. Subotnik, The Journal of Chemical Physics **150**, 164105 (2019).
 [11] Y. Zhang, R. J. Maurer, and B. Jiang, The Journal of Physical Chemistry C (2019), 10.1021/ACS.JPCC.9B01111.
 [12] A. C. Luntz and M. Persson, Journal of Chemical Physics **123**, 074704 (2005).
 [13] A. C. Luntz, M. Persson, S. Wagner, C. Frischkorn, and M. Wolf, Journal of Chemical Physics **124**, 244702 (2006).
 [14] G. Füchsel, T. Klamroth, S. Monturet, and P. Saalfrank, Physical Chemistry Chemical Physics **13**, 8659 (2011).
 [15] S. Monturet and P. Saalfrank, Physical Review B - Condensed Matter and Materials Physics **82** (2010), 041401.
 [16] M. Askerka, R. J. Maurer, V. S. Batista, and J. C. Tully, Physical Review Letters **116**, 217601 (2016).
 [17] C. L. Box, Y. Zhang, R. Yin, B. Jiang, and R. J. Maurer, JACS Au **1**, 164 (2020).
 [18] N. Bode, S. V. Kusminskiy, R. Egger, and F. Von Oppen, Physical Review Letters **107**, 036804 (2011).
 [19] N. Bode, S. V. Kusminskiy, R. Egger, and F. von Oppen, Beilstein Journal of Nanotechnology **3**, 144 (2012).
 [20] J. Daligault and D. Mozyrsky, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics **85**, 041111 (2012).
 [21] W. Dou, G. Miao, and J. E. Subotnik, Physical Review Letters **119**, 046001 (2017).
 [22] M. Born and R. Oppenheimer, Annalen der Physik **389**, 457 (1927).
 [23] M. Born and V. Fock, Zeitschrift für Physik **51**, 165 (1928).
 [24] T. Kato, Journal of the Physical Society of Japan **5**, 435 (1950).
 [25] Henceforth, for notational convenience, we shall omit the n dependence of A_j , $\hat{\pi}_j$, etc. and of the quantum geometric tensor q_{ij} introduced below.
 [26] J. P. Provost and G. Vallee, Communications in Mathematical Physics **76**, 289 (1980).
 [27] M. Berry, Proceedings of the Royal Society of London. A. Mathematics and Physics **49**, 67 (1960).
 [28] It is worth noticing that the transformation $|u_n\rangle \rightarrow e^{-iE_n t/\hbar} |u_n\rangle$ moves the BO PES into the kinetic energy term, since $\hbar A_0 \equiv E_n$ in this *gauge*, and $\hbar A_k \rightarrow \hbar A_k + (\partial_k E_n)t$.
 [29] R. Martinazzo and I. Burghardt, Physical Review Letters **124**, 150601 (2020), arXiv:1907.00841.
 [30] R. Martinazzo and I. Burghardt, (2021), arXiv:2102.12117.
 [31] A. Abedi, N. T. Maitra, and E. K. U. Gross, Physical Review Letters **105**, 123002 (2010).
 [32] A. Abedi, N. T. Maitra, and E. K. U. Gross, The Journal of Chemical Physics **137**, 22A530 (2012).
 [33] We remark that the adiabatic approximation does *not* correspond to $F_k^{\text{NBO}} = 0$ – that would leave the electron corrections to the forces – rather to $F_k^{\text{ED}} = 0$. In a sense, the Born-Oppenheimer approximation incorporates these corrections and thus accounts partially for the electron reaction, thereby removing geometric phase effects and the repulsive *pseudo*-electric forces that become impor-

tant at conical intersections.

- [34] M. D. Kostin, The Journal of Chemical Physics **57**, 3589 (2003).
 [35] R. Resta, Journal of Physics: Condensed Matter **12**, R107 (2000).
 [36] D. Ceresoli, R. Marchetti, and E. Tosatti, Physical Review B **75**, 161101 (2007).

SUPPLEMENTAL MATERIAL

A. Adiabatic approximation

I. Slow variables as external parameters

When the slow variables \mathbf{x} are regarded as parameters that are under the control of the experimenter, the Hamiltonian governing the evolution of the system has a pre-defined time-dependence $H = H(\mathbf{x}(t))$ for any given path $\mathbf{x}(t)$ of the parameters. The adiabatic approximation can be recast as a variational approximation

$$i\hbar \frac{d}{dt} |\Psi_t\rangle = H(t) |\Psi_t\rangle \quad |\Psi_0\rangle = |u_n(0)\rangle$$

which uses a *time-dependent* variational manifold $\mathcal{V}(t)$ (the n^{th} eigenspace of $H(t)$, here assumed to be non-degenerate) and a complex-analytic representation of the wavefunction. In other words, one describes the wavefunction at any time in the form $|\Psi\rangle = C |u_n(s)\rangle$ where C is the only (complex) *variational* parameter of the problem. The Dirac-Frenkel condition amounts to

$$P_n (i\hbar\partial_t - H(t)) |\Psi_t\rangle = 0$$

where $P_n = P_n(t)$ is the instantaneous eigenprojector on the target manifold and $Q_n = 1 - P_n$ its orthogonal complement. Hence

$$i\hbar |\dot{\Psi}_t\rangle = H_{PP}(t) |\Psi_t\rangle + i\hbar \dot{P}_n |\Psi_t\rangle$$

where $Q_n |\dot{\Psi}_t\rangle \equiv \dot{P}_n |\Psi_t\rangle$ has been used to re-write the time-derivative in terms of the manifold dynamics (contained in \dot{P}_n) and of $H_{PP} \equiv P_n H P_n$. Furthermore, since $P_n \dot{P}_n |\Psi_t\rangle \equiv 0$ the equation of motion can be recast as an effective Schrödinger equation

$$i\hbar \frac{d}{dt} |\Psi_t\rangle = H_n(t) |\Psi_t\rangle$$

involving the self-adjoint effective Hamiltonian

$$H_n(t) = H_{PP}(t) + i\hbar [\dot{P}_n, P_n]$$

Local-in-time-error in case I
(Slow variables as parameters)

The local-in-time error (LITE) accompanying a variational solution [29] takes the general form

$$\varepsilon[\Psi] = \hbar^{-1} \|(i\hbar\partial_t - H) \Psi\|$$

and, in the case of a time-dependent, complex-analytic manifold, can be given in terms of the above introduced variational Hamiltonian as

$$\begin{aligned} \hbar^2 \varepsilon^2[\Psi] &= \|(H_n - H) \Psi\|^2 \\ &\equiv \left\| \left(H - i\hbar \dot{P}_n \right) \Psi \right\|^2 - \left\| \left(H_n - i\hbar \dot{P}_n \right) \Psi \right\|^2 \end{aligned}$$

This result follows upon noticing that, for $|\Psi\rangle = P_n |\Psi\rangle$, we have $(H - H_n)|\Psi\rangle = Q_n(H - i\hbar\dot{P}_n)|\Psi\rangle$ hence

$$\begin{aligned}\hbar^2\varepsilon^2[\Psi] &= \langle\Psi|(H + i\hbar\dot{P}_n)Q_n(H - i\hbar\dot{P}_n)|\Psi\rangle \\ &= \langle\Psi|(H + i\hbar\dot{P}_n)(H - i\hbar\dot{P}_n)|\Psi\rangle + \\ &\quad - \langle\Psi|(H + i\hbar\dot{P}_n)P_n(H - i\hbar\dot{P}_n)|\Psi\rangle \\ &\equiv \|(H - i\hbar\dot{P}_n)\Psi\|^2 - \|H_{PP}\Psi\|^2\end{aligned}$$

which is the same as the above result since in the last expression H_{PP} can be replaced with $H_n - i\hbar\dot{P}_n$.

Now, upon noticing that in the adiabatic problem $H_{PP} \equiv E_n P_n$,

$$\hbar^2\varepsilon^2[\Psi] \equiv \left\| E_n \Psi - i\hbar\dot{P}_n \Psi \right\|^2 - \|E_n \Psi\|^2$$

we find

$$\varepsilon = \|\dot{P}_n \Psi\|$$

where $\langle\Psi|\dot{P}_n|\Psi\rangle = 0$ has been used. This error is a purely geometrical property, *i.e.*, for a given infinitesimal displacement in parameter space it does *not* depend on the evolution time. In fact, using $\|\dot{P}_n \Psi\| = \|\dot{P}_n u_n\| = \|Q_n \dot{u}_n\|$, the LITE is seen to be a property of the moving manifold only and takes the form $\varepsilon^2 = \langle\dot{u}_n|Q_n|\dot{u}_n\rangle$ or, equivalently, introducing the parameter dependence,

$$\varepsilon^2 = \sum_{i,j} \langle\partial_i u_n|Q_n|\partial_j u_n\rangle \dot{x}^i \dot{x}^j = \sum_{i,j} g_{ij} \dot{x}^i \dot{x}^j$$

where $\partial_i \equiv \partial/\partial x^i$ and g_{ij} is the symmetric part of the covariant tensor on the parameter space

$$q = \sum_{i,j} \langle\partial_i u_n|Q_n|\partial_j u_n\rangle dx^i dx^j$$

known as quantum geometric tensor. Such symmetric (real) part g is a quantum metric (a Fubini-Study metric on the tangent bundle), while the antisymmetric (imaginary) part is related to the Berry's curvature. In a single equation,

$$q = g - \frac{i}{2} d\omega$$

where d denotes the exterior derivative and ω is the differential form $\omega = \sum_j A_j dx^j$, *i.e.*,

$$d\omega = \sum_{ij} B_{ij} dx^i dx^j$$

Here, $A_j = i\langle u_n|\partial_j u_n\rangle$ subsumes the Berry's connection on the vector bundle $\pi : E \rightarrow \mathcal{M}$ defined by the parametric dependence of the given eigenspace and $B_{ij} \equiv \partial_i A_j - \partial_j A_i$ (here \mathcal{M} is the nuclear configuration space and $\pi^{-1}(\mathbf{x})$ is the n^{th} eigenspace of the electronic Hamiltonian H_{el}).

II. Slow variables as dynamical variables

If the slow variables are considered as dynamical variables the electron-nuclear wavefunction is written as

$$|\Psi\rangle = \int d\mathbf{x} \psi(\mathbf{x}) |u_n(\mathbf{x})\rangle |\mathbf{x}\rangle$$

The variational manifold is complex and application of the variational principle reduces to the Dirac-Frenkel condition,

$$\int d\mathbf{x}' d\mathbf{x} \delta\psi^*(\mathbf{x}') \langle u_n(\mathbf{x}'), \mathbf{x}' | [i\hbar\partial_t - H] |u_n(\mathbf{x}), \mathbf{x}\rangle \psi(\mathbf{x}) = 0$$

Here

$$\langle u_n(\mathbf{x}'), \mathbf{x}' | H |u_n(\mathbf{x}), \mathbf{x}\rangle = \delta(\mathbf{x} - \mathbf{x}') \left(\langle\hat{T}\rangle_n + E_n(\mathbf{x}) \right)$$

where $E_n(\mathbf{x})$ is the Born - Oppenheimer potential energy surface and $\langle\hat{T}\rangle_n$ is the coordinate-representation of the nuclear kinetic energy operator *averaged* over the electronic state,

$$\langle\hat{T}\rangle_n = \langle u_n(\mathbf{x}) | \hat{T} | u_n(\mathbf{x}) \rangle$$

Setting $H_n = \langle T \rangle_n + E_n(\mathbf{x})$, the variational equation of motion takes the form of a Schrödinger equation for the nuclear wavefunction

$$H_n \psi = i\hbar \frac{\partial \psi}{\partial t}$$

with an effective Hamiltonian specific of the electronic state under consideration. The main difference with respect to the common Born-Oppenheimer Hamiltonian lies in the nuclear kinetic operator which gets *dressed* by the electronic motion: this dressing is the way the *gauge* fields originate from the geometric properties of the adiabatic approximation. In this context, it may be worth noticing that the derived equation of motion, *differently from the BO Hamiltonian*, correctly conserves energy since it is variational, and this occurs upon including the above mentioned *gauge* fields.

Let us now examine the dressed kinetic energy operator. Let T be of the form

$$\hat{T} = \frac{1}{2} \sum_{ij} \xi^{ij} \hat{p}_i \hat{p}_j$$

and notice that

$$\begin{aligned}\langle\hat{p}_i \hat{p}_j\rangle_n &= \hat{p}_i \hat{p}_j - i\hbar \langle u_n | \partial_j u_n \rangle \hat{p}_i \\ &\quad - i\hbar \langle u_n | \partial_i u_n \rangle \hat{p}_j - \hbar^2 \langle u_n | \partial_i \partial_j u_n \rangle\end{aligned}$$

where, as above, $A_j = i\langle u_n | \partial_j u_n \rangle \in \mathbb{R}$ and

$$\begin{aligned}\langle u_n | \partial_i \partial_j u_n \rangle &= \partial_i \langle u_n | \partial_j u_n \rangle - \langle \partial_i u_n | \partial_j u_n \rangle \\ &= -i\partial_i A_j - \langle \partial_i u_n | Q \partial_j u_n \rangle - \langle \partial_i u_n | P | \partial_j u_n \rangle \\ &= -i\partial_i A_j - q_{ij} - A_i A_j\end{aligned}$$

where q_{ij} is the ij^{th} component of the quantum geometric tensor (we omit the index n from these quantities). Hence,

$$\langle \hat{p}_i \hat{p}_j \rangle_n = (\hat{p}_i - \hbar A_i) (\hat{p}_j - \hbar A_j) + \hbar^2 q_{ij}$$

and, since ξ^{ij} is symmetric,

$$\langle T \rangle_n = \frac{1}{2} \sum_{ij} \xi^{ij} (\hat{p}_i - \hbar A_i) (\hat{p}_j - \hbar A_j) + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} g_{ij}$$

Clearly, the *dressed* operator contains terms analogous to a vector (A_i) and scalar (g_{ij}) electromagnetic potential and the latter modify the nuclear dynamics, when a comparison is made with the simpler Born-Oppenheimer one. To see the effect of the *gauge* fields on the nuclear dynamics it is enough to consider the nuclear's velocity

$$\hat{v}^k = \frac{i}{\hbar} [\langle H \rangle_n, \hat{x}^k] = \sum_j \xi^{kj} \hat{\pi}_j$$

where $\hat{\pi}_k = \hat{p}_k - \hbar A_k$ is thus the mechanical momentum for the k^{th} degree of freedom. The latter satisfies the commutation relation

$$[\hat{\pi}_i, \hat{\pi}_j] = i\hbar^2 B_{ij}$$

where $B_{ij} = \partial_i A_j - \partial_j A_i$ is the ij^{th} component of the Berry's curvature, *i.e.*, $d\omega \equiv \sum_{ij} B_{ij} dx^i dx^j$ (with $B_{ij} = -B_{ji}$). Notice that the commutator is *gauge* independent [The *gauge* freedom mentioned here is the arbitrariness in the choice of the electronic frame. A *gauge* transformation $|u_n\rangle \rightarrow e^{-i\varphi} |u_n\rangle$ amounts to adding an *exact* 1-form to ω without altering the scalar potential ($\omega \rightarrow \omega + d\varphi$) and, at the same time, to adding a phase factor to the nuclear wavefunction, $\psi \rightarrow e^{+i\varphi} \psi$. In this respect, the *gauge* transformations involved here are more limited than those allowed for a true electromagnetic potential, since they are always stationary w.r.t. t]. The force is then obtained by the rate of variation of particles' mechanical momentum,

$$\dot{\hat{\pi}}^k = \frac{i}{\hbar} [\langle H \rangle_n, \hat{\pi}^k] = \frac{i}{\hbar} \left[\frac{1}{2} \sum_{ij} \xi^{ij} \hat{\pi}_i \hat{\pi}_j + E'_n, \hat{\pi}^k \right]$$

where $E'_n = E_n + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} g_{ij}$. It is convenient to separate two contributions. The first does *not* involve derivatives of the inverse mass tensor and reads as

$$\begin{aligned} \hat{W}^k &= \frac{i}{\hbar} \left\{ \frac{1}{2} \sum_{ij} \xi^{ij} [\hat{\pi}_i \hat{\pi}_j, \hat{\pi}_k] + [E'_n, \hat{\pi}_k] \right\} \\ &= -\frac{\hbar}{2} \sum_{ij} \xi^{ij} (\hat{\pi}_i B_{jk} + B_{ik} \hat{\pi}_j) - \partial_k E'_n \end{aligned}$$

whereas the second only appears when the mass tensor is coordinate-dependent,

$$\begin{aligned} \hat{U}^k &= \frac{i}{2\hbar} \sum_{ij} [\xi^{ij}, \hat{\pi}_k] \hat{\pi}_i \hat{\pi}_j = \frac{i}{2\hbar} \sum_{ij} [\xi^{ij}, \hat{p}_k] \hat{\pi}_i \hat{\pi}_j \\ &= -\frac{1}{2} \sum_{ij} (\partial_k \xi^{ij}) \hat{\pi}_i \hat{\pi}_j \end{aligned}$$

Henceforth we consider only the first term and write the k^{th} component of the force as

$$F_k = -\partial_k E'_n + \frac{\hbar}{2} \sum_j (\hat{v}^j B_{kj} + B_{kj} \hat{v}^j)$$

that is, $F_k = F_k^{\text{BO}} + F_k^{\text{el}} + F_k^{\text{mag}}$, where the first term represents the Born-Oppenheimer force

$$F_k^{\text{BO}} = -\frac{\partial E_n(\mathbf{x})}{\partial x^k}$$

while the latter two form an effective Lorentz force which comprises both an electric component

$$F_k^{\text{el}} = -\frac{\hbar^2}{2} \sum_{ij} \xi^{ij} \frac{\partial g_{ij}}{\partial x^k}$$

arising from the Fubini-Study metric tensor, and a magnetic component

$$F_k^{\text{mag}} = \frac{\hbar}{2} \sum_j (\hat{v}^j B_{kj} + B_{kj} \hat{v}^j)$$

due to the Berry's curvature. (To check that this indeed represents a *pseudo*-magnetic force one can consider the three-dimensional case and observe that $B_{xy} = H_z$, $B_{xz} = -H_y$, $B_{yz} = H_x$, where \mathbf{H} is the *pseudo*-magnetic field. Hence $\mathbf{F}^{\text{mag}} = \frac{\hbar}{2} [\mathbf{v} \wedge \mathbf{H} - \mathbf{H} \wedge \mathbf{v}]$, which is the correct quantum expression of the magnetic component of the Lorentz force). The two components behave very differently from each other: the *pseudo*-magnetic field may vanish almost everywhere (see below) yet give rise to observable effects, similarly to what happens with the Aharonov-Bohm effect, while the *pseudo*-electric field is ubiquitous (*i.e.*, it does not vanish unless the adiabatic error is uniform over the configuration space sampled by the nuclei) but typically of secondary importance and seldom considered in practice. Amazingly, though, they both arise from one and the same object, namely the quantum geometric tensor, here re-written in a form that makes explicit its connections with the omitted adiabatic states

$$\begin{aligned} q &= \sum_{ij} \sum_m \langle \partial_i u_n | u_m \rangle \langle u_m | \partial_j u_n \rangle dx^i dx^j \\ &= \sum_{ij} \sum_m \frac{\langle u_n | \partial_i H | u_m \rangle \langle u_m | \partial_j H | u_n \rangle}{(E_n - E_m)^2} dx^i dx^j \end{aligned}$$

Local-in-time error in case II
[Slow variables as dynamical variables]

Next we consider the LITE in the “dynamic” adiabatic approximation [29]. To this end, we need the time-derivative of the whole wavefunction in the “standard” gauge $\langle \Psi | \dot{\Psi}^+ \rangle = 0$ (where the superscript + is used for the trajectory $|\Psi\rangle = |\Psi(t)\rangle$ to denote this *gauge*) and in particular its squared norm. By this we mean

$$\hbar^2 \|\dot{\Psi}^+\|^2 = \int d\mathbf{x} \psi^*(\mathbf{x}) \left[\langle \hat{T} \rangle_n + (E_n(\mathbf{x}) - \bar{E}) \right]^2 \psi(\mathbf{x})$$

where $\langle T \rangle_n$ is the dressed kinetic energy operator introduced above

$$\langle \hat{T} \rangle_n = \frac{1}{2} \sum_{ij} \xi^{ij} \hat{\pi}_i \hat{\pi}_j + \phi$$

with

$$\hat{\pi}_i = \hat{p}_i - \hbar A_i \quad \phi = \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} g_{ij}$$

and $\bar{E} = \langle \Psi | H | \Psi \rangle$ is the average total energy. We also need the energy variance

$$\Delta E^2 = \int d\mathbf{x} \psi^*(\mathbf{x}) \left\langle \left[\hat{T} + (H_{\text{el}}(\mathbf{x}) - \bar{E}) \right]^2 \right\rangle_n \psi(\mathbf{x})$$

and the result

$$\begin{aligned} \left\langle \left[\hat{T} + (H_{\text{el}}(\mathbf{x}) - \bar{E}) \right]^2 \right\rangle_n &= \langle \hat{T}^2 \rangle_n + \\ &+ \langle (E_n(\mathbf{x}) - \bar{E})^2 \rangle + 2\Re \langle \hat{T} (E_n(\mathbf{x}) - \bar{E}) \rangle \\ &= \langle \hat{T}^2 \rangle_n + \\ &+ (E_n(\mathbf{x}) - \bar{E})^2 + 2\Re \left[\langle \hat{T} \rangle_n (E_n(\mathbf{x}) - \bar{E}) \right] \end{aligned}$$

Hence, upon taking the difference of the two, we find for the local-in-time error the expression

$$\varepsilon^2[\Psi] = \frac{1}{\hbar^2} \int d\mathbf{x} \psi^*(\mathbf{x}) \left[\langle \hat{T}^2 \rangle_n - \langle \hat{T} \rangle_n^2 \right] \psi(\mathbf{x})$$

which shows explicitly the crucial role played by the nuclear kinetic energy fluctuations in the adiabatic approximation.

This expression can also be put in a form that makes explicit the contributions of electronic transitions. To this end it is worth introducing the kinetic energy operator “reduced” with respect to the electronic coordinates, $\langle \hat{T} \rangle_{nm} = \langle u_n | \hat{T} | u_m \rangle$ (the case $n = m$ reduces to the previous dressed kinetic energy operator $\langle T \rangle_n$). These operators have a hermitian symmetry $\langle T \rangle_{nm}^\dagger = \langle T \rangle_{mn}$ (as can be readily checked by either their definition or a direct calculation) and allow us to write

$$\langle \hat{T}^2 \rangle_{nn} - \langle \hat{T} \rangle_{nn}^2 = \sum_{m \neq n} \langle \hat{T} \rangle_{nm} \langle \hat{T} \rangle_{mn} \equiv \sum_{m \neq n} \langle \hat{T} \rangle_{mn}^\dagger \langle \hat{T} \rangle_{mn}$$

In turn, upon defining

$$\varphi_{m \leftarrow n}(\mathbf{x}) = \langle \hat{T} \rangle_{mn} \psi(\mathbf{x}) \equiv \langle u_m | \hat{T} | u_n \rangle_{\text{el}} \psi(\mathbf{x})$$

we have the error in terms of contributing electronic transitions,

$$\varepsilon_{\mathcal{V}}^2[\Psi] = \frac{1}{\hbar^2} \sum_{m \neq n} \int d\mathbf{x} |\varphi_{m \leftarrow n}(\mathbf{x})|^2$$

where

$$\begin{aligned} \nu_{m \leftarrow n}(\mathbf{x}) &= \frac{1}{\hbar^2} |\varphi_{m \leftarrow n}(\mathbf{x})|^2 = \\ &\frac{1}{\hbar^2} \psi^*(\mathbf{x}) \langle u_n | \hat{T} | u_m \rangle \langle u_m | \hat{T} | u_n \rangle \psi(\mathbf{x}) \end{aligned}$$

is a “transition probability density” which, in this form, is manifestly *gauge*-invariant since $\psi(\mathbf{x}) | u_n \rangle \equiv \langle \mathbf{x} | \Psi \rangle$.

In order to make a closer comparison with the error obtained in the previous section for the “static” adiabatic approximation, we introduce $\nu(\mathbf{x}) = \sum_{m \neq n} \nu_{m \leftarrow n}(\mathbf{x})$ and the total conditional transition probability density $\varrho(\mathbf{x}) = \nu(\mathbf{x}) / |\psi(\mathbf{x})|^2$ such that

$$\varepsilon^2[\Psi] = \int d\mathbf{x} |\psi(\mathbf{x})|^2 \varrho(\mathbf{x})$$

which is well defined provided $\psi(\mathbf{x}) \neq 0$. Clearly, $\varrho(\mathbf{x})$ measures the error *locally* in configuration space (as well as in time), and thus describes the tendency of the system in configuration \mathbf{x} to jump to an electronic state other than n . For a nuclear wavefunction $\psi(\mathbf{x})$ and a frame $|u_n(\mathbf{x})\rangle$ in the vector bundle $\pi : E \rightarrow \mathcal{M}$ we consider $|\psi_n(\mathbf{x})\rangle = \psi(\mathbf{x}) |u_n(\mathbf{x})\rangle$ as a smooth section of E , and the map to the normal bundle $|\psi_n(\mathbf{x})\rangle \rightarrow |\varphi(\mathbf{x})\rangle = Q \hat{T} |\psi_n(\mathbf{x})\rangle$ which gives $\nu(\mathbf{x}) = \hbar^{-2} \langle \varphi(\mathbf{x}) | \varphi(\mathbf{x}) \rangle$. We find

$$\begin{aligned} Q \hat{T} |\psi_n(\mathbf{x})\rangle &= \\ &-\frac{\hbar^2}{2} \sum_{ij} \xi^{ij} [(\partial_i \psi) Q | \partial_j u_n \rangle \\ &+ (\partial_j \psi) Q | \partial_i u_n \rangle + \psi Q | \partial_i \partial_j u_n \rangle] \end{aligned}$$

Then, upon introducing $\hat{\pi}_i = -i\hbar \partial_i - \hbar A_i$,

$$\begin{aligned} Q \hat{T} |\psi_n(\mathbf{x})\rangle &= -i\hbar \sum_{ij} \xi^{ij} (\hat{\pi}_i \psi) Q | \partial_j u_n \rangle \\ &- \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} \psi D_{ij} | u_n \rangle \\ &= -i\hbar \sum_j (\hat{v}^j \psi) Q | \partial_j u_n \rangle \\ &- \frac{\hbar^2}{2} \psi \sum_{ij} \xi^{ij} D_{ij} | u_n \rangle \end{aligned}$$

where \hat{v}^j is the j^{th} component of the velocity operator and

$$D_{ij} | u_n \rangle = i A_i Q | \partial_j u_n \rangle + i A_j Q | \partial_i u_n \rangle + Q | \partial_i \partial_j u_n \rangle$$

The reason why we introduce these two components is that they are separately *gauge*-invariant: under the *gauge* transformation $|u_n\rangle \rightarrow |u_n\rangle e^{-i\varphi}$, $\psi \rightarrow \psi e^{+i\varphi}$ we have $A_i \rightarrow A_i + \partial_i\varphi$ and

$$\begin{aligned} Q|\partial_j u_n\rangle &\rightarrow e^{-i\varphi} Q|\partial_j u_n\rangle \\ \hat{\pi}_i\psi &\rightarrow e^{i\varphi} \hat{\pi}_i\psi \\ D_{ij}|u_n\rangle &= e^{-i\varphi} D_{ij}|u_n\rangle \end{aligned}$$

since (there is no need to verify it with an explicit calculation, since both $|\psi_n(\mathbf{x})\rangle$ and the velocity term are *gauge* invariant)

$$\begin{aligned} &Q|\partial_i\partial_j u_n\rangle \rightarrow \\ &e^{-i\varphi} [Q|\partial_i\partial_j u_n\rangle - i(\partial_i\varphi)Q|\partial_j u_n\rangle - i(\partial_j\varphi)Q|\partial_i u_n\rangle] \end{aligned}$$

Stated differently, the operators $Q\partial_j$, $\hat{\pi}_i$, \hat{v}^i , D_{ij} , etc. are *tensorial* under *gauge* transformations. Hence, upon introducing the (*gauge*-tensorial) residue

$$R|u_n\rangle = \frac{\hbar}{2} \sum_{ij} \xi^{ij} D_{ij}|u_n\rangle$$

and the (complex-valued) quantum velocity fields V^j

$$V^j = \frac{\hat{v}^j\psi}{\psi} = \frac{\psi^* \hat{v}^j\psi}{|\psi|^2} \equiv V^j(\mathbf{x})$$

we find

$$\begin{aligned} \varrho(\mathbf{x}) &= \sum_{ij} (V^i)^* V^j q_{ij} \\ &- i \sum_j (V^j)^* \langle \partial_j u_n | R u_n \rangle + \\ &+ i \sum_j V^j \langle R u_n | \partial_j u_n \rangle \\ &+ \langle R u_n | R u_n \rangle \end{aligned}$$

where $q_{ij} = \langle \partial_i u_n | Q | \partial_j u_n \rangle$ is the quantum geometric tensor and the remaining scalar products contain higher derivatives of the electronic state in a *gauge* invariant form. The first term closely resembles the local-in-time error in the standard adiabatic approximation analyzed in the previous section

$$\varepsilon^2 = \sum_{ij} V^i V^j q_{ij}$$

where V^i is now the classical velocity of the i^{th} parameter. There are though notable differences: when turning the slow variables into quantum variables *both* the real (symmetric) *and* the imaginary (antisymmetric) parts of q_{ij} matter for the error, since

$$\sum_{ij} (V^i)^* V^j q_{ij} = \sum_{ij} K^{ij} g_{ij} + \frac{1}{2} \sum_{ij} Y^{ij} B_{ij}$$

where

$$K^{ij} = \Re \left((V^i)^* (V^j) \right) \quad Y^{ij} = \Im \left((V^i)^* (V^j) \right)$$

and g_{ij} and B_{ij} have been introduced above. It is instructive then to consider their total contribution upon integrating over configuration space. For the first we find

$$\begin{aligned} \int d\mathbf{x} |\psi(\mathbf{x})|^2 \sum_{ij} K^{ij} g_{ij} &= \sum_{ij} \int d\mathbf{x} \Re \left((\hat{v}^i\psi)^* g_{ij} (\hat{v}^j\psi) \right) \\ &= \Re \left(\langle \psi | \sum_{ij} \hat{v}^i g_{ij} \hat{v}^j | \psi \rangle_X \right) \\ &\equiv \langle \psi | \sum_{ij} \hat{v}^i g_{ij} \hat{v}^j | \psi \rangle_X \end{aligned}$$

where the scalar product $\langle \cdot | \cdot \rangle_X$ is that of the Hilbert space $L^2(\mathcal{M})$ describing the nuclear degrees of freedom and where the last equality follows from the fact that the operator $\sum_{ij} \hat{v}^i g_{ij} \hat{v}^j$ is self-adjoint on that space. As for the second we have similarly

$$\begin{aligned} \int d\mathbf{x} |\psi(\mathbf{x})|^2 \sum_{ij} Y^{ij} B_{ij} &= \sum_{ij} \int d\mathbf{x} \Im \left((\hat{v}^i\psi)^* B_{ij} (\hat{v}^j\psi) \right) \\ &= \Im \left(\langle \psi | \sum_{ij} \hat{v}^i B_{ij} \hat{v}^j | \psi \rangle_X \right) \\ &= -i \langle \psi | \sum_{ij} \hat{v}^i B_{ij} \hat{v}^j | \psi \rangle_X \end{aligned}$$

since the operator $\sum_{ij} \hat{v}^i B_{ij} \hat{v}^j$ is anti-hermitian

$$\left(\sum_{ij} \hat{v}^i B_{ij} \hat{v}^j \right)^\dagger = \sum_{ij} \hat{v}^j B_{ij} \hat{v}^i = - \sum_{ij} \hat{v}^i B_{ij} \hat{v}^j$$

Hence, overall, by considering the ‘‘classical’’ contribution only, we find that the LITE in the dynamic adiabatic approximation is just the expectation value of a self-adjoint quantum tensor

$$\hat{q} = \sum_{ij} \hat{v}^i \left(g_{ij} - \frac{i}{2} B_{ij} \right) \hat{v}^j \equiv \sum_{ij} \hat{v}^i q_{ij} \hat{v}^j$$

which is nothing but the quantum version of the quantum geometric tensor. That is, to leading order, we have

$$\varepsilon^2 \approx \langle \psi | \sum_{ij} \hat{v}^i q_{ij} \hat{v}^j | \psi \rangle_X$$

On comparing with the static adiabatic approximation, however, one should also observe that additional terms appear whose physical meaning is far less obvious.

Non-adiabatic transition probability

To understand better the meaning of $\varrho(\mathbf{x})$ and its components $\varrho_{m \leftarrow n}$ we consider the situation in which, during the time evolution, the local-in-time error exceeds a given threshold, thereby suggesting the need of going *beyond* the adiabatic approximation. This can be accomplished dynamically by “spawning”[29] the electronic basis that forms the variational manifold, e.g., by expanding the wavefunction *ansatz* to

$$|\Psi_t\rangle = \int d\mathbf{x} \psi_t(\mathbf{x}) |u_n(\mathbf{x}), \mathbf{x}\rangle + \int d\mathbf{x} \phi_t(\mathbf{x}) |u_s(\mathbf{x}), \mathbf{x}\rangle$$

where $s = n \pm 1$ depending on which gap $|E_n - E_{n\pm 1}|$ is the smallest. Henceforth, we shall first address the simpler situation where a single neighboring state affects the dynamics and later generalize the result to a multitude of electronic states.

At the time of spawning t_s the amplitude $\phi_t(\mathbf{x})$ *must* vanish and its time derivative is determined by the variational equations of motion

$$\begin{cases} i\hbar \frac{\partial \psi}{\partial t} = (\langle T \rangle_{nn} + E_n) \psi + \langle T \rangle_{ns} \phi \\ i\hbar \frac{\partial \phi}{\partial t} = \langle T \rangle_{sn} \psi + (\langle T \rangle_{ss} + E_s) \phi \end{cases}$$

which give, for $t = t_s$,

$$i\hbar \frac{\partial \phi}{\partial t} \Big|_{t_s} = \langle T \rangle_{sn} \psi \equiv \varphi_{s \leftarrow n}$$

(notice that the *gauge* does not affect this off-diagonal term). Thus, we see that the probability

$$\nu_{s \leftarrow n} = \hbar^{-2} \int d\mathbf{x} |\varphi_{s \leftarrow n}|^2$$

represents precisely the *error reduction due to electronic spawning*,

$$\varepsilon_V^2 \rightarrow \varepsilon_{V'}^2 = \varepsilon_V^2 - \nu_{s \leftarrow n} \quad \text{at } t = t_s$$

i.e., the error reduction arising from lifting the adiabatic approximation by allowing non-adiabatic transitions to the state s . On the other hand, the above equation also determines the short-time behaviour of the non-adiabatic transition probability P_s to the state s as

$$P_s \approx \nu_{s \leftarrow n} (t - t_s)^2 \quad t \geq t_s$$

since

$$\frac{d|\phi|^2}{dt} \Big|_{t_s} = \phi^* \frac{d\phi}{dt} + \frac{d\phi^*}{dt} \phi \Big|_{t_s} \equiv 0$$

and

$$\frac{d^2|\phi|^2}{dt^2} \Big|_{t_s} = 2 \frac{d\phi^*}{dt} \frac{d\phi}{dt} \Big|_{t_s} \equiv \frac{2}{\hbar^2} |\langle T \rangle_{sn} \psi|^2$$

[Notice that $\nu_{s \leftarrow n}$ is half the second derivative of the transition probability at $t = t_s$, a result which follows in general from the definition of local-in-time error.] This finding leads to an interesting conclusion: when a single term s dominates the sum, $\nu_{s \leftarrow n}$ is approximately the total squared error in the dynamic adiabatic approximation and we have seen above that this is determined by the quantum geometric tensor (to leading order in the derivatives of the $|u_n\rangle$'s). Hence, turning this argument around, we see that the quantum geometric tensor also determines the early transition probability upon spawning. In other words, we have approximately, up to second order in $\delta t = t - t_s$,

$$P_s \approx \int d\mathbf{x} \sum_{ij} (\delta \hat{x}^i \psi_{t_s})^* (\mathbf{x}) (\delta \hat{x}^j \psi_{t_s}) (\mathbf{x}) q_{ij}(\mathbf{x})$$

with $\delta \hat{x}^i := \hat{v}^i \delta t$

if the most important non-adiabatic channel were suddenly opened at time t_s .

More generally, all the above remains unaltered if the adiabatic approximation is suddenly lifted and the “spawning” process is made virtually complete, *i.e.*, the variational constraint is suddenly removed at $t = t_s$ and the wavefunction is allowed to expand into the whole Hilbert space

$$\begin{aligned} |\Psi_t\rangle &= \int d\mathbf{x} \psi_t(\mathbf{x}) |u_n(\mathbf{x}), \mathbf{x}\rangle \rightarrow \\ |\Psi_t\rangle &= \int d\mathbf{x} \psi_t(\mathbf{x}) |u_n(\mathbf{x}), \mathbf{x}\rangle \\ &+ \sum_{m \neq n} \int d\mathbf{x} \phi_t^{(m)}(\mathbf{x}) |u_m(\mathbf{x}), \mathbf{x}\rangle \end{aligned}$$

Again, we have $\phi^{(m)} \equiv 0$ at the time of spawning,

$$i\hbar \frac{\partial \phi^{(m)}}{\partial t} \Big|_{t_s} = \langle T \rangle_{mn} \psi \equiv \varphi_{m \leftarrow n}$$

holds for any $m \neq n$ and now the local-in-time error is reduced *exactly* to zero upon spawning. Thus, the *total* non-adiabatic transition probability P can be given, up to second order in δt , as

$$P \approx \int d\mathbf{x} \sum_{ij} (\delta \hat{x}^i \psi_{t_s})^* (\mathbf{x}) (\delta \hat{x}^j \psi_{t_s}) (\mathbf{x}) q_{ij}(\mathbf{x})$$

with $\delta \hat{x}^i := \hat{v}^i \delta t$

under the only assumption that the terms involving the second derivatives of the electronic states are negligible. This result relates the geometric properties of the adiabatic problem to the rate of non-adiabatic transitions. In a sense, this is an obvious result since the latter transitions represent precisely the failure of the adiabatic approximation. At a closer look, though, it is rather surprising that the exact dynamics of the system beyond the adiabatic paradigm is determined solely by the geometric properties of the approximation.

B. Exact factorization of the wavefunction

As mentioned in the main text, the exact factorization [31, 32] is an “intermediate” representation which is obtained by introducing a local basis of nuclear states $\{|\mathbf{x}\rangle\}$ to represent the exact wavefunction describing the combined electron-nuclear states, *i.e.*,

$$|\Psi_t\rangle = \int d\mathbf{x} |\mathbf{x}\rangle \langle \mathbf{x} | \Psi_t \rangle$$

where $\langle \mathbf{x} | \Psi_t \rangle$ is yet a vector in the electronic Hilbert space \mathcal{H}_{el} that we write as

$$\langle \mathbf{x} | \Psi_t \rangle = \psi_t(\mathbf{x}) |u_t(\mathbf{x})\rangle$$

upon imposing a normalization condition and choosing a smoothly varying phase for the *local* electronic states $|u_t(\mathbf{x})\rangle$. This gives the wavefunction in the (local) exactly-factorized representation

$$|\Psi_t\rangle = \int d\mathbf{x} \psi_t(\mathbf{x}) |\mathbf{x}\rangle |u_t(\mathbf{x})\rangle$$

Clearly, there is some freedom in choosing $|u_t\rangle$ (and correspondingly in defining the nuclear wavefunction ψ_t) that we may fix by imposing the arbitrary (but real) *gauge* term

$$A_0 = i \langle u | \partial_t u \rangle$$

in the equation of motion, besides the usual Berry’s connection terms $A_k = i \langle u | \partial_k u \rangle$.

Equations of motion

To obtain the equations of motion for the above nuclear wavefunction and the electronic state we write the total Hamiltonian using the coordinate representation for the nuclear variables, *i.e.* in the form

$$\hat{H} = \hat{T} + H_{\text{el}}(\mathbf{x})$$

where \hat{T} is the nuclear kinetic energy operator

$$\hat{T} = \frac{1}{2} \sum_{ij} \xi^{ij} \hat{p}_i \hat{p}_j \quad \text{with } \hat{p}_j = -i\hbar \partial_j$$

and $H_{\text{el}}(\mathbf{x})$ is the electronic operator with the nuclei clamped at a configuration \mathbf{x} .

From the Schrödinger equation

$$\hat{H}(\psi |u\rangle) = i\hbar (\partial_t \psi) |u\rangle + i\hbar \psi |\partial_t u\rangle$$

we immediately obtain the equation of motion for the nuclear wavefunction by projecting the above equation onto $|u\rangle$

$$\left(\langle \hat{H} \rangle_{\text{el}} - \hbar A_0 \right) \psi = i\hbar (\partial_t \psi)$$

where $\langle \hat{H} \rangle_{\text{el}} = \langle \hat{T} \rangle_{\text{el}} + \langle u | H_{\text{el}} | u \rangle$ contains the dressed kinetic energy operator and the “Born-Oppenheimer” average energy $E_{\text{el}} = \langle u | H_{\text{el}} | u \rangle$, and A_0 is the *gauge* potential introduced above. We remark that $\langle \hat{T} \rangle_{\text{el}}$ appearing here is the nuclear kinetic energy operator *averaged* over the time-dependent electronic state and reads explicitly

$$\langle \hat{T} \rangle_{\text{el}} = \frac{1}{2} \sum_{ij} \xi^{ij} \hat{\pi}_i \hat{\pi}_j + \frac{1}{2} \sum_{ij} \xi^{ij} q_{ij}$$

where $\hat{\pi}_j = \hat{p}_j - \hbar A_j$ and $q_{ij} = \langle \partial_i u | Q | \partial_j u \rangle$. We use the hat symbol to remind us the coordinate representation adopted, but notice that $|u\rangle$ is everywhere meant to be the time-dependent electronic state (correspondingly, $P = |u\rangle \langle u|$, $Q = 1 - P$, etc.).

As for the equation governing the electron dynamics we only need its projection onto the “unoccupied” electronic space, since $P \partial_t |u\rangle = |u\rangle \langle u | \partial_t u \rangle$ is known once the *gauge* term A_0 has been fixed. Hence,

$$Q \hat{H}(\psi |u\rangle) = +i\hbar \psi Q |\partial_t u\rangle$$

which gives

$$i\hbar Q |\partial_t u\rangle = \frac{1}{\psi} Q \hat{H}(\psi |u\rangle)$$

or, if we write the equation for $\partial_t |u\rangle$,

$$i\hbar |\partial_t u\rangle = +\hbar A_0 |u\rangle + \frac{1}{\psi} Q \hat{H}(\psi |u\rangle)$$

Here, the effective electronic Hamiltonian operator contains two terms

$$\frac{1}{\psi} Q \hat{H}(\psi |u\rangle) = \frac{1}{\psi} Q \hat{T}(\psi |u\rangle) + Q H_{\text{el}} |u\rangle$$

but only the first depends on ψ since H_{el} is local in nuclear coordinates. The first term, which we denote as $K[\psi] |u\rangle$, is found to be

$$K[\psi] |u\rangle = -i\hbar \sum_j V^j Q |\partial_j u\rangle - \hbar R |u\rangle$$

where $V^j = (\hat{v}^j \psi) / \psi$ is the complex-valued nuclear velocity field, $R |u\rangle = \frac{\hbar}{2} \sum_{ij} \xi^{ij} D_{ij} |u\rangle$, and

$$D_{ij} |u\rangle = iA_i Q |\partial_j u\rangle + iA_j Q |\partial_i u\rangle + Q |\partial_i \partial_j u\rangle$$

We stress that the above decomposition has simple *gauge* transformation properties, since V^j is *gauge* invariant and both $Q \partial_j$ and D_{ij} (hence R) behave tensorially under a *gauge* transformation. Hence,

$$i\hbar Q |\partial_t u\rangle = Q H_{\text{el}} + K[\psi] |u\rangle$$

where, on the r.h.s., the first term describes the electron dynamics with the nuclei clamped at \mathbf{x} and the second term describes the drag effect on the electrons due to the motion of the nuclei.

Equivalence with the formulation of Abedi et al.

The above equations for the nuclear and electronic “wavefunctions” are identical to those given in Ref. [31, 32]. This is evident for the nuclear equation but not for the electronic equation since the authors of Ref. [31, 32] wrote it in a rather different form which, in our notation, would read

$$i\hbar |\partial_t u\rangle = \left[H_{\text{el}} - (\bar{E} - \hbar A_0 + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij}) \right] |u\rangle + \left[\sum_{ij} \frac{\xi^{ij}}{2} (\hat{p}_i - \hbar A_i) (\hat{p}_j + \hbar A_j) + \sum_{ij} \xi^{ij} \left(\frac{\hat{p}_i \psi}{\psi} \right) (\hat{p}_j + \hbar A_j) \right] |u\rangle$$

Here, $(\bar{E} - \hbar A_0 + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij}) = \varepsilon$ is the effective energy introduced by the authors of Ref. [31, 32] and the second bracket, denoted F in the following, contains Hamiltonian momentum terms \hat{p}_i 's rather than $\hat{\pi}_i$'s or \hat{v}^i 's (which are *gauge* tensorial). However, it is only a matter of simple algebra to show that indeed

$$F - \varepsilon = K[\psi] + \hbar A_0 - \bar{E}$$

as required by the equation above or, equivalently,

$$F = K[\psi] + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij}$$

To see this notice that

$$\begin{aligned} F &\equiv \sum_{ij} \xi^{ij} \left(\frac{\hat{\pi}_i}{2} + \frac{\hat{\pi}_i \psi}{\psi} + \hbar A_i \right) (\hat{p}_j + \hbar A_j) \\ &= \sum_{ij} \xi^{ij} \left(\frac{\hat{p}_i + \hbar A_i}{2} + \frac{\hat{\pi}_i \psi}{\psi} \right) (\hat{p}_j + \hbar A_j) \\ &= \sum_{ij} \frac{\xi^{ij}}{2} \hat{p}_i \hat{p}_j + \sum_{ij} \frac{\xi^{ij}}{2} \hbar \hat{p}_i A_j + \sum_{ij} \frac{\xi^{ij}}{2} \hbar A_i \hat{p}_j \\ &\quad + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} A_i A_j + \sum_{ij} \frac{\hat{v}^j \psi}{\psi} (\hat{p}_j + \hbar A_j) \end{aligned}$$

where

$$(\hat{p}_j + \hbar A_j) |u\rangle = -i\hbar (\partial_j - \langle u | \partial_j u \rangle) |u\rangle \equiv -i\hbar Q \partial_j |u\rangle$$

gives

$$\sum_j \frac{\hat{v}^j \psi}{\psi} (\hat{p}_j + \hbar A_j) |u\rangle = -i\hbar \sum_j \frac{\hat{v}^j \psi}{\psi} Q \partial_j |u\rangle$$

and, on the other hand,

$$\begin{aligned} &\sum_{ij} \frac{\xi^{ij}}{2} \hbar (\hat{p}_i A_j + A_i \hat{p}_j) |u\rangle = \\ &-i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (\partial_i A_j) |u\rangle + \\ &-i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (A_i |\partial_j u\rangle + A_j |\partial_i u\rangle) = \\ &= -i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (A_i Q |\partial_j u\rangle + A_j Q |\partial_i u\rangle) + \\ &-i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (\partial_i A_j) |u\rangle - \hbar^2 \sum_{ij} \xi^{ij} A_i A_j |u\rangle \end{aligned}$$

Hence,

$$\begin{aligned} F |u\rangle &= -i\hbar \sum_j \frac{\hat{v}^j \psi}{\psi} Q \partial_j |u\rangle \\ &-i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (A_i Q |\partial_j u\rangle + A_j Q |\partial_i u\rangle) \\ &+ \sum_{ij} \frac{\xi^{ij}}{2} \hat{p}_i \hat{p}_j - i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (\partial_i A_j) |u\rangle \\ &- \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} A_i A_j |u\rangle \end{aligned}$$

Finally, upon observing that

$$\begin{aligned} A_i A_j + q_{ij} &= \langle \partial_i u | u \rangle \langle u | \partial_j u \rangle + \langle \partial_i u | Q \partial_j u \rangle \\ &\equiv \langle \partial_i u | \partial_j u \rangle \equiv \partial_i (\langle u | \partial_j u \rangle) - \langle u | \partial_i \partial_j u \rangle \end{aligned}$$

we write

$$\begin{aligned} &-\frac{\hbar^2}{2} \sum_{ij} \xi^{ij} A_i A_j |u\rangle - i\hbar^2 \sum_{ij} \frac{\xi^{ij}}{2} (\partial_i A_j) |u\rangle \\ &= \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} P |\partial_i \partial_j u\rangle + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij} |u\rangle \end{aligned}$$

and obtain

$$\begin{aligned} F |u\rangle &= -i\hbar \sum_j \frac{\hat{v}^j \psi}{\psi} Q \partial_j |u\rangle \\ &- \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} (i A_i Q |\partial_j u\rangle + i A_j Q |\partial_i u\rangle + Q |\partial_i \partial_j u\rangle) \\ &+ \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij} |u\rangle \end{aligned}$$

where

$$i A_i Q |\partial_j u\rangle + i A_j Q |\partial_i u\rangle + Q |\partial_i \partial_j u\rangle \equiv D_{ij} |u\rangle$$

i.e.,

$$F |u\rangle = -i\hbar \sum_j \frac{\hat{v}^j \psi}{\psi} Q \partial_j |u\rangle - \hbar R |u\rangle + \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} q_{ij} |u\rangle$$

as we intended to show.

*Dynamically corrected pseudo-Lorentz force:
proof of the vanishing of its average*

In the main text, we have mentioned that introducing the time derivative of the electronic state

$$\hbar Q \partial_t |u\rangle = -iQ H_{\text{el}} |u\rangle - iK[\psi_t] |u\rangle$$

in the electron dynamical force

$$F_k^{\text{ED}} = -2\Im \langle \partial_k u | Q | \hbar \partial_t u \rangle$$

one obtains a genuine non-adiabatic term

$$F_k^{\text{nad}} = 2\Re \langle \partial_k u | Q H_{\text{el}} | u \rangle$$

and a correction

$$F_k^{\text{corr}} = 2\Re \langle \partial_k u | K[\psi_t] | u \rangle$$

to the *pseudo*-Lorentz force that makes the latter vanish on average. We give here the details of the calculation, starting from the observation that, with $-iK[\psi_t] |u\rangle = -\hbar \sum_j V^j Q |\partial_j u\rangle + i\hbar R |u\rangle$, we obtain

$$\begin{aligned} F_k^{\text{corr}} &= 2\hbar \sum_j \Im (\langle \partial_k u | Q |\partial_j u\rangle V^j) - 2\hbar \Re \langle \partial_k u | R u \rangle \\ &\equiv 2\hbar \sum_j g_{kj} \Im V^j - \hbar \sum_j B_{kj} \Re V^j - 2\hbar \Re \langle \partial_k u | R u \rangle \end{aligned}$$

where $\Im q_{kj} = -B_{kj}/2$ has been used. The corrected magnetic force is easily identified

$$F_k^{\text{mag,c}} = \frac{\hbar}{2} \sum_j (\hat{v}^j B_{kj} + B_{kj} \hat{v}^j) - \hbar \sum_j B_{kj} \Re V^j$$

and found to have zero average with a simple calculation,

$$\begin{aligned} \langle \psi | F_k^{\text{mag,c}} | \psi \rangle_X &= \frac{\hbar}{2} \sum_j \langle \psi | \hat{v}^j B_{kj} + B_{kj} \hat{v}^j | \psi \rangle_X \\ &\quad - \hbar \sum_j \langle \psi | B_{kj} \Re V^j | \psi \rangle_X \end{aligned}$$

since

$$\frac{1}{2} \langle \psi | \hat{v}^j B_{kj} + B_{kj} \hat{v}^j | \psi \rangle_X = \Re \langle \psi | B_{kj} \hat{v}^j | \psi \rangle_X$$

and, on the other hand,

$$\begin{aligned} \langle \psi | B_{kj} \Re V^j | \psi \rangle_X &= \\ &= \int d\mathbf{x} \psi^*(\mathbf{x}) B_{kj}(\mathbf{x}) \frac{\Re (\psi(\mathbf{x})^* \hat{v}^j \psi(\mathbf{x}))}{|\psi(\mathbf{x})|^2} \psi(\mathbf{x}) \\ &= \int d\mathbf{x} B_{kj}(\mathbf{x}) \Re (\psi(\mathbf{x})^* \hat{v}^j \psi(\mathbf{x})) \\ &= \Re \int d\mathbf{x} B_{kj}(\mathbf{x}) \psi(\mathbf{x})^* \hat{v}^j \psi(\mathbf{x}) \\ &= \Re \langle \psi | B_{kj} \hat{v}^j | \psi \rangle_X \end{aligned}$$

Hence,

$$\langle \psi | F_k^{\text{mag,c}} | \psi \rangle_X \equiv 0$$

for any state of the nuclei. The corrected *pseudo*-electric force reads as

$$\begin{aligned} F_k^{\text{el,c}} &= 2\hbar \sum_j g_{kj} \Im V^j \\ &\quad - \hbar^2 \sum_{ij} \xi^{ij} \Re \langle \partial_k u | D_{ij} u \rangle - \frac{\hbar^2}{2} \sum_{ij} \xi^{ij} \frac{\partial g_{ij}}{\partial x^k} \end{aligned}$$

and can be re-written in a more symmetric form

$$\begin{aligned} F_k^{\text{el,c}} &= 2\hbar \sum_j g_{kj} \Im V^j \\ &\quad - \hbar^2 \sum_{ij} \xi^{ij} (\Re \langle \partial_i u | D_{kj} u \rangle + \Re \langle \partial_k u | D_{ij} u \rangle) \end{aligned}$$

This can be seen by observing that

$$\begin{aligned} \frac{\partial g_{ij}}{\partial x^k} &\equiv \frac{\partial}{\partial x^k} \Re \langle \partial_i u | Q Q \partial_j u \rangle \\ &= \Re \langle \partial_i u | Q \frac{\partial}{\partial x^k} Q \partial_j u \rangle + \Re \langle \partial_j u | Q \frac{\partial}{\partial x^k} Q \partial_i u \rangle \end{aligned}$$

where the (*gauge*-invariant) derivatives

$$\begin{aligned} Q \frac{\partial}{\partial x^k} Q |\partial_j u\rangle &= -Q (\partial_k P) |\partial_j u\rangle + Q |\partial_k \partial_j u\rangle \\ &\equiv +iA_j Q |\partial_k u\rangle + Q |\partial_k \partial_j u\rangle \\ &\equiv D_{kj} |u\rangle - iA_k Q |\partial_j u\rangle \end{aligned}$$

can be used to write

$$\begin{aligned} \frac{\partial g_{ij}}{\partial x^k} &= \Re \langle \partial_i u | D_{kj} u \rangle + A_k \Im q_{ij} + \Re \langle \partial_j u | D_{ki} u \rangle + A_k \Im q_{ji} \\ &\equiv \Re \langle \partial_i u | D_{kj} u \rangle + \Re \langle \partial_j u | D_{ki} u \rangle \end{aligned}$$

(this is of course symmetric w.r.t. exchange of i and j). Now, on taking the average

$$\begin{aligned} \langle \psi | 2\hbar \sum_j g_{kj} \Im V^j | \psi \rangle_X &= 2\hbar \sum_j \Im \int d\mathbf{x} \psi^*(\mathbf{x}) g_{kj}(\mathbf{x}) \hat{v}^j \psi(\mathbf{x}) \\ &\equiv 2\hbar \sum_j \langle \psi | \Im (g_{kj} \hat{v}^j) | \psi \rangle \end{aligned}$$

where

$$\begin{aligned} 2\hbar \Im (g_{kj} \hat{v}^j) &= -i\hbar [g_{kj}, \hat{v}^j] = \\ &= -i\hbar \sum_i \xi^{ij} [g_{kj}, \hat{\pi}_i] = -i\hbar \sum_i \xi^{ij} [g_{kj}, \hat{p}_i] \equiv \hbar^2 \sum_i \xi^{ij} \frac{\partial g_{kj}}{\partial x^i} \end{aligned}$$

hence

$$\begin{aligned} \langle \psi | F_k^{\text{el,c}} | \psi \rangle &= \hbar^2 \sum_{ij} \xi^{ij} \langle \psi | \\ &\quad \left[\frac{\partial g_{ki}}{\partial x^j} - (\Re \langle \partial_i u | D_{kj} u \rangle + \Re \langle \partial_k u | D_{ij} u \rangle) \right] | \psi \rangle \end{aligned}$$

where the operator to be averaged reads as

$$\begin{aligned} & \Re \langle \partial_k u | D_{ij} u \rangle + \Re \langle \partial_i u | D_{kj} u \rangle + \\ & - \Re \langle \partial_j u | D_{ki} u \rangle - \Re \langle \partial_k u | D_{ij} u \rangle \equiv 0 \end{aligned}$$

since $\Re \langle \partial_i u | D_{kj} u \rangle = \Re \langle \partial_j u | D_{ki} u \rangle$. In fact, importantly, we have exploited

$$\Re \langle \partial_k u | D_{ij} u \rangle = \frac{1}{2} \left(\frac{\partial g_{ik}}{\partial x^j} + \frac{\partial g_{kj}}{\partial x^i} - \frac{\partial g_{ij}}{\partial x^k} \right)$$

that shows how $\Re \langle \partial_k u | D_{ij} u \rangle$ is related to the connection ∇^q on the tangent bundle induced by the Fubini-Study metric,

$$\Re \langle \partial_k u | D_{ij} u \rangle = g_{kl} \Gamma_{ij}^l$$

where Γ_{ij}^l is the Christoffel symbol of the connection. Indeed, the zeroing of the average *pseudo*-electric force merely expresses the conservation of the metric by the corresponding Ricci-Levi Civita connection

$$\begin{aligned} \langle \psi | F_k^{\text{el},c} | \psi \rangle &= \hbar^2 \sum_{ij} \xi^{ij} \langle \psi | \frac{\partial g_{ki}}{\partial x^j} - g_{il} \Gamma_{kj}^l - g_{kl} \Gamma_{ij}^l | \psi \rangle \\ &= \hbar^2 \sum_{ij} \xi^{ij} \langle \psi | (\nabla_j^q g)_{ik} | \psi \rangle \end{aligned}$$

where $(\nabla_j^q g)_{ik}$ is the ik^{th} component of the covariant derivative of g taken with the connection ∇^q along the direction j .

Statistical properties

We emphasize here that, despite its role of a marginal probability amplitude, $\psi(\mathbf{x})$ alone cannot determine the full statistical properties of the nuclear subset of particles, not even instantaneously, *i.e.*, at a given instant of time. This is evident from the fact that the (instantaneous) statistical properties require the reduced density operator $\rho_X = \text{tr}_e \rho$ which for pure states and the factorization introduced above reads

$$\begin{aligned} \langle \mathbf{x} | \rho_X | \mathbf{x}' \rangle &= \text{tr}_e (\psi(\mathbf{x}) | u(\mathbf{x}) \rangle \langle u(\mathbf{x}') | \psi^*(\mathbf{x}') \rangle) \\ &= \sigma(\mathbf{x}, \mathbf{x}') \langle u(\mathbf{x}') | u(\mathbf{x}) \rangle \end{aligned}$$

where $\sigma(\mathbf{x}, \mathbf{x}') = \psi(\mathbf{x}) \psi^*(\mathbf{x}')$ is the “apparent” nuclear density matrix. In view of this, we have two different strategies (and interpretative tools) to investigate the statistical properties of nuclear observables. Either we use the true density matrix $\rho_X(\mathbf{x}, \mathbf{x}')$ and bare nuclear observables N

$$\langle N \rangle = \int d\mathbf{x} \int d\mathbf{x}' \rho_X(\mathbf{x}, \mathbf{x}') N(\mathbf{x}', \mathbf{x})$$

or we use the apparent density matrix $\sigma(\mathbf{x}, \mathbf{x}')$ and dressed nuclear observables \tilde{N} ,

$$\langle N \rangle = \int d\mathbf{x} \int d\mathbf{x}' \sigma(\mathbf{x}, \mathbf{x}') \tilde{N}(\mathbf{x}', \mathbf{x})$$

where

$$\tilde{N}(\mathbf{x}, \mathbf{x}') = N(\mathbf{x}, \mathbf{x}') \langle u(\mathbf{x}) | u(\mathbf{x}') \rangle$$

or, equivalently,

$$\tilde{N}(\mathbf{x}, \mathbf{x}') = \langle u(\mathbf{x}) | N(\mathbf{x}, \mathbf{x}') | u(\mathbf{x}') \rangle_{\text{el}}$$

which shows that the dressed observables are “averaged” over the electronic states.

As for the electronic density operator ρ_{el} , it takes the form of a convex combination of electronic density operators $\rho_{\text{el}}(\mathbf{x})$

$$\rho_{\text{el}} = \int_X d\mathbf{x} P(\mathbf{x}) \rho_{\text{el}}(\mathbf{x})$$

where $P(\mathbf{x}) = |\psi(\mathbf{x})|^2$ is the probability density of finding the nuclei at \mathbf{x} and $\rho_{\text{el}}(\mathbf{x})$ is the conditional density operator

$$\rho_{\text{el}}(\mathbf{x}) = \frac{\langle \mathbf{x} | \rho | \mathbf{x} \rangle}{P(\mathbf{x})} \equiv |u(\mathbf{x}) \rangle \langle u(\mathbf{x})|$$

which describes a pure local state, the one defined locally by the exact factorization representation.

For comparison, notice that the results in the adiabatic approximation are very similar to the one given here, the only difference being that $|u\rangle$ is replaced by a stationary state. Therefore, the concept that the adiabatic approximation “artificially” forces the *local* electronic state to be a pure state is misleading, because this is true for an arbitrary wavefunction.

In order to clarify the meaning of observables dressed by the electronic state let us consider in detail the nuclear momentum for the k^{th} nuclear degree of freedom, \hat{p}_k (in the coordinate representation appropriate for the exact factorization). This is first “extended” to an operator $\hat{P}_k = \hat{p}_k \otimes \mathbb{I}_{\text{el}}$ acting on the Hilbert space of the electronic-nuclear system, and then “reduced” to an operator \tilde{p}_k on the nuclear space by averaging over the electronic state

$$\tilde{p}_k = \langle u | \hat{P}_k | u \rangle = \hat{p}_k - i\hbar \langle u | \partial_k u \rangle \equiv \hat{\pi}_k$$

The result is the operator for the mechanical momentum $\hat{\pi}_k$ introduced in the main text, which can thus be considered the canonical momentum dressed by the electronic state. In general, for notational convenience, one does not distinguish \hat{P}_k from \hat{p}_k , and then care is needed in interpreting \hat{p}_k as the “microscopic” operator acting on the electronic-nuclear space or the “averaged” one acting on the nuclear space only. As for the dressed operators, they are always averaged over the electronic state, and thus meant to be operators on the Hilbert space of the nuclei.

It is instructive at this point to re-consider the total force F_k acting on the k^{th} nuclear degree of freedom in light of the above difference between “microscopic” and

“electronically averaged” quantities. On the one hand we have

$$\begin{aligned} \frac{d\langle \hat{p}_k \rangle}{dt} &= \langle \Psi | \frac{i}{\hbar} [H, \hat{p}_k] | \Psi \rangle \\ &= \langle \Psi | -\partial_k H_{\text{el}} | \Psi \rangle \\ &= \int d\mathbf{x} \psi^*(\mathbf{x}) \langle -\partial_k H_{\text{el}} \rangle_{\text{el}} \psi(\mathbf{x}) \end{aligned}$$

where, to avoid confusion, we used the subscript el on the angular bracket to denote the electronic average. This shows that the average total force is the expectation value of the dressed microscopic force $-\partial_k H_{\text{el}}$ acting on the given nuclear degree of freedom. The latter can be equivalently re-written as

$$\begin{aligned} \langle -\partial_k H_{\text{el}} \rangle_{\text{el}} &= F_k^{\text{BO}} + 2\Re \langle \partial_k u | H_{\text{el}} | u \rangle \\ &\equiv F_k^{\text{BO}} + F_k^{\text{NBO}} \end{aligned}$$

since $\Re(\langle \partial_k u | u \rangle \langle u | H_{\text{el}} | u \rangle) = 0$, in order to make evident the Born-Oppenheimer-like contribution. On the other hand, we also have

$$\frac{d\langle \hat{p}_k \rangle}{dt} = 2\Re \langle \Psi | \hat{p}_k | \partial_t \Psi \rangle$$

where the time-derivative of the total wavefunction in the exact factorized form can be written as

$$\partial_t(\psi | u) = [(\partial_t \psi) + \psi \langle u | \partial_t u \rangle] | u \rangle + \psi Q | \partial_t u \rangle$$

Here, the term between square brackets amounts to

$$[(\partial_t \psi) + \psi \langle u | \partial_t u \rangle] = -\frac{i}{\hbar} \langle H \rangle_{\text{el}} \psi$$

and thus

$$\begin{aligned} \frac{d\langle \hat{p}_k \rangle}{dt} &= 2\Re \int d\mathbf{x} \psi^*(\mathbf{x}) \left(-\frac{i}{\hbar} \right) (\hat{\pi}_k \langle H \rangle_{\text{el}}) \psi(\mathbf{x}) + \\ &+ 2\Re \int d\mathbf{x} \psi^*(\mathbf{x}) [i\hbar \langle \partial_k u | Q \partial_t u \rangle] \psi(\mathbf{x}) \end{aligned}$$

Here, for the first line we have used $\langle \hat{p}_k \rangle_{\text{el}} = \hat{\pi}_k$, whereas for the second one we have exploited

$$\begin{aligned} \langle u | \hat{p}_k Q \partial_t u \rangle &= -i\hbar \langle u | (\partial_k Q | \partial_t u \rangle) + \langle u | Q \partial_t u \rangle \hat{p}_k \\ &= -i\hbar \partial_t (\langle u | Q \partial_t u \rangle) + i\hbar \langle \partial_k u | Q \partial_t u \rangle \\ &\equiv i\hbar \langle \partial_k u | Q \partial_t u \rangle \end{aligned}$$

since $\langle u | Q \partial_t u \rangle \equiv 0$ (here the scalar products are all meant to be on the electronic space only). Furthermore, since

$$\begin{aligned} 2\Re \int d\mathbf{x} \psi^*(\mathbf{x}) \left(-\frac{i}{\hbar} \right) (\hat{\pi}_k \langle H \rangle_{\text{el}}) \psi(\mathbf{x}) &= \\ = \int d\mathbf{x} \psi^*(\mathbf{x}) \frac{i}{\hbar} [\langle H \rangle_{\text{el}}, \hat{\pi}_k] \psi(\mathbf{x}) \end{aligned}$$

and the second term is the expectation value of F_k^{ED} introduced in the main text, we finally arrive at

$$\frac{d\langle \hat{p}_k \rangle}{dt} = \int d\mathbf{x} \psi^*(\mathbf{x}) [F_k^{\text{BO}} + F_k^{\text{mag}} + F_k^{\text{el}} + F_k^{\text{ED}}] \psi(\mathbf{x})$$

On comparing with the previous expression and remembering that $F_k^{\text{mag}} + F_k^{\text{el}} + F_k^{\text{ED}} = F_k^{\text{mag},c} + F_k^{\text{el},c} + F_k^{\text{ED}}$ we find,

$$\int d\mathbf{x} \psi^*(\mathbf{x}) [F_k^{\text{mag},c} + F_k^{\text{el},c}] \psi(\mathbf{x}) = 0$$

This is consistent with the result given in the previous section, however the proof given there makes clear that the dynamically corrected *pseudo*-electric and *pseudo*-magnetic forces vanish separately when averaged.

C. Electronic friction

Linear response

Let us consider the integral form of the electronic equation in the spirit of linear response theory, set $\hbar A_0 \equiv E_{\text{el}} = \langle u | H_{\text{el}} | u \rangle$ and assume that $|u(t_0)\rangle = e^{-\frac{i}{\hbar} E_0 t_0} |u_0\rangle$ holds for some initial time t_0 in the infinite past. Let us first take the simpler non-adiabatic term represented by the following impulsive ‘kick’

$$\delta(t - \tau) \tilde{K}[\psi_\tau] |u(\tau)\rangle$$

that acts instantaneously, *i.e.*, only at time $t = \tau$. The electronic state soon after the kick reads

$$|u(\tau^+)\rangle \approx |u(\tau^-)\rangle - \frac{i}{\hbar} K_0[\psi_\tau] |u(\tau^-)\rangle$$

where $|u(\tau^-)\rangle = e^{-\frac{i}{\hbar} E_0 \tau} |u_0\rangle$ is the freely propagating state and $Q \rightarrow Q_0 = 1 - |u_0\rangle \langle u_0|$ has been used for $t = \tau - \epsilon$, $\epsilon > 0$. This follows from the integral version of the equation of motion by shrinking the time interval around the kick time τ . Note that, correspondingly, K has been replaced by K_0 to remind us the use of Q_0 rather than Q , and of the ground-state connection in the velocity operators. Hence, for any time t , we have

$$\begin{aligned} |u(t)\rangle &\approx e^{-\frac{i}{\hbar} E_0 t} |u_0\rangle \\ &- \frac{i}{\hbar} e^{-\frac{i}{\hbar} H_{\text{el}}(t-\tau)} K_0[\psi_\tau] |u_0\rangle e^{-\frac{i}{\hbar} E_0 \tau} \Theta(t - \tau) \end{aligned}$$

where $\Theta(t) = 1$ for $t > 0$ and zero otherwise. Now, when considering the full driving term

$$K[\psi_t] |u(t)\rangle = \int_{-\infty}^{+\infty} d\tau \delta(t - \tau) K[\psi_\tau] |u(\tau)\rangle$$

we have, assuming linear response,

$$\begin{aligned} |u(t)\rangle &\approx e^{-\frac{i}{\hbar} E_0 t} |u_0\rangle \\ &- \frac{i}{\hbar} \int_{-\infty}^{+\infty} e^{-\frac{i}{\hbar} H_{\text{el}}(t-\tau)} K_0[\psi_\tau] |u_0\rangle e^{-\frac{i}{\hbar} E_0 \tau} \Theta(t - \tau) d\tau \end{aligned}$$

hence

$$|\Delta u\rangle \approx -\frac{i}{\hbar} e^{-\frac{i}{\hbar} E_0 t} \int_0^\infty e^{-\frac{i}{\hbar} (H_{\text{el}} - E_0) t'} K_0[\psi_{t-t'}] |u_0\rangle dt'$$

is such that $\langle \Delta u | u_0 \rangle = 0$ since $K_0 = Q_0 K_0$.

Plugging this expression in the genuine non-Born-Oppenheimer force given above we obtain two terms,

$$F_k^{\text{NBO,I}} = -2 \sum_j \Re \int_0^\infty \Gamma_{kj}(\tau) V^j(t - \tau) d\tau$$

with the kernel

$$\Gamma_{kj}(t) = \langle \partial_k u_0 | Q_0 H'_{\text{el}} e^{-\frac{i}{\hbar} H'_{\text{el}} t} | \partial_j u_0 \rangle$$

and

$$F_k^{\text{NBO,II}} = -2 \Im \int_0^\infty \langle \partial_k u_0 | Q_0 H'_{\text{el}} e^{-\frac{i}{\hbar} H'_{\text{el}} \tau} R | u_0 \rangle d\tau$$

where we have set $H'_{\text{el}} = H_{\text{el}} - E_0$. The first is a friction-like term, with $2\Re\Gamma_{kj}(t)$ playing the role of memory kernel in the classical limit where V^j is real. In the Markov limit

$$F_k^{\text{NBO,I}} = - \sum_j \bar{\gamma}_{kj} V^j(t)$$

where

$$\bar{\gamma}_{kj} = 2 \lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon t} \Gamma_{kj}(t) dt$$

with the usual ϵ converging factor included. Later we shall find that $\bar{\gamma}_{kj}$ is better defined as the zero-frequency limit (from above) of the frequency-dependent kernel

$$\bar{\gamma}_{kj}(\omega) = 2 \lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon t} e^{i\omega t} \Gamma_{kj}(t) dt$$

where the excitation energy $\hbar\omega$ can be viewed as a ‘‘running’’ correction to E_0 in the dynamical phase factor $e^{\frac{i}{\hbar} E_0 t}$ appearing in $\Gamma_{kj}(t)$.

Equivalence with DMS friction at $T=0$ K

Let us first prove the equivalence of the DMS expression for the friction (Eq. 2 of the main text) with the one obtained above. When the electronic bath is not carrying any current the steady-state density operator is the canonical one, and in the limit $T \rightarrow 0$ we have $\rho \rightarrow |u_0\rangle \langle u_0| = P_0$ and $\partial_j \rho = |\partial_j u_0\rangle \langle u_0| + |u_0\rangle \langle \partial_j u_0|$. This gives two terms

$$\begin{aligned} \gamma_{kj}^{\text{DMS}} &= - \int_0^\infty \text{tr}_e \left((\partial_k H_{\text{el}}) e^{-\frac{i}{\hbar} H'_{\text{el}} \tau} | \partial_j u_0 \rangle \langle u_0 | \right) d\tau \\ &\quad - \int_0^\infty \text{tr}_e \left((\partial_k H_{\text{el}}) | u_0 \rangle \langle \partial_j u_0 | e^{+\frac{i}{\hbar} H'_{\text{el}} \tau} \right) d\tau \end{aligned}$$

which are the complex conjugate of each other, *i.e.*,

$$\gamma_{kj}^{\text{DMS}} = -2 \Re \int_0^\infty \langle u_0 | (\partial_k H_{\text{el}}) e^{-\frac{i}{\hbar} H'_{\text{el}} \tau} | \partial_j u_0 \rangle d\tau$$

Then, upon noticing that

$$(\partial_k H) | u_0 \rangle = (E_0 - H_{\text{el}}) | \partial_k u_0 \rangle + (\partial_k E_0) | u_0 \rangle$$

and introducing the projector $Q_0 = 1 - P_0$ we find

$$\begin{aligned} \gamma_{kj}^{\text{DMS}} &= 2 \Re \int_0^\infty \langle \partial_k u_0 | Q_0 H'_{\text{el}} e^{-\frac{i}{\hbar} H'_{\text{el}} \tau} | \partial_j u_0 \rangle d\tau \\ &\quad - 2 (\partial_k E_0) \int_0^\infty \Re \langle u_0 | \partial_j u_0 \rangle d\tau \end{aligned}$$

where the first term is precisely the real part of $\bar{\gamma}_{kj}$ introduced above, and the second term vanishes identically since $\langle u_0 | \partial_j u_0 \rangle$ is pure imaginary. Note that the usual converging factor has been here tacitly assumed.

Pseudo-magnetic contribution

Secondly, we show that the memoryless friction

$$\bar{\gamma}_{kj} = 2 \lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon t} \Gamma_{kj}(t) dt$$

contains in fact a *pseudo-magnetic* contribution. To this end we need

$$H'_{\text{el}} e^{-\epsilon t} e^{-\frac{i}{\hbar} H'_{\text{el}} t} = i\hbar \frac{d}{dt} \left(e^{-\epsilon t} e^{-\frac{i}{\hbar} H'_{\text{el}} t} \right) + i\hbar \epsilon \left(e^{-\epsilon t} e^{-\frac{i}{\hbar} H'_{\text{el}} t} \right)$$

and

$$\begin{aligned} H'_{\text{el}} \int_0^\infty e^{-\epsilon t} e^{-\frac{i}{\hbar} H'_{\text{el}} t} dt &= -i\hbar \left(1 + \frac{i\epsilon\hbar}{H'_{\text{el}} - i\epsilon\hbar} \right) \\ &= -i\hbar \left(1 + i\epsilon\hbar \frac{H'_{\text{el}} + i\epsilon\hbar}{(H'_{\text{el}})^2 + \epsilon^2\hbar^2} \right) \\ &\rightarrow -i\hbar (1 + i\pi H'_{\text{el}} \delta(H'_{\text{el}})) \end{aligned}$$

where we have used the common notation $\frac{1}{A}$ for A^{-1} . Notice that in this expression the second term on the r.h.s. $\propto H'_{\text{el}} \delta(H'_{\text{el}})$ would vanish if it were applied to a regular electronic state, but this is not the case here because of the presence of the derivative couplings.

Plugging this identity in the friction expression we find

$$\begin{aligned} \Re \bar{\gamma}_{kj} = \gamma_{kj}^{\text{DMS}} &= 2\hbar \Im \langle \partial_k u_0 | Q_0 | \partial_j u_0 \rangle \\ &\quad + 2\pi\hbar \Re \langle \partial_k u_0 | Q_0 H'_{\text{el}} \delta(H'_{\text{el}}) | \partial_j u_0 \rangle \end{aligned}$$

where the first term

$$2\hbar \Im \langle \partial_k u_0 | Q_0 | \partial_j u_0 \rangle = 2\hbar \Im q_{kj} = -\hbar B_{kj}$$

gives a magnetic component $+\hbar \sum_j B_{kl} V^j$ that, when evaluating the force, precisely cancels the magnetic correction introduced in the main text. This term does not

appear in the common case when the electronic states can be taken as real functions of the electron coordinates (as DMS assumed), which is possible in the absence of magnetic fields and for a trivial topology of the ground adiabatic state. It is however necessary when the magnetic field is turned on or if conical intersections exist that can be encircled by the evolving nuclear wavepacket. This corrective effect may thus be viewed physically as restoration of the full adiabatic dynamics in this friction limit: electronic friction cools the nuclear motion and enforces the adiabatic limit (with its *gauge* fields).

To summarize

$$\bar{\gamma}_{kj} = -2i\hbar q_{kj} + \gamma_{kj}$$

where the ‘‘corrected’’ friction kernel γ_{kj} takes the form of the real part of the expression

$$\gamma_{kj} = 2\pi\hbar \langle \partial_k u_0 | Q_0 H'_{\text{el}} \delta(H'_{\text{el}}) | \partial_j u_0 \rangle$$

or, equivalently,

$$\gamma_{kj} = 2i\hbar \lim_{\epsilon \rightarrow 0^+} \epsilon \int_0^\infty e^{-\epsilon t} \langle \partial_k u_0 | Q_0 e^{-\frac{i}{\hbar} H'_{\text{el}} t} | \partial_j u_0 \rangle$$

Running correction to E_0

The friction kernel γ_{kj} defined above is, strictly speaking, ill-defined, as it is apparent from the presence of both Q_0 and $\delta(H'_{\text{el}})$. A more appropriate definition is obtained by taking the zero-frequency limit (from above) of the frequency-dependent kernel

$$\gamma_{kj}(\omega) = 2i\hbar \lim_{\epsilon \rightarrow 0^+} \epsilon \int_0^\infty e^{-\epsilon t} e^{i\omega t} \langle \partial_k u_0 | Q_0 e^{-\frac{i}{\hbar} H'_{\text{el}} t} | \partial_j u_0 \rangle$$

The physical motivation for introducing a small (eventually vanishing in the end) positive frequency ω is that the evolving ground-electronic state has an energy slightly above E_0 , *i.e.* $E_0 + \hbar\omega$ for $\hbar\omega \rightarrow 0^+$, right because of excitations of $e - h$ pairs into the substrate. Thus the replacement

$$\exp\left(\frac{i}{\hbar} E_0 t\right) \rightarrow \exp(i\omega t) \exp\left(\frac{i}{\hbar} E_0 t\right)$$

is needed to correct the LRT result for this effect.

Before addressing this issue in detail, let us first derive some relationships needed to handle the derivative couplings, and useful to derive different equivalent expressions for the friction kernel. Let first E_0 be a discrete, non-degenerate energy eigenvalue of the electronic Hamiltonian H_{el} for some value of the nuclear coordinates $\mathbf{x} = (x^1, x^2, \dots, x^k, \dots)$. Upon taking the derivative of the electronic Schrödinger equation w.r.t. x^k , $(\partial_k H_{\text{el}}) |u_0\rangle = (\partial_k E_0) |u_0\rangle + (E_0 - H_{\text{el}}) |\partial_k u_0\rangle$, and projecting with Q_0 one easily finds

$$Q_0 |\partial_k u_0\rangle = G_0(E_0) Q_0 (\partial_k H_{\text{el}}) |u_0\rangle$$

where $G_0(\lambda) = (\lambda - H_{\text{el}}^0)^{-1}$ is the resolvent of the *restriction* of H_{el} to $Q_0 \mathcal{H}_{\text{el}}$, *i.e.* the operator $H_{\text{el}}^0 = Q_0 H_{\text{el}} = H_{\text{el}} Q_0$ defined in the subspace $Q_0 \mathcal{H}_{\text{el}}$ (here \mathcal{H}_{el} represents the Hilbert space of the electronic system). More generally, for $\lambda \in \mathbb{C}$

$$Q_0 |\partial_k u_0\rangle = (1 + (E_0 - \lambda)G(\lambda))^{-1} G(\lambda) Q_0 (\partial_k H_{\text{el}}) |u_0\rangle$$

and thus

$$Q_0 |\partial_k u_0\rangle = (1 + (\lambda - E_0)G(\lambda) + (\lambda - E_0)^2 G(\lambda)^2 + \dots) \times G(\lambda) Q_0 (\partial_k H_{\text{el}}) |u_0\rangle$$

provided λ is closer to E_0 than to any other eigenvalue (here the projector Q_0 effectively removes the pole at E_0 in the spectral representation of $G(\lambda)$). If E_0 is part of the continuous spectrum we shall use

$$Q_0 |\partial_k u_m\rangle = G^+(E_0) Q_0 (\partial_k H_{\text{el}}) |u_0\rangle$$

which amounts to defining the eigenvectors $|u_0\rangle$ through a limiting procedure. Specifically, given H_{el} , E_0 and $|u_0\rangle$ at some point \mathbf{x} , in order to fix $|u_0\rangle$ at a neighboring geometry $\mathbf{x}' = \mathbf{x} + d\mathbf{x}$ one first defines the family of vectors $|u_0^\lambda\rangle$ through the solutions of

$$[\lambda - (H_{\text{el}} + \Delta H)] |u_0^\lambda\rangle = (\lambda - H_{\text{el}}) |u_0\rangle$$

for $\Delta H = (\partial_k H_{\text{el}} - \partial_k E_0) dx^i$. This gives, for infinitesimal displacements of the nuclear coordinates,

$$Q_0 |\partial_k u_0^\lambda\rangle \approx G(\lambda) Q_0 (\partial_k H) |u_0\rangle$$

from which the above result follows upon taking the limit $\lambda \rightarrow E_0$ for $\Im\lambda > 0$.

Consider now the frequency-dependent friction kernel. In the limit $\omega \rightarrow 0$ we can use the above expression for $\lambda = E_0 + \hbar(\omega + i\epsilon)$ to write

$$\gamma_{kj}(\omega) = 2i\hbar \lim_{\epsilon \rightarrow 0^+} \epsilon \int_0^\infty dt e^{-\epsilon t} e^{i\omega t} \times \langle u_0 | (\partial_k H_{\text{el}}) Q_0 G(\lambda^*) e^{-\frac{i}{\hbar} H'_{\text{el}} t} | \partial_j u_0 \rangle$$

$$= -2\hbar^2 \lim_{\epsilon \rightarrow 0^+} \epsilon \langle u_0 | (\partial_k H_{\text{el}}) Q_0 G(\lambda^*) G(\lambda) | \partial_j u_0 \rangle$$

and thus, to leading order in ω ,

$$\gamma_{kj}(\omega) = -2\pi\hbar \langle u_0 | (\partial_k H_{\text{el}}) \delta(E_0 + \hbar\omega - H_{\text{el}}) | \partial_j u_0 \rangle$$

where $\lim_{\epsilon \rightarrow 0} \hbar\epsilon G(E - i\hbar\epsilon) G(E + i\hbar\epsilon) = \pi\delta(E - H_{\text{el}})$ has been used (here Q_0 could be removed since the δ term projects onto states with energy above E_0). This is precisely Eq.10 appearing in the main text.

Equivalently, in the same limit as above, we can also make explicit the role of $\partial_j H_{\text{el}}$

$$\gamma_{kj}(\omega) = -2\pi\hbar \langle u_0 | (\partial_k H_{\text{el}}) \delta(E_0 + \hbar\omega - H_{\text{el}}) G_p(E_0) \partial_j H_{\text{el}} | u_0 \rangle$$

where $G_p(E_0)$ is the principal part of $G^+(E_0)$, that is, $G_p(E_0) = \lim_{\eta \rightarrow 0} \Re G(E_0 + i\eta)$, and $\Im G^+(E_0) = -\pi\delta(E_0 - H_{\text{el}})$ has been neglected since it gives a vanishing contribution. [Here and in the following $\Re A = (A + A^\dagger)/2$ and $\Im A = (A - A^\dagger)/2i$]. In the limit $\omega \rightarrow 0$ in which we are interested we can replace the above expression with

$$\gamma_{kj}(\omega) = -2\pi\hbar \langle u_0 | (\partial_k H_{\text{el}}) \delta(E_0 + \hbar\omega - H_{\text{el}}) G_p(E_0 + \hbar\omega) \partial_j H_{\text{el}} | u_0 \rangle$$

and thus write the friction kernel as the zero-frequency limit of

$$\gamma_{kj}(\omega) = \frac{\partial}{\partial\omega} \langle u_0 | (\partial_k H_{\text{el}}) \delta(E_0 + \hbar\omega - H_{\text{el}}) (\partial_j H_{\text{el}}) | u_0 \rangle$$

where use has been made of

$$\begin{aligned} \delta(E - H_{\text{el}}) G_p(E) &= \frac{1}{2} \lim_{\epsilon \rightarrow 0} \epsilon [G(\lambda^*) G(\lambda)^2 + G(\lambda^*)^2 G(\lambda)] \\ &= -\frac{1}{2\hbar} \frac{\partial}{\partial\omega} \lim_{\epsilon \rightarrow 0} \epsilon G(\lambda^*) G(\lambda) \\ &= -\frac{1}{2\hbar} \frac{\partial}{\partial\omega} \delta(E_0 + \hbar\omega - H_{\text{el}}) \end{aligned}$$

for $\lambda = E + i\epsilon$, with $E = E_0 + \hbar\omega$. For comparison notice that

$$q_{kj}(\omega) = -\frac{1}{\hbar} \frac{\partial}{\partial\omega} \langle u_0 | (\partial_k H_{\text{el}}) G_p(E_0 + \hbar\omega) (\partial_j H_{\text{el}}) | u_0 \rangle$$

gives, in the limit $\omega \rightarrow 0$, the quantum geometric tensor.

Notice further that if the excitation energy $\hbar\omega$ is introduced at the level of the “bare” kernel $\bar{\gamma}_{kj}$, the above conclusions about the *pseudo*-magnetic contribution and the residual friction kernel γ_{kj} remain unaltered. In fact, one finds

$$\begin{aligned} \bar{\gamma}_{kj}(\omega) &= -2i\hbar q_{kj} + \gamma_{kj}(\omega) \\ &\quad + 2\hbar\omega \langle \partial_k u_0 | Q_0 G^+(E_0 + \hbar\omega) | \partial_j u_0 \rangle \end{aligned}$$

where the last term vanishes in the limit we are interested in.

Independent electrons

For independent electrons $\partial_k H_{\text{el}}$ is a mono-electronic operator that we write as $\partial_k h$ and the projector $\delta(E_0 + \hbar\omega - H_{\text{el}})$ can thus be restricted to singly-excited Slater

determinants, *i.e.* $|\Psi_a^b\rangle = c_b^\dagger c_a |\Phi_0\rangle$ where $|\Phi_0\rangle$ is the Hartree-Fock ground-state and c_a (c_b^\dagger) is an annihilation (creation) operator for the single-particle state $|\phi_a\rangle$ ($|\phi_b\rangle$). Here, the single-particle energies are such that $\epsilon_a < \epsilon_F < \epsilon_b$, where ϵ_F is the Fermi level, and $\hbar\omega = \epsilon_b - \epsilon_a = \Delta\epsilon_{ba}$. As a result, Eq. 10 of the main text becomes

$$\begin{aligned} \gamma_{kj}(\omega) &= -2\pi\hbar \sum_a^{\epsilon_a < \epsilon_F} \sum_b^{\epsilon_b > \epsilon_F} D_{ab}^k \langle \phi_b | \partial_j \phi_a \rangle \delta(\hbar\omega - \Delta\epsilon_{ba}) \\ &= -2\pi\hbar \sum_a^{\epsilon_a < \epsilon_F} \sum_b^{\epsilon_b > \epsilon_F} D_{ab}^k D_{ba}^j \frac{f(\epsilon_b) - f(\epsilon_a)}{\epsilon_b - \epsilon_a} \delta(\hbar\omega - \Delta\epsilon_{ba}) \end{aligned}$$

where we have defined $D_{ab}^k = \langle \phi_a | \partial_k h | \phi_b \rangle$ and introduced the electron occupation function $f(\epsilon) = \Theta(\epsilon_F - \epsilon)$. Here, we can set $\epsilon_b = \epsilon_a + \hbar\omega$ in the incremental ratio of f , and taking the limit $\omega \rightarrow 0$ replace it with $-\delta(\epsilon_a - \epsilon_F)$. Hence, upon freeing the sum over the orbitals we obtain

$$\gamma_{kj} = \pi\hbar \sum_{a,b} D_{ab}^k D_{ba}^j \delta(\epsilon_b - \epsilon_F) \delta(\epsilon_a - \epsilon_F)$$

which is the HGT expression, Eq. 1 of the main text. Notice that $\gamma_{kj}(\omega)$ above becomes real in the limit $\omega \rightarrow 0$.

Frictional vector potential

Let us show here how the above results follow, in linear response, by an appropriate modification of the Hamiltonian governing the adiabatic dynamics, in particular of the vector potential entering such Hamiltonian. This is important for introducing friction (*i.e.*, dissipation) into an effective Hamiltonian for the nuclei. As shown below, this turns the corresponding Schrödinger equation into a *non-linear* equation, but this is the price to pay if the energy transfer mechanism has to depend on the system dynamics and it is not due simply to an “external” field.

To this end, we work in a *gauge* where the electronic states evolve according to a zero-averaged Hamiltonian (the “standard” *gauge*), *i.e.*, $\langle u^+ | \partial_t u^+ \rangle = 0$ where $^+$ denotes the chosen *gauge*. Clearly, in linear-response this amounts to reference the electronic Hamiltonian to the ground-state energy, E_0 , and to write $|u^+\rangle = |u_0\rangle + |\Delta u^+\rangle$ where

$$|\Delta u^+\rangle := -\frac{i}{\hbar} \int_0^{+\infty} e^{-\frac{i}{\hbar} H_{\text{el}} t'} K_0[\psi_{t-t'}] |u_0\rangle dt'$$

since $\langle u | H | u \rangle \approx E_0 + 2\Re \langle u_0 | \Delta u \rangle \equiv E_0$ holds thanks to $\langle \Delta u^+ | u_0 \rangle = 0$. This also implies that the nuclear Hamiltonian in this *gauge*

$$H^+ \approx \frac{1}{2} \sum_{ij} \xi^{ij} \hat{\pi}_i^+ \hat{\pi}_j^+ + (E_0(\mathbf{x}) + \phi^+)$$

resembles closely the *adiabatic* Hamiltonian H^0 : the only difference is the presence of $|u\rangle = |u_0\rangle + |\Delta u^+\rangle$ in place of $|u_0\rangle$ in the vector and scalar potentials, *e.g.*,

$$\begin{aligned} A_k^0 &= i \langle u_0 | \partial_k u_0 \rangle \rightarrow \\ A_k &= A_k^0 + i \langle u_0 | \partial_k \Delta u_0 \rangle + i \langle \Delta u_k | \partial_k u_0 \rangle \\ &\equiv A_k^0 + 2\Im \langle \partial_k u_0 | \Delta u_0 \rangle \end{aligned}$$

where $\partial_k (\langle u_0 | \Delta u \rangle) = 0$ has been used in the last line. In fact, it turns out that the main modification is precisely the *time-dependent* term

$$\begin{aligned} \delta A_k &= A_k - A_k^0 \\ &= 2\Im \left(-\frac{i}{\hbar} \right) \int_0^\infty \langle \partial_k u_0 | e^{-\frac{i}{\hbar} H_{\text{el}}^0 t'} K_0[\psi_{t-t'}] | u_0 \rangle dt' \end{aligned}$$

since this is of first order in the spatial derivative of the electronic states and generates a force term of the same order through its *time-derivative*,

$$\begin{aligned} F_k &= \frac{\partial \hat{\pi}_k^+}{\partial t} + \frac{i}{\hbar} [H^+, \hat{\pi}_k^+] = -\hbar \frac{\partial (\delta A_k)}{\partial t} + \frac{i}{\hbar} [H^+, \hat{\pi}_k^+] \\ &\approx -\hbar \frac{\partial (\delta A_k)}{\partial t} + \frac{i}{\hbar} [H^0, \hat{\pi}_k^0] \end{aligned}$$

where $\hat{\pi}_k^0 = \hat{p}_k - \hbar A_k^0$ is the adiabatic momentum. Here, the last line holds if we retain only terms that contain up to three spatial derivatives of the electronic state at a time, *e.g.*, of the form

$$-\partial_k \phi_0 = -\frac{\hbar^2}{2} \sum_{ij} \xi^{ij} \partial_k \Re \langle \partial_i u_0 | Q_0 | \partial_j u_0 \rangle$$

In other words, to this “order” in the spatial derivatives, we have

$$[\hat{\pi}_i^+, \hat{\pi}_j^+] \approx [\hat{\pi}_i^0, \hat{\pi}_j^0] \quad \phi^+ \approx \phi^0$$

which can be summarized by stating that the geometric properties are the same as in the adiabatic limit, $q_{ij}^+ \approx q_{ij}^0$.

Let us then take a closer look at the correction δA_k to the vector potential. From the definition of K_0 we have

$$\begin{aligned} \delta A_k &= -2 \sum_j \Im \int_0^\infty \langle \partial_k u_0 | e^{-\frac{i}{\hbar} H_{\text{el}}^0 t'} Q_0 | \partial_j u_0 \rangle V_{t-t'}^j dt' \\ &\quad + 2\Re \int_0^\infty \langle \partial_k u_0 | e^{-\frac{i}{\hbar} H_{\text{el}}^0 t'} R | u_0 \rangle dt' \end{aligned}$$

where the second term can be neglected in the approximation above since it is time independent and it is of third order, hence contributes to the force only with a fourth order term. Upon introducing the complex-valued “position” fields

$$X^j(\mathbf{x}, t) = \int_{-\infty}^t V^j(\mathbf{x}, t') dt'$$

and integrating by parts we find

$$\begin{aligned} \delta A_k &= -2 \sum_j \Im (q_{kj}^0 X^j) \\ &\quad + \frac{2}{\hbar} \sum_j \Re \left(\int_0^\infty \langle \partial_k u_0 | \Gamma(t') | \partial_j u_0 \rangle X_{t-t'}^j dt' \right) \end{aligned}$$

where $\Gamma(t) = H_{\text{el}}^0 e^{-\frac{i}{\hbar} H_{\text{el}}^0 t} Q_0$ has been introduced. In the Markov limit we have

$$\int_0^\infty \langle \partial_k u_0 | \Gamma(t') | \partial_j u_0 \rangle X_{t-t'}^j dt \approx X_t^j \int_0^\infty \langle \partial_k u_0 | \Gamma(t') | \partial_j u_0 \rangle dt$$

where, as seen above,

$$\lim_{\epsilon \rightarrow 0^+} \int_0^\infty e^{-\epsilon t} \Gamma(t) dt = -i\hbar \left(1 + i\pi H_{\text{el}}^0 \delta(H_{\text{el}}^0) \right) Q_0$$

Hence, in this limit, we find the following simple “frictional correction” to the adiabatic dynamics

$$\delta A_k = 2\pi \sum_j \Re \left(\langle \partial_k u_0 | H_{\text{el}}^0 \delta(H_{\text{el}}^0) | \partial_j u_0 \rangle X_t^j \right)$$

It is not hard to show that these expressions for δA_k give exactly the correction to the forces discussed above. In the Markov limit, for instance, $F_k = F_k^0 + \delta F_k$ where

$$\delta F_k = - \sum_j \gamma'_{kj} \Re V_t^j + \sum_j \gamma''_{kj} \Im V_t^j$$

and γ' and γ'' are, respectively, the real and imaginary parts of the kernel

$$\gamma_{kj} = 2\pi\hbar \langle \partial_k u_0 | (H_{\text{el}}^0 \delta(H_{\text{el}}^0)) | \partial_j u_0 \rangle$$

In the Markov limit, in the simplest case where $\xi^{ij} = \delta_{ij} M^{-1}$, if γ_{kj} can be taken *diagonal* and *uniform* in the configuration space of the system where the dynamics occurs, we can write

$$\delta A_k = \Re(\gamma X_t^k) \approx \partial_k \Re \left(-i\hbar M^{-1} \gamma \int_{-\infty}^t \ln \psi_{t'}(\mathbf{x}) dt' \right)$$

if we neglect the contribution of the vector potential to the velocity field. We thus see that δA_k becomes longitudinal and can be replaced by an appropriate *scalar* field

$$\delta \phi = \hbar M^{-1} \Im(\gamma \ln \psi_t(\mathbf{x}))$$

When setting $\gamma'' \approx 0$ the resulting Hamiltonian $H = H^0 + \delta \phi$ is the Kostin Hamiltonian which is a simple way to introduce dissipation into a Schrödinger-like equation by adding a simple “phase potential” (depending on the phase of the system wavefunction in the position representation).