

“Chemoenzymatic Approach towards the Synthesis of the Antitumor and Antileishmanial Marine Metabolite (+)-Harzialactone A via the Stereoselective, Biocatalyzed Reduction of a Prochiral Ketone”

Angelica Artasensi ¹, Ivan Bassanini ², Erica Elisa Ferrandi ², Lucia Feni ¹, Giulio Vistoli ¹, Laura Fumagalli ^{1,*} and Raffaella Gandolfi ¹

¹Department of Pharmaceutical Sciences, Università degli Studi di Milano; 20133, Milan, Italy, angelica.artasensi@unimi.it; lucia.feni@unimi.it; giulio.vistoli@unimi.it; laura.fumagalli@unimi.it

²Istituto di Scienze e Tecnologie Chimiche “Giulio Natta”, Consiglio Nazionale delle Ricerche, Via Mario Bianco 9, 20131, Milan, Italy,

ivan.bassanini@scitec.cnr.it, erica.ferrandi@scitec.cnr.it

*Correspondence: laura.fumagalli@unimi.it; Tel.: +390250319303.

Abstract: As a rich source of biological active compounds, marine natural products have been increasingly screened as candidates for developing new drugs. Among the several marine products and metabolites, (+)-Harzialactone A has drawn considerable attention for its antitumor and antileishmanial activity. In this work a chemoenzymatic approach has been implemented for the preparation of the marine metabolite (+)-Harzialactone A. The synthesis involved a stereoselective, biocatalyzed reduction of the prochiral ketone 4-oxo-5-phenylpentanoic acid or the corresponding esters, all generated by chemical reactions. A collection of different promiscuous oxidoreductases (both wild-type and engineered) and diverse microorganism strains were investigated to mediate the bioconversions. After co-solvent and co-substrate investigation in order to enhance the bioreduction performance, *T. molischiana* in presence of NADES (choline hydrochloride-glucose) and ADH442 were identified as the most promising biocatalysts, allowing the obtainment of the (*S*)-enantiomer with excellent *ee* (97% to >99% respectively) and good to excellent conversion (88% to 80% respectively). The successful attempt in this study provides a new chemoenzymatic approach for the synthesis of (+)-Harzialactone A.


Keywords: Harzialactone A; marine metabolite; biocatalysis; stereoselective reduction; oxidoreductases; whole cell bioreactor; prochiral ketones.

1. Introduction

Medicinal chemists have frequently been inspired by nature during the scaffold hopping stage albeit the structural complexity of the natural products compared to the synthetic drug-like molecules is often a limit. Indeed, either natural products or their metabolites offer unexplored frameworks for the development of innovative drugs, new leads or even are already pharmacologically active substances.

Fungi play an irreplaceable role within this context since they have exceptional abilities to produce a variety of metabolites characterized by unique structures and diverse biological activities. In particular, fungi coming from the marine environment have attracted a lot of interest because are very distinctive since they grow under pressure, high salinity and low temperature [1,2]. Among these fungi, *Trichoderma* species have gained a lot of attention since they are able to produce plentiful of secondary metabolites which possess

attractive chemical structures and remarkable biological activities. More precisely *Trichoderma harzianum* probably contributed the most secondary metabolites (SMs) originating from *Trichoderma* species [3,4]. Among the SMs generated by *Trichoderma harzianum*, the metabolite (+)-Harzialactone A (Figure 1) has drawn our attention since it possesses a recognized antitumor activity and a promising growth inhibitor activity against *Leishmania amazoniensis* [5–8]. Moreover, Harzialactone A is also structurally characterized by a γ -valerolactone (GVL) core, a valuable chiral building block of which we have already expertise [9–11]. From a synthetic point of view, the stereoisomers of (+)-Harzialactone A have been prepared according to different strategies comprising both chiral pool-based approaches or chemically catalyzed stereoselective reactions [12–16]. In this context, only one example dealing with the lactonase-mediated kinetic resolution of racemic Harzialactone A can be found as a biocatalyzed entry to this bioactive molecule [17]. To the best of our knowledge, in fact, no reports dealing with the preparation of 5-phenyl- γ -valerolactones, the core of the molecular skeleton of (+)-Harzialactone A, *via* the stereoselective biocatalyzed reduction of the corresponding γ -ketoesters or acids can be found in the literature. For all these reasons, we became interested in the possibility of proposing a stereoselective, biocatalyzed synthesis to (+)-Harzialactone A *via* the enantioselective reduction of prochiral ketones.




(+)-Harzialactone

Figure 1. The metabolite (+)-Harzialactone A; depicted in red the GVL moiety.

The enzyme mediated preparation of chiral γ -lactones is a topic that has been already explored in the literature. Some reported examples are, again, based on the kinetic resolution of chemically prepared racemic lactone-precursors catalyzed by different lipases (Figure 2a) [18,19].

More elegant preparations have been achieved *via* stereoselective biocatalyzed syntheses. Specifically, the enzymatic reduction and tandem cyclization of differently decorated γ -ketoesters, regarded as “bulky-bulky” substrates has been reported using both wild-type and engineered enzymes [20–25]. Among the γ -ketoesters investigated, the enantioselective reduction of methyl 4-oxo-4-phenylbutanoate (**8**, Figure 2b), a compound structurally related to the focus of this work *i.e.* methyl 4-oxo-5-phenylpentanoate (**13**, Scheme 1), resulted in the preparation of enantiomerically enriched (*S*)-5-phenyl γ -valerolactone ((*S*)-**10**, Figure 2b) with excellent isolated yield and enantiomeric excess (*ee*) using *Ralstonia* sp. or *Sphingobium yanoikuyae* alcohol dehydrogenase (Ras-ADH and Sy-ADH, respectively) as biocatalysts. More recently, a novel chemoenzymatic entry to 5-aryl or 5-alkylaryl valerolactone derivatives has been proposed by Özgen *et al.* who reported an integrated photo- and biocatalyzed synthesis of chiral γ -lactones using simple aldehydes, acrylates, or unsaturated acid as starting materials [26]. Furthermore, the microbial bioreduction of γ -ketoesters or acids was only accomplished for the preparation of 5-alkyl or 5-phenyl γ -lactones [27].

a) Kinetic Resolution



b) Enantioselective Carbonyl Reduction





Figure 2. Biocatalytic entries to chiral GVLs.

On the basis of these considerations, a novel chemoenzymatic synthesis of the two enantiomers of **16** (Scheme 1), direct precursors of enantiomerically enriched Harzialactone A isomers [15], has been here investigated and successfully validated. Seeking for good conversions and high *ee* in the bioreduction of the “bulky-bulky” substrates **12-15** to the corresponding γ -hydroxyester, different biocatalysts have been screened. Specifically: (1) a small family of ADHs of different nature and substrate scope [28–31]; (2) a collection of promiscuous hydroxysteroid dehydrogenases (HSDHs) [32–34], (3) a commercial kit of ADHs from Evonx® and (4) different strains of *Torulopsis* sp. yeasts, selected among those species that have previously been exploited in the microbial reduction of structurally complex ketones [35].



Scheme 1. Novel biocatalytic approach to enantiomerically enriched 5-benzyltetrahydrofuran-2-one (**16**).

Reagents and conditions: (a) Na, diethyl succinate, EtOH, RT; (b) H₂O, Acetic Acid, Hydrochloric acid 37%, Reflux; (c) PTSA, MeOH, Trimethyl orthoformate, RT; (d) PTSA, EtOH, Triethyl orthoformate, RT; (e) SOCl₂, *i*-PrOH, reflux; (f) 10% aqueous solution of HCl; (g) 2-[(4-Methylphenyl)sulfonyl]-3-phenyloxaziridine, KHMDS, THF, -78 °C [15].

2. Materials and Methods

2.1 General

All chemicals and solvents were purchased from Merck KGaA, Darmstadt, Germany, and TCI and used as commercially distributed. All purifications were performed by flash chromatography using prepacked Biotage Sfär columns or silica gel (particle size 40–63 μm , Merck) on an Isolera (Biotage, Uppsala, Sweden) apparatus. Thin-layer chromatography (TLC) analyses were performed on aluminum plates precoated with silica gel 60 matrix with a fluorescent indicator and visualized in a TLC UV cabinet followed by an appropriate staining reagent. The content of solvents in eluent mixtures is given as v/v percentage. R_f values are given for guidance. ^1H NMR (300 MHz) and ^{13}C NMR (75 MHz) spectra were recorded on a Varian NMR System 300 MHz spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) relative to the residual solvent (CHCl_3 , MeOH, or dimethyl sulfoxide (DMSO)) as an internal standard. Melting points were determined by a Buchi Melting Point B-540 apparatus.

The Alcohol Dehydrogenases Screening Kit (cat. n° evo-1.1.100) was purchased from Evoxx technologies GmbH (Monheim am Rhein, Germany) and applied in the screening according to its technical bulletin. Plasmid pMS470/pEamTA was kindly donated by Prof. W. Kroutil, University of Graz, Austria.

Abbreviations used: NADES = Natural Deep Eutectic Solvent; ChCl = choline hydrochloride; Glu = glucose; Gly = glycerol; *i*-PrOH = *iso*-propanol; EtOAc = ethyl acetate. PB = phosphate buffer; PTSA = *p*-Toluenesulfonic acid; YPD Yeast Extract Peptone Dextrose.

2.2 Synthesis

2.2.1 Preparation of 4-oxo-4-phenylbutanoic acid (**7**)

To a solution of succinic anhydride (500 mg, 4.99 mmol) in benzene (2.5 mL, 28.03 mmol) aluminum chloride was added (1.645 g, 12.34 mmol). The reaction was heated to reflux under stirring for 45 minutes, checking the progress via TLC. Afterward, the mixture was cooled down to room temperature and poured into ice. Then, the residue was diluted in dichloromethane (15 mL) and extracted with 10% aqueous solution of NaHCO_3 (7 mL). The basic extract was acidified with 10% aqueous solution of HCl, and the keto acid was extracted with dichloromethane (3 x 15 mL). The organic phase was dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording the pure product **7** as a white solid (396 mg, 2.22 mmol). Yield: 44.5% Mp: 109 °C. TLC (cyclohexane/ethyl acetate 7:3 + 1% formic acid) R_f : 0.22 ^1H NMR (300 MHz, CDCl_3): δ 8.03-7.94 (m, 2H), 7.62-7.54 (m, 1H), 7.47 (t, J = 7.5 Hz, 2H), 3.32 (t, J = 6.6 Hz, 2H), 2.82 (t, J = 6.6 Hz, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 198.1, 179.1, 136.6, 133.5, 128.8, 128.2, 33.4, 28.3 NMR data matches reported literature data [36].

2.2.2 Preparation of Methyl 4-oxo-4-phenylbutanoate (**8**)

Sulfuric acid (1.1 mL, 20.6 mmol) was added to a solution of **7** (656 mg, 3.68 mmol) in methanol (5 mL) and the mixture was heated at 50 °C for 12 h. The solution was evaporated under reduced pressure, and the residue was partitioned between water (5 mL) and dichloromethane (20 mL). The organic phase was washed with 10% aqueous solution of NaHCO_3 (5 mL) and brine (5 mL), dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **8** (441 mg, 2.29 mmol) as a yellowish oil. Yield: 62.2%. TLC (cyclohexane/ethyl acetate 7:3) R_f : 0.44 ^1H NMR (300 MHz, CDCl_3): δ 8.03-7.94 (m, 2H), 7.61-7.53 (m, 1H), 7.52-7.42 (m, 2H), 3.71 (s, 3H), 3.33 (t, J = 6.6 Hz, 2H), 2.77 (t, J = 6.6 Hz, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 198.2, 173.5, 136.6, 133.4, 128.7, 128.1, 51.9, 33.5, 28.1. NMR data match reported literature data [37].

2.2.3 Preparation of (\pm)-5-phenyltetrahydrofuran-2-one (**10**)

Under nitrogen atmosphere, NaBH₄ (33 mg, 0.85 mmol) was added to a solution of **7** (126 mg, 0.704 mmol) in NaOH 5%. (1.4 mL). The reaction mixture was stirred at room temperature 18 h until TLC indicated the disappearance of the starting material. The reaction mixture was acidified with a 10% aqueous solution of HCl and extracted with dichloromethane (6 mL). The organic phase was washed with brine (2 mL) and dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording of the pure product **10** (82 mg; 0.45 mmol) as a colorless oil. Yield: 64.3%. TLC (cyclohexane/ethyl acetate 1:1 + 1% formic acid) *R*_f: 0.69 ¹H NMR (300 MHz, CDCl₃): δ 7.45-7.27 (m, 5H), 5.52 (dd, *J* = 7.9, 6.3 Hz, 1H), 2.74-2.60 (m, 3H), 2.27-2.09 (m, 1H). ¹³C NMR (75 MHz, CDCl₃) δ 176.9, 138.9, 129.2, 128.5, 125.0, 80.9, 30.8, 28.9. NMR data match reported literature data [38].

2.2.4 Preparation of Ethyl-5-phenyl-5-cianolevulinate (**11**)

Under nitrogen atmosphere, benzyl cyanide (5 mL; 42.68 mmol) and diethyl succinate (11 mL, 65.73 mmol) were added to 21 mL of a freshly prepared sodium ethoxide solution. The resulting mixture was stirred at room temperature overnight, until TLC indicated the disappearance of the starting material. The solution was then diluted with water (50 mL) and washed three times with toluene (3 x 15 mL). The basic aqueous layer was acidified with concentrated HCl and extracted three times with ethyl ether (3 x 20 mL). The organic phase was washed with 10% aqueous solution of NaHCO₃ (5 mL) and brine (5 mL), dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **11** (6.01 g, 24.54 mmol) as a yellow oil. Yield: 57.5%. TLC (cyclohexane/ethyl acetate 7:3 + 1% formic acid) *R*_f: 0.47 ¹H NMR (300 MHz, CDCl₃): δ 7.47-7.36 (m, 5H), 4.80 (s, *J* = 2.0 Hz, 1H), 4.09 (q, *J* = 7.2 Hz, 2H), 2.83 (t, *J* = 6.4 Hz, 3H), 2.61 - 2.53 (m, 2H), 1.21 (t, *J* = 7.2 Hz, 3H).

2.2.5 Preparation of 5-phenyllevulinic acid (**12**)

Compound **11** (6.01 g, 24.5 mmol) was dissolved in a mixture of water (0.85 mL), conc. HCl (22 mL), and glacial acetic acid (29 mL). The reaction was heated to reflux overnight under stirring the progress was checked via TLC. The mixture was diluted with water (30 mL) and extracted with dichloromethane (3 x 10 mL). Then the organic layer was extracted with 10% aqueous solution of NaHCO₃ (15 mL). The basic extract was acidified with aqueous HCl, and the keto acid was extracted with dichloromethane (3 x 15 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **12** (4.038 g, 21.0 mmol) as a yellow solid. Yield: 85.7%. Mp: 57 °C. TLC (cyclohexane/ethyl acetate 7:3 + 1% formic acid) *R*_f: 0.32 ¹H NMR (300 MHz, CDCl₃): δ 7.40-7.13 (m, 5H), 3.74 (s, 2H), 2.75 (t, *J* = 6.6 Hz, 2H), 2.60 (t, *J* = 6.6 Hz, 2H). ¹³C NMR (75 MHz, CDCl₃) δ 206.5, 178.1, 134.0, 129.6, 128.9, 127.3, 50.2, 36.3, 27.9 NMR data match reported literature data [39].

2.2.6 Preparation of Methyl 4-oxo-5-phenylpentanoate (**13**)

Under nitrogen atmosphere, PTSA (25 mg, 0.13 mmol) and trimethyl orthoformate (284 μ L, 2.6 mmol) were added to a solution of **12** in methanol (15 mL). The reaction mixture was stirred at room temperature 18 h until TLC indicated the disappearance of the starting material. Afterward, the solvent was evaporated *in vacuo*, and the crude was dissolved in ethyl acetate (15 mL). The organic phase was washed with 10% aqueous solution of NaHCO₃ (5 mL) and brine (5 mL), dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **13** (441 mg; 2.29 mmol) as a yellowish oil. Yield: 84.7%. TLC (cyclohexane/ethyl acetate 7:3) *R*_f: 0.43 ¹H NMR (300 MHz, CDCl₃): δ 7.40-7.16 (m, 5H), 3.74 (s, 2H), 3.65 (s,

3H), 2.76 (t, $J = 6.6$ Hz, 2H), 2.56 (t, $J = 6.6$ Hz, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 206.40, 173.09, 134.06, 129.45, 128.70, 127.04, 51.72, 49.92, 36.42, 27.77. NMR data match reported literature data [40].

2.2.7 Preparation of Ethyl 4-oxo-5-phenylpentanoate (**14**)

Under a nitrogen atmosphere, PTSA (25 mg, 0.13 mmol) and triethyl orthoformate (540 μL , 3.25 mmol) were added to a solution of **12** in ethanol (15 mL). The reaction mixture was stirred at room temperature 18 h until TLC indicated the disappearance of the starting material. Afterward, the solvent was evaporated *in vacuo*, and the crude was dissolved in ethyl acetate (15 mL). The organic phase was washed with 10% aqueous solution of NaHCO_3 (5 mL) and of brine (5 mL), dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **14** (490 mg, 2.22 mmol) as a yellow oil. Yield: 84.7%. TLC (cyclohexane/ethyl acetate 7:3) R_f : 0.49 ^1H NMR (300 MHz, CDCl_3): δ 7.41–7.12 (m, 5H), 4.11 (q, $J = 7.1$ Hz, 2H), 3.74 (s, 2H), 2.75 (t, $J = 6.6$ Hz, 2H), 2.55 (t, $J = 6.6$ Hz, 2H), 1.23 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (75 MHz, CDCl_3) δ 206.3, 172.5, 134.2, 129.4, 128.6, 126.9, 53.2, 49.8, 36.4, 28.0, 14.1.

2.2.8 Preparation of Isopropyl 4-oxo-5-phenylpentanoate (**15**)

Under a nitrogen atmosphere, SOCl_2 (283 μL , 3.9 mmol) was added to a solution of **12** (500 mg, 2.6 mmol) in *iso*-propanol (15 mL). The reaction mixture was heated to reflux for 18 h until TLC indicated the disappearance of the starting material. Afterward, the solvent was evaporated *in vacuo*, and the crude was dissolved in ethyl acetate (15 mL). The organic phase was washed with 10% aqueous solution of NaHCO_3 (5 mL) and brine (5 mL), dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording compound **15** (536 mg, 2.28 mmol) as a yellow oil. Yield: 87.9%. TLC (cyclohexane/ethyl acetate 7:3) R_f : 0.56 ^1H NMR (300 MHz, CDCl_3): δ 7.46–6.95 (m, 5H), 5.04–4.89 (m, 1H), 3.74 (s, 2H), 2.74 (t, $J = 6.6$ Hz, 2H), 2.52 (t, $J = 6.6$ Hz, 2H), 1.21 (d, $J = 0.8$ Hz, 3H), 1.19 (d, $J = 0.8$ Hz, 3H). ^{13}C NMR (75 MHz, CDCl_3) δ 206.3, 172.0, 134.1, 129.4, 128.6, 127.0, 67.8, 60.1, 49.9, 36.5, 28.3, 21.7.

2.2.9 Preparation of (\pm)-5-benzyltetrahydrofuran-2-one (**16**)

Under nitrogen atmosphere, NaBH_4 (17 mg, 0.45 mmol) was added to a solution of **13** (76.7 mg, 0.37 mmol) in 5% aqueous solution of NaOH (0.8 mL). The reaction mixture was stirred at room temperature 18 h until TLC indicated the disappearance of the starting material. The reaction mixture was acidified with a 10% aqueous solution of HCl and extracted with dichloromethane (3 X 2 mL). The organic phase was washed with brine (2 mL) and dried over anhydrous sodium sulfate, filtered, and evaporated *in vacuo*, affording the pure product **16** (65 mg, 0.33 mmol) as a colorless oil. Yield: 91.5%. TLC (cyclohexane/ethyl acetate 7:3 + 1% formic acid) R_f : 0.50 ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 7.40–7.13 (m, 5H), 4.78–4.61 (m, 1H), 3.02–2.79 (m, 2H), 2.56–2.28 (m, 3H), 2.25–2.08 (m, 1H), 1.97–1.75 (m, 1H). ^1H NMR (300 MHz, CDCl_3) δ 7.34–7.22 (m, 5H), 4.76–4.70 (m, 1H), 3.09–2.90 (m, 2H), 2.51–2.33 (m, 2H), 2.29–2.21 (m, 1H), 2.00–1.90 (m, 1H). ^{13}C NMR (75 MHz, CDCl_3) δ 176.9, 135.9, 129.4, 128.6, 126.9, 80.7, 41.3, 28.6, 27.1. CDCl_3 NMR data match reported literature data [16].

2.3 Enzymatic and microbial bioreduction

2.3.1 Enzymes preparation.

Recombinant expression in *E. coli* and purification by affinity chromatography of the following enzymes was carried out as previously described: MI-ADH [30], Lb-ADH [30] HSDHs [33], Is2-SDR [33], BmGDH [41]

RasADH, ADH-A and SyADH were expressed in *E. coli* BL21 DE3 according to [30,42]. After cell pellet recovery by centrifugation, cells were resuspended in water and lyophilized.

2.3.2 Culture conditions.

The yeasts were routinely maintained on YPD medium slants (18 g/L agar, 10 g/L yeast extract, 10 g/L peptone, 20 g/L glucose, pH = 5.6). The strain, grown on YPD medium slants for 72 h at 28 °C, was inoculated in a 500 mL Erlenmeyer flask containing 100 mL of the same liquid medium (without agar) and incubated on a reciprocal shaker (150 rpm) for 48 h at 28 °C. In the case of solid-solid-liquid growth, the microorganism was grown in solid medium for 24 h and later it was inoculated in 100 mL liquid medium in a 500 mL Erlenmeyer flask. The microorganism was incubated on a reciprocal shaker (150 rpm) for 48 h at 28 °C before testing. Cell growth was evaluated by cells dry weight. 20 mL of broth culture were centrifuged (4500 rpm for 15 min), the pellet was washed twice with milliQ water and re-suspended in milliQ water. The cell suspension was placed in an oven for 12 h at 104 °C.

2.3.3 Enzymatic screening for **8** and **13** bioreduction.

Reactions were run in analytical scales working with a final volume of 0.5 mL. Substrates **8** and **13** were dissolved in PB (pH 7, 50 mM) at a final concentration of 10 mM starting from stock solutions prepared in DMSO (200 mM) reaching a co-solvent percentage of the 5% (v/v). The selected oxidoreductase was then added to the substrate solution based on its activity reaching a final concentration of 1 U/mL for Is2-SDR and all the HSDHs and of 0.5 U/mL in the case of MI-ADH. Lb-ADH was used with a loading of 2 mg/mL. *E. coli*/SyADH, *E. coli*/ADH-A and *E. coli*/RasADH were instead added with concentrations of 15 mg of lyophilized cells per mL. Glucose (40 mM), glucose dehydrogenase BmGDH (0.5 U/mL) and NAD(P)⁺ (0.4 mM) were added to the reaction mixtures which were then incubated for 24-48 h at 25°C and 180 rpm. Stock solutions of cofactor (prepared in MilliQ-water at concentration of 20 mM) and glucose (prepared in PB buffer -pH 7, 50 mM- at a concentration of 100 mM) were used while BmGDH was added accordingly to its activity. In the case of *E. coli*/RasADH, following literature protocols [42], a final concentration of 1 mM of NAPD⁺ and an operational temperature of 30 °C were used. Reactions were followed by TLC analysis *via* micro-extractions (*ca* 20 µL of reaction extracted with 40 µL of EtOAc). After 24h, reaction mixtures were acidified *ca* to pH 2 with 2N HCl_(aq), extracted two times with 500 µL of EtOAc. The organic phases were combined and dried over Na₂SO₄ and were concentrated *in vacuo* to be analyzed by means of ¹H NMR (400 MHz, CDCl₃) and HPLC (Method B).

2.3.4 Microbial biotransformations: general procedure.

Cells obtained by centrifugation (4500 rpm for 15 min) of the culture broth were washed twice with tap water. The pellet was suspended in Tris/HCl buffer containing 50 g/L of glucose (pH 7, 0.1 M) to obtain a doubled cells concentration. Substrates (**8**, **12** or **13**) were dissolved in DMSO or *iso*-propanol added to the biotransformations in a final concentration of 0.5, 1 or 2 g/L; keeping the co-solvent v/v ratio equal to the 5% (DMSO) and the 10% (*iso*-propanol). The reactions were carried out at 28°C under magnetic stirring. After 96h, the biotransformations were acidified with an HCl solution ([HCl]_{biotransformation} = 2 M), left under magnetic stirring for 24 h and then extracted twice with ethyl acetate. The combined organic phases were dried over anhydrous Na₂SO₄ and were evaporated *in vacuo*. The samples were analyzed by chiral HPLC analyses according to Method A (see Supplementary Materials for details).

2.3.5 NADES preparation.

NADES were prepared by mixing choline hydrochloride (ChCl) and glucose (Glu) in a molar ratio of 1.5:1 or glycerol (Gly) and glucose (Glu) in a molar ratio of 2:1. The compounds were placed under magnetic stirring at a temperature of 75 °C until a colorless liquid mixture was obtained.

2.3.6 HPLC analyses.

Molar conversion and *ee* were determined by HPLC equipped with a chiral column according to one of the two different methods reported below.

Method A: Merck-Hitachi L-7100 instrument coupled with a UV6000LP detector and a Chiralcel OD-H chiral column (250 mm×4.6 mm). HPLC conditions: flow rate = 0.8 mL/min, detection λ = 210 nm; temperature = 30 °C, mobile phase: 90% of hexane, 10% of *i*-PrOH and 0.1% of formic acid. Compound, R_t : **8**, 10.9 min; **7**, 12.6 min; (S)- **10**, 19 min; (R)-**10**, 18 min; **13**, 12.45 min; **12**, 13.1 min; (S)-**16**, 18 min; (R)-**16**, 21 min; **14**, 9.9 min; **15**, 8 min.

Method B: Shimadzu LC-20AD high performance liquid chromatography system equipped with a Shimadzu SPD-20 A UV detector and a Phenomenex Lux 3u Cellulose-1 chiral column (250 mm×4.6 mm). HPLC conditions: flow rate = 1 mL/min; detection λ = 280 nm; temperature = 30 °C, mobile phase: 70% of petroleum ether and 30% of *i*-PrOH. Compounds: R_t = **7**, 5.88 min; (S)-**10**, 6.9 min; (R)-**10**, 7.3 min; **13**, 6.27 min; (S)-**16**, 7.4 min; (R)-**16**, 8.3 min.

2.4 Semi-preparative scale preparation of (R)- and (S)-**16**

2.4.1 Preparation of (+)-5-benzyltetrahydrofuran-2-one (S)-**16**

The liquid culture inoculum of *T. molischiana* was prepared by making two passages in solid YPD medium by incubating the plates in both cases at 28 °C for 24h. Cells from 50 mL of broth culture were centrifuged, washed and suspended in Tris/HCl buffer (25 mL, pH 7, 0.1 M) containing 10% v/v ChCl-Glc NADES and 25 mg of compound **12**. After 96 h at 30 °C under magnetic stirring, the biotransformation was acidified with HCl and extracted twice with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and were concentrated *in vacuo* affording crude (S)-**16** which was purified by preparative TLC (Eluent: cyclohexane : ethyl acetate = 7:3). (S)-**16** was obtained as yellowish oil with a 65% isolated yield and an *ee* > 97%. $[\alpha]_D^{25} = +11.37$ (c 1.0, CHCl₃) [43], HRMS (ESI): *m/z* calculated for [M + Na]⁺ C₁₁H₁₂O₂ 199.0735; found 199.0736.

2.4.2 Preparation of (-)-5-benzyltetrahydrofuran-2-one (R)-**16**

Glucose (40 mM), BmGDH (0.5 U/mL) and NADP⁺ (0.4 mM) were added to a solution of **13** (30 mg, 10 mM) prepared in 10% v/v mixture of DMSO in PB buffer (pH 7.0, 50 mM). Reduction was started by adding *E. coli*/SyADH (10 mg/mL of lyophilized *E. coli* cells) and it was controlled by TLC analysis (ethyl acetate/petroleum ether = 7:3 as mobile phase, UV light and phosphomolybdic acid solution to stain) over the course of 48 h. After that, the mixture was extracted with EtOAc (3 X 500 μ L), the organic layers were combined, dried over Na₂SO₄, and were concentrated *in vacuo* affording crude (R)-**16** which was purified by preparative TLC (Eluent: cyclohexane : ethyl acetate = 7:3). (R)-**16** was obtained as yellowish oil with a 70% isolated yield and an *ee* > 99%. (R)-**16** $[\alpha]_D^{25} = -10.98$ (c 1.0, CHCl₃) [44], HRMS (ESI): *m/z* calculated for [M + Na]⁺ C₁₁H₁₂O₂ 199.0735; found 199.0735.

2.5 Computational Methods

Docking simulations were performed by using the resolved structure of the RasADH in complex with NADP⁺ (PDB Id 4I5D) [45] which was prepared by adding the hydrogen atoms according to the physiological pH

equal to 7.4 by using the VEGA suite of programs [46]. The precise arrangement of the substrates was extrapolated by the resolved structure of F12 RasADH in complex with NADPH and A6O (PDB Id: 6IHH) [47]. In detail, the A6O was manually inserted into the catalytic cavity of the first structure by superimposing the common cofactor and the docking searches were focused within a 10 Å radius sphere around the so inserted A6O ligand. Docking simulations were carried out by PLANTS [48]; 10 poses per ligand were generated and ranked by the ChemPLP scoring function with speed equal to 1. The generated complexes were minimized by using Namd and rescored by Rescore⁺ [49].

3. Results & Discussion

3.1 Chemistry

3.1.2 Synthesis of the suitable substrates


The substrate **8** was synthesized starting from benzene that underwent a Friedel-Crafts acylation and a subsequent Fisher esterification as reported by Alazet *et al.* whose procedure was properly adapted [39]. The series of suitable prochiral ketones **12-15** were synthesized as shown in Scheme 1. The synthesis started with the commercially available benzyl cyanide that reacted with diethyl succinate to give compound **11** which was subsequently treated with a mixture of acetic acid/hydrochloric acid to afford the derivative **12**. Then, compound **12** underwent esterification with methanol, ethanol, and *iso*-propanol to obtain compounds **13**, **14**, and **15** respectively. Furthermore, compounds *rac*-**10** and *rac*-**16** were synthesized as references for HPLC analyses.

Copies of the NMR spectra and HPLC chromatograms can be found in the Supplementary Materials.

3.2 Bioreduction of **13**: biocatalysts screening

3.2.1 Wild-type, recombinant oxidoreductases

The biocatalyzed stereoselective reduction of **13** and the one-pot tandem conversion of the obtained γ -hydroxyester into the corresponding γ -valerolactone **16** were investigated using a collection NAD⁺ or NADP⁺ dependent oxidoreductases of different origin, both belonging to the alcohol dehydrogenase (ADHs) and hydroxysteroid dehydrogenases (HSDHs) groups (Table 1). The glucose/glucose dehydrogenase from *Bacillus megaterium* (BmGDH) system was used for cofactor regeneration (Scheme 2), and enzymes were produced as recombinant proteins in *E. coli* according to established literature protocols (Table 1). Compound **8**, whose bioreduction was already reported in the literature, was used as model substrate to compare the performances of the selected biocatalysts' on both the bulky-bulky γ -ketoester substrates. At variance to **8**, compound **13** is in fact characterized by a more flexible benzyl substituent as C α decoration.



Scheme 2. Enzymatic reduction of compounds **8** and **13** and their tandem conversion into the corresponding GVLs (**10** and **16**).

Table 1 summarizes the source and redox cofactor of the screened biocatalysts, all of them characterized by different synthetic features. Specifically, Is2-SDR, a short-chain dehydrogenase discovered by the authors


from metagenomes sampling, the alcohol dehydrogenase from *Micrococcus luteus* (MI-ADH) are enzymes known for their broad substrate promiscuity and ability to reduce bulky or structurally complex substrates [32,50]. The above mentioned *E. coli*/RasADH and *E. coli*/SyADH showed good activity (60-70%) and excellent enantioselectivity in the reduction of **8**, while the alcohol dehydrogenase from *Rhodococcus ruber* (ADH-A) and *Lactobacillus brevis* (Lb-ADH) did not converted this model substrate [21,26,30]. Finally, a small collection of the different hydroxysteroid dehydrogenases (HSDHs) available to SCITEC-CNR laboratories was selected based on their high regio- and stereoselectivity towards steroid skeletons (*i.e.* 7 α , 7 β and 12 α HSDHs) and substrate promiscuity [32,34].

Table 1. Recombinant NAD(P)⁺-dependent oxidoreductases screened for the stereoselective reduction of **8** and **13**.

Enzyme	Source, Reference	Cofactor
Alcohol Dehydrogenases		
Is2-SDR	<i>Metagenome sample</i> , [32]	NADP ⁺
MI-ADH	<i>Micrococcus luteus</i> , [50]	NAD ⁺
<i>E. coli</i>/ADH-A¹	<i>Rhodococcus ruber</i> , [51]	NAP ⁺
Lb-ADH	<i>Lactobacillus brevis</i> , [52]	NADP ⁺
<i>E. coli</i>/RasADH¹	<i>Ralstonia sp.</i> , [42]	NADP ⁺
<i>E. coli</i>/SyADH¹	<i>Sphingobium yanoikuyae</i> , [29,42]	NADP ⁺
Hydroxysteroids Dehydrogenases		
Ca7α-HSDH	<i>Clostridium absonum</i> , [33]	NADP ⁺
Dm7α-HSDH	<i>Deinococcus marmoris</i> , [32]	NAD ⁺
Hh7α-HSDH	<i>Halomonas halodenitrificans</i> , [32]	NAD ⁺
Ngi1_7α-HSDH	<i>Metagenome sample</i> , [32]	NAD ⁺
Bsp7β-HSDH	<i>Brucella sp.</i> , [32]	NAD ⁺
Ca7β-HSDH	<i>Clostridium absonum</i> , [53]	NADP ⁺
Rs7β-HSDH	<i>Rhodobacter sphaeroides</i> , [32]	NAD ⁺
Sc7β-HSDH	<i>Stanieria cyanosphaera</i> , [32]	NAD ⁺
Csp12α-HSDH	<i>Clostridium sp.</i> , [54]	NADP ⁺

¹ Lyophilized recombinant *E. coli* cell pellets were used in the biotransformations [29,42].

All the described enzymes were thus used in analytical scale as biocatalysts for the stereoselective reduction of compounds **8** and **13** conducting their one-pot, tandem transformation into **10** and **16** respectively without isolating the corresponding γ -hydroxyester intermediate. Reactions were run for 24 h (for details of the screening protocol see the *Material and Methods* section); after extraction with ethyl acetate, conversion and *ee* for compounds **10** and **16** were determined *via* HPLC using chiral stationary phase methods already reported in the literature [16,20]. The successful biotransformations obtained during the screening process are summarized in Table 2.

Table 2. Screening results for the bioreduction of compounds **8** and **13**.

		Substrate			
		8		13	
		Product			
Entry	Oxidoreductases	10 ¹		16 ²	
		<i>c</i> (%) ³	<i>ee</i> (%) ³	<i>c</i> (%) ³	<i>ee</i> (%) ³
1	Is2-SDR	31	78 (<i>S</i>)	N.C.	
2	MI-ADH	38	> 99 (<i>R</i>)	N.C.	
3	<i>E. coli</i>/RasADH	60	> 99 (<i>S</i>)	50	66 (<i>R</i>)
4	<i>E. coli</i>/SyADH	70	> 99 (<i>S</i>)	45	> 99 (<i>R</i>)

¹ Method: Lux Cellulose-1, λ 210 nm and 254 nm, mobile phase petroleum ether/*iso*-propanol = 7:3, flow = 1 mL/min. R_t = (*S*)-**10**, 6.9 min; (*R*)-**10**, 7.3 min. 369

² Method: Lux Cellulose-1, λ 210 nm and 254 nm, mobile phase petroleum ether/*iso*-propanol = 7:3, flow = 1 mL/min. R_t = (*S*)-**16**, 7.4min; (*R*)-**16**, 8.3 min. 371

³ Conversion (*c* (%)) and enantiomeric excesses (*ee* (%)) were determined by HPLC analysis on chiral column (see Materials and Methods section). 373

N.C. = no conversion. 375

To our surprise, none of the promiscuous HSDHs, which successfully catalyzed the reduction of β -ketoesters [32,34], were able to convert compounds **8** or **13** to **10** or **16** respectively. 377

Interestingly, both Is2-SDR and MI-ADH reduced **8** with modest conversion (< 40%) and complementary enantioselectivity. The former produced (*S*)-**10** (the Prelog product) with 78% *ee* while the latter catalyzed the formation of the (*R*)-**10** (the *anti*-Prelog product) with an excellent *ee* (> 99%). However, both these biocatalysts did not convert the bulkier substrate **13**. 379

In agreement with previous works [21,26,30], neither Lb-ADH nor ADH-A converted **8**, while both *E. coli*/RasADH and *E. coli*/SyADH reduced **8** with excellent *ee* (> 99%) in favor of the Prelog product (*S*)-**10**. In agreement with what was reported for **8**, compound **13** was not converted by neither Lb-ADH nor ADH-A while it was reduced by *E. coli*/RasADH and *E. coli*/SyADH, though with lower conversions. Moreover, both *E. coli*/RasADH and *E. coli*/SyADH, at variance to the reduction of **8**, formed the *anti*-Prelog product (*R*)-**16** from **13** with a modest to an excellent *ee* (66% and > 99%, respectively). 383

Surprised by the unexpected *anti*-Prelog stereoselectivity of the bioreduction of **13** an *in silico* investigation was performed to compare the docking poses of the two prochiral γ -ketoesters in RasADH active site (PDB Id: 4I5D) [45]. 389

The docking results revealed that both the substrates, namely compounds **8** and **13** can be suitably accommodated within the RasADH catalytic pocket where they assume two specular binding modes thus providing different enantiotopic faces as shown in Figure 3. Both esters arrange the keto group in a pose 392

conductive to the catalysis since the oxygen atom interacts with Ser137, His147 and Tyr150, while the cofactor's nicotinamide ring points to the substrate's carbonyl carbon atom. The differences between the two simulated substrates involve the relative arrangement of the phenyl ring and the ester moiety. Indeed, Figure 3A reveals that the small substrate (**8**) is able to arrange the phenyl ring towards the cofactor with which it can stabilize p-p interactions further reinforced by similar stacking contacts with Phe205 plus hydrophobic interactions with surrounding alkyl-chain residues (e.g. Ile91, Ala162, Ile187 and Ile188). The ester group of compound **8** is facing the exit of the binding pocket where it can only stabilize hydrophobic interactions with Val138, Leu144 and Leu246. In contrast, Figure 3B shows that compound **13** prefers to assume an overturned pose in which the ester group approaches the cofactor, and the phenyl assumes a more superficial arrangement. In detail, the ester moiety is engaged by a H-bond with Gln191 plus hydrophobic contacts with Ile91 and Ile187, while the phenyl ring elicits a clear p-p stacking with Phe205 plus hydrophobic contacts with Val138, Leu144 and Leu246.





Figure 3. Main interactions stabilizing the putative complexes for compounds **8** (A) and **13** (B) within the catalytic pocket of RasADH (PDB Id: 4I5D) [45]. (C) Schematic representation of Prelog and *anti*-Prelog reduction.

3.2.2. The Evoxx® Alcohol Dehydrogenase Kit

As discussed, compound **13** was reduced by *E. coli*/RasADH even though with low conversion and *ee*, and by *E. coli*/SyADH, with an excellent *ee* and a lower conversion when compared to the reduction of compound **8**. Moreover, both *E. coli*/RasADH and *E. coli*/SyADH produced the (*R*)-enantiomer of **16**, which presents the opposite absolute stereochemistry of (+)-Harzialactone A (Figure 1). Thus, seeking for an enantiocomplementary enzyme, a kit of commercially available engineered ADH (EvoXX® Technologies GmbH) whose good performances on bulky and structurally complex natural compounds have been already reported by us [30] was screened for the reduction of substrates **8** and **13**. Among the screened enzymes ADH442 was found to efficiently catalyzed the formation of (*R*)-**16** with excellent conversion and *ee*. Obtained data are reported in detail in the Supplementary Materials (Table S1).

3.2.3. Microbial bioreduction of **13**

After screening isolated enzymes, looking for a low-cost biocatalyst to produce (*S*)-**16**, the core of (+)-Harzialactone A, we focused our attention on the microbial reduction of **13** (Scheme 1) because of no reports can be found in the literature dealing with the biotransformation of this substrate using whole cells as biocatalyst. Twenty different yeast strains selected from our in-house collection and belonging to different species and *genera* were exploited in an analytical scale screening. As planned for isolated enzymes, compound **13** was used as a model substrate (Figure 2b) since, as demonstrated by Forzato *et al.*, its acid and the ethyl ester derivatives were stereoselectively reduced to the corresponding (*S*)-butyrolactones either by *P. etchelsii*, *P. glucozyma*, or *S. cerevisiae* [27].

Screening results (Table 3) are reported as conversion and *ee* of the corresponding lactones determined *via* HPLC (See Materials and Methods section and Supplementary Materials). At variance to the use of isolated enzymes both the substrates (**8** and **13**) underwent competitive hydrolytic processes in the presence of whole cells, thus they were also converted to the corresponding γ -ketoacids (**7** and **12**).

As expected, compound **8** was successfully reduced by *P. etchelsii*, *P. glucozyma*, and *S. cerevisiae* with good conversion and moderate to excellent *ee*; among them *S. cerevisiae* and *L. fabianii* were the best ones in term of enantioselectivity (*ee* \geq 95%). In general, the yeast strains that were found active reduced compound **8** only to (*S*)-**10** with different degrees of enantioselectivity.

Table 3. Microorganisms screened for the bioreduction of compounds **8** and **13**.

Entry	Microorganism	Substrate					
		8			13		
		7	Product		12	Product	
<i>c</i> (%) ¹	<i>c</i> (%) ¹	<i>ee</i> (%) ¹	<i>c</i> (%) ¹	<i>c</i> (%) ¹	<i>ee</i> (%) ¹		
1	<i>Kluyveromyces marxianus</i> CBS 397	71	17	72 (S)	59	41	85 (S)
2	<i>K. marxianus</i> var. <i>lactis</i> CL69	53	41	75 (S)	78	10	37 (S)
3	<i>Lindnera fabianii</i> CBS 5640	5	95	95 (S)	65	35	66 (S)
4	<i>Pachysolen tannophilus</i> CBS 4044	5	75	22 (S)	47	45	33 (R)
5	<i>Pichia capsulata</i> CBS 1993	70	13	60 (S)	73	traces	
6	<i>Pichia etchellsii</i> MIM	24	51	47 (S)	48	traces	
7	<i>Pichia glucozyma</i> CBS 5766	34	30	45 (S)	13	traces	
8	<i>Pichia jadinii</i> CBS 4885	57	25	65 (S)	89	16	64 (R)
9	<i>Pichia pastoris</i> CBS 704	71	traces		80	traces	
10	<i>Rhodotorula rubra</i> MIM 146	79	traces		39	59	87 (R)
11	<i>Rhodotorula rubra</i> MIM 147	63	traces		35	12	55 (R)
12	<i>Saccharomyces cerevisiae</i> ZEUS	47	41	98 (S)	73	traces	
13	<i>Sporidiobolus pararoseus</i> SD2	96	traces		92	11	77 (R)
14	<i>Sporobolomyces holsaticus</i> NCYC 420	76	traces		75	traces	
15	<i>Sporobolomyces salmonicolor</i> MIM	0	0		46	traces	
16	<i>Slooffia tsugae</i> CBS 5038	51	38	98 (S)	73	18	81 (S)
17	<i>Torulopsis magnolie</i> IMAP 4425	8	82	75 (S)	51	32	71 (S)
18	<i>Torulopsis molischiana</i> CBS 837	14	71	92 (S)	68	16	58 (S)
19	<i>Torulopsis pinus</i> IMAP 4573	73	traces		68	13	4 (S)
20	<i>Torulopsis castelli</i> CBS 4332	11	11	50 (S)	67	15	91 (S)

Screening conditions: substrate 2 mg/mL, DMSO 5% v/v, glucose 50 mg/mL, biocatalyst double concentrated compared to the liquid culture, 96 h and 30 °C.

¹ Conversion (*c* (%)) and enantiomeric excesses (*ee* (%)) were determined by HPLC analysis on chiral column (see Materials and Methods section).

As far as the bioreduction of compound **13** concerns (Table 3), this bulkier substrate was generally transformed to the corresponding lactone with lower conversion with respect to compound **8**. Interestingly, the screened yeast strains showed different stereoselectivity in reducing **13** leading to the formation of both the enantiomers of **16** with different degrees of enantioselectivity. In details, five yeast strains (Entries 4, 8, 10, 11, 13) catalyzed the formation of the *anti*-Prelog lactone (*R*)-**16** with up to 87% of *ee*, while all the other microorganisms produce the Prelog product with poor to good *ee*.

This inversion of enantioselectivity in the reduction of **13**, with the respect to **8**, may be caused by different factors. As a matter of fact, several enzymes endowed by different affinity and selectivity towards the two substrates **8** and **13** could be present in the same yeast. In addition, the bulkier and more flexible nature of compound **13**, as previously demonstrated *in silico* in the case of *E. coli*/RasADH, could induce a different

binding pose within the catalytic site of the same reductase thus offering a distinctive prochiral face to the cofactor.

Furthermore, as previously mentioned, in all the biotransformations modest to considerable amounts of **12** were formed attesting the high hydrolytic activity of these microorganisms toward compound **13**.

This preliminary screening (Table 3) allowed us to identify *T. castelli* CBS 4332 as the most promising biocatalyst for the reduction of **13** to the Prelog enantiomer (*S*)-**16** with a good *ee* (> 90%). Interestingly, *Torulopsis* sp. yeasts were already applied for the reduction of bulky substrate [35] and for these reasons *Torulopsis* strains were selected for upcoming experiments for the optimization of the biotransformations. Specifically, the undesired hydrolytic activity shown toward **13** has been studied to address if (1) it could be lowered avoiding competition with the desired reducing activity or if (2) it could be directly exploited to produce the target lactone **16** from **12** (Scheme 2). Also, the reasons at the basis of the modest conversions of **13** into **16** have been investigated in terms of substrate loading, substrate/product inhibitory activity and in-cell availability of redox cofactors via *ad hoc* designed experiments.

At first, we tried to suppress/lower the hydrolytic activity using bulkier esters as substrates, *i.e.* ethyl (**14**) and *iso*-propyl (**15**) derivatives, in the presence of *T. magnolie*, *T. molischiana*, and *T. castelli*. Also, a hexane-buffer biphasic system was tested using **13** as substrate. As shown in Table 4 (Entries 1-3), the use of bulkier esters or a biphasic system did not produce a significant increase in the amount of **16** with the respect to **12**.

Table 4. Studies on the bioreduction of the different prochiral ketones.

Entry	Substrate	<i>T. magnolie</i>			<i>T. molischiana</i>			<i>T. castelli</i>		
		16 (%)	12 (%)	16/12 ¹	16 (%)	12 (%)	16/12 ¹	16 (%)	12 (%)	16/12 ¹
1	14	30	57	0.5	28	51	0.5	10	63	0.1
2	15	16	36	0.4	20	47	0.4	5	41	0.1
3	13-Biphasic ²	0	37	N.C.	0	34	N.C.	0	43	N.C.
4	13, 1 mg/mL	29	67	0.4	51	49	1.0	25	60	0.4
5	13, 2 mg/mL	15	63	0.2	16	67	0.2	15	67	0.2

Cell grown for 72 h in solid medium and 48 h in liquid medium (S-L), 96 h of biotransformation, [substrates] = 2 mg/mL, 5% v/v DMSO, glucose 50 g/L.

¹ Ratio between the HPLC abundance (%) of compounds **16** and **12**.

² Hexane : Tris/HCl buffer (0.1 M, pH 7) = 1:1.

N.C. = not calculated.


Thus, according to the potential inhibitory effects and/or different enzymatic activities above described, we focused our attention on the effects of using different concentrations of **13** on the **16/12** ratio (Table 4, Entries 4, 5). We discovered that increasing the concentration of compound **13** always lowered the **16/12** ratio, *de facto* facilitating the undesired hydrolytic processes.

In addition, based on these results we again hypothesized substrate/product inhibitory effects and/or a competition between hydrolytic and redox enzymes characterized by different kinetics.

Afterwards, a set of glucose-feeding experiments (feed every 24 h 50 g/L) was conducted to investigate if the low conversion of compound **13** into **16** could be ascribed to poor redox cofactor availability in the cells. Percentages of compounds **16** and **12** seemed to be unmodified by this approach as the **16/12** ratio basically did not change in all the experiments (Data not shown).

Finally, considering the high hydrolytic activities of the *Torulopsis* yeasts and the possible alternative reaction pathways reported in Scheme 3 we checked whether the microorganisms could have directly reduced compound **12** with better conversion and enantioselectivity (Table 5).

In the meantime, we could also determine if the reduction of **12** would be enantioconvergent with the bioreduction of **13** using the same biocatalyst.



Scheme 3. Different biotransformation pathways of compound **13** in cells.

Surprisingly, the three microorganisms produced (*S*)-**16** with excellent *ees* (97-98%). These results corroborated the hypothesis of the presence of no-stereoconvergent ADHs operating on **13** and/or **12** that lowered the *ees* of the previously discussed biotransformations. This is of relevance in the case of bioreduction of **12** catalyzed by *T. magnolie* and *T. molischiana* whose *ees* were significantly higher when compared to the results of Table 3.

Table 5. Studies on bioreduction of compound **12**.

Entry	Microorganism	<i>c</i> (%) ¹	<i>ee</i> (%) ¹
1 ₁₃	<i>T. magnolie</i>	44	98 (<i>S</i>)
2 ₁₄	<i>T. molischiana</i>	35	98 (<i>S</i>)
3 ₁₅	<i>T. castelli</i>	17	97 (<i>S</i>)

Cell grown for 72 h in solid medium and 48 h in liquid medium (S-L), 96 h of biotransformation, [**12**] = 1 mg/mL. *T. molischiana* [cells] = 21.2 mg/mL; *T. magnolie* [cells] = 20.4 mg/mL; *T. castelli* [cells] = 15.7 mg/mL.

¹ Conversion (*c* (%)) and enantiomeric excesses (*ee* (%)) were determined by HPLC analysis on chiral column (see Materials and Methods section).

Once we discovered that compound **12** represented a more convenient substrate, we decided to optimize its biotransformation *via* an approach of reaction media engineering. Specifically, we tested: (1) possible inhibitory effects of compound **16**; (2) the concentration of compound **12** and, most importantly; (3) the effects of different co-solvents and co-substrates.

The presence of 0.5 mg/mL of **16** did not produce any inhibitory effects on bioreduction of compound **12**, while high substrate's concentrations sensibly lowered its conversion into **16** (from 22% in the presence of 2.0 mg/mL to the 68% in the presence of 0.5 mg/mL).

The effects of different co-substrates and co-solvents on the bioconversion of compound **12** into (*S*)-**16** are reported in Table 6. The use of increasing concentrations of glucose (Entries 2-3) or glycerol did not produce any appreciable effects on the conversion of compound **12** (Entry 5), while xylose slightly lowered it. At variance to what we discovered before for the bioreduction of bulky ketones mediated by *Torulopsis* sp. [35], the 10% v/v of *iso*-propanol lowered the conversion of compound **12** (Entry 1), highlighting that different ADHs could be involved in the reduction of this specific substrate. Based on these results we decided to use a glucose-based NADES [55] (natural deep eutectic solvents, Entries 6 and 7) since they merged the characteristics of green and biocompatible co-solvent with a convenient co-substrate. This strategy allowed us to strongly increase the conversion of **12** into the target (*S*)-**16**. Additionally, a double growth passage on solid medium and then in liquid medium was found to increase the conversion of **12**.

Table 6. Co-substrates and co-solvents effects on bioreduction of **12**.

Entry	Co-substrate	<i>T. molischiana</i>		<i>T. magnolie</i>	
		<i>c</i> (%) ¹	<i>ee</i> (%) ¹	<i>c</i> (%) ¹	<i>ee</i> (%) ¹
1	<i>iso</i> -Propanol 10% v/v	10	98 (<i>S</i>)	N.C.	
2	Glucose 50 g/L	36	97 (<i>S</i>)	34	96 (<i>S</i>)
3	Glucose 100 g/L	43	93 (<i>S</i>)	33	93 (<i>S</i>)
4	Xylose 50 g/L	41	92 (<i>S</i>)	30	94 (<i>S</i>)
5	Glycerol 50 g/L	25	97 (<i>S</i>)	22	96 (<i>S</i>)
6	NADES (glu-gly) ²	80	93 (<i>S</i>)	79	97 (<i>S</i>)
7	NADES (ChCl-glu) ³	82	97 (<i>S</i>)	60	96 (<i>S</i>)

Cell grown for 48 h (*S*-L), 96 h of biotransformation, **12** 1 mg/mL. *T. molischiana* [cells]= 21.2 mg/mL; *T. magnolie* [cells]= 20.4 mg/mL.

¹ Conversion (*c* (%)) and enantiomeric excesses (*ee* (%)) were determined by HPLC analysis on chiral column (see Materials and Methods section).

² NADES (glu-gly): Glucose-glycerol.

³ NADES (CHCl-glu): Choline chloride-glucose.

N.C. = no conversion.

3.2.4 Semi-preparative scale preparation of (*R*)- and (*S*)-**16**

(*R*)- and (*S*)-**16** were prepared using two different biocatalytic systems.

The *anti*-Prelog enantiomer (*R*)-**16** was prepared using *E. coli*/SyADH as biocatalyst (10 mg/mL of lyophilized *E. coli* cells) using a 10 mM solution of **13** prepared in the presence of 10% v/v DMSO in PB buffer (pH 7.0, 50 mM), glucose (40 mM), BmGDH (0.5 U/mL) and NADP⁺ (0.4 mM). Product was obtained with an *ee* >99% (80% molar conversion) and a 70% isolated yield (see the Material and Methods section).

The Prelog product (*S*)-**16**, the direct precursor of (+)-Harzialactone A [15], was obtained by biotransformation with whole cells of *T. molischiana* (18.9 mg/mL) grown in solid medium 24 h (double step) and liquid medium 48 h. The cells were suspended in Tris/HCl buffer (0.1M, pH 7) containing 10% NADES (ChCl-glu) and 1 g/L of **12**. After 96 h (*S*)-**16** was obtained with a molar conversion of 88%, *ee* >97% and 65% isolated yield (see the Material and Methods section).

4. Conclusions

In this work, after screening libraries of more than 15 wild type, promiscuous oxidoreductases, a commercial kit of ADHs and 20 different yeasts strains as whole-cell biocatalysts, we have successfully designed and realized a chemo-enzymatic entry to both the enantiomer of **16**. To the best of our knowledge this is the first time that a biocatalytic approach is applied for the preparation of (*S*)-**16**, a valuable synthon of the bioactive marine drug (+)-Harzialactone A.

This convenient, facile, and biocatalytic approach to enantiomerically enriched γ -lactones is now being exploited by us for the synthesis of the two enantiomers of the metabolite 5-(3',4'-dihydroxyphenyl)- γ -valerolactone, that we found to be active against *Candida albicans* when tested as a racemate [10].

Supplementary Materials: The supporting information file for the manuscript was uploaded.

Author Contributions: Angelica Artasensi and Ivan Bassanini share the First author.

Funding: This work was supported by Università degli Studi di Milano.

References

1. Pang, X.; Lin, X.; Yang, J.; Zhou, X.; Yang, B.; Wang, J.; Liu, Y. Spiro-Phthalides and Isocoumarins Isolated from the Marine-Sponge-Derived Fungus *Setosphaeria* Sp. SCSIO41009. *J Nat Prod* **2018**, *81*, 1860–1868, doi:10.1021/acs.jnatprod.8b00345. 585
586
587
588
2. Carroll, A.R.; Copp, B.R.; Davis, R.A.; Keyzers, R.A.; Prinsep, M.R. Marine Natural Products. *Nat Prod Rep* **2022**, *39*, 1122–1171, doi:10.1039/D1NP00076D. 589
590
3. Guo, R.; Li, G.; Zhang, Z.; Peng, X. Structures and Biological Activities of Secondary Metabolites from *Trichoderma Harzianum*. *Mar Drugs* **2022**, *20*, 701, doi:10.3390/md20110701. 591
592
4. Zhang, J.-L.; Tang, W.-L.; Huang, Q.-R.; Li, Y.-Z.; Wei, M.-L.; Jiang, L.-L.; Liu, C.; Yu, X.; Zhu, H.-W.; Chen, G.-Z.; et al. *Trichoderma*: A Treasure House of Structurally Diverse Secondary Metabolites With Medicinal Importance. *Front Microbiol* **2021**, *12*, doi:10.3389/fmicb.2021.723828. 593
594
595
5. Amagata, T.; Usami, Y.; Minoura, K.; Ito, T.; Numata, A. Cytotoxic Substances Produced by a Fungal Strain from a Sponge: Physico-Chemical Properties and Structures. *J Antibiot (Tokyo)* **1998**, *51*, 33–40, doi:10.7164/antibiotics.51.33. 596
597
598
6. Souza, A.D.L.; Rodrigues-Filho, E.; Souza, A.Q.L.; Henrique-Silva, F.; Pereira, J.O. A New Guaiane Mannoside from a *Eutypa*-like Fungus Isolated from *Murraya Paniculata* in Brazil. *J Braz Chem Soc* **2008**, *19*, 1321–1325, doi:10.1590/S0103-50532008000700014. 599
600
601
7. Pawar, V.U.; Ghosh, S.; Chopade, B.A.; Shinde, V.S. Design and Synthesis of Harzialactone Analogues: Promising Anticancer Agents. *Bioorg Med Chem Lett* **2010**, *20*, 7243–7245, doi:10.1016/j.bmcl.2010.10.100. 602
603
604
8. Braun, G.H.; Ramos, H.P.; Candido, A.C.B.B.; Pedroso, R.C.N.; Siqueira, K.A.; Soares, M.A.; Dias, G.M.; Magalhães, L.G.; Ambrósio, S.R.; Januário, A.H.; et al. Evaluation of Antileishmanial Activity of Harzialactone a Isolated from the Marine-Derived Fungus *Paecilomyces* Sp. *Nat Prod Res* **2021**, *35*, 1644–1647, doi:10.1080/14786419.2019.1619725. 605
606
607
608
9. Artasensi, A.; Baron, G.; Vistoli, G.; Aldini, G.; Fumagalli, L. (Z)-5-(30,40-Bis(Benzyloxy)Benzylidene)Furan-2(5H)-One. *Molbank* **2021**, *2021*, doi:10.3390/M1193. 609
610
10. Ottaviano, E.; Baron, G.; Fumagalli, L.; Leite, J.; Colombo, E.A.; Artasensi, A.; Aldini, G.; Borghi, E. *Candida Albicans* Biofilm Inhibition by Two *Vaccinium Macrocarpon* (Cranberry) Urinary Metabolites: 5-(3',4'-Dihydroxyphenyl)- γ -Valerolactone and 4-Hydroxybenzoic Acid. *Microorganisms* **2021**, *9*, doi:10.3390/microorganisms9071492. 611
612
613
614
11. Baron, G.; Altomare, A.; Regazzoni, L.; Fumagalli, L.; Artasensi, A.; Borghi, E.; Ottaviano, E.; del Bo, C.; Riso, P.; Allegrini, P.; et al. Profiling *Vaccinium Macrocarpon* Components and Metabolites in Human Urine and the Urine Ex-Vivo Effect on *Candida Albicans* Adhesion and Biofilm-Formation. *Biochem Pharmacol* **2020**, *173*, doi:10.1016/j.bcp.2019.113726. 615
616
617
618
12. Mereyala, H.B.; Gadikota, R.R. A Concise Synthesis of Harzialactone A from D-Glucose and Revision of Absolute Stereochemistry. *Tetrahedron Asymmetry* **1999**, *10*, 2305–2306, doi:10.1016/S0957-4166(99)00245-1. 619
620
621
13. He, L.; Zhang, S.; Wu, Y.; Li, Y. Synthesis of (-)-Harzialactone a from a Readily Accessible Epoxy Chiral Building Block. *Chin J Chem* **2011**, *29*, 2664–2668, doi:10.1002/cjoc.201100373. 622
623
14. Ballaschk, F.; Özkaya, Y.; Kirsch, S.F. Stereocontrolled Synthesis of Harzialactone A and Its Three Stereoisomers by Use of Standardized Polyketide Building Blocks. *European J Org Chem* **2020**, *2020*, 6078–6080, doi:10.1002/ejoc.202001046. 624
625
626
15. Kotkar, S.P.; Suryavanshi, G.S.; Sudalai, A. A Short Synthesis of (+)-Harzialactone A and (R)-(+)-4-Hexanolide via Proline-Catalyzed Sequential α -Aminoxylation and Horner-Wadsworth-Emmons 627
628

- Olefination of Aldehydes. *Tetrahedron Asymmetry* **2007**, *18*, 1795–1798, doi:10.1016/j.tetasy.2007.07.031. 629
16. Shu, C.; Liu, M.Q.; Sun, Y.Z.; Ye, L.W. Efficient Synthesis of γ -Lactones via Gold-Catalyzed Tandem Cycloisomerization/Oxidation. *Org Lett* **2012**, *14*, 4958–4961, doi:10.1021/ol302323a. 630
17. Chen, B.; Yin, H.F.; Wang, Z.S.; Xu, J.H. New Synthesis of Harzialactone A via Kinetic Resolution Using Recombinant *Fusarium Proliferatum* Lactonase. *Tetrahedron Asymmetry* **2010**, *21*, 237–240, doi:10.1016/j.tetasy.2010.01.021. 631
18. Ketterer, C.; Wünsch, B. Lipase-Catalyzed Kinetic Resolution of 2-Phenylethanol Derivatives and Chiral Oxa-Pictet-Spengler Reaction as the Key Steps in the Synthesis of Enantiomerically Pure Tricyclic Amines. *European J Org Chem* **2012**, *2012*, 2428–2444, doi:10.1002/ejoc.201101800. 632
19. Shimotori, Y.; Hoshi, M.; Inoue, K.; Osanai, T.; Okabe, H.; Miyakoshi, T. Preparation of Optically Active 4-Substituted γ -Lactones by Lipase-Catalyzed Optical Resolution. *Heterocycl Comm* **2015**, *21*, 165–174, doi:10.1515/hc-2015-0027. 633
20. Borowiecki, P.; Telatycka, N.; Tataruch, M.; Żądło-Dobrowolska, A.; Reiter, T.; Schühle, K.; Heider, J.; Szaleniec, M.; Kroutil, W. Biocatalytic Asymmetric Reduction of Γ -Keto Esters to Access Optically Active Γ -Aryl- γ -butyrolactones. *Adv Synth Catal* **2020**, *362*, 2012–2029, doi:10.1002/adsc.201901483. 634
21. Díaz-Rodríguez, A.; Borzeicka, W.; Lavandera, I.; Gotor, V. Stereodivergent Preparation of Valuable γ -Or δ -Hydroxy Esters and Lactones through One-Pot Cascade or Tandem Chemoenzymatic Protocols. *ACS Catal* **2014**, *4*, 386–393, doi:10.1021/cs4010024. 635
22. Zhou, J.; Xu, G.; Ni, Y. Stereochemistry in Asymmetric Reduction of Bulky–Bulky Ketones by Alcohol Dehydrogenases. *ACS Catal* **2020**, *10*, 10954–10966, doi:10.1021/acscatal.0c02646. 636
23. Wu, K.; Yang, Z.; Meng, X.; Chen, R.; Huang, J.; Shao, L. Engineering an Alcohol Dehydrogenase with Enhanced Activity and Stereoselectivity toward Diaryl Ketones: Reduction of Steric Hindrance and Change of the Stereocontrol Element. *Catal Sci Technol* **2020**, *10*, 1650–1660, doi:10.1039/c9cy02444a. 637
24. Xu, G.; Dai, W.; Wang, Y.; Zhang, L.; Sun, Z.; Zhou, J.; Ni, Y. Molecular Switch Manipulating Prelog Priority of an Alcohol Dehydrogenase toward Bulky-Bulky Ketones. *Molecular Catalysis* **2020**, *484*, doi:10.1016/j.mcat.2019.110741. 638
25. de Matos, I.L.; Birolli, W.G.; Santos, D. de A.; Nitschke, M.; Porto, A.L.M. Stereoselective Reduction of Flavanones by Marine-Derived Fungi. *Molecular Catalysis* **2021**, *513*, doi:10.1016/j.mcat.2021.111734. 639
26. Özgen, F.F.; Jorea, A.; Capaldo, L.; Kourist, R.; Ravelli, D.; Schmidt, S. The Synthesis of Chiral γ -Lactones by Merging Decatungstate Photocatalysis with Biocatalysis. *ChemCatChem* **2022**, *14*, doi:10.1002/cctc.202200855. 640
27. Forzato, C.; Gandolfi, R.; Molinari, F.; Nitti, P.; Pitacco, G.; Valentin, E. Microbial Bioreductions of γ - and δ -Ketoacids and Their Esters. *Tetrahedron Asymmetry* **2001**, *12*, 1039–1046, doi:10.1016/S0957-4166(01)00184-7. 641
28. Ferrandi, E.E.; Bassanini, I.; Bertuletti, S.; Riva, S.; Tognoli, C.; Vanoni, M.; Monti, D. Functional Characterization and Synthetic Application of Is2-SDR, a Novel Thermostable and Promiscuous Ketoreductase from a Hot Spring Metagenome. *Int J Mol Sci* **2022**, *23*, doi:10.3390/ijms232012153. 642
29. Kędziora, K.; Bisogno, F.R.; Lavandera, I.; Gotor-Fernández, V.; Montejo-Bernardo, J.; García-Granda, S.; Kroutil, W.; Gotor, V. Expanding the Scope of Alcohol Dehydrogenases towards Bulkier 643

- Substrates: Stereo- and Enantioselectivity for α,α -Dihalogenated Ketones. *ChemCatChem* **2014**, *6*, 1066–1072, doi:10.1002/cctc.201300834. 672
30. Nasti, R.; Bassanini, I.; Ferrandi, E.E.; Linguardo, F.; Bertuletti, S.; Vanoni, M.; Riva, S.; Verotta, L.; Monti, D. Stereoselective Biocatalyzed Reductions of Ginger Active Components Recovered from Industrial Wastes. *ChemBioChem* **2022**, *23*, doi:10.1002/cbic.202200105. 673
31. Lavandera, I.; Kern, A.; Resch, V.; Ferreira-Silva, B.; Glieder, A.; Fabian, W.M.F.; de Wildeman, S.; Kroutil, W. One-Way Biohydrogen Transfer for Oxidation of *Sec*-Alcohols. *Org Lett* **2008**, *10*, 2155–2158, doi:10.1021/ol800549f. 674
32. Bertuletti, S.; Ferrandi, E.E.; Marzorati, S.; Vanoni, M.; Riva, S.; Monti, D. Insights into the Substrate Promiscuity of Novel Hydroxysteroid Dehydrogenases. *Adv Synth Catal* **2020**, *362*, 2474–2485, doi:10.1002/adsc.202000120. 675
33. Ferrandi, E.E.; Bertuletti, S.; Monti, D.; Riva, S. Hydroxysteroid Dehydrogenases: An Ongoing Story. *European J Org Chem* **2020**, *2020*, 4463–4473, doi:10.1002/ejoc.202000192. 676
34. Bertuletti, S.; Bayout, I.; Bassanini, I.; Ferrandi, E.E.; Bouzemi, N.; Monti, D.; Riva, S. Biocatalytic Approaches to the Enantiomers of Wieland–Miescher Ketone and Its Derivatives. *European J Org Chem* **2021**, *2021*, 3992–3998, doi:10.1002/ejoc.202100174. 677
35. Gandolfi, R.; Coffetti, G.; Facchetti, G.; Rimoldi, I. Double Approaches for Obtaining an Asymmetric One-Pot Addition/Reduction Reaction. *Molecular Catalysis* **2022**, doi:10.1016/j.mcat.2022.112716. 678
36. Zhou, J.; Jia, M.; Song, M.; Huang, Z.; Steiner, A.; An, Q.; Ma, J.; Guo, Z.; Zhang, Q.; Sun, H.; et al. Chemoselective Oxyfunctionalization of Functionalized Benzylic Compounds with a Manganese Catalyst. *Angewandte Chemie International Edition* **2022**, *61*, doi:10.1002/anie.202205983. 679
37. Santra, S.K.; Szpilman, A.M. Visible-Spectrum Solar-Light-Mediated Benzylic C-H Oxygenation Using 9,10-Dibromoanthracene As an Initiator. *Journal of Organic Chemistry* **2021**, *86*, 1164–1171, doi:10.1021/acs.joc.0c01720. 680
38. Zhao, M.; Li, W.; Li, X.; Ren, K.; Tao, X.; Xie, X.; Ayad, T.; Ratovelomanana-Vidal, V.; Zhang, Z. Enantioselective Ruthenium(II)/Xyl-SunPhos/Daipen-Catalyzed Hydrogenation of γ -Ketoamides. *Journal of Organic Chemistry* **2014**, *79*, 6164–6171, doi:10.1021/jo5008916. 681
39. Alazet, S.; Le Vaillant, F.; Nicolai, S.; Courant, T.; Waser, J. Divergent Access to (1,1) and (1,2)-Azidolactones from Alkenes Using Hypervalent Iodine Reagents. *Chemistry - A European Journal* **2017**, *23*, 9501–9504, doi:10.1002/chem.201702599. 682
40. Guo, K.; Zhang, Z.; Li, A.; Li, Y.; Huang, J.; Yang, Z. Photoredox-Catalyzed Isomerization of Highly Substituted Allylic Alcohols by C–H Bond Activation. *Angewandte Chemie - International Edition* **2020**, *59*, 11660–11668, doi:10.1002/anie.202000743. 683
41. Crotti, M.; Parmeggiani, F.; Ferrandi, E.E.; Gatti, F.G.; Sacchetti, A.; Riva, S.; Brenna, E.; Monti, D. Stereoselectivity Switch in the Reduction of α -Alkyl- β -Arylenones by Structure-Guided Designed Variants of the Ene Reductase OYE1. *Front Bioeng Biotechnol* **2019**, *7*, doi:10.3389/fbioe.2019.00089. 684
42. Lavandera, I.; Kern, A.; Ferreira-Silva, B.; Glieder, A.; De Wildeman, S.; Kroutil, W. Stereoselective Bioreduction of Bulky-Bulky Ketones by a Novel ADH from *Ralstonia* Sp. *Journal of Organic Chemistry* **2008**, *73*, 6003–6005, doi:10.1021/jo800849d. 685
43. Deng, C.Q.; Liu, J.; Luo, J.H.; Gan, L.J.; Deng, J.; Fu, Y. Proton-Promoted Nickel-Catalyzed Asymmetric Hydrogenation of Aliphatic Ketoacids. *Angewandte Chemie - International Edition* **2022**, *61*, doi:10.1002/anie.202115983. 686

44. Armstrong, A.; Braddock, D.C.; Jones, A.X.; Clark, S. Catalytic Asymmetric Bromolactonization Reactions Using (DHQD) 2PHAL-Benzoic Acid Combinations. *Tetrahedron Lett* **2013**, *54*, 7004–7008, doi:10.1016/j.tetlet.2013.10.043. 715
716
717
45. Lerchner, A.; Jarasch, A.; Meining, W.; Schiefner, A.; Skerra, A. Crystallographic Analysis and Structure-Guided Engineering of NADPH-Dependent *Ralstonia* Sp. Alcohol Dehydrogenase toward NADH Cosubstrate Specificity. *Biotechnol Bioeng* **2013**, *110*, 2803–2814, doi:10.1002/bit.24956. 718
719
720
46. Pedretti, A.; Mazzolari, A.; Gervasoni, S.; Fumagalli, L.; Vistoli, G. The VEGA Suite of Programs: An Versatile Platform for Cheminformatics and Drug Design Projects. *Bioinformatics* **2021**, *37*, 1174–1175, doi:10.1093/bioinformatics/btaa774. 721
722
723
47. Chen, X.; Zhang, H.; Maria-Solano, M.A.; Liu, W.; Li, J.; Feng, J.; Liu, X.; Osuna, S.; Guo, R.-T.; Wu, Q.; et al. Efficient Reductive Desymmetrization of Bulky 1,3-Cyclodiketones Enabled by Structure-Guided Directed Evolution of a Carbonyl Reductase. *Nat Catal* **2019**, *2*, 931–941, doi:10.1038/s41929-019-0347-y. 724
725
726
727
48. Korb, O.; Stütze, T.; Exner, T.E. Empirical Scoring Functions for Advanced Protein–Ligand Docking with PLANTS. *J Chem Inf Model* **2009**, *49*, 84–96, doi:10.1021/ci800298z. 728
729
49. Pedretti, A.; Granito, C.; Mazzolari, A.; Vistoli, G. Structural Effects of Some Relevant Missense Mutations on the MECP2-DNA Binding: A MD Study Analyzed by Rescore+, a Versatile Rescoring Tool of the VEGA ZZ Program. *Mol Inform* **2016**, *35*, 424–433, doi:10.1002/minf.201501030. 730
731
732
50. Song, J.-W.; Jeon, E.-Y.; Song, D.-H.; Jang, H.-Y.; Bornscheuer, U.T.; Oh, D.-K.; Park, J.-B. Multistep Enzymatic Synthesis of Long-Chain α,ω -Dicarboxylic and ω -Hydroxycarboxylic Acids from Renewable Fatty Acids and Plant Oils. *Angewandte Chemie International Edition* **2013**, *52*, 2534–2537, doi:10.1002/anie.201209187. 733
734
735
736
51. Edegger, K.; Stampfer, W.; Seisser, B.; Faber, K.; Mayer, S.F.; Oehrlein, R.; Hafner, A.; Kroutil, W. Regio- and Stereoselective Reduction of Diketones and Oxidation of Diols by Biocatalytic Hydrogen Transfer. *European J Org Chem* **2006**, *2006*, 1904–1909, doi:10.1002/ejoc.200500839. 737
738
739
52. Niefind, K.; Müller, J.; Riebel, B.; Hummel, W.; Schomburg, D. The Crystal Structure of R-Specific Alcohol Dehydrogenase from *Lactobacillus Brevis* Suggests the Structural Basis of Its Metal Dependency. *J Mol Biol* **2003**, *327*, 317–328, doi:10.1016/S0022-2836(03)00081-0. 740
741
742
53. Ferrandi, E.E.; Bertolesi, G.M.; Polentini, F.; Negri, A.; Riva, S.; Monti, D. In Search of Sustainable Chemical Processes: Cloning, Recombinant Expression, and Functional Characterization of the 7 α - and 7 β -Hydroxysteroid Dehydrogenases from *Clostridium Absonum*. *Appl Microbiol Biotechnol* **2012**, *95*, 1221–1233, doi:10.1007/S00253-011-3798-X/FIGURES/5. 743
744
745
746
54. Schmid, R.; Braun, M.; Liu, L.; Aigner, A.; Weuster-Botz, D. 7 α -Hydroxysteroid Dehydrogenase Knockout Mutants and Use Therefor **2015**. 747
748
55. Annunziata, F.; Guaglio, A.; Conti, P.; Tamborini, L.; Gandolfi, R. Continuous-Flow Stereoselective Reduction of Prochiral Ketones in a Whole Cell Bioreactor with Natural Deep Eutectic Solvents. *Green Chemistry* **2022**, *24*, 950–956, doi:10.1039/d1gc03786b. 749
750
751
752
753
754