



Original article

Clinical characteristics and outcomes of vaccinated patients hospitalised with SARS-CoV-2 breakthrough infection: Multi-IPV, a multicentre study in Northern Italy



Andrea Lombardi ^{ae,af,*}, Simone Villa ^{af}, Marta Colaneri ^{ag}, Giovanni Scaglione ^{ag}, Francesca Bai ^{ah}, Benedetta Varisco ^{ah}, Valeria Bono ^{ah}, Antonio Vena ^{ai,aj}, Chiara Dentone ^{ai,aj}, Chiara Russo ^{ai,aj}, Mauro Tettamanti ^{ak}, Giulia Renisi ^{ae}, Giulia Viero ^{ae}, Cecilia Azzarà ^{ae}, Marco Mantero ^{af,al}, Flora Peyvandi ^{af,am}, Matteo Bassetti ^{ai,aj}, Giulia Marchetti ^{ah}, Antonio Muscatello ^{ae}, Alessandro Nobili ^{ak}, Andrea Gori ^{af,ag}, Alessandra Bandera ^{ae,af}, “COVID 19 NETWORK” (Silvano Bosari ^a, Luigia Scudeller ^a, Giuliana Fusetti ^a, Laura Rusconi ^a, Silvia Dell’Orto ^a, Daniele Prati ^b, Luca Valenti ^b, Silvia Giovannelli ^b, Maria Manunta ^b, Giuseppe Lamorte ^b, Francesca Ferarri ^b, Andrea Gori ^c, Alessandra Bandera ^c, Antonio Muscatello ^c, Davide Mangioni ^c, Laura Alagna ^c, Giorgio Bozzi ^c, Andrea Lombardi ^c, Riccardo Ungaro ^c, Giuseppe Ancona ^c, Marco Mussa ^c, Bianca Veronica Mariani ^c, Matteo Bolis ^c, Nathalie Iannotti ^c, Serena Ludovisi ^c, Agnese Comelli ^c, Giulia Renisi ^c, Simona Biscarini ^c, Valeria Castelli ^c, Emanuele Palomba ^c, Marco Fava ^c, Carlo Alberto Peri ^c, Paola Saltini ^c, Giulia Viero ^c, Teresa Itri ^c, Valentina Ferroni ^c, Valeria Pastore ^c, Roberta Massafra ^c, Arianna Liparoti ^c, Toussaint Muheberimana ^c, Alessandro Giommi ^c, Rosaria Bianco ^c, Grazia Eliana Chitani ^c, Chiara Bobbio ^c, Irene De Matteis ^c, Angelo Bianchi Bonomi ^c, Flora Peyvandi ^d, Roberta Gualtierotti ^d, Barbara Ferrari ^d, Raffaella Rossio ^d, Nadia Boasi ^d, Erica Pagliaro ^d, Costanza Massimo ^d, Michele De Caro ^d, Andrea Giachi ^d, Nicola Montano ^e, Barbara Vigone ^e, Chiara Bellocchi ^e, Angelica Carandina ^e, Elisa Fiorelli ^e, Valerie Melli ^e, Eleonora Tobaldini ^e, Francesco Blasi ^f, Stefano Aliberti ^f, Maura Spotti ^f, Leonardo Terranova ^f, Sofia Misuraca ^f, Alice D’Adda ^f, Silvia Della Fiore ^f, Marta Di Pasquale ^f, Marco Mantero ^f, Martina Contarini ^f, Margherita Ori ^f, Letizia Morlacchi ^f, Valeria Rossetti ^f, Andrea Gramegna ^f, Maria Pappalettera ^f, Mirta Cavallini ^f, Agata Buscemi ^f, Marco Vicenzi ^g, Irena Rota ^g, Giorgio Costantino ^g, Monica Solbiati ^g, Ludovico Furlan ^g, Marta Mancarella ^g, Giulia Colombo ^g, Giorgio Colombo ^g, Alice Fanin ^g, Mariele Passarella ^g, Valter Monzani ^h, Ciro Canetta ^h, Angelo Rovellini ^h, Laura Barbetta ^h, Filippo Billi ^h, Christian Folli ^h, Silvia Accordino ^h, Diletta Maira ⁱ, Cinzia Maria Hu ⁱ, Irene Motta ⁱ, Natalia Scaramellini ⁱ, Anna Ludovica Fracanzani ^j, Rosa Lombardi ^j, Annalisa Cespiati ^j, Matteo Cesari ^k, Tiziano Lucchi ^k, Marco Proietti ^k, Laura Calcaterra ^k, Clara Mandelli ^k, Carlotta Coppola ^k, Arturo Cerizza ^k, Antonio Maria Pesenti ^l, Giacomo Grasselli ^l, Alessandro Galazzi ^l, Alessandro Nobili ^m, Mauro Tettamanti ^m, Igor Monti ^m, Alessia Antonella Galbussera ^m, Ernesto Crisafulli ^{n,o}, Domenico Girelli ^{n,o}, Alessio Maroccia ^{n,o}, Daniele Gabbiani ^{n,o}, Fabiana Busti ^{n,o}, Alice Vianello ^{n,o}, Marta Biondan ^{n,o}, Filippo Sartori ^{n,o}, Paola Faverio ^{p,q}, Alberto Pesci ^{p,q}

* Correspondence to: Department of Pathophysiology and Transplantation, Via Francesco Sforza 35, 20122 Milan, Italy.
E-mail address: andrea.lombardi@unimi.it (A. Lombardi).

Stefano Zucchetti ^{P,q}, Paolo Bonfanti ^r, Marianna Rossi ^r, Ilaria Beretta ^r, Anna Spolti ^r, Sergio Harari ^{q,s}, Davide Elia ^t, Roberto Cassandro ^t, Antonella Caminati ^t, Francesco Cipollone ^{u,v}, Maria Teresa Guagnano ^{u,v}, Damiano D'Ardes ^{u,v}, Ilaria Rossi ^{u,v}, Francesca Vezzani ^{u,v}, Antonio Spanevello ^{w,x}, Francesca Cherubino ^{w,x}, Dina Visca ^{w,x}, Marco Contoli ^{y,z}, Alberto Papi ^{y,z}, Luca Morandi ^{y,z}, Nicholas Battistini ^{y,z}, Guido Luigi Moreo ^{aa,ab}, Pasqualina Iannuzzi ^{aa,ab}, Daniele Fumagalla ^{ac}, Sara Leone ^{ad}

^{ae} Foundation IRCCS Ca' Granda Ospedale Maggiore Policlinico, Infectious Diseases Unit, Milan, Italy

^{af} Department of Pathophysiology and Transplantation, University of Milan, Milano, Italy

^{ag} Department of Infectious Diseases, Ospedale Luigi Sacco, Milan, Italy

^{ah} Department of Health Sciences, Clinic of Infectious Diseases, ASST Santi Paolo e Carlo, Milan, Italy

^{ai} Department of Health Sciences, University of Genoa, Genoa, Italy

^{aj} Infectious Diseases Unit, IRCCS Ospedale Policlinico San Martino, Genoa, Italy

^{ak} Department of Health Policy, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milano, Italy

^{al} Foundation IRCCS Ca' Granda Ospedale Maggiore Policlinico, Respiratory Unit and Cystic Fibrosis Center, Milan, Italy

^{am} Foundation IRCCS Ca' Granda Ospedale Maggiore Policlinico, Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Milan, Italy

^a Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Scientific Direction, Italy

^b Department of Transfusion Medicine and Hematology (Biobank), Italy

^c Infectious Diseases Unit, Italy

^d Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Italy

^e UOC Internal Medicine, Immunology and Allergology, Italy

^f Respiratory Unit and Cystic Fibrosis Center, Italy

^g Cardiology Unit, Italy

^h Acute Internal Medicine, Italy

ⁱ Rare Diseases Center, Italy

^j General Medicine and Metabolic Diseases, Italy

^k Geriatric Unit, Italy

^l Intensive Care Unit, Italy

^m Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Italy

ⁿ Policlinico G.B. Rossi, Verona, Italy

^o UOC di Medicina d'Urgenza, Italy

^p Ospedale San Gerardo, ASST Monza, Italy

^q UOC Pneumologia, Italy

^r Malattie Infettive, Italy

^s San Giuseppe Hospital MultiMedica IRCCS and community health, Università degli Studi di Milano, Italy

^t Unità di Pneumologia e terapia Semi-intensiva respiratoria, Servizio di Fisiopatologia Respiratoria ed Emodinamica Polmonare, Italy

^u Ospedale Clinicizzato "SS. Annunziata", Italy

^v Clinica Medica, Italy

^w ICS Maugeri Tradate, Università Insubria, Italy

^x Pneumologia Riabilitativa, Italy

^y Azienda Ospedaliera Universitaria di Ferrara e Dipartimento di Medicina Traslazionale Università di Ferrara, Italy

^z UO Pneumologia

^{aa} Clinica Polispecialistica San Carlo, Italy

^{ab} UO Medicina Interna, Italy

^{ac} UO Oncologia, Italy

^{ad} UO Chirurgia Generale, Italy

ARTICLE INFO

Article history:

Received 6 June 2023

Received in revised form 29 December 2023

Accepted 31 December 2023

Keywords:

Vaccination

Breakthrough infection

SARS-COV-2

ABSTRACT

Background: Despite the well-known efficacy of anti-COVID-19 vaccines in preventing morbidity and mortality, several vaccinated individuals are diagnosed with SARS-CoV-2 breakthrough infection, which might require hospitalisation. This multicentre, observational, and retrospective study aimed to investigate the clinical characteristics and outcomes of vaccinated vs. non-vaccinated patients, both hospitalised with SARS-CoV-2 infection in 3 major hospitals in Northern Italy.

Methods: Data collection was retrospective, and paper and electronic medical records of adult patients with a diagnosed SARS-CoV-2 infection were pseudo-anonymised and analysed. Vaccinated and non-vaccinated individuals were manually paired, using a predetermined matching criterion (similar age, gender, and date of hospitalisation). Demographic, clinical, treatment, and outcome data were compared between groups differing by vaccination status using Pearson's Chi-square and Mann-Whitney tests. Moreover, multiple logistic regression analyses were performed to assess the impact of vaccination status on ICU admission or intra-hospital mortality.

Results: Data from 360 patients were collected. Vaccinated patients presented with a higher prevalence of relevant comorbidities, like kidney replacement therapy or haematological malignancy, despite a milder clinical presentation at the first evaluation. Non-vaccinated patients required intensive care more often than their vaccinated counterparts (8.8% vs. 1.7%, $p = 0.002$). Contrariwise, no difference in intra-hospital mortality was observed between the two groups (19% vs. 20%, $p = 0.853$). These results were confirmed by multivariable logistic regressions, which showed that vaccination was significantly associated with decreased risk of ICU admission (aOR=0.172, 95%CI: 0.039–0.542, $p = 0.007$), but not of intra-hospital mortality (aOR=0.996, 95%CI: 0.582–1.703, $p = 0.987$).

Conclusions: This study provides real-world data on vaccinated patients hospitalised with COVID-19 in Northern Italy. Our results suggest that COVID-19 vaccination has a protective role in individuals with higher risk profiles, especially regarding the need for ICU admission. These findings contribute to our

understanding of SARS-CoV-2 infection outcomes among vaccinated individuals and emphasise the importance of vaccination in preventing severe disease, particularly in those countries with lower first-booster uptake rates.

© 2024 The Author(s). Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Although there are still questions about the impact of SARS-CoV-2 variants and the duration of the immune response, messenger RNA (mRNA)-based and adenoviral-vectored vaccines have shown an overall efficacy of 70–95%, being the most effective way to prevent morbidity and mortality from SARS-CoV-2 [1].

While breakthrough infections, defined as a positive nasopharyngeal swab at least one week after the vaccine schedule, can be expected [2], more attention should be paid to vaccinated patients who experience a progression of COVID-19 to a severe form that requires hospitalisation after being infected [3]. Some studies have already explored the nature and course of SARS-CoV-2 among vaccinated individuals, guiding us in public health preparedness [4–6]. However, despite Italy being the first European country to be heavily affected by the spread of the SARS-CoV-2 pandemic, with a burden of excess mortality [7], real-life data on clinical, laboratory and sociodemographic parameters of patients hospitalised for SARS-CoV-2 despite being fully or partially vaccinated are still scarce [8].

With this in mind, the main objective of this multicentre retrospective observational study was to investigate risk factors and clinical, immunological, and infection characteristics of vaccinated hospitalised patients with SARS-CoV-2 infection in 3 major hospitals in Northern Italy.

2. Materials and methods

2.1. Study design and participants

This study (Multi-IPV study) was a multicentre, observational, retrospective cohort study of hospitalised patients with a confirmed diagnosis of SARS-CoV-2 conducted in Northern Italy (IRCCS Ospedale Maggiore Policlinico, Milan; ASST Santi Paolo e Carlo, Milan; Ospedale Policlinico San Martino, Genoa) from February 2021 to November 2021. The three centres are all university hospitals with around 1000 beds.

Medical records of all adult patients with clinical and virological diagnosis of SARS-CoV-2 infection were pseudo-anonymised and abstracted on standardised data collection forms. Data were retrospectively extracted from paper and electronic medical records only of the patients with complete information regarding their vaccination status by 1 data entry person per hospital.

2.2. Patients' consent

This research project was carried out following a research plan and according to the current version of the Declaration of Helsinki (<https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects>).

The sponsor (IRCCS Ospedale Maggiore Policlinico, Milan) ensured that approval from a Competent Ethics Committee (CEC) was sought for the study. Before the initiation of treatment and data collection, all participants signed an informed consent. The study was approved by the IRCCS Ospedale Maggiore Policlinico Foundation Institutional Review Board (1191_2021bis).

2.3. Patients' characteristics

The demographic data included sex and age. Clinical data included symptoms at presentation, including the use of a clinical presentation score (i.e., WHO Ordinal Scale [9]), and comorbidities among the following: diabetes, obesity (i.e., body mass index equal to 25 kg/m² or above), chronic obstructive pulmonary disease (COPD), kidney disease, liver disease, dialysis, haematological malignancy, solid-organ transplant.

Treatment data included the use of remdesivir, monoclonal antibodies (mAbs), and invasive and non-invasive respiratory support therapies, such as low-flow oxygen, Venturi masks, continuous positive airway pressure (CPAP) helmets, high flow nasal cannula (HFNC), and mechanical ventilation.

Genotyping analysis of SARS-CoV-2 isolates, the COVID-19 vaccine type, and the immunological status were finally considered.

2.4. Main exposure

Vaccination status was considered the primary exposure of interest. This was defined as being vaccinated or not if the patient received at least one COVID-19 vaccine dose.

2.5. Outcomes

The primary outcome was to describe the clinical characteristics of the patients vaccinated for COVID-19 and admitted to the three considered hospitals with a breakthrough infection.

The secondary outcome was to compare the proportions of admissions to intensive care unit (ICU), intra-hospital mortality and type of respiratory support needed, if any, between vaccinated and non-vaccinated individuals hospitalised for SARS-CoV-2.

Finally, we wanted to assess the impact of the COVID-19 vaccine on ICU admission and intra-hospital mortality.

2.6. Statistics

Qualitative variables were described as counts and percentages of each category. Quantitative variables were summarised as median and interquartile range (IQR).

The primary author manually paired all the vaccinated and non-vaccinated individuals based on predetermined matching criteria. Non-vaccinated individuals were matched to vaccinated ones of a similar age (± 5 years compared to the age of the vaccinated patient), same gender, and hospitalised in the same period (± 7 days concerning the date of hospitalisation of the vaccinated patient).

Demographic, clinical, treatment, and outcome data were compared between groups differing by vaccination status using Pearson's Chi-square (χ^2) tests and Mann-Whitney tests.

Two multiple logistic regression analyses were conducted to examine the relationship between vaccination status, ICU admission, and intra-hospital death. In both cases, the relationship was adjusted for patient gender, age, and the number of comorbidities.

The results are expressed in odds ratio (OR) and 95% confidence interval. A significance threshold of 0.05 was used.

Statistical analyses were conducted using R (version 4.1.2).

3. Results

Overall, 360 patients were included in the study, mainly from ASST Santi Paolo e Carlo, which provided data from 192 patients (53%), while 126 (35%) and 42 (12%) patients were from Ospedale Policlinico San Martino of Genoa and IRCCS Ospedale Maggiore Policlinico of Milan, respectively. Patients' demographic characteristics and comorbidities at admission, stratified by vaccination status, are reported in Table 1.

Most of the patients were males (203, 56%), had a mean age of 68 (IQR: 51–80) years, and had at least one comorbidity (185, 51%). Overall, we observed a high prevalence of comorbidities such as diabetes (23%), chronic obstructive pulmonary diseases (18%), obesity (8.1%) and kidney disease (7.9%). By stratifying the study population for vaccination status, those vaccinated against COVID-19 had a higher frequency of suffering from at least one other disease (58% vs. 45%, $P = 0.011$), especially being on dialysis (3.4% vs. 0.0%, $P = 0.014$) and had a history of haematological malignancy (15% vs 4.7%, $P = 0.047$).

At admission, as presented in Table 2, patients generally presented fever (187, 52%), dyspnoea (147, 41%), and cough (140, 39%). Overall, the clinical presentation at admission was scored 5 (IQR: 4–5) using the WHO Ordinal Scale, and vaccinated patients were generally less affected by dyspnoea (35% vs 46%, $P = 0.030$).

Table 1
Demographic characteristics and comorbidities of enrolled patients at admission.

Baseline characteristics	N	Overall N = 360	Non-vaccinated N = 181	Vaccinated N = 179	p-value
Hospital	360				0.959
PC-MI		42 (12%)	22 (12%)	20 (11%)	
SM-GE		126 (35%)	63 (35%)	63 (35%)	
SPC-MI		192 (53%)	96 (53%)	96 (54%)	
Sex	360				0.989
Male		203 (56%)	102 (56%)	101 (56%)	
Female		157 (44%)	79 (44%)	78 (44%)	
Age (years)	360	68 (51, 80)	67 (51, 79)	68 (50, 81)	0.554
Diabetes	359				0.140
No		278 (77%)	146 (81%)	132 (74%)	
Yes		81 (23%)	35 (19%)	46 (26%)	
Unknown		1	0	1	
Obesity	309				0.321
No		284 (92%)	141 (90%)	143 (93%)	
Yes		25 (8.1%)	15 (9.6%)	10 (6.5%)	
Unknown		51	25	26	
COPD	359				0.451
No		295 (82%)	146 (81%)	149 (84%)	
Yes		64 (18%)	35 (19%)	29 (16%)	
Unknown		1	0	1	
Liver disease	359				> 0.999
No		354 (99%)	178 (98%)	176 (99%)	
Yes		5 (1.4%)	3 (1.7%)	2 (1.1%)	
Unknown		1	0	1	
Kidney disease	359				0.220
No		331 (92%)	170 (94%)	161 (90%)	
Yes		28 (7.8%)	11 (6.1%)	17 (9.6%)	
Unknown		1	0	1	
Dialysis	359				0.014
No		353 (98%)	181 (100%)	172 (97%)	
Yes		6 (1.7%)	0 (0%)	6 (3.4%)	
Unknown		1	0	1	
Haematological malignancy	130				0.047
No		117 (90%)	61 (95%)	56 (85%)	
Yes		13 (10%)	3 (4.7%)	10 (15%)	
Unknown		230	117	113	
Solid-organ transplant	322				0.086
No		305 (95%)	155 (97%)	150 (93%)	
Yes		17 (5.3%)	5 (3.1%)	12 (7.4%)	
Unknown		38	21	17	
Number of comorbidities*					0.011
None		175 (49%)	100 (55%)	75 (42%)	
More than 1		185 (51%)	81 (45%)	104 (58%)	

The clinical presentation at discharge, measured using the WHO ordinal scale, was 1 (IQR: 0–1), with no differences between groups (Table 3). Overall, 19 (5.3%) patients were admitted to ICU, and 69 (19%) died. Non-vaccinated individuals were more frequently admitted to ICU than those vaccinated (8.8% vs. 1.7%, $P = 0.002$). On the other hand, no difference was observed in intra-hospital mortality (19% vs 20%, $P = 0.853$).

Focusing on treatment (Table 4), 251 (70%) patients required respiratory support. Overall, 100 patients (60%) required HFNC. Pharmacological treatment was provided in 108 (30%) and 117 (32%) patients with remdesivir and mABs, respectively. No significant differences were observed among those vaccinated and non-vaccinated.

A small subset of patients was further investigated for their immunological profile and SARS-CoV-2 variant, as described in Table 5. The most common variant detected was the Delta (24/31, 77%), followed by the Omicron (4, 13%), with no difference based on patients' vaccination status. Conversely, the median of anti-S antibody titres at admission was 18 (IQR: 0–1702) U/mL, with those vaccinated with 6380 (IQR: 71–12,500) U/mL against 0 (IQR: 0–5) U/mL among non-vaccinated patients ($P = 0.001$).

In the multivariable logistic models, we observed no relationship between vaccination status and intra-hospital deaths (OR=1.051, 95%CI: 0.621–1.780, $P = 0.853$), also when adjusted for age, gender,

Table 2
Clinical presentation and characteristics of enrolled patients at admission.

Symptoms at presentation	N	Overall N = 360	Non-vaccinated N = 181	Vaccinated N = 179	p-value
Fever	360				0.836
No		173 (48%)	86 (48%)	87 (49%)	
Yes		187 (52%)	95 (52%)	92 (51%)	
Coryza	360				0.724
No		352 (98%)	176 (97%)	176 (98%)	
Yes		8 (2.2%)	5 (2.8%)	3 (1.7%)	
Cough	360				0.343
No		220 (61%)	115 (64%)	105 (59%)	
Yes		140 (39%)	66 (36%)	74 (41%)	
Dyspnoea	360				0.030
No		213 (59%)	97 (54%)	116 (65%)	
Yes		147 (41%)	84 (46%)	63 (35%)	
Loss of smell/taste	359				> 0.999
No		350 (97%)	175 (97%)	175 (98%)	
Yes		9 (2.5%)	5 (2.8%)	4 (2.2%)	
Unknown		1	1	0	
Gastrointestinal symptoms	360				0.704
No		322 (89%)	163 (90%)	159 (89%)	
Yes		38 (11%)	18 (9.9%)	20 (11%)	
WHO Ordinal Scale					
• Admission	168	5.00 (4.00, 5.00)	5.00 (4.00, 5.00)	5.00 (3.50, 5.00)	0.038
		192	96	96	

and the number of comorbidities (aOR= 0.996, 95%CI: 0.582–1.703, $P = 0.987$). Conversely, patients vaccinated were found to have a lower probability of being admitted to the ICU (OR=0.176, 95%CI: 0.040–0.539, $P = 0.007$) and remained so when controlled for confounders (aOR=0.172, 95%CI: 0.039–0.542, $P = 0.007$).

4. Discussion

This study investigated the differences in clinical outcomes between vaccinated and non-vaccinated individuals hospitalised for COVID-19. Based on real-world data, our findings revealed that vaccinated patients had a lower risk of being admitted to the ICU, whereas no relationship was found between vaccination status and intra-hospital death.

Firstly, vaccinated patients were more frequently affected by comorbidities notoriously linked to greater severity of SARS-CoV-2 infection, such as dialysis [10] and haematological malignancy [11]. Nonetheless, these patients presented significantly less dyspnoeic at admission and had reduced median WHO Ordinal Scales scores, potentially implying a lower risk of progression to a severe form of SARS-CoV-2 infection [9]. However, the differences mentioned above in clinical characteristics did not lead to a difference in treatments administered regarding respiratory support and pharmacological therapies.

Secondly, we found a significant difference in the number of patients admitted to ICU, with non-vaccinated patients requiring intensive care more often than their vaccinated counterparts. Differently, no difference in intra-hospital mortality was observed

between the two groups. These results are also confirmed by the multivariable logistic regressions, which showed that vaccination was significantly associated with decreased ICU admission risk but not intra-hospital mortality, as further supported by current evidence [8,12]. However, it is essential to note that the sample in this study was not fully balanced in terms of comorbidities. This imbalance is reasonably expected, as vaccinated individuals mostly had specific recommendations for the COVID-19 vaccine.

However, the increased likelihood of ICU admission for non-vaccinated individuals was often associated with their younger age and fewer comorbidities. As a result, these patients were more likely to be eligible for intensive care and had a higher chance of survival. This partly explains the lack of difference in intra-hospital mortality compared to the vaccinated individuals. A higher burden of comorbidities among vaccinated patients might explain their lower admission to the ICU [13]. Our findings suggest that the COVID-19 vaccine has a protective role in individuals with higher risk profiles, as defined by dyspnoea rates and WHO ordinal scale scores, especially regarding the need for ICU admission. This study adds to the growing body of evidence demonstrating the benefits of COVID-19 vaccination, even without booster vaccine doses, against a very aggressive COVID-19 variant such as Delta [4,14].

However, our study has several limitations. It has a retrospective design and a relatively small sample size. Additionally, Multi-IPV was performed during the Delta-dominant period, characterised by a virulent and aggressive variant and relatively low vaccine coverage in Italy. Most of the Italian population had received only 1 or 2

Table 3
Clinical presentation at discharge and outcomes.

Therapies	N	Overall N = 360	Non-vaccinated N = 181	Vaccinated N = 179	p-value
WHO Ordinal Scale					
• Discharge	167	1.0 (0.0, 1.0)	1.0 (0.0, 1.0)	1.0 (0.0, 1.0)	0.705
		193	96	97	
ICU admission	360				0.002
No		341 (95%)	165 (91%)	176 (98%)	
Yes		19 (5.3%)	16 (8.8%)	3 (1.7%)	
Death	360				0.853
No		291 (81%)	147 (81%)	144 (80%)	
Yes		69 (19%)	34 (19%)	35 (20%)	

Table 4
Respiratory support and pharmacological therapies administered to the enrolled patients.

Therapies	N	Overall N = 360	Non-vaccinated N = 181	Vaccinated N = 179	p-value
Respiratory support^s	358				0.611
No		107 (30%)	56 (31%)	51 (29%)	
Yes		251 (70%)	124 (69%)	127 (71%)	
Unknown		2	1	1	
HFNC	168				0.616
No		68 (40%)	36 (42%)	32 (39%)	
Yes		100 (60%)	49 (58%)	51 (61%)	
Unknown		192	96	96	
Venturi mask	167				0.367
No		162 (97%)	80 (95%)	82 (99%)	
Yes		5 (3.0%)	4 (4.8%)	1 (1.2%)	
Unknown		193	97	96	
CPAP	167				> 0.999
No		166 (99%)	84 (99%)	82 (100%)	
Yes		1 (0.6%)	1 (1.2%)	0 (0%)	
Unknown		193	96	97	
Remdesivir	360				0.765
No		252 (70%)	128 (71%)	124 (69%)	
Yes		108 (30%)	53 (29%)	55 (31%)	
Monoclonal Antibodies	360				0.681
No		243 (68%)	124 (69%)	119 (66%)	
Yes		117 (32%)	57 (31%)	60 (34%)	

Table 5
Immunological profile against SARS-CoV-2's Spike protein and SARS-CoV-2 variants of concern.

Variable	N	Overall N = 42	Non-vaccinated N = 22	Vaccinated N = 20	p-value
Anti-S antibodies (U/mL)	29	18 (0 - 1702)	0 (0 - 5)	6380 (71 - 12,500)	0.001
Missing	13	7	6	13	
VOCs	31				0.521
Beta		1 (3.2%)	0 (0%)	1 (7.1%)	
Delta		24 (77%)	13 (76%)	11 (79%)	
Gamma		2 (6.5%)	2 (12%)	0 (0%)	
Omicron		4 (13%)	2 (12%)	2 (14%)	
Unknown		11	5	6	

vaccine doses, if any. Therefore, our findings cannot be directly compared to more recent studies that evaluated the disease course of patients infected with the Omicron variant or subvariants who had already received booster doses of the COVID-19 vaccine. As we discussed earlier, considering the imbalance in terms of comorbidities between vaccinated and non-vaccinated individuals, the findings of this study may not be generalisable to populations with different comorbidity profiles. Lastly, data on SARS-CoV-2 variants and the prognostic role of anti-S antibody titres were unfortunately incomplete, preventing us from drawing any relevant conclusions.

However, we believe that our findings have important implications for public health and provide further support for vaccination efforts against the SARS-CoV-2 pandemic in Italy, where people who have received only two doses of vaccine are now estimated to be around 2 to 3 million [15]. Similarly, these findings might be relevant

Table 6
Crude odds ratio (OR) (95%CI) for the association with intra-hospital death and adjusted OR (aOR) (95%CI) for potential confounders of the association between intra-hospital death and COVID-19 vaccination status (n = 360).

Variables	Categories	OR (95%CI)	p-value	aOR (95%CI)*	p-value
COVID-19 vaccine status	Non-vaccinated	1	0.850	1	0.987
	Vaccinated	1.051 (0.621-1.780)		0.996 (0.582-1.703)	
Sex	Female	1	0.017	1	0.011
	Male	0.525 (0.307-0.889)		0.494 (0.285-0.847)	
Age (years)	continuous	1.002 (0.988-1.018)	0.749	1.004 (0.989-1.019)	0.646
Number of comorbidities	continuous	1.236 (0.969-1.564)	0.082	1.308 (1.014-1.679)	0.036

Acronyms: CI, confident interval; COVID-19, coronavirus disease 2019; OR, odds ratio.
* Adjusted for all other variables included in the table.

Table 7
Crude odds ratio (OR) (95%CI) for the association with ICU admission and adjusted OR (aOR) (95%CI) for potential confounders of the association between ICU admission and COVID-19 vaccination status (n = 360).

Variables	Categories	OR (95%CI)	p-value	aOR (95%CI)*	p-value
COVID-19 vaccine status	Non-vaccinated	1	0.006	1	0.007
	Vaccinated	0.176 (0.040-0.539)		0.172 (0.039-0.542)	
Sex	Female	1	0.010	1	0.010
	Male	7.083 (1.990-45.117)		7.165 (1.985-45.953)	
Age (years)	continuous	0.986 (0.960-1.011)	0.266	0.986 (0.955-1.018)	0.385
Number of comorbidities	continuous	0.956 (0.574-1.461)	0.849	0.953 (0.556-1.511)	0.847

Acronyms: CI, confident interval; COVID-19, coronavirus disease 2019; OR, odds ratio.
* Adjusted for all other variables included in the table.

in other European countries where the first-booster uptake rates are below 50% [16].

Our study calls for prospective data on patients with SARS-CoV-2 infection who were hospitalised despite having received the COVID-19 vaccination.

5. Conclusions

Although a complete COVID-19 vaccine course is recommended, it does not guarantee immunity against adverse outcomes. Therefore, it is strongly recommended, especially for those with underlying health conditions, to receive vaccinations and booster shots to prevent severe consequences from SARS-CoV-2.

Data Availability

Anonymised participant-level data will be made available upon requests directed to the corresponding author. Proposals will be reviewed and approved by the competing Ethical Committee and investigators based on scientific merit. After approval of a proposal, data can be shared through a secure online platform after signing a data access agreement.

Declaration of Competing Interest

All authors declare no competing interests.

Acknowledgements

None.

References

- [1] Deplanque D, Launay O. Efficacy of COVID-19 vaccines: from clinical trials to real life. *Therapie* 2021;76(4):277–83. <https://doi.org/10.1016/j.therap.2021.05.004>
- [2] Khoury J, Najjar-Debbiny R, Hanna A, et al. COVID-19 vaccine - long term immune decline and breakthrough infections. *Vaccine* 2021;39(48):6984–9. <https://doi.org/10.1016/j.vaccine.2021.10.038>
- [3] Yek C, Warner S, Wiltz JL, et al. Risk Factors for Severe COVID-19 outcomes among persons aged ≥18 years who completed a primary COVID-19 vaccination series - 465 health care facilities, United States, December 2020–October 2021. *MMWR Morb Mortal Wkly Rep* 2022;71(1):19–25. <https://doi.org/10.15585/mmwr.mm7101a4>
- [4] Havers FP, Pham H, Taylor CA, et al. COVID-19-associated hospitalisations among vaccinated and unvaccinated adults 18 years or older in 13 US States, January 2021 to April 2022. *JAMA Intern Med* 2022;182(10):1071–81. <https://doi.org/10.1001/jamainternmed.2022.4299>
- [5] Brosh-Nissimov T, Orenbuch-Harroch E, Chowers M, et al. BNT162b2 vaccine breakthrough: clinical characteristics of 152 fully vaccinated hospitalised COVID-19 patients in Israel. *Clin Microbiol Infect* 2021;27(11):1652–7. <https://doi.org/10.1016/j.cmi.2021.06.036>
- [6] Epaulard O, Abgrall S, Lefebvre M, et al. Impact of vaccination on the symptoms of hospitalised patients with SARS-CoV-2 Delta variant (B.1.617.1) infection. *Clin Microbiol Infect* 2022;28(12):1629–35. <https://doi.org/10.1016/j.cmi.2022.06.019>
- [7] Scortichini M, Schneider Dos Santos R, De' Donato F, et al. Excess mortality during the COVID-19 outbreak in Italy: a two-stage interrupted time-series analysis. *Int J Epidemiol* 2021;49(6):1909–17. <https://doi.org/10.1093/ije/dyaa169>
- [8] Colaneri M, Asperges E, Calia M, et al. Despite vaccination: a real-life experience of severe and life-threatening covid-19 in vaccinated and unvaccinated patients. *Vaccin (Basel)* 2022;10(9). <https://doi.org/10.3390/vaccines10091540>
- [9] WHO Working Group on the Clinical Characterisation and Management of COVID-19 infection. A minimal common outcome measure set for COVID-19 clinical research. *Lancet Infect Dis* 2020;20(8):e192–7. [https://doi.org/10.1016/S1473-3099\(20\)30483-7](https://doi.org/10.1016/S1473-3099(20)30483-7)
- [10] De Meester J, De Bacquer D, Naesens M, et al. Incidence, characteristics, and outcome of COVID-19 in adults on kidney replacement therapy: a regionwide registry study. *J Am Soc Nephrol* 2021;32(2):385–96. <https://doi.org/10.1681/ASN.2020060875>
- [11] Pagano L, Salmanton-García J, Marchesi F, et al. COVID-19 infection in adult patients with hematological malignancies: a European Hematology Association Survey (EPICOVIDEHA). *J Hematol Oncol* 2021;14(1):168. <https://doi.org/10.1186/s13045-021-01177-0>
- [12] Liu Q, Qin C, Liu M, Liu J. Effectiveness and safety of SARS-CoV-2 vaccine in real-world studies: a systematic review and meta-analysis. *Infect Dis Poverty* 2021;10(1):132. <https://doi.org/10.1186/s40249-021-00915-3>
- [13] Whittaker R, Bråthen Kristofferson A, Valcarcel Salamanca B, et al. Length of hospital stay and risk of intensive care admission and in-hospital death among COVID-19 patients in Norway: a register-based cohort study comparing patients fully vaccinated with an mRNA vaccine to unvaccinated patients. *Clin Microbiol Infect* 2022;28(6):871–8. <https://doi.org/10.1016/j.cmi.2022.01.033>
- [14] Havers FP, Pham H, Taylor CA, et al. COVID-19-associated hospitalisations among vaccinated and unvaccinated adults 18 years or older in 13 US States, January 2021 to April 2022. *JAMA Intern Med* 2022;182(10):1071–81. <https://doi.org/10.1001/jamainternmed.2022.4299>
- [15] Il vaccino anti covid in Italia in tempo reale | Il Sole 24 ORE.
- [16] Fan G, Song H, Yip S, Zhang T, He D. Impact of low vaccine coverage on the resurgence of COVID-19 in Central and Eastern Europe. *One Health* 2022;14:100402. <https://doi.org/10.1016/j.onehlt.2022.100402>