On the underestimation of dust mass in protoplanetary disks: Effects of disk structure and dust properties *

Yao Liu¹, Hendrik Linz², Min Fang¹, Thomas Henning², Sebastian Wolf³, Mario Flock², Giovanni P. Rosotti⁴, Hongchi Wang¹, and Dafa Li^{1,5}

¹ Purple Mountain Observatory & Key Laboratory for Radio Astronomy, Chinese Academy of Sciences, 10 Yuanhua Road, Nanjing 210023, China;

e-mail: yliu@pmo.ac.cn

² Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

³ Institut für Theoretische Physik und Astrophysik, Christian-Albrechts-Universität zu Kiel, Leibnizstr. 15, 24118 Kiel, Germany

⁴ DAMTP, University of Cambridge, CMS, Wilberforce Road, Cambridge CB3 0WA, UK

⁵ School of Astronomy and Space Science, University of Science and Technology of China, 96 Jinzhai Road, Hefei 230026, China

Preprint online version: October 17, 2022

ABSTRACT

The total amount of dust grains in protoplanetary disks is one of the key properties that characterize the potential for planet formation. With (sub-)millimeter flux measurements, literature studies usually derive the dust mass using an analytic form under the assumption of optically thin emission, which may lead to substantial underestimation. In this work, we conduct a parameter study with the goal of investigating the effects of disk structure and dust properties on the underestimation through self-consistent radiative transfer models. Different dust models, scattering modes and approaches for dust settling are considered and compared. The influences of disk substructures, such as rings and crescents, on the mass derivation are investigated as well. The results indicate that the traditional analytic method can underestimate the mass by a factor of a few to hundreds, depending on the optical depth along the line of sight set mainly by the true dust mass, disk size and inclination. As an application, we perform a detailed radiative transfer modeling of the spectral energy distribution of DoAr 33, one of the observed DSHARP disks. When the DSHARP dust opacities are adopted, the most probable dust mass returned from the Bayesian analysis is roughly 7 times higher than the value given by the analytic calculation. Our study demonstrates that estimating disk dust masses from radiative transfer modeling is one solution for alleviating the problem of insufficient mass for planet formation raised in the ALMA era.

Key words. circumstellar matter – planetary systems: protoplanetary disks – radiative transfer – facilities: Herschel

1. Introduction

Dust and gas material in protoplanetary disks are the building blocks of planetary systems. Within the time scale for disk dispersal, typically a few Myrs (Williams & Cieza 2011), micronsized dust grains coagulate and grow up by more than ten orders of magnitude until the assembly of large entities, such as pebbles and planetesimals (Testi et al. 2014; Birnstiel et al. 2016; Drazkowska et al. 2022). The gas-phase component either dissipates through a combination of complex mechanisms or can be captured by the growing giant planets (Balog et al. 2008; Machida et al. 2010; Ercolano & Pascucci 2017; Picogna et al. 2019; Pascucci et al. 2022). The total amount of material available in the disk plays a crucial role in determining whether planets can eventually form and what their nature is (e.g., Mordasini et al. 2012). The determination of protoplanetary disk masses can be accomplished through gas line or (sub-)millimeter continuum observations, though limitations exist for both diagnoses (Bergin & Williams 2017; Miotello et al. 2022).

Determining the disk gas mass is a difficult task, mainly because the most dominant component, H_2 , has no dipole moment, and it does not emit from the main disk mass reservoir. Carbon monoxide (CO) is an alternative tracer for gas masses. However, in addition to the requirement for a $CO-H_2$ abundance ratio appropriate for protoplanetary disk environments, mass determinations from CO (and its isotopologues) lines need to be corrected for the effects of CO freeze-out in the cold midplane, which may be more effective than previously thought (Powell et al. 2022), and CO photodissociation in the hot surface layers as well as the isotope-selective processes (Williams & Best 2014; Miotello et al. 2014, 2016). Hydrogen deuteride is believed to be a better probe, but to date its J = 1–0 rotational transition line has been detected for only three disks with the Herschel/PACS spectrometer (Bergin et al. 2013; McClure et al. 2016).

Measuring dust continuum emission is less expensive compared to gas line observations. With the advent of the Atacama Large Millimeter/submillimeter Array (ALMA), several nearby star formation regions have been surveyed, resulting in a large sample of disks (of the order of 1,000) with flux measurements at (sub-)millimeter wavelengths (e.g., Ansdell et al. 2016; Barenfeld et al. 2016; Pascucci et al. 2016; Cazzoletti et al. 2019; Grant et al. 2021). Under optically thin conditions, (sub-)millimeter flux densities F_v can be converted into dust masses via the analytic equation

$$M_{\rm dust.ana} = \frac{F_{\nu}D^2}{\kappa_{\nu}B_{\nu}(T_{\rm dust})},\tag{1}$$

where $B_{\nu}(T_{\text{dust}})$ refers to the Planck function given at the observed frequency ν and dust temperature T_{dust} , and the dis-

^{*} *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

tance to the object is denoted as *D*. At a reference frequency of v = 230 GHz (or wavelength $\lambda \sim 1.3 \text{ mm}$), a value of $2.3 \text{ cm}^2/\text{g}$ has been frequently adopted for the dust absorption opacity κ_v (e.g., Beckwith et al. 1990; Andrews et al. 2013). For the dust temperature, 20 K is a common choice. However, Andrews et al. (2013) suggested a stellar-luminosity (L_{\star}) dependent dust temperature $T_{\text{dust}} = 25 (L_{\star}/L_{\odot})^{0.25} \text{ K}$. The scaling of T_{dust} with L_{\star} was further investigated for disks around low-mass stars and brown dwarfs (e.g., Daemgen et al. 2016; van der Plas et al. 2016; Hendler et al. 2017), and generally the relation is found to be flatter than the prescription by Andrews et al. (2013).

A comparison between the statistic distribution of the analytic dust mass $M_{dust,ana}$ and the minimum-mass solar nebula shows that only few disks have enough material to produce our solar system or its counterparts in the exoplanet population (e.g., Najita & Kenyon 2014; Manara et al. 2018; Andrews 2020; Mulders et al. 2021). Explanations for such a discrepancy are generally provided by three different scenarios. The first scenario is that planet formation might be more efficient than what has been postulated, namely that in the protoplanetary disk evolutionary stage dust grains have already grown up to pebbles, planetesimals or even planetary embryos that are insensitive to (sub-)millimeter observations (e.g., Tychoniec et al. 2020). Supporting evidence is the prevalence of disk substructures, such as rings and gaps (e.g., ALMA Partnership et al. 2015; Long et al. 2018; Andrews et al. 2018a), even in protostellar disks with relatively young ages (Segura-Cox et al. 2020; Sheehan et al. 2020). These fine-scale features are likely created by (proto-)planets that are still embedded in their natal disk (Kley & Nelson 2012). In the second scenario, the disk acts as a conveyor belt that transports material from the environment to the central star (Manara et al. 2018). Therefore, the disk is replenished with material from the surrounding remnant envelope, part of which can contribute to the formation of planets. The third proposal concerns the reliability of mass estimation from Eq. 1 that is based on an optically thin assumption. Detailed radiative transfer analysis of millimeter data have demonstrated that protoplanetary disks, particularly in the inner regions, are not necessarily optically thin at millimeter wavelengths (e.g., Wolf et al. 2008; Pinte et al. 2016; Liu et al. 2019; Ueda et al. 2020). The measured fluxes only reflect dust material residing above the $\tau = 1$ surface. Consequently, the calculated $M_{dust,ana}$ is merely a lower limit. Indeed, such an underestimation has been found in previous studies, but mostly on limited number of objects (e.g., Maucó et al. 2018; Ballering & Eisner 2019; Carrasco-González et al. 2019; Ribas et al. 2020; Villenave et al. 2020; Macías et al. 2021; Guidi et al. 2022). Given the fact that the optical depth along the line of sight increases with the viewing angle, the analytic dust mass is expected to further deviate from the true dust mass of disks with high inclinations.

Moreover, deriving dust masses using Eq. 1 does not take into account the dust density distribution in the disk. The accumulating ALMA data show that in some cases, dust grains are quite concentrated within substructures like rings and crescents. Examples are the disks around Oph IRS 48 (van der Marel et al. 2013), HD143006 (Pérez et al. 2018), HD 135344B (Cazzoletti et al. 2018), and MWC 758 (Dong et al. 2018). Whether, and to which level, these configurations affect the mass estimation are not clear. In addition to a direct impact of dust opacities (therefore dust emissivities at millimeter wavelengths) on the mass determination, the choice for the dust model will also influence the resulting dust temperature, which in turn affects the mass determination. Such parameter coupling is not considered in Eq. 1. In this work, using self-consistent radiative transfer models, we

Table 1. Parameters of the fiducial model.

Parameter	Value	Note
$M_{\star} [M_{\odot}]$	0.5	stellar mass
$T_{\rm eff}$ [K]	4000	effective temperature
$L_{\star}[L_{\odot}]$	0.92	stellar luminosity
$R_{\rm in}$ [AU]	0.1	disk inner radius
R _{out} [AU]	100	disk outer radius
γ	1.0	surface density gradient
β	1.15	flaring index
$H_{100.a_{\min}}$ [AU]	10	reference scale height
$M_{\rm dust} \left[M_\odot \right]$	3×10^{-4}	disk dust mass
ξ	0.14	degree of dust settling
<i>i</i> [0]	41.8	disk inclination
$a_{\min} \left[\mu \mathbf{m} \right]$	0.01	minimum grain size
a_{\max} [mm]	1	maximum grain size
Dust model	DSHARP	Mie theory
Scattering mode	Isotropic	RADMC-3D setup

performed a detailed parameter study with the goal of systematically evaluating the extent of dust mass underestimation and its dependence on the disk structure and dust properties. The setup of the fiducial model and the method used to quantify the mass underestimation are described in Sect. 2. In Sect. 3, effects of various model parameters and assumptions on the results are investigated. As an application, we modelled the spectral energy distribution (SED) of the DoAr 33 disk in Sect. 4. The paper ends up with a summary in Sect. 5.

2. The fiducial disk model

In this section, we devise a fiducial radiative tranfer model, and introduce the method used to quantify the degree of underestimation of the dust mass. Starting from the fiducial model, we can alter the assumption, and therefore investigate the effects of various parameters on the result. The well-tested code RADMC-3D¹ is invoked to conduct the radiative transfer simulation (Dullemond et al. 2012). Table 1 gives an overview of parameters together with their adopted values used to describe the fiducial model.

2.1. Dust density distribution

We employ a flared disk surrounding a T Tauri star with stellar luminosity and temperature of $L_{\star} = 0.92 L_{\odot}$ and $T_{\rm eff} = 4000$ K, respectively. The disk is assumed to be passively heated by stellar irradiation. The dust density is assumed to follow a Gaussian vertical profile

$$\rho(R, z, a) = \frac{\Sigma(R, a)}{\sqrt{2\pi} h(R, a)} \exp\left[-\frac{1}{2} \left(\frac{z}{h(R, a)}\right)^2\right],\tag{2}$$

where R is the radial distance from the central star measured in the disk midplane. The surface density is described as a power law

$$\Sigma(R,a) = \Sigma_0(a) \left(\frac{R}{100 \,\text{AU}}\right)^{-\gamma},\tag{3}$$

with the proportionality factor $\Sigma_0(a)$ determined by normalizing the mass of dust grains in a certain grain size (a) bin. We set

¹ https://www.ita.uni-heidelberg.de/ dullemond/software/radmc-3d/.



Fig. 1. Vertical density profile of the fiducial model at a radius of R = 10 AU. The model includes 32 grains sizes logarithmically spaced from $a_{\min} = 0.01 \,\mu\text{m}$ to $a_{\max} = 1 \,\text{mm}$. The results are shown here for dust grains with three representative sizes, i.e. $0.01 \,\mu\text{m}$ (dotted line), $10 \,\mu\text{m}$ (dashed line) and 1 mm (solid line).

the disk inner radius R_{in} to 0.1 AU which is close to the dust sublimation radius. The disk outer radius R_{out} is chosen to be 100 AU, meaning that we truncate the disk at R_{out} .

Theoretical studies show that dust grains gradually decouple from the underlying gas distribution, and tend to settle towards the midplane (Dullemond & Dominik 2004). Such a process has been confirmed by high resolution observations, in which millimeter dust grains are found to be confined within scale heights of an order of a few AU (e.g., Gräfe et al. 2013; Pinte et al. 2016; Villenave et al. 2020) which is smaller than the typical gas pressure scale height (> 10 AU, Law et al. 2021; Rich et al. 2021). To account for dust settling, we assume that the dust scale height *h* varies with *R* and the grain size *a*

$$h(R,a) = H_{100}(a) \left(\frac{R}{100 \,\text{AU}}\right)^{\beta},\tag{4}$$

where β is the flaring index, and $H_{100}(a)$ stands for the scale height for a certain grain size at R = 100 AU. Using a power law, we parameterize the reference dust scale height as

$$H_{100}(a) = H_{100.a_{\min}} \left(\frac{a}{a_{\min}}\right)^{-\xi}.$$
 (5)

The parameter $H_{100,a_{min}}$ denotes the scale height of the smallest dust grains that are expected to be well coupled with the gas. The quantity ξ is used to characterize the degree of dust settling. This simple prescription of dust settling has been used for multi-wavelength modeling of protoplanetary disks (Pinte et al. 2008; Liu et al. 2017). We include 32 grain size bins that are logarithmically distributed from a fixed minimum grain size $a_{min} = 0.01 \,\mu\text{m}$ to a maximum grain size a_{max} . The radiative transfer is performed using these 32 grain sizes as 32 different dust species, each of which has its own dust opacity (see Sect. 2.2). Moreover, they are assumed to be thermally decoupled in RADMC-3D. For the fiducial model, we adopt $a_{max} = 1 \,\text{mm}$. The scale height for each of the 32 grain sizes is described by Eq. 4 and 5. Given $\xi = 0.14$ and $H_{100,a_{min}} = 10 \,\text{AU}$ for the fiducial model, the scale height of the 1 mm dust grain equals to 2 AU, which is consistent with observational constraints (Pinte et al.



Fig. 2. Mass absorption coefficient κ_v at $\lambda = 1.3$ mm as a function of grain size *a*. The blue line shows the result by using the DSHARP dust model, whereas the case for the DIANA dust model is indicated with the black line. The horizontal dashed line marks the value of $2.3 \text{ cm}^2/\text{g}$ that is commonly adopted to derive the analytic dust mass $M_{\text{dust,ana}}$ in the literature.

2016; Villenave et al. 2020; Liu et al. 2022). When the dust and gas are perfectly mixed, $\xi = 0$, and $H_{100}(a) = H_{100.a_{\min}}$ becomes independent of the grain size and is equal to the gas scale height, the disk turns to be the well-mixed case.

The fraction of dust mass in each grain size bin is calculated via

$$f(a_j) = \frac{\int_{a_0}^{a_1} \frac{4\pi}{3} \rho_{\text{grain}} n(a) a^3 \, \mathrm{d}a}{\int_{a_{\text{lower}}}^{a_{\text{upper}}} \frac{4\pi}{3} \rho_{\text{grain}} n(a) a^3 \, \mathrm{d}a},\tag{6}$$

where $a_0 = \exp \left[\ln(a_j) - \Delta/2 \right]$, $a_1 = \exp \left[\ln(a_j) + \Delta/2 \right]$, $a_{\text{lower}} = \exp \left[\ln(a_{\min}) - \Delta/2 \right]$, $a_{\text{upper}} = \exp \left[\ln(a_{\max}) + \Delta/2 \right]$, and Δ is the step width in the logarithmic scale used to sample the 32 grain sizes. The vertically integrated grain size distribution is assumed to be $n(a) \propto a^{-3.5}$, and ρ_{grain} is the bulk density of the dust ensemble. Therefore, once the total dust mass (M_{dust}) in the entire disk is given, we know the dust mass for each grain size bin, therefore can derive the proportionality factor $\Sigma_0(a)$ in Eq. 3. Figure 1 shows the vertical density profile of the fiducial model at a radius of R = 10 AU for three representative grain sizes.

2.2. Dust properties

For the dust composition, we directly make use of the model by Birnstiel et al. (2018), which is composed of water ice, astronomical silicates, troilite, and refractory organic material, with volume fractions being 36%, 17%, 3% and 44%, respectively. The bulk density of the dust mixture is $\rho_{\text{grain}} = 1.675 \text{ g cm}^{-3}$. This dust model was used for the interpretation of the ALMA data from the Disk Substructures at High Angular Resolution Project (DSHARP, Andrews et al. 2018a). We use the Mie theory to calculate the mass absorption/scattering coefficients for the sampled 32 grain sizes. The blue solid line in Figure 2 shows the absorption coefficient at $\lambda = 1.3 \text{ mm}$ as a function of grain size. There are clear fluctuations for $a \sim 0.2 - 0.7 \text{ mm}$ due to the resonances when the size of the particle is comparable with $\lambda/2\pi$.



Fig. 3. Upper panel: 2D density distribution of the fiducial model with parameters given in Table 1. The blue and black lines draw the contour of $\tau_{1.3 \text{ mm}} = 1$ when the disk is viewed at $i = 41.8^{\circ}$, and includes the DSHARP and DIANA dust opacities, respectively. *Bottom panel:* SEDs of the fiducial model when the DSHARP (blue line) or DIANA (black line) opacities are adopted in the simulation. The dashed curve stands for the input photospheric spectrum with $T_{\text{eff}} = 4000 \text{ K}$, assuming $\log g = 3.5$ and solar metallicity (Kurucz 1994).

Dust scattering is expected to significantly affect the emission level and millimeter spectral index of optically thick disks (Zhu et al. 2019; Liu 2019). Consequently, whether or not this mechanism is taken into account will influence the inferred dust temperature, optical depth, dust mass and grain size, see Appendix C of Carrasco-González et al. (2019) for instance. We switch on isotropic scattering in both the thermal Monte Carlo simulation for computing the dust temperature and the subsequent step for SED generation. A more realistic treatment for dust scattering is considered and compared in Sect. 3.2.

Multi-wavelength observations have shown that millimeter spectral slopes differ among disks even in the same star formation region (e.g., Ricci et al. 2010; Pinilla et al. 2017; Andrews 2020). The scatter of the observed millimeter spectral slopes can be attributed to different levels of dust processing or to different optical depths. Without a detailed analysis of multi-wavelength data, the grain size distribution is not known, which could have a direct impact on the mass determination with Eq. 1. Moreover, the composition of dust grains is also found to vary for different disks (e.g., Juhász et al. 2010). Therefore, we also consider another dust model that is introduced in the DiscAnalysis project (DIANA, Woitke et al. 2016), and investigate the influence of a different choice for dust properties on the result, see Sect. 3.4.

2.3. Properties of the fiducial model

The upper panel of Figure 3 shows the dust density distribution of the fiducial model. The blue curve indicates the location of the $\tau_{1.3 \text{ mm}} = 1$ surface considering that the disk is viewed at $i = 41.8^{\circ}$. Within $R \leq 40$ AU, disk interior layers are optically thick. We run the thermal Monte Carlo simulation using 3×10^7 photons in order to obtain a smooth temperature structure. The SED of the fiducial model is simulated and scaled to the distance of 140 pc, which is indicated as the blue line in the bottom panel of Figure 3.

The midplane dust temperature ranges from ~1300 K to ~10 K, roughly following a power law $T_{\text{midplane}} \propto R^{-0.5}$. To give a representative dust temperature characterizing millimeter continuum emission, we calculate the mass-averaged dust temperature via

$$T_{\rm dust} = \frac{1}{M_{\rm dust}} \sum_{i=1}^{N_{\rm cells}} \sum_{j=1}^{N=32} m_{\rm dust}(i,j) t_{\rm dust}(i,j),$$
(7)

where $m_{dust}(i, j)$ and $t_{dust}(i, j)$ are the dust mass and temperature in each cell (*i*) for each grain size (*j*), respectively. Similarly, we derive the mass-averaged dust opacity from

$$\overline{\kappa_{\nu}} = \frac{1}{M_{\text{dust}}} \sum_{i=1}^{N_{\text{cells}}} \sum_{j=1}^{N=32} m_{\text{dust}}(i,j) \,\kappa_{\nu}(i,j),\tag{8}$$

where $\kappa_{\nu}(i, j)$ are the dust opacity in each cell for each grain size. The fiducial model features a $T_{\text{dust}} = 16.3 \text{ K}$ (see Table 2) and $\overline{\kappa_{1.3\text{mm}}} = 1.9 \text{ cm}^2/\text{g}$.

The simulated flux density at $\lambda = 1.3$ mm is $F_{\nu} = 80.7$ mJy. Because F_{ν} , T_{dust} and $\overline{\kappa_{\nu}}$ of the fiducial model are exactly known, we can use Eq. 1 to derive a corresponding analytic dust mass required to produce the same amount of continuum in the optically thin regime. This gives us $M_{dust,ana} = 2.2 \times 10^{-4} M_{\odot}$. The true dust mass imported into the fiducial model is $M_{dust} = 3 \times 10^{-4} M_{\odot}$. We define the degree of mass underestimation as

$$\Lambda_{1.3\text{mm}} = \frac{M_{\text{dust}}}{M_{\text{dust,ana}}} = 1.4,$$
(9)

where the subscript is used to state that the calculation is based on 1.3 mm flux densities. Similar factors can be derived using data points at other wavelengths. The difference between M_{dust} and $M_{dust,ana}$ basically reflects dust material hidden below the $\tau = 1$ surface (see Figure 3), because uncertainties on the dust temperature and opacity are eliminated in our definition.

Effects of model parameters and assumptions on the results

3.1. Effects of model parameters

With the methodology used to quantify the underestimation factor outlined in Sect. 2.3, we then change the values for each model parameter to investigate their effects on the results. Parameters considered in the exploration are $R_{\rm in}$, $R_{\rm out}$, γ , $M_{\rm dust}$, $H_{100.a_{\rm min}}$, β , $H_{100}(1$ mm dust) or equivalent to ξ , $a_{\rm max}$, and *i*. Calculating the underestimation factor always utilizes the mass-averaged dust opacity and temperature of the corresponding model. Figure 4 shows the flux densities at 0.88 mm and 1.3 mm, mass-averaged temperature and underestimation factors.

As can be seen, although the stellar properties are fixed, there is a broad range for the mass-averaged temperature.



Fig. 4. The effect of model parameters on the flux densities at 0.88 and 1.3 mm ($F_{0.88\text{mm}}$, $F_{1.3\text{mm}}$, top row on the left Y axis), massaveraged dust temperature (T_{dust} , top row on the right Y axis), and underestimation factors ($\Lambda_{0.88\text{mm}}$, $\Lambda_{1.3\text{mm}}$, bottom row). When exploring one particular parameter, all the rest parameters are fixed to their fiducial values listed in Table 1. The red dashed lines show the underestimation factors $\Lambda_{1.3\text{mm}}$ that are calculated by assuming a constant $T_{\text{dust}} = 25 (L_{\star}/L_{\odot})^{0.25} = 24.5 \text{ K}$ and $\kappa_{1.3\text{mm}} = 2.3 \text{ cm}^2/\text{g}$.

Particularly, the variation in T_{dust} with R_{out} can be up to a factor of 4, from ~ 32 K for compact disks with $R_{out} = 20 \text{ AU}$ to ~ 8 K for large disks with $R_{out} = 800 \text{ AU}$. Of the nine parameters, only R_{in} , a_{amax} and *i* have minor impacts on T_{dust} .

Therefore, a simple presumption on the temperature according to $T_{\text{dust}} = 25 (L_{\star}/L_{\odot})^{0.25} = 24.5 \text{ K}$ will lead to differences in the determination of $M_{\text{dust.ana}}$.

 Table 2. Comparison of results between models with isotropic and anisotropic scattering.

$M_{\rm dust} \ [M_{\odot}]$	i [0]	<i>F</i> _{1.3 mm} [mJy]	T _{dust} [K]	$\Lambda_{1.3mm}$
3×10^{-5}	41.8	12.2 (12.6)	17.5 (18.0)	1.00 (1.00)
3×10^{-4}	41.8	80.7 (86.7)	16.3 (16.9)	1.38 (1.35)
3×10^{-3}	41.8	168.7 (195.5)	15.7 (16.2)	6.25 (5.62)
3×10^{-4}	8.1	101.3 (105.9)	16.3 (16.9)	1.10 (1.10)
3×10^{-4}	63.1	51.5 (57.0)	16.3 (16.9)	2.16 (2.04)
3×10^{-4}	79.3	21.6 (24.0)	16.3 (16.9)	5.16 (4.86)

Notes. Numbers given outside parentheses refer to the result when isotropic scattering is assumed, while the ones in the parentheses are the case when full scattering matrices are taken into account.

The underestimation factor Λ ranges from a few to hundreds. Higher dust masses, inclinations, or smaller disk sizes result in the greatest underestimation in the dust mass, because they work together to achieve a higher optical depth along the line of sight. Other parameters that aggravate the underestimation include smaller R_{in} , steeper γ , lower β and a_{amax} that yields a higher κ . As the optical depth decreases with increasing wavelength, underestimations calculated based on 1.3 mm fluxes are systematically lower than those derived using the 0.88 mm fluxes. Therefore, long wavelength observations are required to better probe the dust mass in protoplanetary disks. The red dashed lines in Figure 4 show the underestimation factors $\Lambda_{1.3mm}$ that are calculated by assuming a constant $\kappa_{1.3\text{mm}} = 2.3 \text{ cm}^2/\text{g}$ and $T_{\text{dust}} = 25 (L_{\star}/L_{\odot})^{0.25} = 24.5 \text{ K}$ (hotter than most of the models). As can be seen, they are generally larger than those obtained by using the mass-averaged dust opacity and temperature of the models, directly demonstrating the importance of taking disk structure and dust properties into account in the task of mass estimation.

Figure A.1 and A.2 show how Λ varies with different model parameters when the disk is viewed at two extreme inclinations, i.e., $i = 0^{\circ}$ (face on) and $i = 90^{\circ}$ (edge-on). The behavior of faceon disks is quite similar to the fiducial inclined disks. For edgeon disks, the underestimation factor becomes more sensitive to most of the explored parameters. This tendency can be clearly identified from the $\Lambda - R_{out}$, $\Lambda - M_{dust}$, $\Lambda - H_{100}(1 \text{ mm dust})$ and $\Lambda - a_{max}$ profiles. As a consequence, mass determination for edge-on disks suffers more difficulties than for disks with low inclinations, because the accuracy of the result highly depends on how well many other parameters are constrained.

3.2. Effects of dust scattering mode

When dust grains grow up to millimeter sizes, scattering coefficients at millimeter wavelengths become significantly larger than absorption coefficients, resulting in a very high albedo. For instance, the albedo of the fiducial dust model is 0.9 at 1.3 mm. In this case, scattering of thermal reemission radiation on the SED should be taken into account. Theoretical studies have found that dust scattering can considerably reduce the emission from optically thick regions, with the reduction level depending on the inclination (e.g., Zhu et al. 2019; Sierra & Lizano 2020). Thus, an optically thick disk with scattering can be misidentified as an optically thin disk if scattering is ignored.

We compared the models when different modes of dust scattering are turned on. The results are summarized in Table 2. The difference in dust temperature between isotropic and anisotropic



Fig. 5. Scale height of 1 mm dust as a function of *R*. The blue line shows the fiducial model with dust settling prescribed by Eq. 5. The black and red lines stand for models when dust settling is parameterized according to Eq. 10 under different turbulence strengths α_{turb} .

 Table 3. Comparison of models using different prescriptions for dust settling.

Settling approach	ξ or α_{turb}	$F_{1.3 \text{ mm}}$	$T_{\rm dust}$	$\Lambda_{1.3mm}$
		[Jy]	[K]	
Fiducial model	$\xi = 0.14$	80.7	16.3	1.38
Well-mixed model	$\xi = 0$	124.4	22.2	1.34
Settling by Eq. 10	$\alpha_{turb} = 3e - 3$	108.5	20.6	1.40
	$\alpha_{turb} = 3e - 4$	86.6	18.1	1.47

Notes. Other parameters of these models are fixed to the fiducial values.

scattering models is merely ~ 0.5 K, regardless of the total dust mass (hence, intrinsic optical thickness). Anisotropic scattering always yields stronger emission. However, the flux discrepancy between the two modes does not exceed 16%, with larger differences generally for more massive or more inclined disks. Therefore, once dust scattering is included, this does not have a significant impact on the underestimation factor.

3.3. Effects of dust settling approach

We also consider different approaches for dust settling that basically determines the millimeter dust scale height. In the fiducial model, we use only three free parameters ($H_{100.a_{min}}$, ξ and β) to parameterize the scale height, see Eq. 4 and 5. The scale height of 1 mm dust grain is indicated as the blue line in Figure 5. Theoretically, the vertical distribution of dust grains is a consequence of equilibrium between dust settling and vertical stirring induced by turbulent motions. Therefore, the scale heights of dust grains with different sizes are related to the turbulence strength α_{turb}

$$h(R,a) = H_{\text{gas}} \left(1 + \frac{\text{St}}{\alpha_{\text{turb}}} \frac{1 + 2 \,\text{St}}{1 + \text{St}} \right)^{-1/2},\tag{10}$$

where St = $\frac{\pi}{2} \frac{\rho_{\text{grain}} a}{\Sigma_{\text{gas}(R)}}$ is the Stokes number (e.g., Dubrulle et al. 1995; Schräpler & Henning 2004; Woitke et al. 2016; Birnstiel et al. 2016). Since the smallest dust grain (0.01 μ m in our case)



Fig. 6. A comparison of $\Lambda_{1.3mm}$ (squares) and T_{dust} (filled circles) between models that use the DSHARP (blue symbols) and DIANA (black symbols) dust opacities in the simulation, respectively. Details can be found in Sect. 3.4.

is expected to be well-mixed with the gas, we assume that the gas pressure scale height H_{gas} follows

$$H_{\rm gas} = H_{100.a_{\rm min}} \left(\frac{R}{100\,\rm AU}\right)^{\beta}.$$
 (11)

This settling approach has four free parameters: $H_{100.a_{\min}}$, α_{turb} , β and gas surface density Σ_{gas} . For Σ_{gas} , we simply scaled the dust surface density by a constant gas-to-dust mass ratio of 100. For α_{turb} , we consider 3×10^{-4} and 3×10^{-3} that are consistent with observational constraints from gas line observations (e.g., Flaherty et al. 2017; Teague et al. 2018; Flaherty et al. 2020). The α_{turb} /St ratios for the 0.2 mm dust grain calculated at $R \sim 80$ AU are ~ 0.03 and ~ 0.3 for $\alpha_{turb} = 3 \times 10^{-4}$ and 3×10^{-3} , respectively. These ratios are comparable with the results derived from analyzing the widths of dust rings in several DSHARP disks (Dullemond et al. 2018). The dust scale heights are shown with the black and red lines in Figure 5. The model with $\alpha_{turb} = 3 \times 10^{-3}$ features a larger scale height than the fiducial setup across the entire radius. Consequently, it has a higher temperature and emits more continuum (see Table 3). A weaker turbulence of 3×10^{-4} makes the disk cooler and fainter, and now T_{dust} and $F_{1.3 \text{ mm}}$ are comparable with those of the fiducial model.

In Table 3, we also provide the results when dust grains and gas are well coupled, which can be considered as a limiting case of the fiducial settled disk. When the settling degree ξ (see Eq. 5) tends towards zero, the settled disk model turns into the well-mixed case, and becomes the brightest at millimeter wavelengths among the models under comparison. It is naturally understood that using well-mixed radiative transfer models to fit observed SEDs, one would need the least amount of dust grains. Hence, dust mass determinations from radiative transfer analysis based on the well-mixed assumption are probably still lower limits.

3.4. Effects of dust properties

To investigate how the dust properties affect the results, we make use of the standard dust properties introduced by the DIANA project (Woitke et al. 2016). The dust grains consist of 60% silicate ($Mg_{0.7}Fe_{0.3}SiO_3$, Dorschner et al. 1995), 15% amorphous carbon (BE–sample, Zubko et al. 1996), and 25% porosity. These percentages are volume fractions that are used to derive

Table 4. Ring and crescent parameters.

Parameter	1st ring	2nd ring	3rd ring	Crescent
$R_{\rm ring}$ or $R_{\rm cres}$ [AU]	10	35	65	70
$\sigma_{\rm ring}$ or $\sigma_{R_{\rm cres}}$ [AU]	5	5	5	5
$\theta_{\rm cres}$ [0]	-	-	-	45
$\sigma_{\theta_{ m cres}}$ [0]	-	-	-	20

the effective refractory indices by applying the Bruggeman mixing rule (Bruggeman 1935). Porous grains are considered because porosity, and its evolution during the collisional growth, helps to overcome the radial-drift barrier (e.g., Ormel et al. 2007; Garcia & Gonzalez 2020). We assumed a distribution of hollow spheres with a maximum hollow volume ratio of 0.8, and calculated the opacities using the OpacityTool package (Toon & Ackerman 1981; Min et al. 2005).

The black line in Figure 2 shows $\kappa_{1.3 \text{ mm}}$ as a function of grain size. The DIANA dust absorption coefficients are systematically larger than the DSHARP values. The choice for the material termed "organic" in the DSHARP model has a significant impact on the absorption coefficient, see Appendix B in Birnstiel et al. (2018). When using the amorphous carbon (BE-sample, Zubko et al. 1996) instead, the absorption coefficients increase by a factor of ~5. The SEDs using different dust opacities are compared in the lower panel of Figure 3, with $F_{1.3mm} = 165.2 \text{ mJy}$ and 80.7 mJy for the DIANA and DSHARP dust being imported, respectively. Models using the DIANA opacity are generally hotter, but the differences in the mass-averaged temperature are within ~ 2 K, even for very massive disks, see Figure 6. Given the disk is more optically thick when adopting the DIANA dust model, correspondingly, the underestimation factors are systematically higher, as shown in Figure 6.

3.5. Effects of the presence of disk substructures

In the above modeling procedure, dust surface densities are assumed to follow a smooth power law, without sharp density fluctuations, see Eq. 3. However, high resolution multi-wavelength observations have revealed that many protoplanetary disks possess rings and crescents (e.g., van Boekel et al. 2017; Avenhaus et al. 2018; Fedele et al. 2018; Long et al. 2018; Andrews et al. 2018a; Cieza et al. 2021), which are believed to be the favorable place for planet formation (Andrews 2020). In some cases, these substructures, even located at large radial distances from the host central star, appear very bright in the millimeter continuum, indicating that a large amount of dust grains are concentrated (e.g., ALMA Partnership et al. 2015; Pérez et al. 2018; Dong et al. 2018; Cazzoletti et al. 2018).

The overdensities in disk substructures will increase the optical depth locally, which in principle can affect the mass determination. Detailed radiative transfer analysis confirms that some substructures are optically thick at ALMA wavelengths (e.g., Pinte et al. 2016; Liu et al. 2017; Ohashi & Kataoka 2019; Sierra et al. 2021; Liu et al. 2022). In order to investigate how the redistribition of dust grains in disks with substructures affect the mass determination, we consider four scenarios: one ring, two rings, three rings, and three rings plus a crescent. The surface densities of the substructures are defined as

$$\Sigma_{\rm ring} = \Sigma_0 \exp^{-(R - R_{\rm ring})^2 / 2\sigma_{\rm ring}^2},\tag{12}$$

$$\Sigma_{\text{crescent}} = 3\Sigma_0 \exp^{-(R-R_{\text{cres}})^2/2\sigma_{R_{\text{cres}}}^2} \exp^{-(\theta-\theta_{\text{cres}})^2/2\sigma_{\theta_{\text{cres}}}^2},$$
(13)



Fig. 7. Effects of disk substructures on the mass underestimation. *Left column:* representative 1.3 mm dust continuum image for each scenario of disk substructures. The raw radiative transfer images are convolved with a Gaussian beam of 35 mas in size (~ 5 AU at the distance of 140 pc, indicated with a white filled circle in the lower left corner of each image). The disk is seen face-on merely for a better presentation of the disk configuration. *Middle column:* 1.3 mm flux density ($F_{1.3mm}$) and mass-averaged temperature (T_{dust}) as a function of total dust mass (M_{dust}). When simulating the SED, the fiducial inclination of 41.8° is adopted. *Right column:* underestimation factor as a function of M_{dust} . The green filled circles refer to the underestimation factors for smooth disks that have an equivalent R_{out} to that of the disks with substructures, see Sect. 3.5.

where R_{ring} or R_{cres} refers to the radial location of the feature, and σ_{ring} or $\sigma_{R_{\text{cres}}}$ stands for the radial width. For the crescent, θ_{cres} and $\sigma_{\theta_{\text{cres}}}$ are used to control the location and width in the azimuthal direction. The proportionality constant is set to be 3 times larger than that for the rings, under which the crescent can feature a high millimeter flux contrast to its neighboring ring. Table 4 lists the adopted parameter values. These values are chosen to be representative among the results that are derived by analyzing the brightness profiles of rings/crescents revealed by ALMA (e.g., Long et al. 2018; Huang et al. 2018; Pérez et al. 2018). If all the features are included, integrating Eq. 12 and 13 results in the mass fractions of 8%, 28%, 50%, and 14% for the 1st, 2nd, 3rd ring and crescent, respectively.

We gradually increase M_{dust} , therefore to let the substructures turn to be optically thick. Dust opacities and settling approach are kept to the fiducial assumptions. The results for each scenario are presented in Figure 7. We found that the tendencies of $F_{1.3mm}$, T_{dust} , and the overall level of $\Lambda_{1.3mm}$ are generally similar to those for smooth disks. The maximum underestimation is achieved by only introducing the innermost ring. In this case, dust grains are confined within the smallest disk radius. When the outer substructures are added, the underestimation becomes less severe. Such an effect is mainly driven by the actual R_{out} . In other words, R_{out} has a stronger influence on the mass underestimation than the presence/absence of substructures. In order to testify the hypothesis, we run additional simulations for smooth disks. The surface densities for $R \le R_f$ are all fixed to the surface density at $R = R_f$, where R_f is the location of the outermost substructure (e.g., $R_f = 65 \text{ AU}$ for the case of three rings, see Table 4), and for $R > R_f$ we set the surface densities according to Eq. 12. In the case of three rings plus a crescent, we set the surface densities for $R > R_f$ according to an azimuthally averaged profile described by a combination of Eq. 12 and Eq. 13. Under such a circumstance, the dust surface density does not vary within R_f (i.e., no rings/gaps/crescent), and follows a same profile to that of the disk with substructures in the outermost region, which would lead to an equivalent R_{out} . The green filled circles shown in the right column of Figure 7 refer to the underestimation factors, which are similar to those of the disks with substructures.

4. Application to the DoAr 33 disk

We have shown that due to the optical depth effect, calculating disk dust masses using the analytic approach can lead to substantial underestimation, even if the dust temperature (T_{dust}) and opacity (κ) are known. The disk outer radius (R_{out}), inclination (*i*) and true dust mass (M_{dust}) are most important to create such optically thick regions. Fortunately, R_{out} and *i* can be derived from high resolution (sub-)millimeter images that are nowadays available for a large number of disks thanks to ALMA. In this section, we conduct a detailed radiative transfer modeling of the SED for DoAr 33 in order to constrain the total dust mass in the disk, and then compare the result with the analytic dust mass.

4.1. DoAr 33 and its SED

DoAr 33 is a T Tauri star located in the Ophiuchus star formation region at a distance of 140 pc (Gaia Collaboration et al. 2018). The spectral type K5.5 derived by Wilking et al. (2005) translates to an effective temperature of 4160 K using the SpT - T_{eff} conversion reported by Herczeg & Hillenbrand (2014). To compile the SED, on one hand we collected data points at various wavelengths from the literature. On the other hand, through the Chinese Telescope Access Program, we conducted new CSO/SHARC II observations at $350 \,\mu\text{m}$, see Sect. B in the Appendix for details about the observations and data reduction. Herschel photometry for DoAr 33 in the literature show a surprisingly steep decline toward submillimeter wavelengths (Rebollido et al. 2015). Therefore, we re-reduced the Herschel/PACS and SPIRE data, and performed the photometry based on a point-spread function fitting, see Sect. C. These efforts allow us to build the SED with an excellent wavelength coverage. The measurements from optical to millimeter domains are summarized in Table 5, and shown with red dots in panel (a) of Figure 8. Fitting the optical photometry with the Kurucz atmosphere models yields a stellar luminosity of $L_{\star} = 0.97 L_{\odot}$.

4.2. A large grid of radiative transfer model SEDs

As one of the 20 disks selected in the DSHARP project, DoAr 33 was observed by ALMA with an angular resolution of ~ 35 mas (Andrews et al. 2018a). The disk is inclined by $i = 41.8^{\circ}$, and appears smooth in the 1.3 mm continuum image. Huang et al. (2018) defined the outer boundary of the millimeter disk to be the radius at which the enclosed flux is equal to 95% of the to-

Table 5. The photometry of DoAr 33.

λ [μm]	F_{ν} [mJy]	Instrument / Filter	Reference
0.505	4.5 ± 0.039	GAIA/bp	1
0.623	16.1 ± 0.039	GAIA/g	1
0.773	38.8 ± 0.239	GAIA/rp	1
0.64	15.5 ± 1.548	R	2
0.79	37.7 ± 3.765	Ι	2
1.235	175 ± 3.88	2MASS/J	3
1.662	332 ± 11	2MASS/H	3
2.159	348 ± 5.5	2MASS/K	3
3.55	212 ± 15.3	Spitzer/IRAC 1	3
4.493	169 ± 12.3	Spitzer/IRAC 2	3
5.731	193 ± 11.9	Spitzer/IRAC 3	3
7.872	222 ± 11.7	Spitzer/IRAC 4	3
12	200.4 ± 3.8	WISE 3	4
22	225 ± 9.5	WISE 4	4
23.68	211 ± 19.5	Spitzer/MIPS 1	3
70	469 ± 33	Herschel/PACS	this work
100	537 ± 38	Herschel/PACS	this work
160	539 ± 38	Herschel/PACS	this work
250	416 ± 23	Herschel/SPIRE	this work
350	260 ± 14	Herschel/SPIRE	this work
350	212 ± 21.2	CSO/SHARC II	this work
850	79 ± 7	JCMT	5
880	80 ± 2	SMA	6
870	76.4 ± 1.4	ALMA	7
1254	35 ± 0.017	ALMA	8
1300	33.6 ± 0.47	ALMA	3
3300	3.7 ± 0.3	ATCA	9

Notes. References: (1) Gaia Collaboration et al. (2018); (2) Wilking et al. (2005); (3) Cieza et al. (2019); (4) Cutri & et al. (2013); (5) Andrews & Williams (2007); (6) Andrews et al. (2010); (7) Cox et al. (2017); (8) Andrews et al. (2018a); (9) Ricci et al. (2010)

tal flux. That is 27 AU, making DoAr 33 to be the smallest disk among the 18 single-star disk systems targeted by DSHARP. We checked the ¹²CO J = 2-1 line data, and found that the emission of ¹²CO is more extended, with a 95% enclosing flux radius of 70 AU. A more compact millimeter dust disk than the CO disk has also been found for other targets (e.g., Ansdell et al. 2018; Long et al. 2022), and can be naturally explained by radial drift of dust particles (Trapman et al. 2019; Toci et al. 2021).

We built a large grid of radiative transfer models for DoAr 33. The model configuration is a slight variant of the fiducial setup. Dust grains with sizes ranging from $a_{\min} = 0.01 \,\mu\text{m}$ to $10\,\mu\text{m}$ are assumed to be distributed from 0.1 AU to 70 AU. Larger dust grains (i.e., $a > 10 \,\mu$ m) have already drifted to the location where the DSHARP continuum data probes, and therefore we truncated them at R = 27 AU. For the most interesting parameter M_{dust} , we sampled 13 grid points that are logarithmically distributed from 3×10^{-6} to $3 \times 10^{-3} M_{\odot}$. For a_{max} , we took 7 logarithmically spaced points from 0.1 mm to 100 mm. There are 8 points for $H_{100.a_{\min}}$ linearly spaced between 4 and 18 AU, and 9 linear points for β from 1.05 to 1.25. For the power-law index (γ) of the surface density, we chose a value of 0.5 which is close to the slope of a power law used to fit the azimuthallyaveraged surface brightness of the DSHARP image. Parameters not mentioned here were fixed to the fiducial values, see Table 1. In total, there are 6,552 models in the grid, and we separately ran the simulation with the DSHARP and DIANA dust opacities.



Fig. 8. *Panel (a):* SEDs of DoAr 33. The best-fit radiative transfer models using the DSHARP and DIANA opacities are shown with blue and black solid lines, respectively. The dashed line refers to the input photospheric spectrum, while observational data points are overlaid with red dots. *Panels (b)–(e):* Bayesian probability distributions for $\text{Log}_{10}(M_{\text{dust}}/M_{\odot})$, $\text{Log}_{10}(a_{\text{max}}/\text{mm})$, $H_{100.a_{\text{min}}}$ and β . The triangles indicate the parameter values of the best-fit radiative transfer model. The vertical dotted line in panel (b) marks the analytic dust mass $M_{\text{dust,ana}}$ that is derived by using the flux density measured at 1.3 mm, $T_{\text{dust}} = 24.8$ K, and $\kappa_{1.3\text{mm}} = 2.3 \text{ cm}^2/\text{g}$, see Sect. 4.3.

4.3. Result and discussion

The best-fit model SEDs, identified as the ones with the lowest reduced χ^2 , are shown in panel (a) of Figure 8. Following the procedure described in Pinte et al. (2008), a Bayesian analysis is conducted by using the 6,552 SED models and assuming flat priors because there is no preliminary information about the parameters. The resulting marginalized probability distributions are presented in panels (b)–(e) of Figure 8. The triangles indicate the parameter values of the best-fit radiative transfer model, all of which are less than one bin away from the most probable values. The relatively wide distribution of the probabilities, especially for a_{max} , implies that there are degeneracies between model parameters in fitting the SED. The most probable dust masses are $M_{\text{dust,RT}} = 3 \times 10^{-4} M_{\odot} (9.5 \times 10^{-5}, 1.7 \times 10^{-3})$ and $9.5 \times 10^{-5} M_{\odot} (5.3 \times 10^{-5}, 5.3 \times 10^{-4})$ when the DSHARP and DIANA dust opacities are used, respectively, where the numbers given in parenthesis correspond to the 68% confidence in-



Fig. 9. Upper panel: $\Lambda_{1.3mm}$ as a function of $F_{1.3mm}$ and R_{out} . Black dots stand for the DSHARP disks. The black star represents the DoAr 33 disk. Smooth disks and disks with substructures observed in Long et al. (2019) are indicated with grey dots and grey rings, respectively. All the flux densities are scaled to the distance of 140 pc. *Bottom panel:* T_{dust} distribution in the $F_{1.3mm} - R_{out}$ space. The dashed line draws the contour level of 24.8 K, the analytic temperature simply given by $25 (L_*/L_{\odot})^{0.25}$ K.

terval. To calculate the analytic dust mass $M_{dust.ana}$, we adopted $\kappa_{1.3mm} = 2.3 \text{ cm}^2/\text{g}$ and $T_{dust} = 25 (L_{\star}/L_{\odot})^{0.25} = 24.8 \text{ K}$ for consistency with literature studies. We took the integrated flux of 35 mJy from the DSHARP project (see Table 5), and derived $M_{dust.ana} = 4.1 \times 10^{-5} M_{\odot}$ that is indicated with the vertical dotted line in panel (b) of Figure 8. Dust masses from radiative transfer modeling are 7.3 and 2.3 times higher than the analytic dust mass, when the DSHARP and DIANA dust opacities are adopted in the simulation, respectively.

Taking the most probable model with the inclusion of the DSHARP opacity as an illustration, regions below the $\tau_{1.3mm} = 1$ surface occupy $\sim 70\%$ of the total mass, which partly explains the large difference between $M_{dust.ana}$ and $M_{dust.RT}$. The massaveraged dust temperature is $T_{dust} = 19.6 \text{ K}$ that is cooler than the analytic dust temperature 24.8 K. This also contributes to the mass difference. To reproduce the shallow millimeter spectral slope $\alpha_{\rm mm} = 2.1$, the most probable maximum grain size achieves $a_{\text{max}} = 32 \text{ mm}$, yielding an absorption coefficient of $\overline{\kappa_{1.3\text{mm}}} = 0.7 \text{ cm}^2/\text{g}$. The discrepancy in $\kappa_{1.3\text{mm}}$ accounts for a factor of 2.3/0.7 = 3.3 difference in the dust mass. The millimeter spectral slope of DoAr 33 is typically found in other disks around both brown dwarfs (BDs) and T Tauri/Herbig stars (e.g., Ricci et al. 2010, 2014; Pinilla et al. 2017), implying that dust grains have grown to millimeter sizes at least. A multi-wavelength analysis, such as self-consistent radiative transfer modeling implemented in this study, is necessary to recover the grain size distribution, which is very important to characterize dust emissivities, and therefore obtain a reliable determination of dust masses. This information is basically missing in the traditionally analytic approach.

The large underestimation of the dust mass obtained for DoAr 33 may not be a peculiarity of that source.. With a thorough analysis of millimeter observations, Macías et al. (2021) recovered 4.5 - 5.9 times more mass for the TW Hydrae disk than with the usual analytic methodology. Similar underestimation factors have also been found for some disks in the Taurus star formation region (Ballering & Eisner 2019; Ribas et al. 2020). To give a rough assessment on how frequently a large underestimation would be encountered, Figure 9 shows $\Lambda_{1.3mm}$ and T_{dust} as a function of $F_{1.3\text{mm}}$ and R_{out} . When making the plot, we include models in which only M_{dust} and R_{out} are varied, because as discussed in Sect. 3.1 they have the most significant impact on the underestimation factor. Other parameters including the disk inclination are fixed to their fiducial values given in Table 1. Disks from the DSHARP program and ALMA survey conducted by Long et al. (2019) are overlaid. We excluded disks in multiple systems and disks around Herbig stars. For simplicity, we directly took the 95% enclosing flux radius from the literature to be R_{out} . A majority of disks are located in parameter spaces that yield a relatively large underestimation (e.g., $\Lambda_{1.3mm} \gtrsim 3$). There are few outliers even brighter than model predictions at the highest level. The maximum dust mass sampled in our grid is $M_{\rm dust} = 3 \times 10^{-3} M_{\odot}$. Under the 2D density distribution describe in Sect. 2.1, such a large amount of material concentrated in a narrow radius range makes the disk quite optically thick, any further increase in M_{dust} does not contribute much more to the emergent flux. However, stellar properties, power-law index for the dust surface density, flaring index, scale height, and degree of dust settling are all fixed in generating the plot. A combination of adjustments to (some of) these assumptions is able to induce a higher dust temperature T_{dust} . Alternatively, lowering the disk inclination would increase the solid angle towards the observer. Both possibilities can bring the millimeter flux up to the observed level.

Correcting for the underestimation on disk dust masses will certainly ease the mass budget problem for planet formation (e.g., Ueda et al. 2020; Andrews 2020). It is also expected to have an impact on the $M_{dust} - M_{\star}$ relation. Millimeter surveys for protoplanetary disks have found that the measured fluxes strongly correlate to the stellar masses $F_{\rm mm} \propto M_{\star}^{\sim 1.7}$ (e.g., Andrews et al. 2013; Pascucci et al. 2016; Andrews 2020). A correlation between the dust mass and stellar mass will be naturally identified, when converting $F_{mm}(s)$ to dust masses with Eq. 1 and using a constant absorption coefficient κ_{mm} and dust temperature T_{dust} . If the measured fluxes fully reflect the total amount of dust material (i.e., not considering the effects of $\kappa_{\rm mm}$ and $T_{\rm dust}$), T Tauri/Herbig disks are expected to be more massive than BD disks, because of the $F_{\rm mm} - M_{\star}$ relation. This means that the optical depth of T Tauri/Herbig disks is systematically higher than for BD disks if all disks have similar sizes. In such a circumstance, corrections for true dust masses are larger in disks around higher mass stars, which in principle will steepen the $M_{dust} - M_{\star}$ relation. However, the size of the millimeter disk $R_{\rm disk,mm}$ has been found to scale with the stellar mass $R_{\rm disk,mm} \propto M_{\star}^{0.6}$, though such a relation is less pronounced (Andrews et al. 2018b; Andrews 2020), and whether or not it extends down to the BD regime needs more spatially resolved observations for further confirmation (e.g., Ricci et al. 2013, 2014; Testi et al. 2016). As shown in Sect 3.1, the disk size

significantly affects the mass derivation, with more severe underestimation for smaller disks. In this regard, correction factors for the analytic dust masses of BD disks are higher than those for their higher mass counterparts, which conversely will shallow the $M_{dust} - M_{\star}$ relation. Moreover, there is a large scatter on the millimeter spectral slope α_{mm} for each individual stellar mass bin (see for instance Figure 2 in Pinilla et al. (2017) and Figure 9 in Andrews (2020)), pointing to inhomogeneities of the dust properties and/or a dispersion in optical depths. Given the complexity and parameter coupling, a dedicated investigation on the relation between M_{dust} and M_{\star} , and its evolution with time (Pascucci et al. 2016; Rilinger & Espaillat 2021), requires a homogeneous radiative transfer analysis.

Spatially resolved observations at various wavelengths provide the radial and vertical brightness distributions, which are directly linked to the product of the dust emissivity and density distribution at different locations in the disk. Therefore, spatially resolved data, when available, should be taken into account (e.g., Gräfe et al. 2013; Menu et al. 2014; Pinte et al. 2016; Liu et al. 2017). If the disk is completely optically thick at all the observed millimeter wavelengths, having multi-wavelength observations would still not be enough to provide a good measurement of the dust mass. Observations at longer wavelengths (even with the Very Large Array) are also necessary in this situation (e.g., Macías et al. 2018; Carrasco-González et al. 2019; Macías et al. 2021; Guidi et al. 2022).

5. Summary

Dust mass in protoplanetary disks is an important parameter characterizing the potential for planet formation. Literature studies usually derive dust masses using an analytic approach based on the optically thin assumption. Statistic analysis of dust masses obtained in this way reveals that only few disks are able to form our solar system or its analogs. However, protoplanetary disks, particularly in the inner regions, are probably optically thick, which might lead to substantial underestimation on the true dust mass.

Using self-consistent radiative transfer models, we conducted a detailed parameter study to investigate to which degree dust masses can be underestimated, and how the underestimation is influenced by disk and dust properties. Our results show that mass underestimations can be a few times up to hundreds depending on the optical depth along the line of sight. The most significant impacts on the underestimation are produced by the disk outer radius, inclination and the true dust mass. We also compared models with different dust settling approaches that control the millimeter dust scale height. The results show that the underestimation is not significantly affected, but the well-mixed model produces the strongest millimeter emisson, and consequently it needs the lowest M_{dust} to fit the data. Different dust scattering modes are also tested, but the differences in the millimeter flux, dust temperature, and therefore the underestimation are generally small. Given the prevalence of disk substructures revealed by ALMA, we also included these small-scale features in the models. The results show that their impacts on the mass underestimation is weaker than that induced by the disk outer radius and true dust mass.

As an application, we also conducted a detailed SED modeling for DoAr 33 that is one of the 20 disks observed by the DSHARP project. In the modeling procedure, we fixed the disk outer radius and disk inclination to the constraints set by the ALMA observations. The most probable dust masses are 7.3 and 2.3 times higher than the analytic dust mass, when the DSHARP and DIANA dust opacities are adopted in the radiative transfer simulation, respectively.

A homogeneous radiative transfer modeling is a more appropriate way to determine the disk dust mass, and investigate its dependence on the host stellar properties. In addition, in the analysis, multi-wavelength spatially resolved data are required for better reliability.

Acknowledgements. We thank the anonymous referee for the constructive comments that highly improved the manuscript. Y.L. acknowledges the financial support by the National Natural Science Foundation of China (Grant No. 11973090), and the science research grants from the China Manned Space Project with NO. CMS-CSST-2021-B06. T.H. acknowledges support from the European Research Council under the Horizon 2020 Framework Program via the ERC Advanced Grant Origins 832428. M.F. acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 757957). G.R. acknowledges support from the Netherlands Organisation for Scientific Research (NWO, program number 016.Veni.192.233) and from an STFC Ernest Rutherford Fellowship (grant number ST/T003855/1). H.W. acknowledges the support by NSFC grant 11973091.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

PACS has been developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany); INAF-IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); IAC (Spain). This development has been supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy), and CICYT/MCYT (Spain). SPIRE has been developed by a consortium of institutes led by Cardiff University (UK) and including Univ. Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); and Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC, UKSA (UK); and NASA (USA).

ALMA is a partnership of ESO (representing its member states), NSF (USA), and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO, and NAOJ.

References

- ALMA Partnership, Brogan, C. L., Pérez, L. M., et al. 2015, ApJ, 808, L3
- André, P., Men'shchikov, A., Bontemps, S., et al. 2010, A&A, 518, L102
- Andrews, S. M. 2020, ARA&A, 58, 483
- Andrews, S. M., Huang, J., Pérez, L. M., et al. 2018a, ApJ, 869, L41
- Andrews, S. M., Rosenfeld, K. A., Kraus, A. L., & Wilner, D. J. 2013, ApJ, 771, 129
- Andrews, S. M., Terrell, M., Tripathi, A., et al. 2018b, ApJ, 865, 157
- Andrews, S. M. & Williams, J. P. 2007, ApJ, 671, 1800
- Andrews, S. M., Wilner, D. J., Hughes, A. M., Qi, C., & Dullemond, C. P. 2010, ApJ, 723, 1241
- Ansdell, M., Williams, J. P., Trapman, L., et al. 2018, ApJ, 859, 21
- Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, ApJ, 828, 46
- Avenhaus, H., Quanz, S. P., Garufi, A., et al. 2018, ApJ, 863, 44
- Ballering, N. P. & Eisner, J. A. 2019, AJ, 157, 144
- Balog, Z., Müller, T., Nielbock, M., et al. 2014, Experimental Astronomy, 37, 129
- Balog, Z., Rieke, G. H., Muzerolle, J., et al. 2008, ApJ, 688, 408
- Barenfeld, S. A., Carpenter, J. M., Ricci, L., & Isella, A. 2016, ApJ, 827, 142
- Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Guesten, R. 1990, AJ, 99, 924
- Bergin, E. A., Cleeves, L. I., Gorti, U., et al. 2013, Nature, 493, 644
- Bergin, E. A. & Williams, J. P. 2017, in Astrophysics and Space Science Library, Vol. 445, Formation, Evolution, and Dynamics of Young Solar Systems, ed. M. Pessah & O. Gressel, 1
- Beuther, H. & Steinacker, J. 2007, ApJ, 656, L85
- Birnstiel, T., Dullemond, C. P., Zhu, Z., et al. 2018, ApJ, 869, L45
- Birnstiel, T., Fang, M., & Johansen, A. 2016, Space Sci. Rev., 205, 41 Bruggeman, D. 1935, Ann. Phys., 416, 636
- Carrasco-González, C., Sierra, A., Flock, M., et al. 2019, ApJ, 883, 71
- Cazzoletti, P., Manara, C. F., Liu, H. B., et al. 2019, A&A, 626, A11
- Cazzoletti, P., van Dishoeck, E. F., Pinilla, P., et al. 2018, A&A, 619, A161

- Cieza, L. A., González-Ruilova, C., Hales, A. S., et al. 2021, MNRAS, 501, 2934
- Cieza, L. A., Ruíz-Rodríguez, D., Hales, A., et al. 2019, MNRAS, 482, 698
- Cox, E. G., Harris, R. J., Looney, L. W., et al. 2017, ApJ, 851, 83
- Cutri, R. M. & et al. 2013, VizieR Online Data Catalog, 2328
- Daemgen, S., Natta, A., Scholz, A., et al. 2016, A&A, 594, A83
- Diolaiti, E., Bendinelli, O., Bonaccini, D., et al. 2000, A&AS, 147, 335
- Dong, R., Liu, S.-y., Eisner, J., et al. 2018, ApJ, 860, 124
- Dorschner, J., Begemann, B., Henning, T., Jaeger, C., & Mutschke, H. 1995, A&A, 300, 503
- Dowell, C. D., Allen, C. A., Babu, R. S., et al. 2003, in Proc. SPIE, Vol. 4855, Millimeter and Submillimeter Detectors for Astronomy, ed. T. G. Phillips & J. Zmuidzinas, 73-87
- Drazkowska, J., Bitsch, B., Lambrechts, M., et al. 2022, arXiv e-prints, arXiv:2203.09759
- Dubrulle, B., Morfill, G., & Sterzik, M. 1995, Icarus, 114, 237
- Dullemond, C. P., Birnstiel, T., Huang, J., et al. 2018, ApJ, 869, L46
- Dullemond, C. P. & Dominik, C. 2004, A&A, 421, 1075
- Dullemond, C. P., Juhasz, A., Pohl, A., et al. 2012, RADMC-3D: A multi-
- purpose radiative transfer tool, Astrophysics Source Code Library Ercolano, B. & Pascucci, I. 2017, Royal Society Open Science, 4, 170114
- Fedele, D., Tazzari, M., Booth, R., et al. 2018, A&A, 610, A24
- Flaherty, K., Hughes, A. M., Simon, J. B., et al. 2020, ApJ, 895, 109
- Flaherty, K. M., Hughes, A. M., Rose, S. C., et al. 2017, ApJ, 843, 150
- Fruchter, A. S. & Hook, R. N. 2002, PASP, 114, 144
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1
- Garcia, A. J. L. & Gonzalez, J.-F. 2020, MNRAS, 493, 1788
- Graciá-Carpio, J., Wetzstein, M., & Roussel, H. 2015, ArXiv e-prints
- Gräfe, C., Wolf, S., Guilloteau, S., et al. 2013, A&A, 553, A69
- Grant, S. L., Espaillat, C. C., Wendeborn, J., et al. 2021, ApJ, 913, 123
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, A&A, 518, L3
- Guidi, G., Isella, A., Testi, L., et al. 2022, arXiv e-prints, arXiv:2207.01496
- Hendler, N. P., Mulders, G. D., Pascucci, I., et al. 2017, ApJ, 841, 116
- Herczeg, G. J. & Hillenbrand, L. A. 2014, ApJ, 786, 97
- Huang, J., Andrews, S. M., Dullemond, C. P., et al. 2018, ApJ, 869, L42
- Juhász, A., Bouwman, J., Henning, T., et al. 2010, ApJ, 721, 431
- Kley, W. & Nelson, R. P. 2012, ARA&A, 50, 211
- Kovács, A. 2008, in Proc. SPIE, Vol. 7020, Millimeter and Submillimeter Detectors and Instrumentation for Astronomy IV, 70201S
- Kurucz, R. 1994, Solar abundance model atmospheres for 0,1,2,4,8 km/s. Kurucz CD-ROM No. 19. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1994., 19
- Law, C. J., Teague, R., Loomis, R. A., et al. 2021, ApJS, 257, 4
- Liu, H. B. 2019, ApJ, 877, L22
- Liu, Y., Bertrang, G. H. M., Flock, M., et al. 2022, arXiv e-prints, arXiv:2208.09230
- Liu, Y., Dipierro, G., Ragusa, E., et al. 2019, A&A, 622, A75
- Liu, Y., Henning, T., Carrasco-González, C., et al. 2017, A&A, 607, A74
- Long, F., Andrews, S. M., Rosotti, G., et al. 2022, ApJ, 931, 6
- Long, F., Herczeg, G. J., Harsono, D., et al. 2019, ApJ, 882, 49
- Long, F., Pinilla, P., Herczeg, G. J., et al. 2018, ApJ, 869, 17
- Machida, M. N., Kokubo, E., Inutsuka, S.-I., & Matsumoto, T. 2010, MNRAS, 405, 1227
- Macías, E., Espaillat, C. C., Ribas, Á., et al. 2018, ApJ, 865, 37
- Macías, E., Guerra-Alvarado, O., Carrasco-González, C., et al. 2021, A&A, 648, A33
- Manara, C. F., Morbidelli, A., & Guillot, T. 2018, A&A, 618, L3
- Maucó, K., Briceño, C., Calvet, N., et al. 2018, ApJ, 859, 1
- McClure, M. K., Bergin, E. A., Cleeves, L. I., et al. 2016, ApJ, 831, 167
- Menu, J., van Boekel, R., Henning, T., et al. 2014, A&A, 564, A93
- Min, M., Hovenier, J. W., & de Koter, A. 2005, A&A, 432, 909
- Miotello, A., Bruderer, S., & van Dishoeck, E. F. 2014, A&A, 572, A96
- Miotello, A., Kamp, I., Birnstiel, T., Cleeves, L. I., & Kataoka, A. 2022, arXiv e-prints, arXiv:2203.09818
- Miotello, A., van Dishoeck, E. F., Kama, M., & Bruderer, S. 2016, A&A, 594, A85
- Mordasini, C., Alibert, Y., Klahr, H., & Henning, T. 2012, A&A, 547, A111
- Mulders, G. D., Pascucci, I., Ciesla, F. J., & Fernandes, R. B. 2021, ApJ, 920, 66 Najita, J. R. & Kenyon, S. J. 2014, MNRAS, 445, 3315
- Ohashi, S. & Kataoka, A. 2019, ApJ, 886, 103
- Ormel, C. W., Spaans, M., & Tielens, A. G. G. M. 2007, A&A, 461, 215
- Pascucci, I., Cabrit, S., Edwards, S., et al. 2022, arXiv e-prints, arXiv:2203.10068
- Pascucci, I., Testi, L., Herczeg, G. J., et al. 2016, ApJ, 831, 125
- Pearson, C., Lim, T., North, C., et al. 2014, Experimental Astronomy, 37, 175
- Pérez, L. M., Benisty, M., Andrews, S. M., et al. 2018, ApJ, 869, L50
- Picogna, G., Ercolano, B., Owen, J. E., & Weber, M. L. 2019, MNRAS, 487,
- 691
- Pinilla, P., Quiroga-Nuñez, L. H., Benisty, M., et al. 2017, ApJ, 846, 70

- Pinte, C., Dent, W. R. F., Ménard, F., et al. 2016, ApJ, 816, 25
- Pinte, C., Padgett, D. L., Ménard, F., et al. 2008, A&A, 489, 633
- Powell, D., Gao, P., Murray-Clay, R., & Zhang, X. 2022, Nature Astronomy
- Ragan, S., Henning, T., Krause, O., et al. 2012, A&A, 547, A49
- Rebollido, I., Merín, B., Ribas, Á., et al. 2015, A&A, 581, A30
- Ribas, Á., Espaillat, C. C., Macías, E., & Sarro, L. M. 2020, A&A, 642, A171
- Ricci, L., Isella, A., Carpenter, J. M., & Testi, L. 2013, ApJ, 764, L27
- Ricci, L., Testi, L., Natta, A., et al. 2010, A&A, 512, A15
- Ricci, L., Testi, L., Natta, A., et al. 2014, ApJ, 791, 20
- Rich, E. A., Teague, R., Monnier, J. D., et al. 2021, ApJ, 913, 138
- Rilinger, A. M. & Espaillat, C. C. 2021, ApJ, 921, 182
- Roussel, H. 2013, PASP, 125, 1126
- Schräpler, R. & Henning, T. 2004, ApJ, 614, 960
- Segura-Cox, D. M., Schmiedeke, A., Pineda, J. E., et al. 2020, Nature, 586, 228 Sheehan, P. D., Tobin, J. J., Federman, S., Megeath, S. T., & Looney, L. W. 2020,
- ApJ, 902, 141
- Sierra, A. & Lizano, S. 2020, ApJ, 892, 136
- Sierra, A., Pérez, L. M., Zhang, K., et al. 2021, ApJS, 257, 14
- Teague, R., Henning, T., Guilloteau, S., et al. 2018, ApJ, 864, 133 Testi, L., Birnstiel, T., Ricci, L., et al. 2014, in Protostars and Planets VI, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 339
- Testi, L., Natta, A., Scholz, A., et al. 2016, A&A, 593, A111
- Toci, C., Rosotti, G., Lodato, G., Testi, L., & Trapman, L. 2021, MNRAS, 507, 818
- Toon, O. B. & Ackerman, T. P. 1981, Appl. Opt., 20, 3657
- Trapman, L., Facchini, S., Hogerheijde, M. R., van Dishoeck, E. F., & Bruderer, S. 2019, A&A, 629, A79
- Tychoniec, Ł., Manara, C. F., Rosotti, G. P., et al. 2020, A&A, 640, A19
- Ueda, T., Kataoka, A., & Tsukagoshi, T. 2020, ApJ, 893, 125
- van Boekel, R., Henning, T., Menu, J., et al. 2017, ApJ, 837, 132
- van der Marel, N., van Dishoeck, E. F., Bruderer, S., et al. 2013, Science, 340, 1199
- van der Plas, G., Ménard, F., Ward-Duong, K., et al. 2016, ApJ, 819, 102
- Villenave, M., Ménard, F., Dent, W. R. F., et al. 2020, A&A, 642, A164
- Wilking, B. A., Meyer, M. R., Robinson, J. G., & Greene, T. P. 2005, AJ, 130, 1733
- Williams, J. P. & Best, W. M. J. 2014, ApJ, 788, 59
- Williams, J. P. & Cieza, L. A. 2011, ARA&A, 49, 67
- Woitke, P., Min, M., Pinte, C., et al. 2016, A&A, 586, A103
- Wolf, S., Schegerer, A., Beuther, H., Padgett, D. L., & Stapelfeldt, K. R. 2008, ApJ, 674, L101
- Zhu, Z., Zhang, S., Jiang, Y.-F., et al. 2019, ApJ, 877, L18
- Zubko, V. G., Mennella, V., Colangeli, L., & Bussoletti, E. 1996, MNRAS, 282, 1321

Appendix A: Underestimation factors for face-on and edge-on disks

In this section, we explore how the underestimation factor changes with different model parameters for face-on $(i = 0^{\circ})$ and edge-on $(i = 90^\circ)$ disks. The results are shown in Figure A.1 and A.2.

The degree of underestimation, and its variation with different mode parameters, are broadly comparable between the faceon and fiducial inclined $(i = 41.8^{\circ})$ disks. However, the dependencies of Λ on most of the explored parameters are found to be tighter for edge-on disks. For instance, when the disk is viewed face-on, Λ decreases with increasing disk outer radius R_{out} , and it becomes insensitive to R_{out} when $R_{out} \gtrsim 100$ AU. However, for edge-on disks, a strong correlation between Λ and R_{out} exists for a wide range of R_{out} , i.e., $10 \le R_{out} \le 100$ AU. Moreover, different dust scale heights ($H_{100,a_{\min}}$ or $H_{100}(1 \text{ mm dust})$) generally result in a similar underestimation for disks with low inclinations. But, edge-on disks show clear dependencies between Λ and both parameters.

Appendix B: CSO/SHARC II observations

Through the Chinese Telescope Plan, DoAr 33 was observed with the SHARCII bolometer array (Dowell et al. 2003) on the 10.4 m Caltech Submillimeter Observatory (CSO) on Mauna Kea. SHARC II contains a 12×32 array of pop-up bolometers with a > 90% filling factor over the field. The field-of-view of the SHARCII maps was $\sim 2' \times 0.6'$, and the beam size at 350 μ m is about 8.5".

The data at $350\,\mu\text{m}$ were acquired on June 05 2015. The conditions during the observations were excellent with $\tau_{225 \text{ GHz}}$ = 0.03. The observing sequence consists of a series of target scans bracketed by scans of Pollas and cal_1629m2422, serving as both absolute flux calibration and pointing calibration measurements. We used the sweep mode for the observations, in which the telescope moves in a Lissajous pattern that keeps the central regions of the maps fully sampled. Typical individual target scan time was 300 s, while 120 s was sufficient for the calibrators. We obtained a total of 15 scans on the science target. The data analysis consists of reduction of raw data with the CRUSH pipeline (Kovács 2008), flux calibration and aperture photometry. Figure B.1 shows the reduced image. The flux density of DoAr 33 is measured to be 0.212 Jy. The total uncertainty on the flux ranges from $\sim 10-20\%$ mainly due to the uncertainties of the calibrator's flux.

Appendix C: Revisiting the Herschel data

DoAr 33 was observed with Herschel within the framework of a large survey of circumstellar disks in the Ophiuchus cloud (André et al. 2010). We re-reduced the data and performed the photometry based on point-spread function (PSF) fitting although literature aperture photometry have been reported on PACS and SPIRE data (Rebollido et al. 2015). We work on the same observations as Rebollido et al. (2015). The PACS 70 μ m and 160 μ m data are from program OT1_pabraham_3, while the SPIRE data at 250, 350, and 500 μ m are from the program KPGT_pandre_1. In addition, we newly report on PACS $100 \,\mu\text{m}$ photometry, also from the KPGT_pandre_1 program. We downloaded Level2.5 data (scan and cross-scan combined) from the Herschel Science Archive (HSA), based on the recent bulk reprocessing which had employed the pipeline versions SPG 14.2.0 for the PACS data, and SPG v14.2.1 for the SPIRE data. This iteration of the data reduction includes improved versions of the pointing products which can refine the sharpness of the PSFs especially at short wavelengths (70 and 100 μ m). The maps for PACS are JSCANAM maps (Graciá-Carpio et al. 2015), which are an implementation of the SCANAMORPHOS suite of algorithms devised by Roussel (2013). These standard maps have been constructed using the pixfrac=0.1 setting for the DRIZZLE algorithm (see Fruchter & Hook 2002). For the SPIRE data, we used the maps from the standard mapper, with the point-source calibration intact (map units Jy/beam).

For PSF references, we used the data released by the PACS and SPIRE instrument teams, i.e., maps of asteroid Vesta² and of planet Neptune³, respectively. For the PACS PSFs, appropriate files were available for the pixfrac, scan instrument angle, and scan speed values of the science data. We just had to regrid the fine PSF data to the somewhat coarser pixel scale of the science maps. The SPIRE PSFs were available in a pixel scale adapted to the science data. These reference PSFs were obtained using the standard scan speed for SPIRE (30"/s, sampling 18.2 Hz), while the science data have been taken with the fast scan speed of the

² ftp://ftp.sciops.esa.int/pub/hsc-calibration/PACS/PSF/

PACSPSF_PICC-ME-TN-033_v2.2.tar.gz

³ http://herschel.esac.esa.int/twiki/bin/view/Public/SpirePhotometer BeamProfile2



Fig. A.1. Same as in Figure 4, but the disk inclination is fixed to $i = 0^{\circ}$ for all models.

SPIRE /PACS parallel mode (60"/s, sampling 10 Hz). Potential differences in the PSF shape would remain a second-order effect (see Griffin et al. 2010, and especially the Parallel-Mode Observers' Manual⁴, Section 2.1). All reference PSFs were rotated in order to match the PSF orientation in the individual science maps, in dependence of the satellite position angles, taken from the science map fits headers.

The actual PSF photometry was performed employing the STARFINDER package (Diolaiti et al. 2000), which has proven to

successfully cope with the circumstances in *PACS* maps with extended emission (e.g. Ragan et al. 2012). For *SPIRE*, several photometry algorithms are implemented in the *Herschel* data reduction environment HIPE, but their performance has been mainly tested on clean stellar or extragalactic fields (Pearson et al. 2014). For our circumstances – a point source embedded in rivalling extended emission especially at the longest *Herschel* wavelengths – we also employ the STARFINDER PSF photometry for the *SPIRE* maps. DoAr 33 was detected with high confidence in all six bands (see also the high correlation coefficients

⁴ http://herschel.esac.esa.int/Docs/PMODE/html/parallel_om.html



Fig. A.2. Same as in Figure 4, but the disk inclination is fixed to $i = 90^{\circ}$ for all models.

mentioned in Table C.1). During the photometry setup, we explicitly truncated the PSF maps radially. For PACS, we work on Jy/pixel maps and are interested in the pixel-integrated flux densities. Therefore, we applied an appropriate aperture correction as tabulated in the encircled-energy-fraction files coming with the aforementioned PSF release (see also Balog et al. 2014). For the *SPIRE* maps (in Jy/beam) we are interested in the peak fluxes of the fitted PSFs when the extended background is eventually removed. STARFINDER produces such maps as a side product which just contain the appropriately scaled copies of the

PSFs, depending on the brightness of the identified sources in the field. We took these maps and fitted a 2-dimensional Gaussian to the source representing DoAr 33 (without extended background emission) in order to better recover the true peak flux, partly mitigating the relatively coarse pixel resolution of the SPIRE data. For a true point source, the value of the peak flux (in Jy/beam) will be identical to the integrated flux of the source.

Finally, a colour correction has to be applied. The *PACS* and *SPIRE* instrument teams provide correction factors for different types of SEDs, in particular for the situation of modified

Table	C.1.	Observational	details of	Herschel	data and	derived	photometry
-------	------	---------------	------------	----------	----------	---------	------------

Wavelength [µm]	70	100	160	250	350	500
abaida	1342238816 +	1342227148 +	1342238816 +	1342205093 +	1342205093 +	1342205093 +
obsids	1342238817	1342227149	1342238817	1342205094	1342205094	1342205094
scan speed ["/s]	20	20	20	60	60	60
parallel mode [y/n]	n	n	n	у	У	у
map pixel size ["]	1.6	1.6	3.2	6.0	10.0	14.0
PSF correlation coeff.	0.991	0.985	0.984	0.982	0.945	0.892
PSF radius used ["]	32	32	57.6	_	-	-
PSF integrated flux [Jy]	0.407	0.474	0.495	-	_	-
Aperture corrected	0.455	0.521	0.543			
flux density [Jy]	0.455	0.551	0.545	-	—	-
PSF peak flux [Jy/beam]	-	-	_	0.422	0.269	0.130
Colour-corrected flux density [Jy]	0.469 ± 0.033	0.537 ± 0.038	0.539 ± 0.038	0.416 ± 0.023	0.260 ± 0.014	0.121 ± 0.007



Fig. B.1. The CSO/SHARC II image of DoAr 33.

black-body SEDs $(F_{\nu} \sim B(T) \cdot \nu^{\beta})$. We therefore fitted the 70– 500 μ m SED with a simple but robust SED fitting tool (Beuther & Steinacker 2007). The SED of DoAr 33 in the FIR cannot be approximated with one temperature component, but the inclusion of two temperature components (fit values: 12.6 K and 32.5 K) gave satisfactory results when adopting $\beta = +2$ (a case explicitly listed in the colour correction tables). For all six Herschel wavelengths, the tool gave the relative flux contributions for both temperature components. We applied the colour correction terms (interpolated to the above-mentioned temperatures) to the respective fractions of the measured flux values, and added up the two corrected values per wavelength. The finally derived flux densities for DoAr 33 are given in Table C.1. The formal fitting errors from the PSF photometry are quite small. As overall uncertainties we therefore quote the sum of instrument calibration uncertainties and a second term arising from uncertainties in the underlying models of the flux calibration standards. These uncertainties are (2.0 + 5.0)% for PACS and (1.5 + 4.0)% for SPIRE, respectively.

On the whole, we can confirm the *PACS* 70 μ m and 160 μ m photometry derived by Rebollido et al. (2015, cf. our Table C.1). However, we derive *SPIRE* fluxes 3–4 times higher than reported by their study. To assume that the photometry for the wrong *SPIRE* source was given in the paper (and in the associated Vizier catalogue) remains speculation. Therefore, we have to conclude that their automatised aperture photometry did not handle the subtraction of the extended background well in this case and led to a strong over-subtraction of flux.