



EUROPEAN
HEMATOLOGY
ASSOCIATION

haematologica

Journal of the European Hematology Association
Published by the Ferrata Storti Foundation

**XV Congress of the Italian Society of Experimental Hematology
Rimini, Italy, October 18-20, 2018**

ABSTRACT BOOK

ISSN 0392-26

Volume 103
OCTOBER
2018 | **s3**

XV Congress of the Italian Society of Experimental Hematology

Rimini, Italy, October 18-20, 2018

COMITATO SCIENTIFICO

Mario LUPPI, Presidente
Mauro KRAMPERA, Vice Presidente
Massimo MASSAIA, Past President
Antonio CURTI
Fabio FORGHIERI
Sara GALIMBERTI
Nicola GIULIANI
Luca MALCOVATI
Pellegrino MUSTO
Stefano SACCHI
Paolo VIGNERI

SEGRETERIA SIES

Via Marconi, 36 - 40122 Bologna
Tel. 051 6390906 - Fax 051 4219534
E-mail: segreteriasies@ercongressi.it
www.siesonline.it

SEGRETERIA ORGANIZZATIVA

Studio ER Congressi
Via Marconi, 36 - 40122 Bologna
Tel. 051 4210559 - Fax 051 4210174
E-mail: ercongressi@ercongressi.it
www.ercongressi.it

ABSTRACT BOOK

XV Congress of the Italian Society of Experimental Hematology

Rimini, Italy, October 18-20, 2018

Contributors

ABBVIE
CELGENE
NOVARTIS

GILEAD SCIENCES
JAZZ PHARMACEUTICALS

AMGEN
BRISTOL-MYERS SQUIBB
JANSSEN ITALIA
MUNDIPHARMA PHARMACEUTICALS
ROCHE
TAKEDA

DIASORIN
MSD
PFIZER
SHIRE

MATTIOLI 1885 - CASA EDITRICE
INCYTE BIOSCIENCES
WERFEN

XV Congress of the Italian Society of Experimental Hematology

Rimini, Italy, October 18-20, 2018

Main Program

..... 1

Best Abstracts

..... 14

Oral Communications

Session	1.	C001-C008	Acute Leukemia 1	18
Session	2.	C009-C016	Monoclonal Gammopathies and Multiple Myeloma 1	22
Session	3.	C017-C024	Chronic Lymphocytic Leukemia and Chronic Lymphoproliferative Disorders	26
Session	4.	C025-C032	Benign Hematology (Red Cell Disease Thrombosis and Hemostasis)	31
Session	5.	C033-C040	Stem Cell Transplantation	35
Session	6.	C041-C048	Molecular Hematology	39
Session	7.	C049-C056	Myelodysplastic Syndromes	43
Session	8.	C057-C064	Myeloproliferative Disorders and Chronic Myeloid Leukemia	47
Session	9.	C065-C072	Monoclonal Gammopathies and Multiple Myeloma 2	51
Session	10.	C073-C080	Acute Leukemia 2	55
Session	11.	C081-C088	Lymphomas	59
Session	12.	C089-C096	Stem Cells and Growth Factors	63

Posters

Session	1.	P001-P014	Acute Leukemia 1	68
Session	2.	P015-P025	Chronic Lymphocytic Leukemia and Chronic Lymphoproliferative Disorders 1	74
Session	3.	P026-P037	Monoclonal Gammopathies and Multiple Myeloma 1	80
Session	4.	P038-P043	Benign Hematology (Red Cell Disease Thrombosis and Hemostasis)	86
Session	5.	P044-P061	Myeloproliferative Disorders and Chronic Myeloid Leukemia	89
Session	6.	P062-P065	Myelodysplastic Syndromes	98
Session	7.	P066-P079	Acute Leukemia 2	100
Session	8.	P080-P090	Chronic Lymphocytic Leukemia and Chronic Lymphoproliferative Disorders 2	107
Session	9.	P091-P101	Monoclonal Gammopathies and Multiple Myeloma 2	113
Session	10.	P102-P111	Lymphomas	117
Session	11.	P112-P118	Molecular Hematology	122
Session	12.	P119-P124	Immunotherapy and cell therapy	126

CK2 inactivation, which could account for GATA1 instability. However, GATA-1 and CK2 β also colocalized in the nucleus in presence of the β -estradiol suggesting a possible direct mechanism in CK2-mediated control of this transcription factor.

Conclusions: CK2 β is essential for erythroid maturation, regulating STAT5, AKT and GATA-1. Further research will clarify if CK2 β subunit could influence these factors in an indirect or direct mechanism.

CO045

COMPORAMENTO DELLE MUTAZIONI DRIVER E NON DRIVER NEI PAZIENTI CON MIELOFIBROSI IN TRATTAMENTO CON RUXOLITINIB

S. Galimberti¹, S. Balducci¹, F. Guerrini¹, S. Grassi^{1,2}, M.G. Massantini¹, F. Ricci¹, D. Cattaneo³, A. Iurlo³, P. Vigneri⁴, M. Petrini¹, G.A. Palumbo⁵, C. Baratè¹

¹UO Ematologia, Dipartimento Medicina Clinica e Sperimentale, Università di Pisa; ²GENOME scuola di dottorato, Università di Siena; ³UO Ematologia, IRCCS Ca' Granda, Milano; ⁴Oncologia, Università di Catania; ⁵UO Ematologia, Università di Catania, Italy

Background: Introduction of ruxolitinib changed the outcome of patients with myelofibrosis (MF), offering longer survivals. Nevertheless, 50% of patients loss response; in some cases, this phenomenon has been ascribed to driver and non-driver mutations, but with conflicting results: Patel *et al.* (Blood 2015) reported that having >3 mutations well correlated with shorter time to discontinuation and overall survival, whereas in the COMFORT-II study the MF-associated mutations did not correlate with response, survival or discontinuation probability (Guglielmelli, Blood 2014).

Aims: in order to investigate if ruxolitinib could play any role in changing the mutational landscape in MF, we assessed the 3 driver and 8 non-driver mutations in 36 MF patients; all were assessed a diagnosis, 19 also after 12 months of ruxolitinib, and other 4 after hydroxyurea.

Methods: In addition to assessment of the driver mutations (JAK2, CALR, MPL), a PCR plate with pre-spotted primers able to detect 8 non-driver mutations was designed (Custom qBiomarker Somatic Mutation PCR Array® - Qiagen, Italy). ASXL1, EZH2, DNMT3A, IDH1, IDH2, SRSF2, TET2, TP53, for total 38 hot-spot sites, were assessed.

Results: JAK2 was mutated in 70% of cases, CALR in 20%, whereas 10% of cases were triple-negative. The median OS was significantly longer for primary MF (160 months) vs post-ET (80 months) or post-PV MF (35 months)(p=0.03), and for CALR- vs JAK2-mutated patients. At the last follow-up, 4 patients (11%) progressed to AML, and 12 (33%) died. The non-driver mutations were found at diagnosis in 33% of cases receiving ruxolitinib and in one/4 patients treated with hydroxyurea. Considering both driver and non-driver mutations, 24 cases (67%) were mutated, with 16 cases carrying one, and 10 two mutations. The most frequently detected mutations belonged to the methylation pathway (DNMT3A, IDH, TET2 = 75%), followed by TP53 (17%), SRSF2 (8%), ASXL1 (8%), and EZH2 (8%). During treatment, JAK2 allele burden remained stable, whereas non-driver mutations changed in 13 cases: 9 acquired a new mutation (DNMT3A in 5, IDH2 in one, and TP53 in another one) while other 4 lost mutation. None of the CALR-mutated cases carried non-driver mutations. In the 4 cases treated with hydroxyurea, one acquired the TP53 and another one the DNMT3A mutation. On the other hand, 4 cases in the group of ruxolitinib, and none in the group of hydroxyurea lost mutations present at diagnosis (TP53, IDH2, ASXL1, DNMT3A). Presence/absence of non-driver mutations, their number (>1), the molecular subgroup (methylation, splicing, chromatin) did not significantly condition OS.

Conclusions: In this work, even if on a small series of patients, we showed that during ruxolitinib about the half of cases developed non-driver mutations, a percentage overlapping to that observed in cases receiving hydroxyurea. Interestingly, ruxolitinib and not hydroxyurea allowed disappearance of mutations in one third of cases.

CO046

DEVELOPMENT OF A COMBINATION STRATEGY BASED ON ER AND OXIDATIVE STRESS TO TARGET ACUTE MYELOID LEUKEMIA

S. Masciarelli¹, E. Capuano¹, T. Ottone², M. Divona², S. Lavorgna², A. Picardi³, M.T. Voso², F. Lo Coco², F. Fazi¹

¹Department of Anatomical, Histological, Forensic & Orthopedic Sciences, Section of Histology & Medical Embryology, Sapienza University of Rome, laboratory affiliated to Istituto Pasteur Italia-Fondazione Cenci Bolognetti, Rome; ²Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome; ³Stem Cell Transplant Unit, Rome Transplant Network, Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy

Introduction: Acute Myeloid Leukemia (AML) is a heterogeneous disease caused by different molecular genetic aberrations. These result in the expression of fusion or mutant proteins that cause impaired differentiation and enhanced proliferation and survival. We previously showed that APL cell lines and primary blasts induced to differentiate by Retinoic Acid (RA) become highly sensitive to amounts of ER stress not detrimental for the same cells in the absence of RA. Furthermore the same cells resulted sensitive to a combination of ER stress inducers with Arsenic Trioxide (ATO) that generates oxidative stress. Importantly we observed that ER stress caused increased amounts of disulphide-bound high molecular weight aggregates of PML-RAR α and PML, exacerbating the alteration of cellular proteostasis already generated by induction of ER stress. This observation provides the rationale to translate the findings we observed in APL to other types of AML characterized by fusion or mutant proteins. The presence of mutant proteins that are easily prone to aggregation or mis-folding, because of their mutant structure or because of mis-localization, could render the cells sensitive to levels of ER and oxidative stress that could be recovered in their absence.

Methods: We treated AML cell lines and AML primary leukemic blasts with RA and ER and oxidative stress inducers, evaluating cell proliferation and death, activation of the ER/oxidative stress responses, localization and possible aggregation of the mutant proteins by confocal microscopy, colony forming capacity.

Results: We first tested a panel of AML cell lines characterized by different oncogenic fusion or mutant proteins and we found that ML-2 cells, bearing the MLL-AF6 fusion protein, and MV-4-11 cells, expressing the fusion protein MLL-AF4 and FLT3-ITD are highly sensitive to the combination of sub-lethal amounts of RA, Tm and ATO. In the cells undergoing ER and oxidative stress in combination, we found prolonged activation of the antioxidant response and of the unfolded protein response (UPR), activated by ER stress, as indicated by the expression of HMOX, CHOP, BiP and sXBP1. The antioxidant agent N-acetyl-cysteine and the inhibitor of the UPR player GADD34 determine resistance of the cells to the treatments. Furthermore, an inhibitor of the PERK branch of the UPR dramatically exacerbates the sensitivity to the combination of ER and oxidative stress pointing to this pathway as a possible new therapeutic molecular target. Importantly, the combination of ER and oxidative stress significantly reduces the colony forming capacity of primary leukemic blasts isolated from the bone marrow of FLT3-ITD positive patients.

Conclusions: Altogether our data suggest that the combination of low levels of ER and oxidative stress leads to apoptosis rather than recovery, achieved instead when the same stresses are induced alone.