



# Parental Somatic Mosaicism Uncovers Inheritance of an Apparently *De Novo* GFAP Mutation

Alice Grossi<sup>1</sup>, Federico Morelli<sup>1†</sup>, Marco Di Duca<sup>2</sup>, Francesco Caroli<sup>1†</sup>, Isabella Moroni<sup>3</sup>, Davide Tonduti<sup>4</sup>, Tiziana Bachetti<sup>1,5†</sup> and Isabella Ceccherini<sup>1\*</sup>

## OPEN ACCESS

### Edited by:

Ioannis Dragatsis,  
University of Tennessee Health  
Science Center (UTHSC),  
United States

### Reviewed by:

Albee Messing,  
University of Wisconsin-Madison,  
United States  
Michael Brenner,  
University of Alabama at Birmingham,  
United States

### \*Correspondence:

Isabella Ceccherini  
isabellaceccherini@gaslini.org

### †Present Address:

Federico Morelli,  
uniQure N.V., Paasheuvelweg 25a, 1105  
BP, Amsterdam, The Netherlands.  
Francesco Caroli,  
UOC Genetica Medica, IRCCS Istituto  
Giannina Gaslini, 16148, Genova, Italy.  
Tiziana Bachetti,  
UO Proteomica e Spettrometria di  
Massa, IRCCS Ospedale Policlinico San  
Martino, Genova, Italy

### Specialty section:

This article was submitted to  
Neurogenomics,  
a section of the journal  
Frontiers in Genetics

Received: 19 July 2021

Accepted: 02 November 2021

Published: 07 December 2021

### Citation:

Grossi A, Morelli F, Di Duca M, Caroli F,  
Moroni I, Tonduti D, Bachetti T and  
Ceccherini I (2021) Parental Somatic  
Mosaicism Uncovers Inheritance of an  
Apparently *De Novo* GFAP Mutation .  
Front. Genet. 12:744068.  
doi: 10.3389/fgene.2021.744068

<sup>1</sup>UOSD Laboratory of Genetics and Genomics of Rare Diseases, IRCCS Istituto Giannina Gaslini, Genoa, Italy, <sup>2</sup>Laboratory of Molecular Nephrology, IRCCS Istituto Giannina Gaslini, Genoa, Italy, <sup>3</sup>Department of Pediatric Neurosciences, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy, <sup>4</sup>Unit of Pediatric Neurology - C.O.A.L.A (Center for Diagnosis and Treatment of Leukodystrophies), V. Buzzi Children's Hospital, Milan, Italy, <sup>5</sup>Laboratory of Developmental Neuro-Biology, DISTAV, University of Genoa, Genoa, Italy

Alexander disease is a leukodystrophy caused by heterozygous mutations of *GFAP* gene. Recurrence in siblings from healthy parents provides a confirmation to the transmission of variants through germinal mosaicism. With the use of DNA isolated from peripheral blood, next-generation sequencing (NGS) of *GFAP* locus was performed with deep coverage ( $\geq 500\times$ ) in 11 probands and their parents (trios) with probands heterozygous for apparently *de novo* *GFAP* mutations. Indeed, one parent had somatic mosaicism, estimated in the range of 8.9%–16%, for the mutant allele transmitted to the affected sibling. Parental germline mosaicism deserves attention, as it is critical in assessing the risk of recurrence in families with Alexander disease.

**Keywords:** central nervous system diseases, genetic counseling, human genetics, DNA sequence analysis, *GFAP* gene, Alexander disease, germline mosaicism, somatic mosaicism

## INTRODUCTION

Alexander disease (AxD) is an extremely rare, untreatable, and usually fatal neurodegenerative disorder (OMIM #203450), classified among leukodystrophies due to white matter deficits (Messing and Brenner, 2020). It is estimated to affect 1:2.7 million people in Japan (Yoshida et al., 2011). The disease presents at different ages of onset, with distinct symptoms and prognosis: in neonates and early childhood (type I) and later, though not restricted to adulthood (type II) (Prust et al., 2011). AxD is caused by heterozygous mutations of glial fibrillary acidic protein (*GFAP*) gene, which eventually lead to the formation of aggregates, also containing alphaB-crystallin, HSP27, ubiquitin, and proteasome components (Quinlan et al., 2007). To date, a broad spectrum of pathogenic *GFAP* variants accounts for more than 90% of patients. Mutations occur either *de novo* or through transmission from the parental generation. A recurrent occurrence of the same disease-causing *GFAP* mutation in siblings from parents who tested negative for the variant strongly suggests the presence of a germinal mosaicism (Melchionda et al., 2013) (two affected siblings were also reported by Namekawa et al. (2002), but the parents were not examined). Indirect evidence for germinal mosaicism in *de novo* AxD cases has also been provided by studies finding that the *de novo* mutations predominantly arise on the paternal chromosome (Li et al., 2006; Zang et al., 2013). Such a condition may be associated with somatic mosaicism, a circumstance nevertheless unproven so far (Messing, 2018). In the case of AxD, the risk of transmitting a *GFAP* mutation to a second child by germline mosaicism has been estimated as less than 1% (Messing, 2018); however, when significant somatic mosaicism is observed in a parent, the risk of recurrence could be substantially higher.

In the initial diagnostic screening of patients affected with AxD, and their parents when available, the entire coding region of *GFAP* gene was examined by Sanger sequencing. To test the hypothesis that during our diagnostic workflow the mosaicism of somatic *GFAP* mutations in AxD families may have been under-recognized, 11 pairs of asymptomatic parents of AxD patients heterozygous for *GFAP* causative and apparently *de novo* mutations were selected and then subjected to targeted in-depth sequencing.

## MATERIALS AND METHODS

As the next-generation sequencing (NGS) technology can facilitate the detection of low-level mosaicism, undetectable by conventional Sanger sequencing (Qin et al., 2016), deep coverage ( $\geq 500\times$ ) NGS, targeting *GFAP* locus, was therefore performed, as already described (Rusmini et al., 2016), using DNA isolated from peripheral blood samples from 11 pairs of asymptomatic parents without previously detected *GFAP* mutations, whose siblings are affected with AxD and heterozygous for a causative *GFAP* mutation already detected in the past by Sanger sequencing. In particular, the samples to be re-evaluated were just plugged into a new diagnostic pipeline making use of an NGS-based gene panel setup more recently.

The genomic region included in the experimental design and sequenced is at chr17:4,292,001–42,997,000 (GRCh37/hg19), contains regulatory regions and UTRs, all nine exons, and flanking regions (*GFAP* reference transcript NM\_002,055.5), is divided into 59 amplicons of 264 bases on average and has a final coverage of 95%, with 752 bases missing, mostly intronic. Our attention was focused on two amplicons of *GFAP* gene containing variants under analysis, namely, p.E75K, p.L76P, p.R79C/H (chr17:42,992,581–42,992,886, according to GRCh37 and corresponding to exon 1), and p.R239C (chr17:42,990,593–42,990,894, according to GRCh37 and corresponding to exon 4). The parameters for the variant calling were configured in the custom mode recommended for detecting mosaicisms (Torrent Suite Software, Thermo Fisher®, Thermo Fisher Scientific, Waltham, MA, USA).


Single-nucleotide primer extension was performed using a specific kit (SNaPshot Multiplex System, Thermo Fisher®) and oligos suitable for the mutation site (p.R79C) on both strand (forward primer: 5'-GAGATGATGGAGCTCAATGACC-3'; reverse primer: 5'-CTTCTCGATGTAGCTGGCAAAG-3'), followed by capillary electrophoresis on automated DNA sequencer (ABI PRISM® 3130XL), according to a protocol already reported (Matera et al., 2013).

The PCR product of exon 1 was cloned using the genomic DNA of ID#110M subject as a template, into the TOPO TA vector (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. The plasmid DNA isolated from 51 clones was directly sequenced using the Sanger protocol, as already reported (Matera et al., 2013).

## RESULTS

**Supplementary Table S1** reports details on the in-depth NGS analysis of AxD-causing *GFAP* mutations in families with

180A_50%	180A_25%	180A_12.5%	180A_6.25%	110M_unknown	
720	656	739	760	1239	TOT (X)
371	549	661	730	1088	WT (X)
349	107	78	30	151	MUT (X)
48.47	16.31	10.55	3.95	12.18	MUT (%)




**FIGURE 1 |** Coverage ratios for mutant and wild-type (WT) alleles obtained from known dilutions of a defined heterozygous control (ID#180A) are compared with those of ID#110M parent. Bottom: WT and mutant allele reads shown above are reported as bars of a histogram.

apparently *de novo* occurrence, all representing single instances of sibs having a mutation. In particular, the coverage of the amplicons containing the mutant codons is reported for each indicated patient's parents (mother or father). All parents could be evaluated for possible mosaicism at the mutation site of corresponding patients, except the father of proband #147 (ID#147F), who had to be removed from the set because the coverage of the region, affected by the mutation present in his son, did not reach the minimum threshold established by the experimental design ( $500\times$ ).

The mother of proband #110 (ID#110M) revealed a mosaicism of *GFAP* mutation already detected in her son, the c.236G>A (p.R79H). The ratios between mutant and total allele reads amount to 12.18%, which can be regarded as the extension of the mosaicism in blood cells for this mother. However, when the 12.18% value was compared with ratios obtained previously with known dilutions of the mutant allele of a heterozygous control, ID#180A (the patient of family #180) carrying the mutations *GFAP* c.227T>C (p.L76P), an estimate of the extension of mosaicism of 16% was achieved (Figure 1), a figure higher than expected and likely affected by technical biases and experimental variability.

The presence of a small proportion of mutant allele was confirmed through the single-nucleotide primer extension



approach, evaluating the amount of the wild-type and mutant alleles obtained from ID#110 families by using both the F- and R-primers (Figure 2A) and then comparing the ratios of peak height and of peak area obtained from ID#110M DNA sample (right-hand yellow column) with a reference curve constructed using DNA mixtures containing known fractions of the two alleles (Figure 2B). This allowed us to estimate an 8.9% proportion of mutant allele in this mosaic mother.

Given the low and uncertain proportion of the expanded alleles in this latter parent, we sought to confirm her mosaic status by cloning the PCR product of exon 1, containing *GFAP* mutation c.236G>A (p.R79H). Overall, 51 colonies were obtained, 46 of which showed the presence of a wild-type G allele at the mutation site after Sanger sequencing. This allowed us to estimate a ratio of clones carrying mutant and wild-type inserts of 10% (Figure 2C).

Based on the different approaches applied and the comparably similar estimates obtained (8.9%, 10%, 12.18%, and 16%), it could therefore be concluded that ID#110M mother shows a rather low degree of mosaicism, although not irrelevant for *GFAP* mutation c.236G>A (p.R79H).

A retrospective reanalysis, carried out on the Sanger sequences initially obtained from this mother's DNA, showed a very small second/mutant peak allele underneath the reference allele, which however was not attributed to the mutant allele but rather to the background noise (data not shown).

## DISCUSSION

Somatic mosaicism, occurring as a result of postzygotic mutations, can affect human health and diseases (Acuna-Hidalgo et al., 2015; D’Gama and Walsh, 2018). Indeed, as far as AxD is concerned, one patient was reported as a somatic mosaic for a *GFAP* mutation that was detected in buccal DNA (presumably present also in brain), but not in blood DNA, clearly arising in the embryo rather than in the parental germline (Flint et al., 2012). Moreover, the combination of both somatic and germline mosaicism has been postulated to account also for another mechanism recently suggested in AxD, where the presence of a *GFAP* mutation only in a subset of cells in the parental generation could attenuate disease severity and delay the

age of onset, thus leading to an apparent anticipation of symptoms in the offspring expressing the mutation ubiquitously. Unfortunately, authors have been unable to test this hypothesis that should be considered anyway in future instances of inherited AxD (Hunt et al., 2021). Finally, rare repetitive occurrences of AxD were already described in families, in association with severe GFAP mutations likely transmitted from parents who did not have symptoms and/or had tested negative for the mutation; however, no germline mosaicism has yet been demonstrated (Namekawa et al., 2002; Melchionda et al., 2013). Mosaic mutations in parents, and particularly mosaicism due to mutations that occurred at an earlier stage and therefore present in both the germline and somatic cells, could underlie a proportion of these cases, otherwise attributable to *de novo* germline mutations. Here, we have confirmed AxD transmission through germline mosaic GFAP mutation, made evident by the demonstration of somatic mosaicism, a circumstance that requires specific and careful workout especially for rare diseases, as already described (Bachetti et al., 2011; Brewer et al., 2020).

Recognition of a mosaicism can be achieved with different methods, each with a different resolution. A major obstacle to observing mosaicism is that the testing sample should harbor the mutation, given the surprisingly variable levels of mosaicism potentially detectable between tissues and organs and even within the same embryonic lineages (Campbell et al., 2015).

A roughly 12% mosaic condition for GFAP c.236G>A (p.R79H) mutation, a genetic defect already shown in association with AxD and recurring in unrelated patients (Rodriguez et al., 2001; Asahina et al., 2006; Zang et al., 2013), has been demonstrated in blood cells of an AxD patient's mother, and for this reason, it must have been present to an unknown extent in the lineage of the germ cells. Unfortunately, we did not have the opportunity to test DNA from tissues other than whole blood, such as exfoliation cells found in oral mucosa swabs, urinary sediments, skin hair bulbs, or other tissues, to investigate the extent of somatic mosaicism. Depending on when the somatic mutation occurs, we expect a different involvement of different tissues that can carry heterogeneous cell populations together with tissues characterized instead by a homogeneous wild-type cell population. This is also consistent with the possible onset of the disease phenotype, whenever mutant cells are found within tissues that can be affected by the presence of the mutation. Subject ID#110M, identified as a mosaic in the present study, is asymptomatic, and this can be explained by postulating that the mutation is absent in the astrocyte cell line of the central nervous system, the target of the neurodegeneration typical of AxD and in which GFAP is expressed. Nevertheless, both the presence of the mosaicism at a tolerable low level in astrocytes and a late onset of a mild disease cannot be ruled out at this stage.

Finding a GFAP mutation in an AxD patient and following its segregation within the family is crucial both to confirm a diagnosis and to assess the risk of disease recurrence. Actually, families with *de novo* mutations are generally considered as at risk for possible germline mosaicism and should be counseled accordingly. Given the high probability of germline mosaicism (Li et al., 2006), and indeed genetic counseling and possibly prenatal testing during a

second pregnancy are already recommended for families at risk of AxD recurrence, the discovery that the patient's mother ID#110M is actually a mosaic has represented a confirmation for parents of their actual transmission hazard. Indeed, the discovery of germline mosaicism exposes this family to a much higher but unquantifiable risk of AxD recurrence: though the possibility that 100% of oocyte cells carrying this variant cannot be excluded, the real risk of recurrence in this family is presumably lower than the 50% risk typical of germline carriers (Gajecka, 2016; Machiela and Chanock, 2017).

Although the search for parental somatic, and therefore germline, mosaicisms should make use, whenever possible, of tissues relevant to the disease, NGS with deep coverage at target sites of even less appropriate DNA sources may represent a fruitful approach to this type of investigation. Obviously, the test becomes informative only if a somatic mutation is found, while for subjects with negative results, mosaicism cannot be definitely excluded (Gajecka, 2016; Machiela and Chanock, 2017).

The tools provided by NGS, and in particular the possibility of an in-depth analysis, allow for the detection of possible mosaicisms, thus providing safer and more informative genetic counseling, an opportunity that should be considered for many rare diseases, including AxD.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are all included in the article. The novel observation of the p.R79H GFAP variant found in a mosaic parent is deposited in the ClinVar database and it has been assigned ID = SCV001786708. Further inquiries can be directed to the corresponding author.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Comitato Etico Regionale Liguria. The patients/participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

AG, TB, and IC contributed to conception and design of the study. AG, FM, MD, and FC contributed to the acquisition and analysis of data. IM and DT contributed to patient clinical data. AG, MD, and IC drafted the manuscript text and prepared the figures. All the authors revised and approved the final manuscript draft.

## FUNDING

This study was supported by the Associazione "Più Unici che Rari" (the Italian Family Association for Alexander disease) and by the Italian Ministry of Health "Ricerca Corrente" and "5 x mille" (to Gaslini Institute).

## ACKNOWLEDGMENTS

We would like to thank all patients and their families for participating. IM is a member of the European Reference Network for Rare Neurological Disorders (ERN-RND).

## REFERENCES

- Acuna-Hidalgo, R., Bo, T., Kwint, M. P., van de Vorst, M., Pinelli, M., Veltman, J. A., et al. (2015). Post-Zygotic point Mutations Are an Underrecognized Source of *De Novo* Genomic Variation. *Am. J. Hum. Genet.* 97, 67–74. doi:10.1016/j.ajhg.2015.05.008
- Asahina, N., Okamoto, T., Sudo, A., Kanazawa, N., Tsujino, S., and Saitoh, S. (2006). An Infantile-Juvenile Form of Alexander Disease Caused by a R79H Mutation in GFAP. *Brain Develop.* 28, 131–133. doi:10.1016/j.braindev.2005.05.004
- Bachetti, T., Parodi, S., Di Duca, M., Santamaria, G., Ravazzolo, R., and Ceccherini, I. (2011). Low Amounts of PHOX2B Expanded Alleles in Asymptomatic Parents Suggest Unsuspected Recurrence Risk in Congenital central Hypoventilation Syndrome. *J. Mol. Med.* 89, 505–513. doi:10.1007/s00109-010-0718-y
- Brewer, C. J., Gillespie, M., Fierro, J., Scaringe, W. A., Li, J., Lee, C.-Y., et al. (2020). The Value of Parental Testing by Next-Generation Sequencing Includes the Detection of Germline Mosaicism. *J. Mol. Diagn.* 22, 670–678. doi:10.1016/j.jmoldx.2020.02.001
- Campbell, I. M., Shaw, C. A., Stankiewicz, P., and Lupski, J. R. (2015). Somatic Mosaicism: Implications for Disease and Transmission Genetics. *Trends Genet.* 31, 382–392. doi:10.1016/j.tig.2015.03.013
- Caroli, F., Biancheri, R., Seri, M., Rossi, A., Pessagno, A., Bugiani, M., et al. (2007). GFAP Mutations and Polymorphisms in 13 Unrelated Italian Patients Affected by Alexander Disease. *Clin. Genet.* 72, 427–433. doi:10.1111/j.1399-0004.2007.00869.x
- D’Gama, A. M., and Walsh, C. A. (2018). Somatic Mosaicism and Neurodevelopmental Disease. *Nat. Neurosci.* 21, 1504–1514. doi:10.1038/s41593-018-0257-3
- Flint, D., Li, R., Webster, L. S., Naidu, S., Kolodny, E., Percy, A., et al. (2012). Splice Site, Frameshift, and chimericGFAPmutations in Alexander Disease. *Hum. Mutat.* 33, 1141–1148. doi:10.1002/humu.22094
- Gajicka, M. (2016). Unrevealed Mosaicism in the Next-Generation Sequencing Era. *Mol. Genet. Genomics.* 291, 513–530. doi:10.1007/s00438-015-1130-7
- Hunt, C. K., Al Khleifat, A., Burchill, E., Ederle, J., Al-Chalabi, A., and Sreedharan, J. (2021). Does Genetic Anticipation Occur in Familial Alexander Disease? *Neurogenetics* 22, 215–219. doi:10.1007/s10048-021-00642-9
- Li, R., Johnson, A. B., Salomons, G. S., van der Knaap, M. S., Rodriguez, D., Boespflug-Tanguy, O., et al. (2006). Propensity for Paternal Inheritance of *De Novo* Mutations in Alexander Disease. *Hum. Genet.* 119 (1–2), 137–144. doi:10.1007/s00439-005-0116-7
- Machiela, M. J., and Chanock, S. J. (2017). The Ageing Genome, Clonal Mosaicism and Chronic Disease. *Curr. Opin. Genet. Develop.* 42, 8–13. doi:10.1016/j.gde.2016.12.002
- Matera, I., Musso, M., Griseri, P., Rusmini, M., Di Duca, M., So, M.-t., et al. (2013). Allele-Specific Expression at theRETLocus in Blood and Gut Tissue of Individuals Carrying Risk Alleles for Hirschsprung Disease. *Hum. Mutat.* 34, 754–762. doi:10.1002/humu.22302
- Melchionda, L., Fang, M., Wang, H., Fugnanesi, V., Morbin, M., Liu, X., et al. (2013). Adult-Onset Alexander Disease, Associated with a Mutation in an Alternative GFAP Transcript, May Be Phenotypically Modulated by a Non-

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2021.744068/full#supplementary-material>

- Neutral HDAC6 Variant. *Orphanet J. Rare Dis.* 8, 66. doi:10.1186/1750-1172-8-66
- Messing, A. (2018). Alexander Disease. *Handb. Clin. Neurol.* 148, 693–700. doi:10.1016/B978-0-444-64076-5.00044-2
- Messing, A., and Brenner, M. (2020). GFAP at 50. *ASN Neuro.* 12, 1759091420949680. doi:10.1177/1759091420949680
- Namekawa, M., Takiyama, Y., Aoki, Y., Takayashiki, N., Sakoe, K., Shimazaki, H., et al. (2002). Identification ofGFAP Gene Mutation in Hereditary Adult-Onset Alexander’s Disease. *Ann. Neurol.* 52, 779–785. doi:10.1002/ana.10375
- Prust, M., Wang, J., Morizono, H., Messing, A., Brenner, M., Gordon, E., et al. (2011). GFAP Mutations, Age at Onset, and Clinical Subtypes in Alexander Disease. *Neurology* 77, 1287–1294. doi:10.1212/WNL.0b013e3182309f72
- Qin, L., Wang, J., Tian, X., Yu, H., Truong, C., Mitchell, J. J., et al. (2016). Detection and Quantification of Mosaic Mutations in Disease Genes by Next-Generation Sequencing. *J. Mol. Diagn.* 18, 446–453. doi:10.1016/j.jmoldx.2016.01.002
- Quinlan, R. A., Brenner, M., Goldman, J. E., and Messing, A. (2007). GFAP and its Role in Alexander Disease. *Exp. Cel Res.* 313, 2077–2087. doi:10.1016/j.yexcr.2007.04.004
- Rodriguez, D., Gauthier, F., Bertini, E., Bugiani, M., Brenner, M., N’guyen, S., et al. (2011). Infantile Alexander Disease: Spectrum of GFAP Mutations and Genotype-Phenotype Correlation. *Am. J. Hum. Genet.* 69, 1134–1140. doi:10.1086/323799
- Rusmini, M., Federici, S., Caroli, F., Grossi, A., Baldi, M., Obici, L., et al. (2016). Next-Generation Sequencing and its Initial Applications for Molecular Diagnosis of Systemic Auto-Inflammatory Diseases. *Ann. Rheum. Dis.* 75, 1550–1557. doi:10.1136/annrheumdis-2015-207701
- Yoshida, T., Sasaki, M., Sasaki, M., Yoshida, M., Namekawa, M., Okamoto, Y., et al. (2011). Nationwide Survey of Alexander Disease in Japan and Proposed New Guidelines for Diagnosis. *J. Neurol.* 258, 1998–2008. doi:10.1007/s00415-011-6056-3
- Zang, L., Wang, J., Jiang, Y., Gu, Q., Gao, Z., Yang, Y., et al. (2013). Follow-up Study of 22 Chinese Children with Alexander Disease and Analysis of Parental Origin of *De Novo* GFAP Mutations. *J. Hum. Genet.* 58, 183–188. doi:10.1038/jhg.2012.152

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher’s Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Grossi, Morelli, Di Duca, Caroli, Moroni, Tonducci, Bachetti and Ceccherini. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.