

---

# Pushing athletic performance boundaries with anodal transcranial direct current stimulation: an exploratory, pilot, randomized controlled study on strength and inflammation in elite weightlifters

---

Received: 12 July 2025

Accepted: 8 June 2026

Published online: 12 June 2026

Cite this article as: Donato R., Prenassi M., Guidetti M. *et al.* Pushing athletic performance boundaries with anodal transcranial direct current stimulation: an exploratory, pilot, randomized controlled study on strength and inflammation in elite weightlifters. *BMC Sports Sci Med Rehabil* (2026). <https://doi.org/10.1186/s13102-026-01803-9>

**Renato Donato, Marco Prenassi, Matteo Guidetti, Clara Tambussi, Roberta Ferrucci, Alessandra Vezzoli, Cinzia Dellanoce, Stefano Pastore, Alessandro Brunello, Alberto Priori, Sara Marceglia & Simona Mrakic-Sposta**

---

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

**Pushing athletic performance boundaries with anodal transcranial direct current stimulation: an exploratory, pilot, randomized controlled study on strength and inflammation in elite weightlifters**

Renato de Donato<sup>1</sup>, Marco Prenassi<sup>2,3</sup>, Matteo Guidetti<sup>4</sup>, Clara Tambussi<sup>4</sup>, Roberta Ferrucci<sup>4,5,6,7</sup>, Alessandra Vezzoli<sup>8</sup>, Cinzia Dellanoce<sup>8</sup>, Stefano Pastore<sup>3</sup>, Alessandro Brunello<sup>9,†</sup>, Alberto Priori<sup>4,6</sup>, Sara Marceglia<sup>3,4\*</sup>, Simona Mrakic-Sposta<sup>8</sup>.

<sup>1</sup>Department of Neurosciences, Biomedicine and Movement Sciences; University of Verona, Verona, Italy.

<sup>2</sup>Institute Research, Innovation and Enterprises, Area Science Park, Trieste, Italy.

<sup>3</sup>Dipartimento di Ingegneria e Architettura, Università degli studi di Trieste, Trieste, Italy.

<sup>4</sup>“Aldo Ravelli” Center for Neurotechnology and Experimental Brain Therapeutics, Department of Health Sciences, University of Milan, Via Antonio di Rudinì 8, 20142, Milan, MI, Italy.

<sup>5</sup>Foundation IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, 20122, Italy.

<sup>6</sup>ASST Santi Paolo e Carlo, University Hospital, Milan, 20142, Italy.

<sup>7</sup>Department of Oncology and Emato-Oncology, University of Milan, Milan 20122, Italy.

<sup>8</sup>Institute of Clinical Physiology, National Research Council (CNR), Milan, Italy.

<sup>9</sup>Associazione Sportiva Dilettantistica Alex Club, Federazione Italiana Pesistica (FIPE), Via Privata Mottarone, 2, 20155 Milano, Italy.

<sup>†</sup>Deceased

### **Corresponding Authors:**

Prof. Sara Marceglia, PhD

“Aldo Ravelli” Center for Neurotechnology and Experimental Brain Therapeutics, Department of Health Sciences, University of Milan, Via Antonio di Rudinì 8, 20142, Milan, MI, Italy.

E-mail: smarceglia@units.it; ORCID: 0000-0002-0456-866X

### **ABSTRACT**

**Background:** Transcranial direct current stimulation (tDCS) has been proposed as a non-invasive neuromodulatory strategy to enhance motor performance and modulate inflammatory responses. However, its effects on strength capacity and physiological stress markers during complex resistance exercises in elite athletes remain underexplored.

**Methods:** In this exploratory, randomized, sham-controlled pilot study, ten elite male weightlifters (n = 10; active tDCS = 5, sham = 5) received either a single 20-min session of anodal tDCS (2 mA) over the motor cortex or sham stimulation immediately before performing high-load back squats (85% 1RM). Outcomes included repetition number, barbell kinematics, Borg Rating of Perceived Exertion (Borg-RPE), Visual Analogue Scale (VAS) measures, and metabolic and oxy-inflammatory biomarkers (blood glucose, lactate, antioxidant capacity; salivary ROS; urinary IL-6, 8-iso-PGF2 $\alpha$ , 8-OH-

dG, creatinine, and neopterin). Given the exploratory design and small sample size, intervention effects were analyzed using Mann-Whitney U tests. For outcomes assessed both before and after stimulation, between-group comparisons were performed on change scores ( $\Delta = \text{post} - \text{pre}$ ). In contrast, mechanical exercise-related variables, which were only collected after stimulation, were directly compared between the active and sham groups.

**Results:** Athletes receiving active tDCS performed a significantly higher number of back squat repetitions and exhibited shorter lift times compared with the sham group ( $p = 0.024$  and  $p = 0.011$ , respectively). No significant between-group differences were observed in Borg-RPE, VAS-based subjective measures, or biochemical markers (all  $p > 0.05$ ). No significant differences were found for the remaining mechanical parameters (all  $p > 0.05$ ).

**Conclusions:** A single session of anodal tDCS applied before high-load back squat exercise may be associated with improvements in repetition capacity and lifting efficiency in elite weightlifters. No significant effects were observed on perceived exertion, biomechanical stability measures, or metabolic, oxidative, inflammatory, and renal biomarkers. These preliminary findings suggest that tDCS may represent a promising neuromodulatory adjunct to strength training, warranting confirmation in larger and methodologically powered studies.

**Trial registration**

None (pilot study).

**Keywords:** non-invasive brain stimulation, oxidative stress, inflammation, transcranial direct current stimulation, back squat, athletes, strength and conditioning.

## **Background**

Muscular strength is a complex and multifaceted construct [1] that encompasses both neurophysiological [2] and psychological factors [3]. In weightlifting, a discipline where performance is centered on achieving maximal force output in compound movements such as the squat, bench press, and deadlift, athletes are trained to progressively increase their one-repetition maximum (1RM) in multi-joint exercises [4]. This process demands a comprehensive approach that encompasses musculoskeletal components like motor control, inter- and intra-muscular coordination, lifting technique, and psychological dimensions like emotion regulation, motivation, and attentional control [5]. Accordingly, weightlifting training can be conceptualized as a means to express the maximum strength possible, prompting the exploration of novel training strategies and tools. Within this performance context, exercise-induced imbalances in oxidative stress and inflammation have also been recognized as relevant physiological concerns, given their interdependence and potential for long-term detrimental effects [6,7]. An increase in oxidative stress (OxS) has been observed following both aerobic and anaerobic exercise of sufficient intensity [8,9], with the magnitude of the response being dependent on

exercise type, duration, and load [10,11]. OxS arises from an imbalance between reactive oxygen species (ROS) and the body's antioxidant defense systems. Numerous studies have demonstrated that acute exercise leads to elevated ROS production [12-14], thereby intensifying oxidative pressure on cellular structures [15,16]. When excessive, this stress may impair physical performance and hinder skeletal muscle recovery [17] by altering contractile and regulatory proteins and disrupting calcium handling [18]. These alterations result in reduced force production and increased fatigue, particularly at submaximal intensities [18]. Excessive ROS may impair mitochondrial efficiency through damage to respiratory chain components and membrane lipids [19]. This phenomenon manifests as higher oxygen consumption at a given workload, shorter time to exhaustion, and greater perceived exertion in closely repeated sessions [20]. Moreover, muscle damage has been associated with post-exercise elevations in interleukin-6, a response with practical implications for recovery and subsequent performance [21]. Considering these mechanisms, quantifying biomarkers of oxidative stress and inflammation is essential, including ROS, 8-iso PGF<sub>2</sub> $\alpha$ , 8-OH-dG, and IL-6, which reflect oxidative cascades, lipid peroxidation, DNA damage, and muscle inflammation, respectively. Alongside these biochemical considerations, transcranial direct current stimulation (tDCS) has shown promising results as a non-invasive neuromodulation technique in sports neuroscience to enhance sport performance [22]. tDCS involves the application of low-intensity direct

current to the scalp to modulate cortical excitability [23]. Typically, anodal stimulation (atDCS) increases excitability, while cathodal stimulation (ctDCS) decreases it [24]. In particular, one of the most used electrode placements in tDCS involves an active electrode over the brain region of interest (e.g., motor cortex, M1) and a reference electrode often over a distant anatomical area (i.e., extracranial, such as the shoulder or deltoid) to complete the electrical circuit [25]. This approach has demonstrated the ability to make the stimulation more focused on the target area, avoiding "unwanted" effects due to the proximity of the electrical fields and possibly affecting subcortical circuits [25]. Emerging evidence suggests that tDCS may influence neuroinflammatory processes by modulating microglial activation and cytokine expression, potentially reducing pro-inflammatory markers in both preclinical and clinical models [26–31]. It should be noted that the inflammatory processes described in neurological disorders differ substantially from the transient systemic inflammatory responses induced by high-intensity exercise. In the context of resistance exercise, inflammatory signaling is primarily driven by peripheral muscle stress and metabolic perturbations rather than by central neuroinflammatory mechanisms [32]. Although tDCS is primarily known for modulating cortical excitability [33], changes in central motor drive, perception of effort, and autonomic regulation may indirectly interact with systemic physiological responses during intense exercise [34]. Through these central mechanisms, tDCS could potentially influence performance outcomes while interacting

with metabolic and inflammatory processes associated with strenuous muscular activity. In the sports domain, despite heterogeneity in protocols and populations studied, increasing evidence supports the use of atDCS over the M1 to improve endurance, power performance, and explosive force in various disciplines such as cycling and ski jumping [35,36]. Additional studies have shown that atDCS can enhance muscle strength [37,38], improve cognitive function [39], and reduce both central and peripheral fatigue [35] and pain perception [40]. However, most investigations on tDCS and muscular performance have focused on time to exhaustion during isolated and simple tasks, such as elbow flexion [38,41], often overlooking more ecologically valid performance conditions that integrate cognitive, emotional, and motivational demands [41] - e.g., sport-specific [42], high-load conditions [43].

To date, few studies have examined the effects of tDCS on complex multi-joint strength performance while simultaneously assessing performance outcomes together with physiological responses in elite athletes. In the present study, we explored the impact of monopolar atDCS applied bilaterally over the M1, compared to sham stimulation, on athletic performance during high-load back squat (BS) exercises in elite male weightlifters, with a specific focus on the physio-biological effects of the intervention in terms of metabolic (glucose, lactate), oxidative-inflammatory (ROS, total antioxidant capacity, lipid peroxidation, DNA oxidation, IL-6), and renal biomarkers (creatinine, neopterin). The results are intended to

inform future studies aimed at investigating the effects of neurostimulation targeting motor regions on performance, as well as on inflammatory and neurobiological profiles.

## **Methods**

### *Study design and Setting*

In this prospective, sham-controlled, exploratory pilot clinical study, 10 elite weightlifting athletes were randomly assigned to either an active anodal tDCS group (active tDCS,  $n = 5$ ) or a sham stimulation group (sham tDCS,  $n = 5$ ) (Fig.1). Each participant, blinded to the treatment group, received a single 20-minute session of tDCS (active or sham) immediately prior to performing a high-load BS exercise. Performance was evaluated by measuring the total number of BS repetitions completed, alongside a set of biomechanical parameters, including gyroscope-based oscillation analysis, angular momentum energy estimation, oscillation count during the lift phase, and horizontal sway as tracked by a motion capture camera. Subjective perceptions of effort and well-being were assessed using the Borg Rating of Perceived Exertion (Borg-RPE) and a Visual Analogue Scale (VAS), respectively. Additionally, physiological responses were assessed through blood, saliva, and urine biomarkers, collected at two timepoints: immediately before (T0, throughout the text referred to as “pre”) and after (T1, throughout the text referred to as “post”) the BS exercise. Assessors were not blinded to participant allocation. This study was designed as an exploratory pilot trial aimed at assessing feasibility and estimating

preliminary effect sizes rather than providing confirmatory hypothesis testing. Given the limited sample size, statistical analyses were considered exploratory, and p-values were interpreted descriptively rather than as confirmatory evidence of population-level effects. In this sense, analyses could inform future power calculations and lay the foundations for further, powered and more methodologically solid studies. All procedures were conducted in accordance with the Declaration of Helsinki and were approved by the Ethics Committee of the IRCCS Ca' Granda Foundation—Maggiore Policlinico Hospital of Milan (approval number J274-2022). All subjects gave written informed consent before participation.

### *Participants*

Participants were recruited via invitation from the Milan section of the Italian Weightlifting Federation (Federazione Italiana Pesistica - FIPE). Inclusion criteria were: (I) age  $\geq 18$  years; (II) official registration with FIPE as “expert” athletes in the “Senior” category across different weight classes [44]; (III) ability to understand and perform the study procedures; and (IV) capacity to provide written informed consent. Exclusion criteria included: (I) history or presence of major neurological, neuropsychological, or psychiatric disorders, as determined by clinical history; and (II) the presence of implanted medical devices such as pacemakers, intracranial metallic components, or spinal cord stimulators. To minimize potential confounding effects of prior physical exertion, all athletes were instructed

to abstain from exercise for at least 24 hours prior to the study session, following the guidance of a certified FIPE trainer (A.B.).

### *tDCS protocol*

tDCS, either active or sham, was administered in a single session immediately prior to the BS exercise. Stimulation was delivered using a constant-current stimulator (HDCStim, Newronika, Italy) connected to silicone rubber electrodes (1 mm thickness) with an area of 35 cm<sup>2</sup> (7 × 5 cm) for the anodes and 48 cm<sup>2</sup> (8 × 6 cm) for the cathode. The anodal electrodes were positioned bilaterally over M1 (corresponding to C3 and C4 according to the international 10–20 EEG system), while the cathode was placed over the right deltoid muscle [25]. Indeed, as demonstrated in previous literature, this montage induces more focal and deeper stimulation [25]. Conductive gel was applied to ensure stable and low-impedance electrode–skin contact throughout the stimulation. For the active condition, direct current was applied at 2 mA for 20 minutes, with a 30-second ramp-up and ramp-down period. The resulting charge density (68.57 mC/cm<sup>2</sup>) was within established safety limits [45,46]. In the sham condition, the device delivered only the initial 30-second ramp-up followed by a ramp-down, mimicking the somatosensory sensations of active stimulation without inducing lasting neuromodulatory effects [47], thereby maintaining participant blinding [48]. To verify blinding, subjects were asked at the end of the session whether they thought to have received active or sham stimulation.

### *BS task*

Participants were instructed to perform the maximum number of repetitions of the BS exercise with a barbell loaded at 85% of their 1RM, as assessed before the experiment. The exercise involved lowering the body with the barbell positioned on the upper back, between the seventh cervical vertebra and the scapular spine, until reaching a squat depth where the tops of the thighs were parallel to the floor ("parallel" position). Execution of the high-load BS required the athlete to control multiple performance-related variables, including barbell trajectory, joint angles at the knees and ankles, postural adjustments, foot positioning, and ground contact. This complex motor task allowed for the integration of both neuromuscular and subjective (cognitive-emotional) factors involved in strength execution. To prevent performance from being influenced by habitual training routines, the actual barbell load was concealed using opaque covers. This approach encouraged athletes to rely on perceived exertion rather than prior expectations about the load, thereby promoting a more authentic maximal-effort performance. Movement analysis focused on capturing barbell trajectory and detecting imbalances throughout the lift. A visual motion tracking system was used to assess lateral sway (Fig.2), and to enhance motion analysis, inertial measurement units (IMUs) were affixed to the barbell to measure rotational dynamics along its central axis (Fig.2). Further details on biomechanical outcome measures are provided in the biomechanical outcomes.

### *Clinical outcomes*

Perceived exertion was assessed using the Borg Rating of Perceived Exertion (Borg-RPE), a 15-point scale that includes standardized verbal descriptors to quantify subjective effort, breathlessness, and fatigue across a range of physical tasks and individuals [49].

Subjective symptoms were further evaluated using a 10-centimeter Visual Analogue Scale (VAS), in which the left anchor (score = 0) represented the absence of a symptom, and the right anchor (score = 10) represented the most intense symptom imaginable [50]. Participants were asked to indicate the intensity of their perceived symptoms at the given moment by marking a point along the line. VAS assessments included mood, general wellness (e.g., happy/unhappy, rest/tired), general sensations (e.g., hot/cold, calm/agitated), and pain (with anatomical location specified).

#### *Biochemical outcomes*

Biochemical outcomes were obtained through capillary blood, saliva, and urine samples. Capillary blood was collected from a fingertip for the determination of blood lactate concentration, glycemia, and total antioxidant capacity (TAC). Saliva was collected in Salivette devices (Sarstedt, Nümbrecht, Germany). Urine samples were collected by voluntary voiding in a sterile container. All saliva and urine samples were stored in multiple aliquots at -20°C until assayed and thawed only once before analysis, which was performed within two weeks of collection.

- *Metabolic markers*: Blood lactate concentration ([La]b) was assessed using an enzymatic method (Lactate Scout 4; EKF Diagnostic for Life, Germany)

with 0.2  $\mu\text{L}$  of capillary blood collected from the fingertip. Blood glucose levels were measured using an electronic glucometer with Accu-Chek Active test strips (Roche Diagnostics) on a fingertip blood sample. Each test was conducted in duplicate for all participants [13].

- *ROS production, antioxidant capacity, and oxidative damage:* ROS production rate was assessed in saliva by Electron Paramagnetic Resonance spectroscopy (EPR), operating X-band (E-Scan- Bruker BioSpin, GmbH, MA USA) as previously described [15,51]. Sample temperatures were stabilized at 37 °C by a temperature controller ("Bio III" unit, Noxygen Science Transfer & Diagnostics GmbH, Germany), interfaced with E-Scan, and CMH spin trapping molecules (also referred to as probes), 1-hydroxy-3-methoxycarbonyl-2,2,5,5-tetramethylpyrrolidine (CMH, Noxygen Science Transfer & Diagnostics, Germany) were used. The EPR signal reflects the number of unpaired electrons and can be translated into absolute production rates ( $\mu\text{mol}\cdot\text{min}^{-1}$ ). The stable CP (3-Carboxy-2,2,5,5-tetramethyl-1-pyrrolidinyloxy) radical was measured independently and served as a reference standard. TAC was measured by a redox sensor using a commercial EDEL potentiostat electrochemical analyser (Edel Therapeutics, Switzerland) in a three-electrode arrangement, in 10  $\mu\text{L}$  of capillary blood [52,53].

- *Oxidative damage and inflammatory markers:* The levels of 8-isoprostane (8-iso-PGF $2\alpha$ ), 8-hydroxy-2'-deoxyguanosine (8-OH-dG), and interleukin-6 (IL-6) were determined in urine using ELISA assays (Enzyme-Linked

Immunosorbent Assay). All determinations were carried out using a microplate reader spectrophotometer (InfiniteM200, Tecan, Austria). All the assessments were duplicated, and the interassay coefficient of variation was within the range indicated by the kit's manufacturer. Briefly, lipid peroxidation and DNA damage were assessed using commercial kits (Cayman Chemical, Ann Arbor, Michigan, USA, Item No. 516351 and Item No. 89320 respectively) as previously described [13,54]; samples were read at a wavelength of 412 nm, and concentration were determined using standard curves as previously described. The coefficient of variation (CV) was indicated by the manufacturer: inter-assay CV 11.2% and 10.7% and intra-assay CV 14.6% and 11.6%, respectively, for 8-iso-PGF $2\alpha$  and 8-OH-dG [53]. IL-6 in urine (IL-6-U) was determined by Human IL-6 Immunoassay (Fine Test, Wuhan, China, Cat No.: EH0201) as previously described [13,54]. The samples' concentrations were determined at 450 nm. The CV was indicated by the manufacturer: inter-assay CV 4.62%, and intra-assay CV 5.35%; urinary lipid peroxidation was assessed by competitive immunoassay of 8-iso-PGF $2\alpha$  (Cayman Chemical, USA) as previously described [13,54]; 8-OH-dG was quantified using a commercially available enzyme-linked immunosorbent assay (EIA) kit (Cayman Chemical, Ann Arbor, MI, USA, Item No.589320). The sample 8-OH-dG concentration was determined using a standard curve as previously described [53]. Neopterin concentrations were measured in duplicate using an isocratic high-pressure liquid

chromatography (HPLC) method. The calibration curves were linear for creatinine (1.25–10 mmol/L), and neopterin (0.125–1  $\mu$ mol/L). The inter-assay and intra-assay variation coefficients were below 5%. The methods have been previously described [53].

- *Renal function*: Bilirubin, urobilinogen, ketones, blood proteins, pH, and leukocytes, as well as specific gravity/density, were assessed semi-quantitatively using Urine Test Strips (Combi screen 11sys PLUS, GIMA, Gessate, Milan, Italy) in duplicate for each subject [13]. Creatinine assessment followed the one previously described for neopterin.

#### *Biomechanical outcomes*

Biomechanical variables and analyses were customized for this pilot study. Briefly, to quantify rotational instability of the barbell, we employed a differential signal analysis from the two IMUs units mounted on the barbell (Fig. 3). This allowed us to minimize structural deformation artifacts and isolate true instability in barbell rotation. As shown in Fig. 3, the difference between the two-sensor signal was calculated to attenuate the deformations of the barbell and to amplify the actual balancing act on the barbell to characterize the gesture of the athlete.

The signal was band-pass filtered (0.5–4 Hz) to remove noise related to gravity or high-frequency peaks. Within the concentric phase, two measures were computed: I) the angular momentum energy, normalized to the weight lifted, to estimate the magnitude of oscillation; and II) the number of oscillations during the lift phase, calculated via peak detection, as a proxy

for motion regularity and control. The latter was particularly useful for identifying instability during the initial concentric stage. To count the number of oscillations, a simple peak detection algorithm was applied on the post-processed signal, using the total number of peaks detected as an oscillation estimate (Fig. 4).

In addition, the horizontal sway of the barbell was assessed from the x-axis displacement in the lateral video recordings. This metric was normalized to each athlete's lifted weight. Since all athletes started from the same position and camera setup remained fixed, we used pixel displacement in the vertical (y) axis as a consistent reference. To extract the sway feature we calculated the maximum extensions of the horizontal offset during each single phase (Fig. 5). These measurements were made for all the repetitions during the exercise of the athletes. To better account for the effects of fatigue on motor control, repetitions were further classified based on their velocity. For each athlete, the mean duration of the eccentric-concentric cycle was used as a threshold. Repetitions faster than the mean were labeled as "fast" (typically occurring at the beginning of the set), while slower repetitions (generally appearing later as fatigue accumulated) were categorized as "slow." This classification allowed us to explore changes in movement quality across the course of the task.

### *Statistical analysis*

Data distribution was assessed using the Shapiro-Wilk test. Given the small sample size ( $n = 5$  per group) and violations of normality in several

variables, non-parametric tests were adopted throughout the analysis. Between-group comparisons were performed using Mann-Whitney U tests, with effect size expressed as rank-biserial correlation where applicable. For variables assessed both before and after stimulation, including clinical outcomes, biochemical outcomes, and total repetition number, between-group comparisons were conducted on change scores ( $\Delta = \text{post} - \text{pre}$ ) to account for baseline interindividual variability. For the remaining mechanical exercise-related variables, comparisons were performed on post-stimulation values only (Active vs Sham), because the maximal exercise task could not be repeated before and after stimulation. Descriptive statistics are reported as mean  $\pm$  standard deviation (SD). Statistical significance was set at  $p < 0.05$  (two-tailed). All statistical analyses were performed using Jamovi (version 2.6).

## **Results**

Ten male weightlifting athletes were included (age  $27.3 \pm 5.27$  years). All of them were expert weightlifters based on training history ( $6.0 \pm 5.5$  years) and high relative load performance ( $\sim 85\%$  1RM;  $174.05 \pm 22.28$  kg).

Demographic and anthropometric characteristics of the sample are summarized in Table 1. Anthropometric parameters were obtained using bipolar bioelectrical impedance analysis (TBF-300A Body Composition Analyzer; Tanita Corporation, Arlington Heights, IL, USA), a method that may present some limitations in the assessment of body composition [55]. All participants were in good health and reported no history of smoking.

Additionally, the training history of each athlete was recorded to better contextualize individual performance profiles. When participants were asked to identify their treatment condition in order to assess the success of blinding, they were unable to determine which study arm they had been assigned to.

Descriptive statistics, together with Mann-Whitney U test results, are reported in Table 2. The table specifies whether each comparison was conducted on change scores ( $\Delta = \text{post} - \text{pre}$ ) or on post-stimulation values only. No significant differences were observed between the active tDCS and sham groups for perceived exertion (Borg-RPE) or for any VAS dimension, including well-being, tiredness, heat, restlessness, happiness, and pain (all  $p > 0.05$ ). For the mechanical exercise-related variables (total repetitions, lift time, fast repetitions, slow repetitions, angular momentum energy, rotational oscillations in the concentric phase, and normalized horizontal sway), significant between-group differences were detected for total repetitions ( $p = 0.024$ ) and lift time ( $p = 0.011$ ), whereas no significant differences were observed for the remaining mechanical variables (all  $p > 0.05$ ). Similarly, no significant between-group differences were observed for blood glucose, blood lactate, ROS production, total antioxidant capacity, 8-iso-PGF $2\alpha$ , 8-OH-dG, urinary IL-6, creatinine, neopterin, bilirubin, urobilinogen, ketones, erythrocytes, urinary pH, leukocytes, or specific gravity (for all,  $p > 0.05$ ).

## **Discussion**

In this randomized, sham-controlled, exploratory study, we explored the acute effects of atDCS on biomechanical, clinical, and biochemical responses during high-load BS performance in elite weightlifting athletes. Athletes who received active stimulation demonstrated a significant increase in the total number of repetitions, performed at a faster pace, compared to the sham group. No differential effect emerged between the fast- and slow-repetition subgroups, suggesting an overall improvement in task performance under the tested condition. These findings support the hypothesis that a single session of anodal tDCS may enhance both physical performance and motor control during high-intensity resistance exercise. However, despite the higher performance observed in the active stimulation group, no measurable differences in perceived exertion or exercise-related oxidative/inflammatory biomarkers were detected in the present pilot sample, with potential implications for neuromodulatory strategies in elite training protocols.

To our knowledge, this is the first study to investigate the interaction between tDCS and repeated high-load BS using a multimodal assessment approach, including micro-invasive sampling of capillary blood, saliva, and urine to evaluate systemic oxy-inflammatory responses. This methodology allows for a more comprehensive characterization of physiological adaptations to acute neuromodulation in elite athletes, a population where physiological precision and training efficiency are critical [54,56–58].

From a physical performance perspective, athletes who received active stimulation performed a higher number of repetitions with reduced lift time. While no significant differences emerged in fast repetitions, slow repetitions, angular momentum energy, rotational oscillations in the concentric phase, or normalized horizontal sway, the increased total performance supports the idea that tDCS can facilitate more efficient neuromuscular activation during complex strength tasks [59], even in the absence of overt changes in conventional biomechanical parameters. Previous research has demonstrated that anodal tDCS applied over the M1 can increase force generation, enhancing performance in maximal voluntary contraction tasks [60] and improving motor control [61-63]. These effects are thought to arise from increased corticospinal tract excitability [64,65], ranging from cortical neuron activation [66] to motor unit recruitment [67], ultimately strengthening the descending drive from cortex to muscle and resulting in more efficient muscle activation [68]. In parallel, atDCS has been shown to promote more synchronized activation of agonist muscles and more effective inhibition of antagonist muscles [66]. Nonetheless, its effects on muscular endurance remain inconclusive [60]. Furthermore, tDCS may reduce perceived exertion [59,69] and delay neuromuscular fatigue, thereby supporting performance gains in both endurance and strength domains [61-63]. Interestingly, our results occurred in absence of differences in between-group comparisons of perceived exertion (Borg-RPE). This suggests that, despite similar perceived effort, the active tDCS

group achieved greater performance. These findings are consistent with a randomized controlled trial by Dos Santos et al. (2023), which demonstrated that motor anodal tDCS acutely improved 5,000 m running performance, yielding faster times and higher average velocity compared with sham, without changes in subjective RPE [70]. Such effects may arise because tDCS might act primarily through increased neural excitability and motor unit recruitment rather than through sensory or cognitive pathways underlying the perception of effort [71].

Biochemically, no significant between-group differences emerged in metabolic markers (blood lactate and blood glucose), oxidative stress parameters (ROS production and TAC), inflammatory markers (8-iso-PGF2 $\alpha$ , 8-OH-dG, IL-6), or indices of renal function (creatinine, bilirubin, urobilinogen, ketones, blood proteins, pH, leukocytes, and specific gravity/density). For most of these biomarkers, direct comparison with the existing literature is challenging, as no prior studies have specifically investigated these outcomes in relation to tDCS; therefore, conclusions can only be drawn indirectly from broader investigations. To date, no evidence has been reported regarding a potential direct effect of tDCS on renal function or on inflammatory biomarkers such as 8-iso-PGF2 $\alpha$  and 8-OH-dG. However, a study conducted in healthy adults with obesity undergoing anodal tDCS applied to the dorsolateral prefrontal cortex for three consecutive mornings demonstrated a stimulation-induced reduction in plasma IL-6 concentrations compared with the sham condition [72]. These

results are only partially comparable, given the differences in both population and stimulation protocol. Nevertheless, they provide useful insight into the potential biochemical effects of tDCS, considering that the significant increase in physical effort occurred with no significant increase in inflammatory response between groups. The available literature investigating the anti-inflammatory effects of stimulation through peripheral biomarkers has primarily focused on clinical populations rather than healthy individuals. For example, studies conducted in knee osteoarthritis [73] and stroke [74] have assessed inflammatory cytokines such as IL-6, TNF- $\alpha$ , and IL-10, although the results remain inconclusive. Similarly, the literature on tDCS and oxidative stress largely concerns clinical populations or animal models and suggests that tDCS may modulate oxidative stress pathways [26]. For instance, a randomized trial in stroke survivors demonstrated that tDCS combined with physiotherapy increased serum superoxide dismutase, an antioxidant enzyme involved in scavenging oxygen radicals [26]. In addition, Li et al. [75] showed that anodal tDCS applied over the frontal cortex of mice reduced levels of malondialdehyde, a key marker of lipid peroxidation [76], potentially through a tDCS-induced upregulation of brain-derived neurotrophic factor (BDNF), which is known to exert antioxidant effects [77].

In contrast, metabolic markers have been more extensively investigated. A crossover study showed that a single session of anodal tDCS promoted systemic glucose uptake and improved systemic glucose tolerance as

measured with the euglycemic-hyperinsulinemic clamp [78]. A similar effect was observed when tDCS was applied twice in series [79]. In addition, in healthy subjects, stimulation resulted in blood glucose concentrations that were distinctly lower under the tDCS condition compared with sham as early as day 1, with persistence of the effect at day 8 [80]. However, in all these studies the experimental conditions did not include any form of athletic performance or physical exertion, unlike in the present study. This warrants particular caution in making comparisons, as blood glucose is known to increase during exercise as a well-established acute physiological response to increased energy demand [81]. Regarding blood lactate, healthy individuals have been assessed primarily following sport-related efforts. In one study, anodal stimulation over the left dorsolateral prefrontal cortex increased time to exhaustion during cycling and, at exhaustion, was associated with significantly higher blood lactate accumulation compared with the sham condition [82]. Furthermore, a systematic review conducted in healthy adults concluded that, during cycling protocols, tDCS was associated in some studies with increased blood lactate accumulation, together with improvements in time to task failure and reductions in perceived exertion [83]. However, findings regarding lactate are not consistent. A 2024 triple-blind crossover study in recreational runners reported no differences between anodal tDCS and sham in blood lactate, nor in torque, motor-evoked potentials (MEP), or perceived exertion [84]. These observations suggest that, when lactate changes are observed, they

may primarily reflect greater tolerance to effort or differences in task-related motor dynamics rather than a direct and robust effect of tDCS on the systemic regulation of lactate under resting conditions [83].

This study has several limitations. First, the small, unstratified, all-male sample size, although justified by the elite status of the athletes, limits statistical power and generalizability. In addition, the absence of blind assessors may have increased the risk of biased assessments. Second, the study examined only the acute effects of a single tDCS session, without assessing potential long-term or cumulative adaptations (e.g., tolerance or chronic adaptations), and lacked direct neurophysiological measures (e.g., motor-evoked potentials, EEG, TMS-derived metrics) to clarify the underlying cortical mechanisms. Third, potential confounding factors such as nutritional intake, sleep patterns, and circadian timing were not standardized, and the timing of biomarker assessment (“immediately post” exercise) may have missed peak fluctuations in IL-6 and oxidative stress responses.

## **Conclusion**

This randomized, sham-controlled, exploratory study provides preliminary evidence that a single session of anodal tDCS, applied prior to high-load BS exercise, may enhance neuromuscular endurance and athletic performance, without detectable increases in the measured systemic oxidative and inflammatory biomarkers. In this small pilot sample, no significant between-group differences were detected in metabolic markers, oxidative stress

parameters, inflammatory markers, or indices of renal function. These findings highlight the potential of tDCS as a neuromodulatory adjunct to conventional strength training, particularly in elite athletic settings. Future studies involving larger samples and repeated stimulation protocols are warranted to confirm these results and to investigate the long-term benefits and mechanisms of action of tDCS in strength-based sports.

### **Abbreviations**

BS: back squat; 1RM: one-repetition maximum; tDCS: transcranial direct current stimulation; atDCS: anodal transcranial direct current stimulation; ctDCS: cathodal transcranial direct current stimulation; OxS: Oxidative Stress; ROS: Reactive Oxygen Species; FIPE: Federazione Italiana Pesistica, Italian Weightlifting Federation; IL-6-U: Interleukin 6 in Urine; 8-iso-PGF<sub>2</sub>α: 8-isoprostane concentration; HPLC: high-pressure liquid chromatography; Borg-RPE: BORG Rate of Perceived Exertion scale; VAS: Visual Analogue Scale

### **Declaration**

#### *Ethics approval and consent to participate*

The study was conducted in accordance with the Declaration of Helsinki and approved by the institutional review board of the Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico di Milano (approval number J274-2022). Written informed consent was obtained from all participants prior to study enrolment.

#### *Consent for publication*

All participants provided written informed consent for the publication of potentially identifying images and/or personal or clinical information in this article.

#### *Availability of data and materials*

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

#### *Competing interests*

S.M. was partially supported by the European Union under the Italian National Recovery and Resilience Plan (NRRP) of NextGenerationEU, partnership on “Telecommunications of the Future” (PE00000001 - program “RESTART”). S.M., R.F. and A.P. are founders and shareholders of Newronika Spa, Italy. All the other authors declare no competing interests

#### *Funding*

This research received no external funding. The research was funded by internal funding from Institute of Clinical Physiology - Milan, National Research Council (IFC-CNR).

#### *Authors' contributions*

R.dD. contributed to Conceptualization, Methodology, Validation, Investigation, Writing - Original Draft. M.P. contributed to Conceptualization, Methodology, Software, Data Curation, Visualization, Writing - Original Draft. M.G. contributed to Data Curation, Formal analysis, Writing - Original Draft, Writing - Review & Editing. C.T. contributed to Data Curation, Formal analysis, Writing - Original Draft,

Writing - Review & Editing. R.F. contributed to Methodology and Data Curation. A.V. contributed to Methodology, Formal analysis. C. D. contributed to Methodology, Formal analysis. S. P. contributed to Formal analysis, Visualization. A.B. contributed to Validation and Investigation. A.P. contributed to Validation, Supervision, Writing - Review & Editing. S.M. contributed to Conceptualization, Software, Data Curation, Writing - Original Draft, Writing - Review & Editing. S.M.S. contributed to Conceptualization, Data Curation, Visualization, Writing - Original Draft, Supervision. All authors, except A.B. (deceased) read and approved the submitted version of the manuscript.

### *Acknowledgements*

The authors acknowledge the athletes of Weightlifting Italian Federation.

**Fig. 1.** Flowchart diagram depicting the flow of participants through study.

**Fig. 2.** Experiment setup and squat motion. A: Front view, B: Side view.

**Fig. 3.** Gyroscope sensor position on the barbell and signal deformation attenuation using the differential of the two recording units.

**Fig. 4:** Peak detection algorithm applied to the post-processed differential module of the two barbell gyroscopes in the subject 1, during the first lift phase. In this example there are 8 oscillations.

**Fig. 5.** Camera tracking trajectory sway segmentation and fast/slow repetitions segmentation example, A) Horizontal camera tracked barbell motion, B) Vertical camera tracked barbell motion.

**Table 1.** Demographic and anthropometric characteristics of the sample, divided by group.

<b>Active tDCS Group</b>										
Subject	Age	Kg	Height	Body fat	FFM	TBW	PMM	BMI	Repetitions number Basal assessment	Weights [kg] (85% of maximum)
1	26	144.8	177	40.9	85.5	62.4	81.3	46.22	5	178.5
2	36	94.8	190	14.5	80.2	57.6	76.3	26.26	9	161.5
4	21	88.7	184	12.7	76	59.2	72.2	26.2	7	135
7	22	76.4	165	11.8	64.6	60.2	61.8	28.06	5	210
8	34	108.4	190	14.7	93.7	58.6	89.1	30.03	5	187
<b>mean</b>	<b>27.8</b>	<b>102.6</b>	<b>181.2</b>	<b>22.6</b>	<b>80</b>	<b>56.1</b>	<b>76.1</b>	<b>31.35</b>	<b>6.2</b>	<b>174.4</b>
<b>SD</b>	<b>6.87</b>	<b>26.24</b>	<b>10.52</b>	<b>20.55</b>	<b>10.8</b>	<b>6.20</b>	<b>10.2</b>	<b>8.46</b>	<b>1.78</b>	<b>28.11</b>
		<b>2</b>			<b>5</b>	<b>6</b>	<b>4</b>	<b>4</b>		
<b>Sham tDCS Group</b>										
Subject	Age	Kg	Height	Body fat	FFM	TBW	PMM	BMI	Repetitions number Basal assessment	Weights [kg] (85% of maximum)
3	24	116.4	191	20.8	95.6	57	90.9	31.91	6	178.5
5	27	80.3	175	8.6	71.7	62	68.1	26.22	5	150
6	31	88.8	168	21.4	67.4	54.4	64.1	31.46	6	170
9	30	86.8	172	18.3	68.5	55.5	65.1	29.34	5	170
10	22	84.6	175	11.5	73.1	60.5	69.5	27.62	5	200
<b>mean</b>	<b>26.8</b>	<b>91.38</b>	<b>176.2</b>	<b>16.12</b>	<b>75.2</b>	<b>57.8</b>	<b>71.5</b>	<b>29.31</b>	<b>5.4</b>	<b>173.7</b>
<b>SD</b>	<b>3.83</b>	<b>14.34</b>	<b>8.76</b>	<b>5.75</b>	<b>11.6</b>	<b>3.26</b>	<b>11.0</b>	<b>2.44</b>	<b>0.55</b>	<b>18.05</b>
	<b>4</b>				<b>0</b>		<b>4</b>			

FFM = Fat-Free Mass; TBW = Total Body Water; PMM = Predicted Muscle Mass; BMI = Body Mass Index.

**Table 2.** Summary of the outcomes considered in the study.

Variable	Comparison metric	Active tDCS (mean $\pm$ SD)	Sham tDCS (mean $\pm$ SD)	U	p-value	Rank-biserial r
----------	-------------------	-----------------------------	---------------------------	---	---------	-----------------

<b>Clinical</b>						
Borg-RPE (perceived exertion)	$\Delta$ (post – pre)	16.80 $\square$ 2.48	17.80 $\square$ 1.30		> 0.05	
VAS well-being	$\Delta$ (post – pre)	3.06 $\square$ 1.79	3.9 $\square$ 1.62		> 0.05	
VAS tiredness	$\Delta$ (post – pre)	4.5 $\square$ 1.26	6.94 $\square$ 1.76		> 0.05	
VAS heat	$\Delta$ (post – pre)	1.64 $\square$ 2.22	1.76 $\square$ 1.12		> 0.05	
VAS restlessness	$\Delta$ (post – pre)	5.14 $\square$ 2.70	3.62 $\square$ 2.57		> 0.05	
VAS happiness	$\Delta$ (post – pre)	1.6 $\square$ 0.90	2 $\square$ 0		> 0.05	
VAS pain	$\Delta$ (post – pre)	2.2 $\square$ 2.70	1 $\square$ 1		> 0.05	
<b>Mechanical</b>						
Total repetitions (n)	$\Delta$ (post – pre)	2.40 $\pm$ 1.52	0.80 $\pm$ 0.45	2.00	<b>0.024*</b>	-0.84 0
Fast repetitions (n)	Post only	2.2 $\pm$ 0.4	2.6 $\pm$ 0.49		> 0.05	
Slow repetitions (n)	Post only	3 $\pm$ 0.63	5 $\pm$ 1.79		> 0.05	
Angular momentum energy ( $^{\circ}\text{s}^{(-1)} \text{Hz}^{(-0.5)}$ )	Post only	0.0048 $\pm$ 0.0071	0.0201 $\pm$ 0.0245		> 0.05	
Rotational oscillations in concentric phase (n)	Post only	10.53 $\pm$ 2.89	12.44 $\pm$ 3.64		> 0.05	
Lift time (sec)	Post only	1.66 $\pm$ 0.08	1.97 $\pm$ 0.12	1.00	<b>0.011*</b>	+0.92 0
Normalized horizontal sway	Post only	0.363 $\pm$ 0.175	0.435 $\pm$ 0.225		> 0.05	
<b>Biochemical</b>						
Blood glucose (mg/dL)	$\Delta$ (post – pre)	92.67 $\pm$ 10.79	81.75 $\pm$ 12.97		> 0.05	
Blood lactate ([La]b(mM))	$\Delta$ (post – pre)	9.26 $\pm$ 2.45	12.54 $\pm$ 2.82		> 0.05	
ROS production ( $\mu\text{mol}\cdot\text{min}^{-1}$ )	$\Delta$ (post – pre)	0.19 $\pm$ 0.01	0.20 $\pm$ 0.01		> 0.05	
Antioxidant capacity (TAC) (mM)	$\Delta$ (post – pre)	3.25 $\pm$ 0.13	3.42 $\pm$ 0.25		> 0.05	

8-iso-PGF2 $\alpha$ (ng·mg <sup>-1</sup> creatinine)	$\Delta$ (post – pre)	0.72 $\pm$ 0.40	0.73 $\pm$ 0.21		> 0.05	
8-OH-dG (ng/mL)	$\Delta$ (post – pre)	3.72 $\pm$ 1.11	4.81 $\pm$ 1.17		> 0.05	
Urinary IL-6 (pg. mL <sup>-1</sup> )	$\Delta$ (post – pre)	8.87 $\pm$ 3.30	7.64 $\pm$ 2.85		> 0.05	
Creatinine (g.L <sup>-1</sup> )	$\Delta$ (post – pre)	0.60 $\pm$ 0.10	0.66 $\pm$ 0.30		> 0.05	
Neopterin ( $\mu$ mol.mol <sup>-1</sup> creatinine)	$\Delta$ (post – pre)	277.60 $\pm$ 72.21	306.40 $\pm$ 108.20		> 0.05	
Bilirubin ( $\mu$ mol.L <sup>-1</sup> )	$\Delta$ (post – pre)	4.4 $\pm$ 2.5	3.2 $\pm$ 2.7		> 0.05	
Urobilinogen ( $\mu$ mol.L <sup>-1</sup> )	$\Delta$ (post – pre)	0.8 $\pm$ 0.7	0.3 $\pm$ 0.4		> 0.05	
Ketones (mmol.L <sup>-1</sup> )	$\Delta$ (post – pre)	4.0 $\pm$ 2.2	4.2 $\pm$ 3.2		> 0.05	
Erythrocytes (Ery. $\mu$ L <sup>-1</sup> )	$\Delta$ (post – pre)	3.6 $\pm$ 4.9	5.4 $\pm$ 4.9		> 0.05	
pH	$\Delta$ (post – pre)	6.6 $\pm$ 0.9	6.2 $\pm$ 0.8		> 0.05	
Leukocytes (Leuko. $\mu$ L <sup>-1</sup> )	$\Delta$ (post – pre)	3.0 $\pm$ 6.7	0.0 $\pm$ 0.0		> 0.05	
Specific gravity	$\Delta$ (post – pre)	1.02 $\pm$ 0.0	1.02 $\pm$ 0.0		> 0.05	

RPE = Rating of Perceived Exertion; VAS = Visual Analogue Scale; ROS = Reactive Oxygen Species; TAC = Total Antioxidant Capacity; 8-iso-PGF2 $\alpha$  = 8-iso-Prostaglandin F2 $\alpha$ ; 8-OH-dG = 8-hydroxy-2'-deoxyguanosine; IL-6 = Interleukin-6; pH = potential of Hydrogen. Values are reported as mean  $\pm$  standard deviation. Between-group comparisons were performed using Mann-Whitney U tests. For variables assessed both before and after stimulation, including clinical outcomes, biochemical outcomes, and total repetitions, analyses were conducted on change scores ( $\Delta$  = post – pre). For the remaining mechanical exercise-related variables, analyses were conducted on post-stimulation values only, as the maximal exercise task could not be repeated before and after stimulation. Mann-Whitney U statistics and rank-biserial correlations are reported for statistically significant between-group comparisons. Statistically significant differences are indicated in bold and with an asterisk (\*p < 0.05).

## References

- [1] Frontera WR, Meredith CN, O'Reilly KP, Knuttgen HG, Evans WJ. Strength conditioning in older men: skeletal muscle hypertrophy and improved function.

- Journal of Applied Physiology 1988;64:1038-44.  
<https://doi.org/10.1152/jappl.1988.64.3.1038>.
- [2] Kraemer WJ, Looney DP. Underlying Mechanisms and Physiology of Muscular Power. *Strength & Conditioning Journal* 2012;34:13.  
<https://doi.org/10.1519/SSC.0b013e318270616d>.
- [3] Lochbaum M, Stoner E, Hefner T, Cooper S, Lane AM, Terry PC. Sport psychology and performance meta-analyses: A systematic review of the literature. *PLOS ONE* 2022;17:e0263408. <https://doi.org/10.1371/journal.pone.0263408>.
- [4] Ferland P-M, Comtois AS. Classic Powerlifting Performance: A Systematic Review. *The Journal of Strength & Conditioning Research* 2019;33:S194.  
<https://doi.org/10.1519/JSC.0000000000003099>.
- [5] Park R. The Mentality of Strength: A Study on the Effects of Grit, Self-Control, and Conscientiousness on Rate of Progress in Powerlifting. Senior Independent Study Theses 2017.
- [6] Forrester SJ, Kikuchi DS, Hernandez MS, Xu Q, Griendling KK. Reactive Oxygen Species in Metabolic and Inflammatory Signaling. *Circ Res* 2018;122:877-902.  
<https://doi.org/10.1161/CIRCRESAHA.117.311401>.
- [7] Powers SK, Ji LL, Kavazis AN, Jackson MJ. REACTIVE OXYGEN SPECIES: IMPACT ON SKELETAL MUSCLE. *Compr Physiol* 2011;1:941-69.  
<https://doi.org/10.1002/cphy.c100054>.
- [8] Bloomer RJ, Goldfarb AH. Anaerobic exercise and oxidative stress: a review. *Can J Appl Physiol* 2004;29:245-63. <https://doi.org/10.1139/h04-017>.
- [9] König D, Wagner KH, Elmalfa I, Berg A. Exercise and oxidative stress: significance of antioxidants with reference to inflammatory, muscular, and systemic stress. *Exerc Immunol Rev* 2001;7:108-33.
- [10] Gussoni M, Moretti S, Vezzoli A, Genitoni V, Giardini G, Balestra C, et al. Effects of Electrical Stimulation on Delayed Onset Muscle Soreness (DOMS): Evidences from Laboratory and In-Field Studies. *JFMK* 2023;8:146.  
<https://doi.org/10.3390/jfink8040146>.
- [11] Steinbacher P, Eckl P. Impact of Oxidative Stress on Exercising Skeletal Muscle. *Biomolecules* 2015;5:356-77. <https://doi.org/10.3390/biom5020356>.
- [12] Hudson MB, Hosick PA, McCaulley GO, Schrieber L, Wrieden J, McAnulty SR, et al. The effect of resistance exercise on humoral markers of oxidative stress. *Med Sci Sports Exerc* 2008;40:542-8.  
<https://doi.org/10.1249/MSS.0b013e31815daf89>.
- [13] Mrakic-Sposta S, Gussoni M, Moretti S, Pratali L, Giardini G, Tacchini P, et al. Effects of Mountain Ultra-Marathon Running on ROS Production and Oxidative Damage by Micro-Invasive Analytic Techniques. *PLOS ONE* 2015;10:e0141780.  
<https://doi.org/10.1371/journal.pone.0141780>.
- [14] Radak Z, Zhao Z, Koltai E, Ohno H, Atalay M. Oxygen Consumption and Usage During Physical Exercise: The Balance Between Oxidative Stress and ROS-Dependent Adaptive Signaling. *Antioxidants & Redox Signaling* 2013;18:1208-46. <https://doi.org/10.1089/ars.2011.4498>.
- [15] Mrakic-Sposta S, Gussoni M, Vezzoli A, Dellanoce C, Comassi M, Giardini G, et al. Acute Effects of Triathlon Race on Oxidative Stress Biomarkers. *Oxidative Medicine and Cellular Longevity* 2020;2020:e3062807.  
<https://doi.org/10.1155/2020/3062807>.
- [16] Vezzoli A, Dellanoce C, Mrakic-Sposta S, Montorsi M, Moretti S, Tonini A, et al. Oxidative Stress Assessment in Response to Ultraendurance Exercise: Thiols Redox Status and ROS Production according to Duration of a Competitive Race.

- Oxidative Medicine and Cellular Longevity 2016;2016:e6439037.  
<https://doi.org/10.1155/2016/6439037>.
- [17] Sunemi SM, Silva FA, Antonio EL, Tucci PJF, Serra AJ. Photobiomodulation: Newly Discovered Actions in Resistance Exercise. *Reactive Oxygen Species* 2019;7:148-53.
- [18] Lamb GD, Westerblad H. Acute effects of reactive oxygen and nitrogen species on the contractile function of skeletal muscle. *The Journal of Physiology* 2011;589:2119-27. <https://doi.org/10.1113/jphysiol.2010.199059>.
- [19] Steinbacher P, Eckl P. Impact of Oxidative Stress on Exercising Skeletal Muscle. *Biomolecules* 2015;5:356-77. <https://doi.org/10.3390/biom5020356>.
- [20] Mrakic-Sposta S, Gussoni M, Porcelli S, Pugliese L, Pavei G, Bellistri G, et al. Training Effects on ROS Production Determined by Electron Paramagnetic Resonance in Master Swimmers. *Oxidative Medicine and Cellular Longevity* 2015;2015:804794. <https://doi.org/10.1155/2015/804794>.
- [21] Pedersen BK, Steensberg A, Schjerling P. Muscle-derived interleukin-6: possible biological effects. *The Journal of Physiology* 2001;536:329-37. <https://doi.org/10.1111/j.1469-7793.2001.0329c.xd>.
- [22] Moreira A, da Silva Machado DG, Moscaleski LA, Baptista AF, Li LM, Morya E, et al. tDCS in Exercise, Sport Performance, and Recovery Process. In: Brunoni AR, Nitsche MA, Loo CK, editors. *Transcranial Direct Current Stimulation in Neuropsychiatric Disorders: Clinical Principles and Management*, Cham: Springer International Publishing; 2021, p. 413-32. [https://doi.org/10.1007/978-3-030-76136-3\\_18](https://doi.org/10.1007/978-3-030-76136-3_18).
- [23] Maiorana N, Guidetti M, Dini M, Priori A, Ferrucci R. Cerebellar tDCS as Therapy for Cerebellar Ataxias. *Cerebellum* 2022;21:755-61. <https://doi.org/10.1007/s12311-021-01357-1>.
- [24] Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *Journal of Physiology* 2000;527:633-9. <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x>.
- [25] Guidetti M, Maria Bianchi A, Parazzini M, Maiorana N, Bonato M, Ferrara R, et al. Monopolar tDCS might affect brainstem reflexes: A computational and neurophysiological study. *Clinical Neurophysiology* 2023;155:44-54. <https://doi.org/10.1016/j.clinph.2023.08.011>.
- [26] Guidetti M, Bertini A, Pirone F, Sala G, Signorelli P, Ferrarese C, et al. Neuroprotection and Non-Invasive Brain Stimulation: Facts or Fiction? *International Journal of Molecular Sciences* 2022, Vol 23, Page 13775 2022;23:13775. <https://doi.org/10.3390/IJMS232213775>.
- [27] Guo T, Fang J, Tong ZY, He S, Luo Y. Transcranial Direct Current Stimulation Ameliorates Cognitive Impairment via Modulating Oxidative Stress, Inflammation, and Autophagy in a Rat Model of Vascular Dementia. *Frontiers in Neuroscience* 2020;14:28. <https://doi.org/10.3389/FNINS.2020.00028/BIBTEX>.
- [28] Zhang K, Guo L, Zhang J, Rui G, An G, Zhou Y, et al. tDCS Accelerates the Rehabilitation of MCAO-Induced Motor Function Deficits via Neurogenesis Modulated by the Notch1 Signaling Pathway. *Neurorehabilitation and Neural Repair* 2020;34:640-51. <https://doi.org/10.1177/1545968320925474>.
- [29] Regner GG, Torres ILS, de Oliveira C, Pflüger P, da Silva LS, Scarabelot VL, et al. Transcranial direct current stimulation (tDCS) affects neuroinflammation parameters and behavioral seizure activity in pentylentetrazole-induced kindling in rats. *Neuroscience Letters* 2020;735:135162. <https://doi.org/10.1016/J.NEULET.2020.135162>.

- [30] Zhang H, Zheng X, Zhang B. Anti-Inflammatory Pathways Mediating tDCS's Effects on Neuropathic Pain. *Biology (Basel)* 2025;14:892. <https://doi.org/10.3390/biology14070892>.
- [31] Straudi S, Antonioni A, Baroni A, Bonsangue V, Lavezzi S, Koch G, et al. Anti-Inflammatory and Cortical Responses after Transcranial Direct Current Stimulation in Disorders of Consciousness: An Exploratory Study. *J Clin Med* 2023;13:108. <https://doi.org/10.3390/jcm13010108>.
- [32] Zhang Y, Li H, Yang J, Ma H. Physical activity and neuroinflammation: a bibliometric analysis of research progress and future perspectives. *Front Aging Neurosci* 2025;17:1602724. <https://doi.org/10.3389/fnagi.2025.1602724>.
- [33] Guidetti M, Arlotti M, Bocci T, Bianchi AM, Parazzini M, Ferrucci R, et al. Electric Fields Induced in the Brain by Transcranial Electric Stimulation: A Review of In Vivo Recordings. *Biomedicines* 2022;10:2333. <https://doi.org/10.3390/biomedicines10102333>.
- [34] Angius L, Hopker J, Mauger AR. The Ergogenic Effects of Transcranial Direct Current Stimulation on Exercise Performance. *Front Physiol* 2017;8:90. <https://doi.org/10.3389/fphys.2017.00090>.
- [35] Vitor-Costa M, Okuno NM, Bortolotti H, Bertollo M, Boggio PS, Fregni F, et al. Improving Cycling Performance: Transcranial Direct Current Stimulation Increases Time to Exhaustion in Cycling. *PLOS ONE* 2015;10:e0144916. <https://doi.org/10.1371/journal.pone.0144916>.
- [36] Lu P, Hanson NJ, Wen L, Guo F, Tian X. Transcranial Direct Current Stimulation Enhances Muscle Strength of Non-dominant Knee in Healthy Young Males. *Front Physiol* 2021;12. <https://doi.org/10.3389/fphys.2021.788719>.
- [37] Cogiamanian F, Marceglia S, Ardolino G, Barbieri S, Priori A. Improved isometric force endurance after transcranial direct current stimulation over the human motor cortical areas. *European Journal of Neuroscience* 2007;26:242-9. <https://doi.org/10.1111/j.1460-9568.2007.05633.x>.
- [38] Krishnan C, Ranganathan R, Kantak SS, Dhaher YY, Rymer WZ. Anodal Transcranial Direct Current Stimulation Alters Elbow Flexor Muscle Recruitment Strategies. *Brain Stimulation* 2014;7:443-50. <https://doi.org/10.1016/j.brs.2014.01.057>.
- [39] Borducchi DMM, Gomes JS, Akiba H, Cordeiro Q, Borducchi JHM, Valentin LSS, et al. Transcranial Direct Current Stimulation Effects on Athletes' Cognitive Performance: An Exploratory Proof of Concept Trial. *Frontiers in Psychiatry* 2016;7.
- [40] Moshfeghinia R, Shekouh D, Mostafavi S, Hosseinzadeh M, Bahadori AR, Abdollahifard S, et al. The effects of transcranial direct-current stimulation (tDCS) on pain intensity of patients with fibromyalgia: a systematic review and meta-analysis. *BMC Neurology* 2023;23:395. <https://doi.org/10.1186/s12883-023-03445-7>.
- [41] Williams PS, Hoffman RL, Clark BC. Preliminary Evidence That Anodal Transcranial Direct Current Stimulation Enhances Time to Task Failure of a Sustained Submaximal Contraction. *PLOS ONE* 2013;8:e81418. <https://doi.org/10.1371/journal.pone.0081418>.
- [42] Maudrich T, Ragert P, Perrey S, Kenville R. A Single Anodal Transcranial Direct Current Stimulation Session to Enhance Sport-Specific Performance in Trained Individuals? A Systematic Review and Meta-Analysis 2022:2022.06.23.22276798. <https://doi.org/10.1101/2022.06.23.22276798>.

- [43] Zhou Y, Zhai H, Wei H. Acute Effects of Transcranial Direct Current Stimulation Combined with High-Load Resistance Exercises on Repetitive Vertical Jump Performance and EEG Characteristics in Healthy Men. *Life (Basel)* 2024;14:1106. <https://doi.org/10.3390/life14091106>.
- [44] Regolamenti Tecnici - Federpesistica | Federazione Italiana Pesistica. Federpesistica n.d. <https://www.federpesistica.it/regolamenti-gara/> (accessed September 10, 2025).
- [45] Nitsche MA, Liebetanz D, Lang N, Antal A, Tergau F, Paulus W, et al. Safety criteria for transcranial direct current stimulation (tDCS) in humans. *Clinical Neurophysiology : Official Journal of the International Federation of Clinical Neurophysiology* 2003;114:2220–2. [https://doi.org/10.1016/S1388-2457\(03\)00235-9](https://doi.org/10.1016/S1388-2457(03)00235-9).
- [46] Guidetti M, Giannoni-Luza S, Bocci T, Pacheco-Barrios K, Bianchi AM, Parazzini M, et al. Modeling Electric Fields in Transcutaneous Spinal Direct Current Stimulation: A Clinical Perspective. *Biomedicines* 2023;11:1283. <https://doi.org/10.3390/biomedicines11051283>.
- [47] Guidetti M, Ferrucci R, Vergari M, Aglieco G, Naci A, Versace S, et al. Effects of Transcutaneous Spinal Direct Current Stimulation (tsDCS) in Patients With Chronic Pain: A Clinical and Neurophysiological Study. *Frontiers in Neurology* 2021;0:1480. <https://doi.org/10.3389/FNEUR.2021.695910>.
- [48] Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): A tool for double-blind sham-controlled clinical studies in brain stimulation. *Clinical Neurophysiology* 2006;117:845–50. <https://doi.org/10.1016/j.clinph.2005.12.003>.
- [49] Borg GAV. Psychophysical bases of perceived exertion. *Medicine & Science in Sports & Exercise* 1982;14:377–81. <https://doi.org/10.1249/00005768-198205000-00012>.
- [50] Kersten P, Küçükdeveci AA, Tennant A. The use of the Visual Analogue Scale (VAS) in rehabilitation outcomes. *Journal of Rehabilitation Medicine* 2012;44:609–10. <https://doi.org/10.2340/16501977-0999>.
- [51] Mrakic-Sposta S, Vezzoli A, Garetto G, Paganini M, Camporesi E, Giacom TA, et al. Hyperbaric Oxygen Therapy Counters Oxidative Stress/Inflammation-Driven Symptoms in Long COVID-19 Patients: Preliminary Outcomes. *Metabolites* 2023;13:1032. <https://doi.org/10.3390/metabo13101032>.
- [52] Mrakic-Sposta S, Di Santo SG, Franchini F, Arlati S, Zangiacomi A, Greci L, et al. *Frontiers | Effects of Combined Physical and Cognitive Virtual Reality-Based Training on Cognitive Impairment and Oxidative Stress in MCI Patients: A Pilot Study* n.d. <https://doi.org/10.3389/fnagi.2018.00282>.
- [53] Mrakic-Sposta S, Gussoni M, Porcelli S, Pugliese L, Pavei G, Bellistri G, et al. Training Effects on ROS Production Determined by Electron Paramagnetic Resonance in Master Swimmers. *Oxidative Medicine and Cellular Longevity* 2015;2015:e804794. <https://doi.org/10.1155/2015/804794>.
- [54] Brizzolari A, Bosco G, Vezzoli A, Dellanoce C, Barassi A, Paganini M, et al. Seasonal Oxy-Inflammation and Hydration Status in Non-Elite Freeskiing Racer: A Pilot Study by Non-Invasive Analytic Method. *International Journal of Environmental Research and Public Health* 2023;20:3157. <https://doi.org/10.3390/ijerph20043157>.
- [55] Campa F, Toselli S, Mazzilli M, Gobbo LA, Coratella G. Assessment of Body Composition in Athletes: A Narrative Review of Available Methods with Special

- Reference to Quantitative and Qualitative Bioimpedance Analysis. *Nutrients* 2021;13:1620. <https://doi.org/10.3390/nu13051620>.
- [56] Giacom TA, Bosco G, Vezzoli A, Dellanoce C, Cialoni D, Paganini M, et al. Oxidative stress and motion sickness in one crew during competitive offshore sailing. *Sci Rep* 2022;12:1142. <https://doi.org/10.1038/s41598-022-05219-6>.
- [57] Bosco G, Giacom TA, Paolucci N, Vezzoli A, Noce CD, Paganini M, et al. Dopamine/BDNF loss underscores narcosis cognitive impairment in divers: a proof of concept in a dry condition. *Eur J Appl Physiol* 2023;123:143–58. <https://doi.org/10.1007/s00421-022-05055-6>.
- [58] De Donato R, Maiorana NV, Vergari M, De Sandi A, Naci A, Aglieco G, et al. ‘Knock down the brain’: a nonlinear analysis of electroencephalography to study the effects of sub-concussion in boxers. *European Journal of Neurology* 2025;32:e16411. <https://doi.org/10.1111/ene.16411>.
- [59] Lattari E, Andrade ML, Alberto Filho S, Moura AM, Neto GM, Silva JG, et al. Can transcranial direct current stimulation improve the resistance strength and decrease the rating perceived scale in recreational weight-training experience? *The Journal of Strength & Conditioning Research* 2016;30:3381–7.
- [60] Machado S, Jansen P, Almeida V, Veldema J. Is tDCS an adjunct ergogenic resource for improving muscular strength and endurance performance? A systematic review. *Frontiers in Psychology* 2019;10:1127.
- [61] Lattari E, Vieira LA, Oliveira BR, Unal G, Bikson M, de Mello Pedreiro RC, et al. Effects of transcranial direct current stimulation with caffeine intake on muscular strength and perceived exertion. *The Journal of Strength & Conditioning Research* 2019;33:1237–43.
- [62] Vieira LAF, Lattari E, de Jesus Abreu MA, Rodrigues GM, Viana B, Machado S, et al. Transcranial direct current stimulation (tDCS) improves back-squat performance in intermediate resistance-training men. *Research Quarterly for Exercise and Sport* 2022;93:210–8.
- [63] Yu Y, Zhang X, Nitsche MA, Vicario CM, Qi F. Does a single session of transcranial direct current stimulation enhance both physical and psychological performance in national-or international-level athletes? A systematic review. *Frontiers in Physiology* 2024;15:1365530.
- [64] Angius L, Pageaux B, Hopker J, Marcora SM, Mauger AR. Transcranial direct current stimulation improves isometric time to exhaustion of the knee extensors. *Neuroscience* 2016;339:363–75. <https://doi.org/10.1016/j.neuroscience.2016.10.028>.
- [65] Tanaka S, Hanakawa T, Honda M, Watanabe K. Enhancement of pinch force in the lower leg by anodal transcranial direct current stimulation. *Experimental Brain Research* 2009;196:459–65. <https://doi.org/10.1007/s00221-009-1863-9>.
- [66] Rroji O, van Kuyck K, Nuttin B, Wenderoth N. Anodal tDCS over the primary motor cortex facilitates long-term memory formation reflecting use-dependent plasticity. *PLoS One* 2015;10:e0127270.
- [67] Qi S, Cao L, Wang Q, Sheng Y, Yu J, Liang Z. The Physiological Mechanisms of Transcranial Direct Current Stimulation to Enhance Motor Performance: A Narrative Review. *Biology* 2024;13:790. <https://doi.org/10.3390/biology13100790>.
- [68] Saruco E, Di Rienzo F, Nunez-Nagy S, Rubio-Gonzalez MA, Jackson PL, Collet C, et al. Anodal tDCS over the primary motor cortex improves motor imagery benefits on postural control: A pilot study. *Scientific Reports* 2017;7:480. <https://doi.org/10.1038/s41598-017-00509-w>.

- [69] Baharlouei H, Goosheh M, Moore M, Ramezani Ahmadi AH, Yassin M, Jaberzadeh S. The effect of transcranial direct current stimulation on rating of perceived exertion: A systematic review of the literature. *Psychophysiology* 2024;61:e14520.
- [70] dos Santos LF, dos Santos Silva D, de Jesus Alves MD, Moura Pereira EV, do Nascimento HR, de Sousa Fernandes MS, et al. Acute effects of transcranial direct current stimulation (tDCS) on peak torque and 5000 m running performance: a randomized controlled trial. *Scientific Reports* 2023;13:9362. <https://doi.org/10.1038/s41598-023-36093-5>.
- [71] Kristiansen M, Thomsen MJ, Nørgaard J, Aaes J, Knudsen D, Voigt M. Anodal transcranial direct current stimulation increases corticospinal excitability, while performance is unchanged n.d. <https://doi.org/10.1371/journal.pone.0254888>.
- [72] Aydin BN, Stinson EJ, Travis KT, Krakoff J, Rodzevik T, Chang DC, et al. Reduced plasma interleukin-6 concentration after transcranial direct current stimulation to the prefrontal cortex. *Behavioural Brain Research* 2024;474:115201. <https://doi.org/10.1016/j.bbr.2024.115201>.
- [73] Suchting R, Colpo GD, Rocha NP, Ahn H. The Effect of Transcranial Direct Current Stimulation on Inflammation in Older Adults With Knee Osteoarthritis: A Bayesian Residual Change Analysis. *Biological Research For Nursing* 2020;22:57-63. <https://doi.org/10.1177/1099800419869845>.
- [74] da Cunha MJ, Pires Dorneles G, Peres A, Maurer S, Horn K, Souza Pagnussat A. tDCS does not add effect to foot drop stimulator and gait training in improving clinical parameters and neuroplasticity biomarkers in chronic post-stroke: randomized controlled trial. *International Journal of Neuroscience* 2024;134:1518-27. <https://doi.org/10.1080/00207454.2023.2272041>.
- [75] Lu C, Wei Y, Hu R, Wang Y, Li K, Li X. Transcranial Direct Current Stimulation Ameliorates Behavioral Deficits and Reduces Oxidative Stress in 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine-Induced Mouse Model of Parkinson's Disease. *Neuromodulation: Technology at the Neural Interface* 2015;18:442-7. <https://doi.org/10.1111/ner.12302>.
- [76] Mas-Bargues C, Escrivá C, Dromant M, Borrás C, Viña J. Lipid peroxidation as measured by chromatographic determination of malondialdehyde. Human plasma reference values in health and disease. *Archives of Biochemistry and Biophysics* 2021;709:108941. <https://doi.org/10.1016/j.abb.2021.108941>.
- [77] Fritsch B, Reis J, Martinowich K, Schambra HM, Ji Y, Cohen LG, et al. Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning. *Neuron* 2010;66:198-204. <https://doi.org/10.1016/j.neuron.2010.03.035>.
- [78] Binkofski F, Loebig M, Jauch-Chara K, Bergmann S, Melchert UH, Scholand-Engler HG, et al. Brain energy consumption induced by electrical stimulation promotes systemic glucose uptake. *Biol Psychiatry* 2011;70:690-5. <https://doi.org/10.1016/j.biopsych.2011.05.009>.
- [79] Wardzinski EK, Friedrichsen L, Dannenberger S, Kistenmacher A, Melchert UH, Jauch-Chara K, et al. Double transcranial direct current stimulation of the brain increases cerebral energy levels and systemic glucose tolerance in men. *J Neuroendocrinol* 2019;31:e12688. <https://doi.org/10.1111/jne.12688>.
- [80] Kistenmacher A, Manneck S, Wardzinski EK, Martens JC, Gohla G, Melchert UH, et al. Persistent blood glucose reduction upon repeated transcranial electric stimulation in men. *Brain Stimul* 2017;10:780-6. <https://doi.org/10.1016/j.brs.2017.03.011>.

- [81] Adams OP. The impact of brief high-intensity exercise on blood glucose levels. *Diabetes Metab Syndr Obes* 2013;6:113-22.  
<https://doi.org/10.2147/DMSO.S29222>.
- [82] Angius L, Santarnecchi E, Pascual-Leone A, Marcora SM. Transcranial Direct Current Stimulation over the Left Dorsolateral Prefrontal Cortex Improves Inhibitory Control and Endurance Performance in Healthy Individuals. *Neuroscience* 2019;419:34-45.  
<https://doi.org/10.1016/j.neuroscience.2019.08.052>.
- [83] Xu Z, Shen B, Xiao S, Zhang C, Zhan J, Li J, et al. The Effect of Transcranial Direct Current Stimulation on Lower-Limb Endurance Performance: A Systematic Review. *Bioengineering (Basel)* 2024;11:1088.  
<https://doi.org/10.3390/bioengineering11111088>.
- [84] Uehara L, Coelho DB, Baptista AF, Santana L, Moreira RJD, Zana Y, et al. Does Transcranial Direct Current Stimulation reduce central and peripheral muscle fatigue in recreational runners? A triple-blind, sham-controlled, randomized, crossover clinical study. *Braz J Phys Ther* 2024;28:101088.  
<https://doi.org/10.1016/j.bjpt.2024.101088>.

**Enrollment**

Assessed for eligibility (n=10)

Excluded (n=0)

Randomized (n=10)

**Allocation**

Allocated to intervention (n=5)  
· Received allocated intervention (n=5)  
· Did not receive allocated intervention (give reasons) (n=0)

Allocated to intervention (n=5)  
· Received allocated intervention (n=5)  
· Did not receive allocated intervention (give reasons) (n=0)

**Analysis**

Analysed (n=5)  
· Excluded from analysis (n=0)

Analysed (n=5)  
· Excluded from analysis (n=0)







