

ANALYSIS & PDE

Volume 16

No. 6

2023

FABIO CAVALLETTI, ANDREA MONDINO AND DANIELE SEMOLA

QUANTITATIVE OBATA'S THEOREM

QUANTITATIVE OBATA'S THEOREM

FABIO CAVALLETTI, ANDREA MONDINO AND DANIELE SEMOLA

We prove a quantitative version of Obata's theorem involving the shape of functions with null mean value when compared with the cosine of distance functions from single points. The deficit between the diameters of the manifold and of the corresponding sphere is bounded likewise. These results are obtained in the general framework of (possibly nonsmooth) metric measure spaces with curvature-dimension conditions through a quantitative analysis of the transport-ray decompositions obtained by the localization method.

| | |
|---|------|
| 1. Introduction | 1389 |
| 2. Background material | 1395 |
| 3. One-dimensional estimates | 1402 |
| 4. Quantitative Obata's theorem on the diameter | 1413 |
| 5. Quantitative Obata's theorem on almost optimal functions | 1416 |
| Acknowledgements | 1428 |
| References | 1429 |

1. Introduction

One of the core topics in geometric analysis is the deep connection between the geometry of a domain (in a possibly curved space) and spectral properties of the Laplacian defined on it.

The present paper focuses on the first eigenvalue λ_1 of the Laplacian (with Neumann boundary conditions, in case the domain has nonempty boundary). Since the Poincaré(–Wirtinger) inequality plays an important role in analysis and since a lower bound of the first eigenvalue gives an upper bound of the constant in the Poincaré(–Wirtinger) inequality, it is extremely useful to have a good lower estimate of λ_1 .

For domains in the Euclidean space, classical estimates of the first eigenvalue of the Laplacian (under Dirichlet or Neumann boundary conditions) date back to Lord Rayleigh [1877], Faber [1923], Krahn [1925], Pólya and Szegő [1951], and Payne and Weinberger [1960], among others. For curved spaces, two major results are due to Lichnerowicz [1958] and Obata [1962]:

Theorem 1.1. *Let (M, g) be an N -dimensional Riemannian manifold with $\text{Ric}_g \geq (N - 1)g$. Then $\lambda_1 \geq N$ (Lichnerowicz spectral gap [1958]).*

Moreover, $\lambda_1 = N$ if and only if (M, g) is isometric to the unit sphere \mathbb{S}^N (Obata's theorem [1962]).

MSC2020: 53C23, 58J50.

Keywords: quantitative inequalities, Obata's theorem, Ricci curvature, spectral gap.

Remark 1.2. On \mathbb{S}^N , the first eigenvalue $\lambda_1 = N$ has multiplicity $N + 1$. The corresponding eigenspace is spanned by the restriction to \mathbb{S}^N of affine functions of \mathbb{R}^{N+1} (i.e., an L^2 -orthogonal basis is composed of the standard coordinate functions $\{x^1, x^2, \dots, x^{N+1}\}$ of \mathbb{R}^{N+1}). Equivalently, a function $u : \mathbb{S}^N \rightarrow \mathbb{R}$ is a first eigenfunction normalized as $\|u\|_{L^2(\mathbb{S}^N)} = 1$ if and only if there exists $P \in \mathbb{S}^N$ such that $u = \sqrt{N + 1} \cos d_P$, where we denote by d_P the Riemannian distance from the point P .

Our main result is a *quantitative spectral gap* involving the shape of the eigenfunctions (or, more generally, of functions with almost optimal Rayleigh quotient), when compared with the eigenfunctions of the model space \mathbb{S}^N (as in Remark 1.2). In detail, we show that if $\text{Ric}_g \geq (N - 1)g$ and $u : M \rightarrow \mathbb{R}$ is a first eigenfunction with $\|u\|_{L^2(M)} = 1$, then there exists $P \in M$ such that

$$\|u - \sqrt{N + 1} \cos d_P\|_{L^2(M)} \leq C(N)(\lambda_1 - N)^{O(1/N)}. \tag{1-1}$$

More generally, the same conclusion holds for every Lipschitz function $u : M \rightarrow \mathbb{R}$ with null mean value and $\|u\|_{L^2(M)} = 1$, provided λ_1 on the right-hand-side is replaced by the Dirichlet energy $\int_M |\nabla u|^2 d \text{vol}_g$.

We will prove (1-1) with tools of optimal transport tailored to study (possibly nonsmooth) metric measure spaces satisfying Ricci curvature lower bounds and dimensional upper bounds in the synthetic sense, the so-called $\text{CD}(K, N)$ spaces introduced in [Sturm 2006a; 2006b; Lott and Villani 2009]. For the sake of this introduction, a metric measure space (m.m.s. for short) is a triple (X, d, \mathfrak{m}) , where (X, d) is a compact metric space and \mathfrak{m} is a Borel probability measure, playing the role of reference volume measure. A $\text{CD}(K, N)$ space should be roughly thought of as a possibly nonsmooth metric measure space having Ricci curvature bounded below by $K \in \mathbb{R}$ and dimension bounded above by $N \in (1, \infty)$ in the synthetic sense. The basic idea of the synthetic approach of Lott, Sturm and Villani is to analyze weighted convexity properties of suitable entropy functionals along geodesics in the space of probability measures endowed with the quadratic transportation (also known as Kantorovich–Wasserstein) distance. An important technical assumption throughout the paper is the essentially nonbranching (“e.n.b.” for short) property [Rajala and Sturm 2014], which roughly corresponds to requiring that the L^2 -optimal transport between two absolutely continuous (with respect to the reference volume measure \mathfrak{m}) probability measures is performed along geodesics which do not branch (for the precise definitions see Sections 2A and 2B). Notable examples of spaces satisfying e.n.b. $\text{CD}(K, N)$ include (geodesically convex domains in) smooth Riemannian manifolds with Ricci bounded below by K and dimension bounded above by N , their measured Gromov–Hausdorff limits (i.e., the so-called “Ricci limits”) and more generally $\text{RCD}(K, N)$ spaces (i.e., $\text{CD}(K, N)$ spaces with linear Laplacian; see Remark 2.4 for more details), finite-dimensional Alexandrov spaces with curvature bounded below, and Finsler manifolds endowed with a strongly convex norm. A standard example of a space failing to satisfy the essentially nonbranching property is \mathbb{R}^2 endowed with the L^∞ norm. Later in the introduction, when discussing the main steps of the proof, we will mention how the essentially nonbranching assumption is used in our arguments.

We will establish our results directly on the more general class of e.n.b. $\text{CD}(N - 1, N)$ metric measure spaces. For an m.m.s. (X, d, \mathfrak{m}) we define the nonnegative real number $\lambda_{(X,d,\mathfrak{m})}^{1,2}$ as

$$\lambda_{(X,d,\mathfrak{m})}^{1,2} := \inf \left\{ \frac{\int_X |\nabla u|^2 \mathfrak{m}}{\int_X |u|^2 \mathfrak{m}} : u \in \text{Lip}(X) \cap L^2(X, \mathfrak{m}), u \neq 0, \int_X u \mathfrak{m} = 0 \right\}, \tag{1-2}$$

where $|\nabla u|$ is the slope (also called local Lipschitz constant) of the Lipschitz function u given by

$$|\nabla u|(x) = \begin{cases} \limsup_{y \rightarrow x} |u(x) - u(y)|/d(x, y) & \text{if } x \text{ is not isolated,} \\ 0 & \text{otherwise.} \end{cases}$$

It is well known that, in case (X, d, \mathfrak{m}) is the m.m.s. corresponding to a smooth compact Riemannian manifold (possibly with boundary) $\lambda_{(X,d,\mathfrak{m})}^{1,2}$ coincides with the first eigenvalue of the problem $-\Delta u = \lambda u$ with Neumann boundary conditions.

Considering the extension of (1-1) to e.n.b. $\text{CD}(N - 1, N)$ spaces is natural: indeed a sequence (M_j, g_j) of Riemannian N -manifolds with $\text{Ric}_{g_j} \geq (N - 1)g_j$ where the right-hand side of (1-1) converges to zero as $j \rightarrow \infty$ may develop singularities and admits a limit (up to subsequences) in the measured Gromov–Hausdorff sense to a possibly nonsmooth e.n.b. $\text{CD}(N - 1, N)$ space (actually the limit is, more strongly, $\text{RCD}(N - 1, N)$).

In the enlarged class of e.n.b. $\text{CD}(N - 1, N)$ spaces (actually already for $\text{RCD}(N - 1, N)$ spaces), Obata’s rigidity theorem must be modified:

- First of all, $N \in (1, \infty)$ is a (possibly noninteger) real number.
- Even in the case when N is an integer, the round sphere \mathbb{S}^N is not anymore the only case of equality in the Lichnerowicz spectral gap as the spherical suspensions achieve equality as well [Ketterer 2015].

A key geometric property of the spherical suspensions is that they have diameter π , thus saturating Bonnet–Myers diameter upper bound. The first part of our main result is a quantitative control of how close to π the diameter must be, in terms of the spectral gap deficit. The second part of the statement is an L^2 -quantitative control of the shape of functions with almost optimal Rayleigh quotient. We can now state our main theorem.

Theorem 1.3 (quantitative Obata’s theorem for e.n.b. $\text{CD}(N - 1, N)$ -spaces). *For every real number $N > 1$ there exists a real constant $C(N) > 0$ with the following properties: If (X, d, \mathfrak{m}) is an essentially nonbranching metric measure space satisfying the $\text{CD}(N - 1, N)$ condition and $\mathfrak{m}(X) = 1$ with $\text{supp}(\mathfrak{m}) = X$, then*

$$\pi - \text{diam}(X) \leq C(N)(\lambda_{(X,d,\mathfrak{m})}^{1,2} - N)^{1/N}. \tag{1-3}$$

Moreover, for any Lipschitz function $u : X \rightarrow \mathbb{R}$ with $\int_X u \, \mathfrak{m} = 0$ and $\int_X u^2 \, \mathfrak{m} = 1$, there exists a distinguished point $P \in X$ such that

$$\|u - \sqrt{N + 1} \cos d_P\|_{L^2(X,\mathfrak{m})} \leq C(N) \left(\int_X |\nabla u|^2 \, \mathfrak{m} - N \right)^\eta, \quad \eta = \frac{1}{6N + 4}. \tag{1-4}$$

Remark 1.4. Although Theorem 1.3 is formulated for e.n.b. $\text{CD}(N - 1, N)$ spaces, a statement for e.n.b. $\text{CD}(K, N)$ spaces with $K > 0$ is easily obtained by scaling. Indeed, (X, d, \mathfrak{m}) satisfies $\text{CD}(K, N)$ if and only if, for any $\alpha, \beta \in (0, \infty)$, the scaled metric measure space $(X, \alpha d, \beta \mathfrak{m})$ satisfies $\text{CD}(\alpha^{-2}K, N)$; see [Sturm 2006b, Proposition 1.4].

Let us compare Theorem 1.3 with related results in the literature. Under the standing assumption that (M, g) is a smooth Riemannian N -manifold without boundary and with $\text{Ric}_g \geq (N - 1)g$:

(1) It follows from Cheng's comparison theorem [1975] that if $\lambda_{(M,g)}^{1,2}$ is close to N then the diameter of M must be close to π . Conversely, Croke [1982] proved that if the diameter is close to π then $\lambda_{(M,g)}^{1,2}$ must be close to N . Bérard, Besson and Gallot [Bérard et al. 1985] sharpened the diameter estimate of Cheng by proving an estimate very similar to (1-3).

(2) Bertrand [2007] established the following stability result for eigenfunctions (see also [Petersen 1999]): for every $\epsilon > 0$ there exists $\delta > 0$ such that if $\lambda_1 \leq N + \delta$ and u is an eigenfunction relative to λ_1 normalized so that $\int_M u^2 d \text{vol}_g = \text{vol}_g(M)$, then there exists a point $P \in M$ such that $\|u - \sqrt{N+1} \cos d_P\|_{L^\infty(X,m)} \leq \epsilon$.

Theorem 1.3 sharpens and extends the above results in various ways:

- The estimate (1-3) extends [Bérard et al. 1985] to e.n.b. $\text{CD}(N-1, N)$ spaces. These spaces are nonsmooth a priori and may have (convex) boundary. Actually, as the reader will realize, the claim (1-3) will be proved in Section 4 along the way to proving the much harder (1-4), to which the entire Section 5 is devoted.
- The estimate (1-4) extends Bertrand's stability [2007] to the more general class of e.n.b. $\text{CD}(N-1, N)$ spaces and to arbitrary functions (a priori not eigenfunctions) with Rayleigh quotient close to N . The fact that u is an eigenfunction was key in [Bertrand 2007] in order to apply maximum principle and gradient estimates in the spirit of [Li and Yau 1980]. Let us stress that our methods are completely different and work for an arbitrary Lipschitz function satisfying a *small energy* condition but *no PDE* a priori.

Inequality (1-4) naturally fits in the framework of quantitative functional/geometric inequalities. A basic result in this context is the quantitative Euclidean isoperimetric inequality proved by Fusco, Maggi and Pratelli [Fusco et al. 2008] (see also [Figalli et al. 2010; Cicalese and Leonardi 2012] for different proofs) stating that for every Borel set $E \subset \mathbb{R}^n$ of positive and finite volume there exists $\bar{x} \in \mathbb{R}^n$ such that

$$\frac{|E \Delta B_{r_E}(\bar{x})|}{|E|} \leq C(N) \left(\frac{P(E)}{P(B_{r_E}(\bar{x}))} - 1 \right)^{1/2}, \quad (1-5)$$

where r_E is such that $|B_{r_E}(\bar{x})| = |E|$. Quantitative results involving the spectrum of the Laplacian have been proved for domains in \mathbb{R}^n , among others, by Hansen and Nadirashvili [1994] in dimension 2, by Melas [1992] for convex bodies, by Fusco, Maggi and Pratelli [Fusco et al. 2009] and Brasco, De Philippis and Velichkov [Brasco et al. 2015] regarding quantitative forms of the Faber–Krahn inequality and by Brasco and Pratelli [2012] regarding quantitative versions of the Krahn–Szegő and Szegő–Weinberger inequalities. More recently, a quantitative version of the Lévy–Gromov isoperimetric inequality was proved for essentially nonbranching $\text{CD}(N-1, N)$ metric measure spaces in [Cavalletti et al. 2019], and a quantitative isoperimetric inequality in the setting of smooth Riemannian manifolds was considered in [Chodosh et al. 2023].

Taking variations in the broad context of metric measure spaces makes the prediction on the sharp exponent η in (1-4) a hard task. Even formulating a conjecture is a challenging question and it could actually be that $\eta = O(1/N)$ as $N \rightarrow \infty$ is already sharp. In the direction of this guess, we notice that the exponent $1/N$ in (1-3) is indeed optimal in the class of metric measure spaces, as a direct computation on the model one-dimensional space $([0, D], |\cdot|, \sin^{N-1}(\cdot) \mathcal{L}^1)$ shows.

Before discussing the main steps in the proof of [Theorem 1.3](#), it is worth recalling remarkable examples of spaces fitting in the assumptions of the result. Let us stress that our main theorem seems new in all of them. The class of essentially nonbranching $CD(N - 1, N)$ spaces includes many notable families of spaces, among them:

- *Geodesically convex domains* in (resp. weighted) Riemannian N -manifolds satisfying $\text{Ric}_g \geq (N - 1)g$ (resp. N -Bakry–Émery Ricci curvature bounded below by $N - 1$).
- *Measured Gromov Hausdorff limits of Riemannian N -manifolds satisfying $\text{Ric}_g \geq (N - 1)g$* (so-called “Ricci limits”) and more generally the class of $RCD(N - 1, N)$ spaces. Indeed Ricci limits are examples of $RCD(N - 1, N)$ spaces (see for instance [\[Gigli et al. 2015\]](#)) and $RCD(N - 1, N)$ spaces are essentially nonbranching $CD(N - 1, N)$ (see [\[Rajala and Sturm 2014\]](#)).
- *Alexandrov spaces with curvature ≥ 1* . Petrunin [\[2011\]](#) proved that the synthetic curvature lower bound in the sense of comparison triangles is compatible with the optimal transport lower bound on the Ricci curvature of Lott, Sturm and Villani (see also [\[Zhang and Zhu 2010\]](#)). Moreover geodesics in Alexandrov spaces with curvature bounded below do not branch. It follows that Alexandrov spaces with curvature bounded from below by 1 and Hausdorff dimension at most N are nonbranching $CD(N - 1, N)$ spaces.
- *Finsler manifolds with strongly convex norm, and satisfying Ricci curvature lower bounds*. More precisely we consider a C^∞ -manifold M , endowed with a function $F : TM \rightarrow [0, \infty]$ such that $F|_{TM \setminus \{0\}}$ is C^∞ and for each $x \in M$ it holds that $F_x := T_x M \rightarrow [0, \infty]$ is a strongly convex norm, i.e.,

$$g_{ij}^x(v) := \frac{\partial^2(F_x^2)}{\partial v^i \partial v^j}(v) \quad \text{is a positive definite matrix at every } v \in T_x M \setminus \{0\}.$$

Under these conditions, it is known that one can write the geodesic equations and the geodesics do not branch: in other words these spaces are nonbranching. We also assume (M, F) to be geodesically complete and endowed with a C^∞ probability measure m in such a way that the associated m.m.s. (X, F, m) satisfies the $CD(N - 1, N)$ condition. This class of spaces has been investigated by Ohta [\[2009\]](#), who established the equivalence between the curvature dimension condition and a Finsler version of the Bakry–Émery N -Ricci tensor bounded from below.

An overview of the proof. The starting point of the proof of [Theorem 1.3](#) is the metric-measure version of the classical *localization technique*. First introduced by Payne and Weinberger [\[1960\]](#) for establishing a sharp Poincaré–Wirtinger inequality for convex domains in \mathbb{R}^n , the localization technique has been developed into a general dimension-reduction tool for geometric inequalities in symmetric spaces by Gromov and Milman [\[1987\]](#), Lovász and Simonovits [\[1993\]](#) and Kannan, Lovász and Simonovits [\[Kannan et al. 1995\]](#). More recently, Klartag [\[2017\]](#) used optimal transportation tools in order to extend the range of applicability of the technique to general Riemannian manifolds. The extension to the metric setting was finally obtained in [\[Cavalletti and Mondino 2017b\]](#); see [Section 2D](#).

Given a function $u \in L^1(X, m)$ with $\int_X u \, m = 0$, the localization theorem ([Theorem 2.10](#)) gives a decomposition of X into a family of one-dimensional sets $\{X_q\}_{q \in Q}$ formed by the transport rays of a Kantorovich potential associated to the optimal transport from the positive part of u (i.e., $\mu_0 := \max\{u, 0\} \, m$)

to the negative part of u (i.e., $\mu_1 := \max\{-u, 0\} \mathfrak{m}$); each X_q is in particular isometric to a real interval. A first key property of such a decomposition is that each ray X_q carries a natural measure \mathfrak{m}_q (given by the disintegration theorem) in such a way that

$$(X_q, d, \mathfrak{m}_q) \text{ is a } CD(N - 1, N) \text{ space and } \int_{X_q} u \mathfrak{m}_q = 0, \tag{1-6}$$

so that both the geometry of the space and the null mean value constraint are *localized* into a family of one-dimensional spaces. An important ingredient used in the proof of such a decomposition is the essentially nonbranching property which, coupled with $CD(N - 1, N)$ (actually the weaker measure contraction would suffice here), guarantees that the rays form a partition of X (up to an \mathfrak{m} -negligible set).

In order to exploit (1-6), as a first step, in Section 3 we prove the one-dimensional counterparts of Theorem 1.3. More precisely, given a one-dimensional $CD(N - 1, N)$ space $(I = [0, D], |\cdot|, \mathfrak{m})$ we show that (Proposition 3.3)

$$\pi - D \leq C(N)(\lambda_{(I, |\cdot|, \mathfrak{m})}^{1,2} - N)^{1/N}, \tag{1-7}$$

and that, if $u \in \text{Lip}(I)$ satisfies $\int u \mathfrak{m} = 0$ and $\int u^2 \mathfrak{m} = 1$, then (Theorem 3.11)

$$\min\{\|u - \sqrt{N + 1} \cos(\cdot)\|_{L^2(\mathfrak{m})}, \|u + \sqrt{N + 1} \cos(\cdot)\|_{L^2(\mathfrak{m})}\} \leq C \left(\int |u'|^2 \mathfrak{m} - N \right)^{\min\{1/2, 1/N\}}. \tag{1-8}$$

Combining (1-6) and (1-7), it is not hard to prove (see Theorem 4.3) the first claim (1-3) of Theorem 1.3. Actually, calling Q_ℓ (for “ Q long”) the set of indices for which $|X_q| \simeq \pi$, we aim to show that $\mathfrak{q}(Q_\ell) \simeq 1$ (i.e., “most rays are long”). As we will discuss in a few lines, this is far from being trivial (in particular, it needs new ideas when compared with [Cavalletti et al. 2019]).

A second crucial property of the decomposition $\{X_q\}_{q \in Q}$, inherited by the variational nature of the construction, is the so-called cyclical monotonicity. This was key in [Cavalletti et al. 2019] for showing that, for $q \in Q_\ell$, the transport ray X_q has its starting point close to a fixed “south pole” P_S , and ends up near a fixed “north pole” P_N (in particular, the distance between P_S and P_N is close to π) (Proposition 5.1).

Then we observe that (1-8) forces, for $q \in Q_\ell$, the fiber $u_q := u|_{X_q}$ (that is the restriction of u to the corresponding one-dimensional element of the partition) to be L^2 close to a multiple of the cosine of the arclength parametrization along the ray X_q , i.e.,

$$u_q(\cdot) \simeq c_q \sqrt{N + 1} \cos(\cdot) \text{ along } X_q, \text{ where } c_q = \|u_q\|_{L^2(\mathfrak{m}_q)} \text{ for } q \in Q_\ell \text{ (see (5-13)).} \tag{1-9}$$

The difficulties in order to conclude the proof are mainly two, and are strictly linked:

- (1) Show that $Q_\ell \ni q \mapsto c_q$ is almost constant.
- (2) Show that $\mathfrak{q}(Q_\ell) \simeq 1$.

Let us stress that at this stage the only given information is that $\int_{Q_\ell} c_q^2 \mathfrak{q} \simeq 1$. The intuition why (1) and (2) should hold is that an oscillation of c_q would correspond to an oscillation of u “orthogonal to the transport rays”, which would be expensive in terms of Dirichlet energy of u . The proofs of the two claims are the most technical part of the work and correspond respectively to Propositions 5.2 and 5.3.

Let us mention that the two difficulties (1) and (2) were not present in the proof of the quantitative Lévy–Gromov inequality in [Cavalletti et al. 2019], where it was sufficient to work with characteristic functions (which have a fixed scale, i.e., they are either 0 or 1).

2. Background material

The goal of this section is to fix the notation and to recall the basic notions/constructions used throughout the paper: in Section 2A we review geodesics in the Wasserstein distance, in Section 2B curvature-dimension conditions, in Section 2C some basics of $CD(K, N)$ densities on segments of the real line, and in Section 2D the decomposition of the space into transport rays (localization).

2A. Geodesics in the L^2 -Kantorovich–Wasserstein distance. Let (X, d) be a compact metric space and m a Borel probability measure over X . The triple (X, d, m) is called metric measure space, m.m.s. for short.

The space of all Borel probability measures over X will be denoted by $\mathcal{P}(X)$. We define the L^2 -Kantorovich–Wasserstein distance W_2 between two measures $\mu_0, \mu_1 \in \mathcal{P}(X)$ as

$$W_2(\mu_0, \mu_1)^2 = \inf_{\pi} \int_{X \times X} d^2(x, y) \pi(dx dy), \tag{2-1}$$

where the infimum is taken over all $\pi \in \mathcal{P}(X \times X)$ with μ_0 and μ_1 as the first and the second marginal, i.e., $(P_1)_\# \pi = \mu_0$, $(P_2)_\# \pi = \mu_1$. Of course P_i , $i = 1, 2$, denotes the projection on the first and second factors, respectively, and $(P_i)_\#$ is the corresponding push-forward map on measures. As (X, d) is complete, $(\mathcal{P}(X), W_2)$ is also complete.

The space of geodesics of (X, d) is denoted by

$$\text{Geo}(X) := \{\gamma \in C([0, 1], X) : d(\gamma_s, \gamma_t) = |s - t|d(\gamma_0, \gamma_1) \text{ for every } s, t \in [0, 1]\}.$$

A metric space (X, d) is said to be a *geodesic space* if and only if for each $x, y \in X$ there exists $\gamma \in \text{Geo}(X)$ such that $\gamma_0 = x$, $\gamma_1 = y$. A basic fact of W_2 geometry is that if (X, d) is geodesic then $(\mathcal{P}(X), W_2)$ is geodesic as well. For any $t \in [0, 1]$, let e_t denote the evaluation map:

$$e_t : \text{Geo}(X) \rightarrow X, \quad e_t(\gamma) := \gamma_t.$$

Any geodesic $(\mu_t)_{t \in [0, 1]}$ in $(\mathcal{P}(X), W_2)$ can be lifted to a measure $\nu \in \mathcal{P}(\text{Geo}(X))$, called a *dynamical optimal plan*, such that $(e_t)_\# \nu = \mu_t$ for all $t \in [0, 1]$. Given $\mu_0, \mu_1 \in \mathcal{P}(X)$, we denote by $\text{OptGeo}(\mu_0, \mu_1)$ the space of all $\nu \in \mathcal{P}(\text{Geo}(X))$ for which $(e_0, e_1)_\# \nu$ realizes the minimum in (2-1). Here as usual $\#$ indicates the push-forward operation. If (X, d) is geodesic, then the set $\text{OptGeo}(\mu_0, \mu_1)$ is nonempty for any $\mu_0, \mu_1 \in \mathcal{P}(X)$.

A set $F \subset \text{Geo}(X)$ is a *set of nonbranching geodesics* if and only if for any $\gamma^1, \gamma^2 \in F$, it holds

$$\text{there exists } \bar{t} \in (0, 1) \text{ such that, for all } t \in [0, \bar{t}], \gamma_t^1 = \gamma_t^2 \implies \gamma_s^1 = \gamma_s^2 \text{ for all } s \in [0, 1].$$

A measure μ on a measurable space (Ω, \mathcal{F}) is said to be *concentrated* on $F \subset \Omega$ if there exists $E \subset F$ with $E \in \mathcal{F}$ so that $\mu(\Omega \setminus E) = 0$. With this terminology, we next recall the definition of *essentially nonbranching space* from [Rajala and Sturm 2014].

Definition 2.1. A metric measure space (X, d, m) is *essentially nonbranching* if and only if for any $\mu_0, \mu_1 \in \mathcal{P}(X)$, with μ_0, μ_1 absolutely continuous with respect to m , any element of $\text{OptGeo}(\mu_0, \mu_1)$ is concentrated on a set of nonbranching geodesics.

2B. Curvature-dimension conditions for metric measure spaces. The L^2 -transport structure described in Section 2A allows us to formulate a generalized notion of Ricci curvature lower bound coupled with a dimension upper bound in the context of possibly nonsmooth metric measure spaces. This corresponds to the $\text{CD}(K, N)$ condition introduced in the seminal works of Sturm [2006a; 2006b] and Lott and Villani [2009], which here is reviewed only for a compact m.m.s. (X, d, m) with $m \in \mathcal{P}(X)$ and in the case $K > 0$, $1 < N < \infty$ (the basic setting of the present paper).

For $N \in (1, \infty)$, the N -Rényi relative-entropy functional $\mathcal{E}_N : \mathcal{P}(X) \rightarrow [0, 1]$ is defined as

$$\mathcal{E}_N(\mu) := \int \rho^{1-1/N} dm,$$

where $\mu = \rho m + \mu^{\text{sing}}$ is the Lebesgue decomposition of μ with $\mu^{\text{sing}} \perp m$.

Given $K \in (0, \infty)$, $N \in (1, \infty)$, and $t \in [0, 1]$, define $\sigma_{K,N}^{(t)} : [0, \infty) \rightarrow [0, \infty]$ as

$$\begin{cases} \sigma_{K,N}^{(t)}(0) := t, \\ \sigma_{K,N}^{(t)}(\theta) := \sin(t\theta\sqrt{K/N})/\sin(\theta\sqrt{K/N}) & \text{if } 0 < \theta < \pi/\sqrt{K/N}, \\ \sigma_{K,N}^{(t)}(\theta) := +\infty & \text{otherwise.} \end{cases} \tag{2-2}$$

Set also

$$\tau_{K,N}^{(t)}(\theta) := t^{1/N} \sigma_{K,N-1}^{(t)}(\theta)^{1-1/N}. \tag{2-3}$$

Definition 2.2 ($\text{CD}(K, N)$). An m.m.s. (X, d, m) is said to satisfy $\text{CD}(K, N)$ if for all $\mu_0, \mu_1 \in \mathcal{P}(X)$ absolutely continuous with respect to m there exists $\nu \in \text{OptGeo}(\mu_0, \mu_1)$ so that for all $t \in [0, 1]$ it holds $\mu_t := (e_t)_{\#} \nu \ll m$ and

$$\mathcal{E}_{N'}(\mu_t) \geq \int_{X \times X} (\tau_{K,N'}^{(1-t)}(d(x_0, x_1))\rho_0^{-1/N'}(x_0) + \tau_{K,N'}^{(t)}(d(x_0, x_1))\rho_1^{-1/N'}(x_1)) \pi(dx_0, dx_1) \tag{2-4}$$

for all $N' \geq N$, where $\pi = (e_0, e_1)_{\#}(\nu)$ and $\mu_i = \rho_i m$, $i = 0, 1$.

If (X, d, m) satisfies the $\text{CD}(K, N)$ condition then the same is valid for $(\text{supp}(m), d, m)$; hence we directly assume $X = \text{supp}(m)$.

For the general definition of $\text{CD}(K, N)$ see [Lott and Villani 2009; Sturm 2006a; 2006b].

Remark 2.3 (case of a smooth Riemannian manifold). It is worth recalling that if (M, g) is a Riemannian manifold of dimension n and $h \in C^2(M)$ with $h > 0$ then, denoting by d_g and vol_g the Riemannian distance and volume measure, the m.m.s. $(M, d_g, h \text{vol}_g)$ satisfies $\text{CD}(K, N)$ with $N \geq n$ if and only if (see [Sturm 2006b, Theorem 1.7])

$$\text{Ric}_{g,h,N} \geq Kg, \quad \text{Ric}_{g,h,N} := \text{Ric}_g - (N - n) \frac{\nabla_g^2 h^{1/(N-n)}}{h^{1/(N-n)}},$$

in other words if and only if the weighted Riemannian manifold $(M, g, h \text{vol}_g)$ has N -Bakry-Émery Ricci tensor bounded below by K . Note that if $N = n$, the Bakry-Émery Ricci tensor $\text{Ric}_{g,h,N} = \text{Ric}_g$ makes sense only if h is constant. □

Remark 2.4 ($CD^*(K, N)$, $RCD^*(K, N)$ and $RCD(K, N)$). The lack of the local-to-global property of the $CD(K, N)$ condition (for $K/N \neq 0$) led Bacher and Sturm [2010] to introduce the reduced curvature-dimension condition, denoted by $CD^*(K, N)$. The $CD^*(K, N)$ condition asks for the same inequality (2-4) of $CD(K, N)$ to hold but the coefficients $\tau_{K,N}^{(s)}(d(\gamma_0, \gamma_1))$ are replaced by the slightly smaller $\sigma_{K,N}^{(s)}(d(\gamma_0, \gamma_1))$. Let us explicitly notice that, in general, $CD^*(K, N)$ is weaker than $CD(K, N)$. A subsequent breakthrough in the theory was obtained with the introduction of the Riemannian curvature dimension condition $RCD(K, N)$: in the infinite-dimensional case $N = \infty$ this condition was introduced in [Ambrosio et al. 2014] (for finite measures m , and in [Ambrosio et al. 2015] for σ -finite ones). The finite-dimensional refinements $RCD(K, N)/RCD^*(K, N)$ with $N < \infty$ were subsequently studied in [Gigli 2015; Erbar et al. 2015; Ambrosio et al. 2019]. We refer to these articles as well as to the survey papers [Ambrosio 2018; Villani 2019] for a general account on the synthetic formulation of Ricci curvature lower bounds, in particular of the latter Riemannian-type. Here we only briefly recall that it is a stable [Gigli et al. 2015] strengthening of the (resp. reduced) curvature-dimension condition: an m.m.s. satisfies $RCD(K, N)$ (resp. $RCD^*(K, N)$) if and only if it satisfies $CD(K, N)$ (resp. $CD^*(K, N)$) and the Sobolev space $W^{1,2}(X, m)$ is a Hilbert space (with the Hilbert structure induced by the Cheeger energy).

To conclude we recall also that recently, the first author together with E. Milman [Cavalletti and Milman 2021] proved the equivalence of $CD(K, N)$ and $CD^*(K, N)$, together with the local-to-global property for $CD(K, N)$, in the framework of essentially nonbranching m.m.s. having $m(X) < \infty$. As we will always assume the aforementioned properties to be satisfied by our ambient m.m.s. (X, d, m) , we will use both formulations with no distinction. It is worth also mentioning that an m.m.s. satisfying $RCD^*(K, N)$ is essentially nonbranching (see [Rajala and Sturm 2014, Corollary 1.2]), implying also the equivalence of $RCD^*(K, N)$ and $RCD(K, N)$ (see [Cavalletti and Milman 2021] for details). \square

We shall always assume that the m.m.s. (X, d, m) is essentially nonbranching and satisfies $CD(K, N)$ for some $K > 0$, $N \in (1, \infty)$ with $\text{supp}(m) = X$. It follows that (X, d) is a geodesic and compact metric space. More precisely: note we assumed from the beginning (X, d) to be compact for the sake of simplicity; however, such an assumption could have been replaced by completeness and separability throughout Sections 2A and 2B, but compactness would have been now a consequence of $CD(K, N)$ for some $K > 0$, $N \in (1, \infty)$.

A useful property of essentially nonbranching $CD(K, N)$ spaces is the validity of a weak local Poincaré inequality.

Proposition 2.5 (weak local Poincaré inequality). *Let (X, d, m) be an essentially nonbranching $CD(K, N)$ space for some $K \geq 0$, $N > 1$. For every $u \in \text{Lip}(X)$ it holds*

$$\int_{B_r(x)} \left| u - \int_{B_r(x)} u \right| m \leq 2^{N+2} r \int_{B_{2r}(x)} |\nabla u| m. \tag{2-5}$$

More generally, for every $p \geq 1$ there exists $C_{p,N}$ such that

$$\int_{B_r(x)} \left| u - \int_{B_r(x)} u \right|^p m \leq C_{p,N} r^p \int_{B_{10r}(x)} |\nabla u|^p m. \tag{2-6}$$

Proof. It is well known that, in essentially nonbranching $CD(K, N)$ spaces, the W_2 geodesic connecting two absolutely continuous probability measures is unique (indeed, it holds more generally for essentially nonbranching $MCP(K, N)$ spaces [Cavalletti and Mondino 2017a, Theorem 1.1]). Thus, (X, d, m) as in the assumptions enters the framework of [Rajala 2012, Corollary 1] and (2-5) follows.

Recalling that by the Bishop–Gromov inequality [Sturm 2006b, Theorem 2.3] it holds

$$\frac{m(B_\rho(x_0))}{m(B_1(x_0))} \geq C_N \rho^N$$

for every $\rho \in [0, 1]$, $x_0 \in X$, the second claim (2-6) is a consequence of (2-5) and [Hajlasz and Koskela 2000, Theorem 5.1]. □

2C. $CD(K, N)$ densities on segments of the real line. We will use several times the following terminology: recalling the coefficients σ from (2-2), a nonnegative function h defined on an interval $I \subset \mathbb{R}$ is called a $CD(K, N)$ density on I , for $K \in \mathbb{R}$ and $N \in (1, \infty)$, if for all $x_0, x_1 \in I$ and $t \in [0, 1]$

$$h(tx_1 + (1 - t)x_0)^{1/(N-1)} \geq \sigma_{K, N-1}^{(t)}(|x_1 - x_0|)h(x_1)^{1/(N-1)} + \sigma_{K, N-1}^{(1-t)}(|x_1 - x_0|)h(x_0)^{1/(N-1)}. \quad (2-7)$$

The link with the definition of $CD(K, N)$ for an m.m.s. can be summarized as follows (see for instance [Cavalletti and Milman 2021, Theorem A.2]): if h is a $CD(K, N)$ density on an interval $I \subset \mathbb{R}$ then the m.m.s. $(I, |\cdot|, h(t) dt)$ satisfies $CD(K, N)$; conversely, if the m.m.s. $(\mathbb{R}, |\cdot|, \mu)$ satisfies $CD(K, N)$ and $I = \text{supp}(\mu)$ is not a point, then $\mu \ll \mathcal{L}^1$ and there exists a representative of the density $h = d\mu/d\mathcal{L}^1$ which is a $CD(K, N)$ density on I .

A $CD(K, N)$ density h defined on an interval $I \subset \mathbb{R}$ satisfies the following properties:

- h is lower semicontinuous on I and locally Lipschitz continuous in its interior (this is easily reduced to the corresponding statement for concave functions on I).
- h is strictly positive in the interior of I whenever it does not vanish identically (this follows directly from the definition (2-7)).
- h is locally semiconcave in the interior of I , i.e., for all x_0 in the interior of I , there exists $C_{x_0} \in \mathbb{R}$ so that $x \mapsto h(x) - C_{x_0}(x - x_0)^2$ is concave in a neighborhood of x_0 . In particular, h is twice differentiable in I with at most countably many exceptions.

As proven in [Cavalletti and Milman 2021, Lemma A.5], if h is a $CD(K, N)$ density on an interval I then at any point x in the interior where it is twice differentiable (thus up to at most countably many exceptions) it holds

$$(\log h)''(x) + \frac{1}{N-1} ((\log h)'(x))^2 = (N-1) \frac{(h^{1/(N-1)})''(x)}{h^{1/(N-1)}(x)} \leq -K. \quad (2-8)$$

Also the converse implication holds; see [Cavalletti and Milman 2021, Lemma A.6] for the proof and the precise statement.

We next recall some estimates on $CD(N - 1, N)$ densities, which will turn out to be useful in the paper. Let h_N be the model density for the $CD(N - 1, N)$ condition given by

$$h_N(t) := \frac{1}{\omega_N} \sin^{N-1}(t) \quad \text{for } t \in [0, \pi], \tag{2-9}$$

where $\omega_N := \int_0^\pi \sin^{N-1}(t) dt$. Let $\epsilon := \pi - D$ and $\lambda_D := \int_0^D h_N(t) dt$ for any $D \in [0, \pi]$.

For a proof of the next proposition see for instance [Cavalletti et al. 2019, Proposition A.3].

Proposition 2.6. *Let $h : [0, D] \rightarrow [0, +\infty)$ be a $CD(N - 1, N)$ density which integrates to 1 on $[0, D]$. Then, for any $t \in (0, D)$, it holds*

$$\left(\frac{\omega_N}{\omega_N \lambda_D + \epsilon} \right) \min\{h_N(t), h_N(t + \epsilon)\} \leq h(t) \leq \left(\frac{\omega_N}{\omega_N - \epsilon} \right) \max\{h_N(t), h_N(t + \epsilon)\}. \tag{2-10}$$

Corollary 2.7. *Under the assumptions of Proposition 2.6, there exist a constant $C = C(N) > 0$ and $\epsilon_0 > 0$ with the following property: if $\epsilon \in [0, \epsilon_0]$ then for any $t \in (0, D)$ it holds*

$$|h(t) - h_N(t)| \leq C\epsilon. \tag{2-11}$$

Moreover, for $r \in (0, \frac{1}{10})$ and $\epsilon \in (0, \frac{1}{10}r)$ the following improved estimate holds:

$$|h(t) - h_N(t)| \leq Cr^{N-2}\epsilon \quad \text{for all } t \in ([0, r] \cup [\pi - r, D]). \tag{2-12}$$

Proof. The validity of (2-11) follows from (2-10) taking into account the Lipschitz continuity of h_N and the asymptotic expansions of

$$\frac{\omega_N}{\omega_N \lambda_D + \epsilon} \quad \text{and} \quad \frac{\omega_N}{\omega_N - \epsilon},$$

as $\epsilon \rightarrow 0$. The improved estimate (2-12) on $([0, r] \cup [\pi - r, D])$ follows analogously from (2-10) and the mean value theorem. □

Armed with Corollary 2.7 we can prove that, if $D \in (0, \pi)$ is close to π , then the integrals of the functions \sin and \cos (and of any bounded function, more in general) with respect to a $CD(N - 1, N)$ density h defined on $[0, D]$ do not differ much from the value of the corresponding integrals computed with respect to the model density h_N .

Corollary 2.8. *Let $f : [0, \pi] \rightarrow [-1, 1]$ be Borel measurable. Define $\mathfrak{m}(dt) := h(t)\mathcal{L}^1(dt)$ and $\mathfrak{m}_N(dt) := h_N(t)\mathcal{L}^1(dt)$. Under the assumptions of Proposition 2.6, there exist a constant $C = C(N) > 0$ and $\epsilon_0 > 0$ with the following property: if $\epsilon \in [0, \epsilon_0]$ then*

$$\left| \int_0^D f(t) \mathfrak{m}(dt) - \int_0^\pi f(t) \mathfrak{m}_N(dt) \right| \leq C\epsilon. \tag{2-13}$$

Moreover, for any $r \in (0, \frac{1}{10})$ and $\epsilon \in (0, \frac{1}{10}r)$ the following improved estimate holds

$$\left| \int_0^r f(t) \mathfrak{m}(dt) - \int_0^r f(t) \mathfrak{m}_N(dt) \right| + \left| \int_{\pi-r}^D f(t) \mathfrak{m}(dt) - \int_{\pi-r}^\pi f(t) \mathfrak{m}_N(dt) \right| \leq C\epsilon r^{N-1}. \tag{2-14}$$

Proof. The conclusion follows from [Corollary 2.7](#) just by integrating on $[0, D]$ and taking into account that $|\int_D^\pi f \, m_N| \leq C\epsilon^N$. \square

2D. Localization and L^1 -optimal transportation. The localization technique has its roots in a work of Payne and Weinberger [\[1960\]](#) and has been developed by Gromov and Milman [\[1987\]](#), Lovász and Simonovits [\[1993\]](#) and Kannan, Lovász and Simonovits [\[Kannan et al. 1995\]](#) in the setting of Euclidean spaces, spheres and Hilbert spaces. The basic idea is to reduce an n -dimensional problem, via tools of convex geometry, to lower-dimensional problems which are easier to handle. In the aforementioned papers, the symmetries of the spaces were heavily used to obtain such a dimensional reduction, typically via iterative bisections. Recently Klartag [\[2017\]](#) found a bridge between L^1 -optimal transportation problems and the localization technique yielding the localization theorem in the framework of smooth Riemannian manifolds. Inspired by this approach, the first and the second author in [\[Cavalletti and Mondino 2017b\]](#) proved a localization theorem for essentially nonbranching metric measure spaces satisfying the $CD(K, N)$ condition. Before stating the result it is worth recalling some basics about the disintegration of a measure associated to a partition (for a comprehensive treatment see the monograph [\[Fremlin 2006\]](#); for a discussion closer to the spirit of this paper see [\[Bianchini and Caravenna 2009\]](#); for a one-page summary see [\[Cavalletti et al. 2019, Appendix B\]](#)).

Given a measure space (X, \mathcal{X}, m) , suppose a *partition* of X into *disjoint* sets is given by $\{X_q\}_{q \in Q}$ so that $X = \bigcup_{q \in Q} X_q$. Here Q is the set of indices and $\Omega : X \rightarrow Q$ is the quotient map, i.e.,

$$q = \Omega(x) \iff x \in X_q.$$

We endow Q with the *push forward σ -algebra* \mathcal{Q} of \mathcal{X} :

$$C \in \mathcal{Q} \iff \Omega^{-1}(C) \in \mathcal{X},$$

i.e., the biggest σ -algebra on Q such that Ω is measurable. Moreover, the push forward measure $q := \Omega_\# m$ defines a natural measure q on (Q, \mathcal{Q}) . The triple (Q, \mathcal{Q}, q) is called *the quotient measure space*.

Definition 2.9 (consistent and strongly consistent disintegration). A *disintegration* of m consistent with the partition is a map

$$Q \ni q \mapsto m_q \in \mathcal{P}(X, \mathcal{X})$$

such that the following requirements hold:

- (1) For all $B \in \mathcal{X}$, the map $q \mapsto m_q(B)$ is q -measurable.
- (2) For all $B \in \mathcal{X}$ and $C \in \mathcal{Q}$, the following consistency condition holds:

$$m(B \cap \Omega^{-1}(C)) = \int_C m_q(B) \, q(dq).$$

A disintegration of m is called *strongly consistent* if in addition:

- (3) For q -a.e. $q \in Q$, m_q is concentrated on $X_q = \Omega^{-1}(q)$.

In the next theorem, for q -a.e. $q \in Q$, the equivalence class X_q is a geodesic in X . With a slight abuse of notation X_q denotes also the arc-length parametrization on a real interval of the corresponding geodesic; i.e., it is a map from a real interval with image X_q . We will use the following terminology: $q \mapsto m_q$ is a $CD(K, N)$ disintegration if, for q -a.e. $q \in Q$, $m_q = h_q \mathcal{H}^1 \llcorner_{X_q}$, where \mathcal{H}^1 denotes the one-dimensional Hausdorff measure and $h_q \circ X_q$ is a $CD(K, N)$ density, in the sense of (2-7).

Theorem 2.10 [Cavalletti and Mondino 2017b]. *Let (X, d, m) be an essentially nonbranching metric measure space satisfying the $CD(K, N)$ condition for some $K \in \mathbb{R}$ and $N \in [1, \infty)$. Let $f : X \rightarrow \mathbb{R}$ be m -integrable such that $\int_X f m = 0$, and assume the existence of $x_0 \in X$ such that $\int_X |f(x)| d(x, x_0) m(dx) < \infty$.*

Then the space X admits a partition $\{X_q\}_{q \in Q}$ and a corresponding (strongly consistent) disintegration of m , $\{m_q\}_{q \in Q}$, such that:

- For any m -measurable set $B \subset \mathcal{T}$ it holds

$$m(B) = \int_Q m_q(B) q(dq),$$

where q is a probability measure over Q defined on the quotient σ -algebra \mathcal{Q} .

- For q -almost every $q \in Q$, the set X_q is a geodesic (possibly of zero length) and m_q is supported on it. Moreover $q \mapsto m_q$ is a $CD(K, N)$ disintegration.
- For q -almost every $q \in Q$, it holds $\int_{X_q} f m_q = 0$.

In Theorem 2.10 we can also distinguish the set of X_α having positive length, whose union forms the so-called *transport set* denoted by \mathcal{T} , from the ones having zero length, i.e., points, whose union we usually denote by Z , so to have a decomposition of X into \mathcal{T} and Z . The last point of Theorem 2.10 implies then that m -a.e. $f \equiv 0$ on Z .

Following the approach of [Klartag 2017], Theorem 2.10 was proven in [Cavalletti and Mondino 2017b] studying the following optimal transportation problem. Let $\mu_0 := f^+ m$ and $\mu_1 := f^- m$, where f^\pm denote the positive and the negative parts of f respectively, and study the L^1 -optimal transport problem associated with it:

$$\inf \left\{ \int_{X \times X} d(x, y) \pi(dx dy) : \pi \in \mathcal{P}(X \times X), (P_1)_\# \pi = \mu_0, (P_2)_\# \pi = \mu_1 \right\}. \tag{2-15}$$

Then the relevant object to study is given by the dual formulation of the previous minimization problem. By the summability properties of f (see the hypotheses of Theorem 2.10), there exists a 1-Lipschitz function $\phi : X \rightarrow \mathbb{R}$ such that π is a minimizer in (2-15) if and only if $\pi(\Gamma) = 1$, where

$$\Gamma := \{(x, y) \in X \times X : \phi(x) - \phi(y) = d(x, y)\}$$

is the naturally associated d -cyclically monotone set; i.e., for any $(x_1, y_1), \dots, (x_n, y_n) \in \Gamma$ it holds

$$\sum_{i=1}^n d(x_i, y_i) \leq \sum_{i=1}^n d(x_i, y_{i+1}), \quad y_{n+1} = y_1,$$

for any $n \in \mathbb{N}$. The set Γ induces a partial order relation whose maximal chains produce a partition (up to an m -negligible subset) of the set $\mathcal{T} \subset X$ appearing in the statement of [Theorem 2.10](#), made of one-dimensional subsets. For a summary of the constructions see [[Cavalletti et al. 2019](#), Section 2.5]; for more details see [[Cavalletti and Mondino 2017b](#); [Cavalletti and Milman 2021](#)].

3. One-dimensional estimates

The goal of this section is to give a self-contained presentation of the one-dimensional estimates we will use throughout the paper.

3A. Bérard–Besson–Gallot explicit lower bound on the model isoperimetric profile. For $N > 1$, let

$$\omega_N := \int_0^\pi (\sin t)^{N-1} dt \quad \text{and} \quad m_N := \frac{1}{\omega_N} (\sin t)^{N-1} \mathcal{L}^1(dt) \llcorner_{[0,\pi]}. \tag{3-1}$$

From now on fix $D \in (0, \pi)$. For $b \in [0, \pi - D]$ and $v \in [0, 1]$, let $R(b, v) \in [b, \pi]$ be uniquely defined by the equation

$$\int_b^{R(b,v)} (\sin t)^{N-1} dt = v \int_b^{b+D} (\sin t)^{N-1} dt. \tag{3-2}$$

Set

$$\mathcal{I}_{N,D}(v) := \inf\{g(b, v) : b \in [0, \pi - D]\}, \tag{3-3}$$

where

$$g(b, v) := \frac{[\sin(R(b, v))]^{N-1}}{\int_b^{b+D} (\sin t)^{N-1} dt}. \tag{3-4}$$

To keep notation short, we also set $\mathcal{I}_N := \mathcal{I}_{N,\pi}$. Notice that \mathcal{I}_N is the isoperimetric profile of \mathbb{S}^N for an integer N . We refer to [Section 4](#) for a brief discussion about the isoperimetric profile; note also that $\mathcal{I}_{N,D}$ is the model isoperimetric profile in the Lévy–Gromov isoperimetric comparison theorem for spaces with Ricci $\geq N - 1$, dimension $\leq N$ and diameter $\leq D$; see [[Gromov 1999](#), Appendix C; [Bérard et al. 1985](#); [Milman 2015](#); [Cavalletti and Mondino 2017b](#)].

The proof of the next lemma is inspired by, but somewhat different from, [[Bérard et al. 1985](#), Appendix 1] and the statement generalizes to arbitrary real $N > 1$ the result stated in the reference for an integer $N \geq 2$.

Lemma 3.1 (Bérard–Besson–Gallot explicit isoperimetric lower bound). *Fix $N > 1$ and $D \in [0, \pi]$, and let $\mathcal{I}_{N,D} : [0, 1] \rightarrow [0, \infty)$ be defined in (3-3). Then*

$$\frac{\mathcal{I}_{N,D}(v)}{\mathcal{I}_N(v)} \geq \left(\frac{\int_0^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{1/N} =: C_{N,D} \geq 1 \quad \text{for all } v \in (0, 1). \tag{3-5}$$

Proof. Let $v' \in (0, 1)$ and $f : [0, \pi - D] \times (0, 1) \rightarrow [0, +\infty)$ be defined by

$$v' := \frac{1}{\omega_N} \int_0^{R(b,v)} (\sin t)^{N-1} dt \quad \text{and} \quad f(b, v) := \frac{g(b, v)}{\mathcal{I}_N(v)}. \tag{3-6}$$

In particular

$$\mathcal{I}_N(v') = \frac{1}{\omega_N} [\sin R(b, v)]^{N-1}, \tag{3-7}$$

and therefore

$$f(b, v) = \omega_N \left(\int_b^{b+D} (\sin t)^{N-1} dt \right)^{-1} \frac{\mathcal{I}_N(v')}{\mathcal{I}_N(v)}. \tag{3-8}$$

Thanks to the explicit expression of the isoperimetric profile \mathcal{I}_N it is possible to compute

$$(\mathcal{I}_N^{N/(N-1)})'' \mathcal{I}_N^{(N-2)/(N-1)} = -N. \tag{3-9}$$

In particular it follows from (3-9) that $\mathcal{I}_N^{N/(N-1)}$ is concave on $(0, 1)$.

We now distinguish two cases: $v' \leq v$ and $v' > v$.

Case 1: $v' \leq v$. First observe that

$$\omega_N v' = \int_0^{R(b,v)} (\sin t)^{N-1} dt \geq \int_b^{R(b,v)} (\sin t)^{N-1} dt = v \int_b^{b+D} (\sin t)^{N-1} dt. \tag{3-10}$$

The concavity observed above, together with (3-10), gives that

$$\frac{\mathcal{I}_N(v')}{\mathcal{I}_N(v)} \geq \left(\frac{v'}{v} \right)^{1-1/N} \geq \left(\omega_N^{-1} \int_b^{b+D} (\sin t)^{N-1} dt \right)^{1-1/N}.$$

Hence, taking into account (3-8), we obtain

$$f(b, v) \geq \omega_N^{1/N} \left(\int_b^{b+D} (\sin t)^{N-1} dt \right)^{-1/N}. \tag{3-11}$$

Case 2: $v' > v$. A change of variables in the definition of R easily yields

$$R(\pi - b - D, 1 - v) = \pi - R(b, v)$$

and therefore

$$f(b, v) = f(\pi - b - D, 1 - v). \tag{3-12}$$

Moreover

$$\int_0^{R(\pi-b-D, 1-v)} (\sin t)^{N-1} dt = \int_{R(b,v)}^{\pi} (\sin t)^{N-1} dt = (1 - v') \omega_N;$$

hence

$$f(\pi - b - D, 1 - v) = \omega_N \left(\int_b^{b+D} (\sin t)^{N-1} dt \right)^{-1} \frac{\mathcal{I}_N(1 - v')}{\mathcal{I}_N(1 - v)}. \tag{3-13}$$

Next we observe that, as in the previous case, the concavity of $\mathcal{I}_N^{N/(N-1)}$ yields

$$\frac{\mathcal{I}_N(1 - v')}{\mathcal{I}_N(1 - v)} \geq \left(\frac{1 - v'}{1 - v} \right)^{1-1/N}. \tag{3-14}$$

Moreover, it holds

$$\omega_N(1 - v') = \int_{R(b,v)}^{\pi} (\sin t)^{N-1} dt \geq \int_b^{b+D} (\sin t)^{N-1} dt = (1 - v) \int_b^{b+D} (\sin t)^{N-1} dt. \tag{3-15}$$

Combining (3-13), (3-14) and (3-15) and taking into account (3-12), we get

$$f(b, v) \geq \omega_N^{1/N} \left(\int_b^{b+D} (\sin t)^{N-1} dt \right)^{-1/N}. \tag{3-16}$$

It is now sufficient to observe that the function $x \mapsto \int_x^{x+D} (\sin t)^{N-1} dt$ attains its maximum at $x = \pi/2 - D/2$ in order to obtain from (3-11), (3-16), (3-6) and (3-3) that

$$\frac{\mathcal{I}_{N,D}(v)}{\mathcal{I}_N(v)} \geq \left(\frac{\omega_N}{\int_{\pi/2-D/2}^{\pi/2+D/2} (\sin t)^{N-1} dt} \right)^{1/N} = C_{N,D} \quad \text{for all } v \in (0, 1).$$

Above, the last identity follows from the expression for $C_{N,D}$ introduced in (3-5) thanks to the identity $\cos(\pi/2 - x) = \sin(x)$ and a change of variables. □

Let us study the behavior of $C_{N,D}$ in the asymptotic $D \rightarrow \pi$.

Lemma 3.2. *It holds that*

$$\lim_{D \rightarrow \pi} \frac{(\pi - D)^N}{C_{N,D}^2 - 1} = 2^{N-1} N^2 \int_0^{\pi/2} (\cos t)^{N-1} dt. \tag{3-17}$$

Hence there exist $\bar{C} = \bar{C}(N) > 0$ and $\bar{D} = \bar{D}(N) < \pi$ such that

$$C_{N,D}^2 - 1 \geq \bar{C}(\pi - D)^N \quad \text{for all } D \in [\bar{D}, \pi]. \tag{3-18}$$

Proof. Recalling the expression of $C_{N,D}$ from (3-5), we have

$$C_{N,D}^2 - 1 = \left(\frac{\int_0^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{2/N} - 1 = \left(1 + \frac{\int_{D/2}^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{2/N} - 1.$$

Now, as $D \rightarrow \pi$, we have the expansion

$$\int_{D/2}^{\pi/2} (\cos t)^{N-1} dt = \int_0^{\pi/2-D/2} (\sin t)^{N-1} dt \sim \int_0^{\pi/2-D/2} s^{N-1} ds \sim \frac{1}{N} \left(\frac{\pi}{2} - \frac{D}{2} \right)^N.$$

Taking into account the asymptotic $(1 + x)^\beta - 1 \sim \beta x$, we obtain (3-17).

The second conclusion in the statement easily follows from the first one. □

3B. Spectral gap and diameter. Building on top of the lower bound of the isoperimetric profile obtained in Lemma 3.1, we next obtain a quantitative spectral gap inequality for Neumann boundary conditions in terms diameters.

The analogous result in the case of smooth Riemannian manifolds was established in [Croke 1982, Theorem B] building upon a quantitative improvement of the Lévy–Gromov inequality and on [Bérard and Meyer 1982] (see also [Bérard et al. 1985, Corollary 17]). The usual strategy to show the improved Neumann spectral gap inequality is based on the observation that a Neumann first eigenfunction of the Laplacian f is a Dirichlet first eigenfunction of the Laplacian on the domains $\{f > 0\}$ and $\{f < 0\}$ (see, for instance, [Matei 2000, Lemma 3.2]). The improved Dirichlet spectral gap inequality is then obtained by rearrangement starting from the isoperimetric inequality.

Proposition 3.3 (1-dimensional quantitative Obata's theorem on the diameter). *Let (I, d_{eucl}, m) be a one-dimensional $CD(N - 1, N)$ m.m.s. with $\text{diam}(I) \leq D$. Then*

$$\frac{\lambda_{(I, d_{\text{eucl}}, m)}^{1,2}}{N} \geq C_{N,D}^2 = \left(\frac{\int_0^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{2/N}, \tag{3-19}$$

where $C_{N,D}$ was defined in (3-5).

In particular, there exists a constant $C_N > 0$ (more precisely one can choose $C_N = \bar{C}N$, where \bar{C} was defined in Lemma 3.2) such that

$$C_N(\pi - \text{diam}(I))^N \leq \lambda_{(I, d_{\text{eucl}}, m)}^{1,2} - N. \tag{3-20}$$

Proof. From [Bakry and Qian 2000] (see also [Cavalletti and Mondino 2017c, Section 4.1] for the regularization procedure) we know that $\lambda_{(I, d_{\text{eucl}}, m)}^{1,2} \geq \lambda_{N,D}^{1,2}$ where $\lambda_{N,D}^{1,2}$ is the first solution $\lambda > 0$ of the eigenvalue problem

$$\ddot{w} + (N - 1) \tan(t)\dot{w} + \lambda w = 0, \tag{3-21}$$

on $[-D/2, D/2]$ with Neumann boundary conditions. The eigenfunction associated to the first eigenvalue in (3-21) is unique, up to a multiplicative constant. Therefore, denoting it by $w_{N,D} : [-D/2, D/2] \rightarrow (-\infty, +\infty)$, it holds $w_{N,D}(-x) = -w_{N,D}(x)$ for any $x \in [-D/2, D/2]$ as a consequence of the symmetry of (3-21). In particular $w_{N,D}(0) = 0$. Let

$$m_{N,D} := \Lambda_{N,D}(\cos t)^{N-1} \mathcal{L}^1 \llcorner_{[-D/2, D/2]},$$

with $\Lambda_{N,D}$ such that $m_{N,D}$ is a probability measure. Note that $([-D/2, D/2], d_{\text{eucl}}, m_{N,D})$ is a $CD(N - 1, N)$ m.m.s. with diameter equal to D and $m_{N,D}([-D/2, 0]) = m_{N,D}([0, D/2]) = \frac{1}{2}$. Hence

$$\lambda_{N,D}^{1,2} = \frac{\int_{-D/2}^{D/2} |w'_{N,D}|^2 m_{N,D}}{\int_{-D/2}^{D/2} |w_{N,D}|^2 m_{N,D}} = \frac{\int_0^{D/2} |w'_{N,D}|^2 m_{N,D}}{\int_0^{D/2} |w_{N,D}|^2 m_{N,D}} \geq \lambda_{N,D}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right),$$

where $\lambda_{N,D}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right)$ is the least first eigenvalue of the Laplacian with Dirichlet boundary conditions on one extremum for intervals of volume $\frac{1}{2}$ in $([-D/2, D/2], d_{\text{eucl}}, m_{N,D})$.

Moreover a coarea argument (see for instance [Bérard et al. 1985, Corollary 17; Mondino and Semola 2020, Proposition 3.13]) using Lemma 3.1 gives

$$\lambda_{N,D}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right) \geq C_{N,D}^2 \lambda_{N,\pi}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right).$$

Recalling that $\lambda_{N,\pi}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right) = \lambda_{N,\pi}^{1,2} = N$ (see for instance [Bakry and Qian 2000]), we conclude that

$$\lambda_{(I, d_{\text{eucl}}, m)}^{1,2} \geq \lambda_{N,D}^{1,2} \geq \lambda_{N,D}^{1,2,\mathcal{D}}\left(\frac{1}{2}\right) \geq N C_{N,D}^2. \tag{3-22}$$

The second part of the statement follows by choosing $D = \text{diam}(I)$ and applying Lemma 3.2. □

A converse of the inequality proved in Proposition 3.3 can be obtained as follows.

Lemma 3.4. *For any $N > 1$ there exists $C = C(N) > 0$ such that if $([0, D], d_{\text{eucl}}, m)$ is a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. with $D \geq \pi - \epsilon$ then*

$$|\lambda_{([0,D],d_{\text{eucl}},m)}^{1,2} - N| \leq C\epsilon.$$

Proof. By the Lichnerowicz spectral gap we already know that $\lambda_{([0,D],d_{\text{eucl}},m)}^{1,2} \geq N$. It is therefore enough to prove the existence of $u \in \text{Lip}([0, D])$ such that

$$\|u\|_{L^2([0,D],m)} = 1, \quad \int_{[0,D]} u \, m = 0, \quad \int_{[0,D]} |u'|^2 \, m \leq N + C_N \epsilon. \tag{3-23}$$

Setting $u_N^*(t) := \sqrt{N+1} \cos(t)$ and using [Corollary 2.8](#) we get

$$\left| \int_{[0,D]} u_N^* \, m \right| \leq C_N \epsilon, \quad \left| 1 - \int_{[0,D]} |u_N^*|^2 \, m \right| \leq C_N \epsilon, \quad \int_{[0,D]} |(u_N^*)'|^2 \, m \leq N + C_N \epsilon. \tag{3-24}$$

Let $v = u_N^* - \int_{[0,D]} u_N^* \, m$ and $c_v := \|v\|_{L^2([0,D],m)}$. Using the estimates [\(3-24\)](#), it is straightforward to check that $u = (1/c_v)v$ satisfies [\(3-23\)](#). □

3C. Spectral gap and shape of eigenfunctions. Next we establish some basic estimates on eigenfunctions which will be useful later.

Given a one-dimensional $\text{CD}(K, N)$ space (I, d_{eucl}, m) , we know that we can write $m(dt) = h\mathcal{L}^1(dt)$ for some $\text{CD}(K, N)$ density h . We start by recalling the definition and basic properties of the Laplace operator Δ . A function $u \in W^{1,2}(I, m)$ is said to be in the domain of Δ , and we write $u \in \text{Dom}(\Delta)$ if for every $\phi \in C_c^\infty(I)$ it holds

$$\left| \int_I u' \phi' \, m \right| \leq C_u \|\phi\|_{L^2(I,m)}$$

for some $C_u \geq 0$ depending on u . In this case, by the Riesz theorem, there exists a function $\Delta u \in L^2(I, m)$ such that

$$- \int_I u' \phi' \, m = \int_I \Delta u \, \phi \, m.$$

It is readily seen that the operator $\text{Dom}(\Delta) \ni u \mapsto \Delta u \in L^2(I, m)$ is linear.

Moreover, using the properties of $\text{CD}(K, N)$ densities recalled at the beginning of the section, it holds that every $u \in \text{Dom}(\Delta)$ is twice differentiable \mathcal{L}^1 -a.e. on I and

$$\Delta u = u'' + (\log h)'u', \quad \mathcal{L}^1\text{-a.e. on } I, \text{ for all } u \in \text{Dom}(\Delta). \tag{3-25}$$

Proposition 3.5. *Let (I, d_{eucl}, m) be a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. Then there exists a constant $C = C(N) > 0$ such that, if u is an eigenfunction of the Laplacian on (I, d_{eucl}, m) associated to an eigenvalue $\lambda \in [N, 2N]$ and with $\|u\|_2 = 1$, then $u \in W_{\text{loc}}^{2,2}(I, d_{\text{eucl}}, \mathcal{L}^1)$ and*

$$\|u'' + u\|_{L^2(m)} \leq C(\lambda - N)^{1/2}. \tag{3-26}$$

Proof. Step 1: We claim that it holds

$$\int_I \left(u'' - \frac{1}{N} \Delta u \right)^2 \, m \leq \int_I \left(\frac{N-1}{N} (\Delta u)^2 - (N-1)(u')^2 \right) \, m. \tag{3-27}$$

Since by assumption $u \in W^{1,2}(I, d_{\text{eucl}}, \mathfrak{m})$ is an eigenfunction we have $-\Delta u \in W^{1,2}(I, d_{\text{eucl}}, \mathfrak{m})$ as well. Thus we can define the Γ_2 operator as

$$\Gamma_2(u; \phi) := \int_I \left(\frac{1}{2}(u')^2 \Delta \phi - (\Delta u)' u' \phi \right) \mathfrak{m} \tag{3-28}$$

for all $\phi \in L^\infty(I, \mathfrak{m})$ with $\Delta \phi \in L^\infty(I, \mathfrak{m})$. Using that h satisfies (2-8), a manipulation via integration by parts gives that for all $\phi \geq 0$ as above it holds:

$$\Gamma_2(u; \phi) \geq \int_I \left[(u'')^2 + (N-1)(u')^2 + \frac{1}{N-1}(\Delta u - u'')^2 \right] \phi \mathfrak{m}. \tag{3-29}$$

By direct computations, one can check that

$$\begin{aligned} & (u'')^2 + (N-1)(u')^2 + \frac{1}{N-1}(\Delta u - u'')^2 \\ &= (N-1)(u')^2 + \left(u'' - \frac{1}{N} \Delta u \right)^2 + \frac{1}{N}(\Delta u)^2 + \frac{1}{N-1} \left(u'' - \frac{1}{N} \Delta u \right)^2 \quad \mathfrak{m}\text{-a.e.} \end{aligned} \tag{3-30}$$

Plugging (3-30) into (3-29) gives

$$\Gamma_2(u; \phi) \geq \int_I \left[(N-1)(u')^2 + \left(u'' - \frac{1}{N} \Delta u \right)^2 + \frac{1}{N}(\Delta u)^2 \right] \phi \mathfrak{m}.$$

Choosing $\phi \equiv 1$ yields (3-27).

Step 2: Inserting the eigenvalue relation $\lambda u = -\Delta u$ into (3-27), we obtain

$$\int_I \left(u'' + \frac{\lambda}{N} u \right)^2 \mathfrak{m} \leq \int_I \left(\frac{N-1}{N} (\lambda u)^2 - (N-1)(u')^2 \right) \mathfrak{m} = \frac{N-1}{N} \lambda (\lambda - N) \int_I u^2 \mathfrak{m}. \tag{3-31}$$

Eventually,

$$\begin{aligned} \int_I (u'' + u)^2 \mathfrak{m} &\leq 2 \int_I \left| u'' + \frac{\lambda}{N} u \right|^2 \mathfrak{m} + 2 \int_I \left| \frac{\lambda - N}{N} u \right|^2 \mathfrak{m} \\ &\leq 2 \left(\frac{N-1}{N} \lambda (\lambda - N) + \frac{(\lambda - N)^2}{N^2} \right) \int_I u^2 \mathfrak{m} \leq C(N) (\lambda - N) \int_I u^2 \mathfrak{m}, \end{aligned}$$

where, in the last estimate, we used the assumption $\lambda \leq 2N$. □

The aim of the remaining part of this section is to prove [Theorem 3.11](#) stating roughly that, on any one-dimensional $\text{CD}(N-1, N)$ m.m.s. $(I, d_{\text{eucl}}, \mathfrak{m})$, a function $u : I \rightarrow \mathbb{R}$ whose 2-Rayleigh quotient is close to N (the optimal one on the model $(N-1, N)$ -space) and with L^2 -norm equal to 1, is $W^{1,2}$ -close to the (normalized) cosine of the distance from one of the extrema of the interval, in quantitative terms.

The conclusion of [Theorem 3.11](#) will be achieved through some intermediate steps. First we estimate the $W^{1,2}$ -closeness of a first eigenfunction u^* for $(I, d_{\text{eucl}}, \mathfrak{m})$ with the cosine of the distance from one of the extremes of the segment, see [Proposition 3.6](#). Then, we bound the $W^{1,2}$ -closeness of the function u from u^* (or $-u^*$), see [Proposition 3.10](#).

Let us observe that

$$\|\cos(\cdot)\|_{L^2(\mathfrak{m}_N)} = \frac{1}{\sqrt{N+1}}, \tag{3-32}$$

and, by symmetry,

$$\int_{[0,\pi]} \cos(t) \mathfrak{m}_N(dt) = 0. \tag{3-33}$$

Proposition 3.6. *For every $N > 1$ there exist constants $C = C(N) > 0$ and $\epsilon_0 = \epsilon_0(N) > 0$ such that for every one dimensional $\text{CD}(N - 1, N)$ m.m.s. $([0, D], d_{\text{eucl}}, \mathfrak{m})$ and every Neumann eigenfunction u^* , with $\|u^*\|_{L^2(\mathfrak{m})} = 1$, of eigenvalue $\lambda \in [N, 2N]$ it holds*

$$\min\{\|u^* - \sqrt{N + 1} \cos(\cdot)\|_{L^2(\mathfrak{m})}, \|u^* + \sqrt{N + 1} \cos(\cdot)\|_{L^2(\mathfrak{m})}\} \leq C\delta^{\min\{1/2, 1/N\}}, \tag{3-34}$$

where $\delta := \int |\nabla u^*|^2 \mathfrak{m} - N < \epsilon_0$. Furthermore the conclusion can be improved to $W^{1,2}$ -closeness:

$$\min\{\|(u^* - \sqrt{N + 1} \cos(\cdot))'\|_{L^2(\mathfrak{m})}, \|(u^* + \sqrt{N + 1} \cos(\cdot))'\|_{L^2(\mathfrak{m})}\} \leq C\delta^{\min\{1/2, 1/N\}}. \tag{3-35}$$

Proof. Let $h : [0, D] \rightarrow [0, +\infty)$ be the density of \mathfrak{m} with respect to \mathcal{L}^1 and let $x_0 \in (0, D)$ be a maximum point of h . In [Cavalletti et al. 2019, Lemma A.4] it is proved that such a maximum point is unique and that h is strictly increasing on $[0, x_0]$ and strictly decreasing on $[x_0, D]$.

Step 1: In this first step we prove that, given $z \in L^2([0, D], \mathfrak{m})$, any solution of $v'' + v = z$ can be written as

$$v(t) = \int_{x_0}^t \sin(t - s)z(s) ds + \alpha \sin(t) + \beta \cos(t) \tag{3-36}$$

for some $\alpha, \beta \in \mathbb{R}$. To this aim, it suffices to prove that

$$v_0(t) := \int_{x_0}^t \sin(t - s)z(s) ds \tag{3-37}$$

solves $v'' + v = z$. First we observe that v_0 is well-defined, since the assumption $z \in L^2((0, D), \mathfrak{m})$ guarantees that $z \in L^1_{\text{loc}}((0, D), \mathcal{L}^1)$ (due to the fact that h is locally bounded from below by a strictly positive constant in the interior of $[0, D]$). The fact that it satisfies $v''_0 + v_0 = z$ follows from an elementary computation.

Step 2: Next, we prove that the function v_0 defined in (3-37) satisfies

$$\|v_0\|_{L^2(\mathfrak{m})} \leq \pi \|z\|_{L^2(\mathfrak{m})}. \tag{3-38}$$

Indeed, taking into account that $|\sin| \leq 1$, applying the Cauchy–Schwarz inequality, Fubini’s theorem and recalling that h is increasing on $[0, x_0]$ and decreasing on $[x_0, D]$, we can compute

$$\begin{aligned} \|v_0\|_{L^2(\mathfrak{m})}^2 &= \int_0^D \left(\int_{x_0}^t \sin(t - s)z(s) ds \right)^2 h(t) dt \leq \pi \int_0^D h(t) \left| \int_{x_0}^t z^2(s) ds \right| dt \\ &= \pi \left(\int_0^{x_0} z^2(s) \int_0^s h(t) dt ds + \int_{x_0}^D z^2(s) \int_s^D h(t) dt ds \right) \\ &\leq \pi^2 \left(\int_0^{x_0} z^2(s)h(s) ds + \int_{x_0}^D z^2(s)h(s) ds \right) = \pi^2 \|z\|_{L^2(\mathfrak{m})}^2. \end{aligned}$$

Let us remark that from (3-38) it follows applying Cauchy–Schwarz inequality that $\|v_0\|_{L^1(\mathfrak{m})} \leq \pi \|z\|_{L^2(\mathfrak{m})}$.

Step 3: Recall from Proposition 3.3 the bound $\pi - D \leq C\delta^{1/N}$. Furthermore we know from (3-26) that if u^* is as in the assumptions of the statement, then $(u^*)'' + u^* = z$ on $[0, D]$ for some function z such that

$\|z\|_{L^2(\mathfrak{m})} \leq C\delta^{1/2}$. Hence, as proved in Step 1, u^* can be written as

$$u^*(t) = \int_{x_0}^t \sin(t-s)z(s) ds + \alpha \sin(t) + \beta \cos(t) \tag{3-39}$$

for some $\alpha, \beta \in \mathbb{R}$. We want to show that there exists $C = C(N) > 0$ such that $|\alpha| + |\beta| \leq C(N)$.

Set $u_0(t) := \int_{x_0}^t \sin(t-s)z(s) ds$ and recall that, from Step 2, it holds $\|u_0\|_{L^2(\mathfrak{m})} \leq C\delta^{1/2}$. Since by assumption u^* has null mean value, integrating (3-39) over $[0, D]$ with respect to \mathfrak{m} gives

$$0 = \alpha \int_{[0,D]} \sin(t)\mathfrak{m}(dt) + \beta \int_{[0,D]} \cos(t)\mathfrak{m}(dt) + \int_{[0,D]} u_0(t)\mathfrak{m}(dt). \tag{3-40}$$

From the last remark in Step 2 and Corollary 2.8, it follows that

$$\left(\int_{[0,\pi]} \sin^N(t) dt + O(\delta^{1/N}) \right) \alpha + O(\delta^{1/N})\beta + O(\delta^{1/2}) = 0,$$

giving that

$$\alpha = O(\delta^{1/N})\beta + O(\delta^{1/2}). \tag{3-41}$$

In order to estimate β , we compute the $L^2(\mathfrak{m})$ -norm squared of both the left- and right-hand sides of (3-39) to obtain

$$1 = \|u_0\|_{L^2(\mathfrak{m})}^2 + \alpha^2 \|\sin(\cdot)\|_{L^2(\mathfrak{m})}^2 + \beta^2 \|\cos(\cdot)\|_{L^2(\mathfrak{m})}^2 + 2\alpha \int u_0(t) \sin(t) \mathfrak{m}(dt) + 2\beta \int u_0(t) \cos(t) \mathfrak{m}(dt) + 2\alpha\beta \int \sin(t) \cos(t) \mathfrak{m}(dt). \tag{3-42}$$

Plugging (3-41) into (3-42), gives

$$(1 + O(\delta)) + O(\delta^{1/N+1/2}) \beta + \left(\int_{[0,\pi]} \cos^2(t) \sin^{N-1}(t) dt + O(\delta^{1/N}) \right) \beta^2 = 0, \tag{3-43}$$

yielding $|\beta| \leq C(N)$ and thus, by (3-41), also $|\alpha| \leq C(N)$.

Step 4: Conclusion. In order to get (3-34), we have to bound $|\alpha|$ and $\min\{|\sqrt{N+1} - \beta|, |\sqrt{N+1} + \beta|\}$ in terms of δ .

From (3-40), Step 3, the last remark in Step 2 and Corollary 2.8 it follows that

$$|\alpha| \leq C(\delta^{1/2} + \delta^{1/N}) \leq C\delta^{\min\{1/N, 1/2\}}, \tag{3-44}$$

up to increasing the value of the constant C in the second inequality. Plugging (3-44) into (3-42) gives

$$1 = O(\delta) + O(\delta^{\min\{1, 2/N\}}) + O(\delta^{1/2}) + O(\delta^{\min\{1/2, 1/N\}}) + \beta^2/(N+1)$$

and therefore

$$\left| 1 - \frac{\beta^2}{N+1} \right| = O(\delta^{\min\{1/2, 1/N\}}). \tag{3-45}$$

From (3-45) we easily obtain that

$$\min\{|\sqrt{N+1} - \beta|, |\sqrt{N+1} + \beta|\} \leq C\delta^{\min\{1/4, 1/(2N)\}}. \tag{3-46}$$

In the case

$$|\sqrt{N+1} - \beta| = \min\{|\sqrt{N+1} - \beta|, |\sqrt{N+1} + \beta|\} \leq C\delta^{\min\{1/4, 1/(2N)\}}$$

(respectively $|\sqrt{N+1} + \beta| = \min\{|\sqrt{N+1} - \beta|, |\sqrt{N+1} + \beta|\} \leq C\delta^{\min\{1/4, 1/(2N)\}}$), it follows that

$$|\sqrt{N+1} + \beta| \geq 2\sqrt{N+1} - C\delta^{\min\{1/4, 1/(2N)\}} \geq \sqrt{N+1} \quad \text{for } \delta \leq \delta_0(N). \tag{3-47}$$

(resp. $|\sqrt{N+1} - \beta| \geq \sqrt{N+1}$). Plugging (3-47) back into (3-45) gives $|\sqrt{N+1} - \beta| \leq C\delta^{\min\{1/2, 1/N\}}$ (resp. $|\sqrt{N+1} + \beta| \leq C\delta^{\min\{1/2, 1/N\}}$). In conclusion, (3-45) and (3-46) can be bootstrapped to give

$$\min\{|\sqrt{N+1} - \beta|, |\sqrt{N+1} + \beta|\} \leq C\delta^{\min\{1/2, 1/N\}}. \tag{3-48}$$

Combining all these ingredients we can eventually estimate the $L^2(\mathfrak{m})$ -distance between the first Neumann eigenfunction and the normalized cosine. Indeed, assuming without loss of generality that $|\sqrt{N+1} - \beta| \leq |\sqrt{N+1} + \beta|$ and taking into account (3-44), (3-48), we obtain

$$\begin{aligned} \|u^* - \sqrt{N+1} \cos(\cdot)\|_{L^2(\mathfrak{m})} &= \|u_0 + \alpha \sin(\cdot) + \beta \cos(\cdot) - \sqrt{N+1} \cos(\cdot)\|_{L^2(\mathfrak{m})} \\ &\leq |\alpha| \|\sin(\cdot)\|_{L^2(\mathfrak{m})} + \|u_0\|_{L^2(\mathfrak{m})} + |\beta - \sqrt{N+1}| \|\cos(\cdot)\|_{L^2(\mathfrak{m})} \\ &\leq C\delta^{\min\{1/2, 1/N\}}. \end{aligned}$$

Finally, we improve the $L^2(\mathfrak{m})$ -closeness to $W^{1,2}(\mathfrak{m})$ -closeness. To this aim, differentiate (3-39) to obtain

$$(u^*)'(t) = \int_{x_0}^t \cos(t-s)z(s) ds + \alpha \cos(t) - \beta \sin(t). \tag{3-49}$$

With computations analogous to the ones used to obtain the bound $\|v_0\|_2 \leq \pi \|z\|_2$ in Step 2, one can prove that, letting $w_0(t) := \int_{x_0}^t \cos(t-s) ds$, it holds $\|w_0\|_2 \leq \pi \|z\|_2$. The sought estimate for

$$\min\{\|(u^* - \sqrt{N+1} \cos(\cdot))'\|_{L^2(\mathfrak{m})}, \|(u^* + \sqrt{N+1} \cos(\cdot))'\|_{L^2(\mathfrak{m})}\}$$

follows taking into account (3-44) and (3-46). □

We isolate the following corollary, which will be useful later in the paper.

Corollary 3.7. *Under the assumptions of Proposition 3.6, setting $r = \delta^{\gamma/N}$ for some $\gamma \in (0, 1)$, it holds*

$$\begin{aligned} \min\{\|u^* - \sqrt{N+1} \cos(\cdot)\|_{W^{1,2}([0,r],\mathfrak{m})}, \|u^* + \sqrt{N+1} \cos(\cdot)\|_{W^{1,2}([0,r],\mathfrak{m})}\} \\ \leq C(N)(\delta^{1/2} + r^{N/2} \delta^{\min\{1/2, 1/N\}}). \end{aligned} \tag{3-50}$$

Moreover, for $\eta \in (0, \frac{1}{10}r)$,

$$\begin{aligned} \min\{\|u^* - \sqrt{N+1} \cos(\cdot)\|_{W^{1,2}([r-\eta, r+\eta],\mathfrak{m})}, \|u^* + \sqrt{N+1} \cos(\cdot)\|_{W^{1,2}([r-\eta, r+\eta],\mathfrak{m})}\} \\ \leq C(N)(\delta^{1/2} + (r^{N-1}\eta)^{1/2} \delta^{\min\{1/2, 1/N\}}). \end{aligned} \tag{3-51}$$

Proof. It is enough to improve the final estimates in Step 4 of the proof of Proposition 3.6 by using (2-14):

$$\begin{aligned} \|u^* - \sqrt{N+1} \cos(\cdot)\|_{L^2([0,r],\mathfrak{m})} &= \|u_0 + \alpha \sin(\cdot) + \beta \cos(\cdot) - \sqrt{N+1} \cos(\cdot)\|_{L^2([0,r],\mathfrak{m})} \\ &\leq \|u_0\|_{L^2([0,r],\mathfrak{m})} + |\alpha| \|\sin(\cdot)\|_{L^2([0,r],\mathfrak{m})} + |\beta - \sqrt{N+1}| \|\cos(\cdot)\|_{L^2([0,r],\mathfrak{m})} \\ &\leq C(\delta^{1/2} + \delta^{\min\{1/2, 1/N\}} (\|\cos(\cdot)\|_{L^2([0,r],\mathfrak{m}_N)} + C\delta^{1/N} r^{N-1})) \\ &\leq C(\delta^{1/2} + r^{N/2} \delta^{\min\{1/2, 1/N\}}). \end{aligned}$$

The improved estimate for the first derivative and for the domain $[r - \eta, r + \eta]$ is analogous. □

Lemma 3.8. *For any $N > 1$ there exist $\bar{D} = \bar{D}(N) < \pi$ and $\alpha = \alpha(N) > 0$ such that the following holds. Let $([0, D], d_{\text{eucl}}, \mathfrak{m})$ be a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. with $D \geq \bar{D}$ and u^* any first Neumann eigenfunction, with $\|u^*\|_{L^2(\mathfrak{m})} = 1$.*

Then for any $v \in L^2([0, D], \mathfrak{m})$ with $\|v\|_{L^2(\mathfrak{m})} = 1$ such that $|\int v u^ \mathfrak{m}| \leq \frac{1}{2}$ we have*

$$N + \alpha \leq \int_{[0, D]} |v'|^2 \mathfrak{m}.$$

Proof. We argue by contradiction.

Suppose there is a sequence of $\text{CD}(N - 1, N)$ measures $\mathfrak{m}_n = h_n \mathcal{L}^1$ with $\text{supp } h_n = [0, D_n]$ and $D_n \uparrow \pi$ satisfying the following: for every n there exists $v_n \in W^{1,2}([0, D_n], d_{\text{eucl}}, \mathfrak{m}_n)$ with $\|v_n\|_{L^2(\mathfrak{m}_n)} = 1$ such that

$$\int_{[0, D_n]} |v'_n|^2 \mathfrak{m}_n \rightarrow N \quad \text{as } n \rightarrow \infty, \quad \text{and} \quad \left| \int v_n u_n^* \mathfrak{m}_n \right| \leq \frac{1}{2}, \tag{3-52}$$

where u_n^* is a first Neumann eigenfunction on $([0, D_n], d_{\text{eucl}}, h_n \mathcal{L}^1)$, i.e.,

$$\int_{[0, D_n]} |u_n^*|^2 \mathfrak{m}_n = 1, \quad \int_{[0, D_n]} |(u_n^*)'|^2 \mathfrak{m}_n = \lambda_n \rightarrow N, \tag{3-53}$$

where in the last identity we used (3-25), and the convergence of λ_n to N follows from Lemma 3.4.

From Corollary 2.7, the fact that $\text{supp } h_n = [0, D_n]$ with $D_n \uparrow \pi$ implies that (h_n) (extended to the constant $h(D_n)$ on $[D_n, \pi]$) converges uniformly to the model one-dimensional $\text{CD}(N - 1, N)$ -density $h_N = (1/c'_N) \sin^{N-1}$ on $[0, \pi]$. In particular, for every $\eta \in (0, \pi/2)$ the densities h_n restricted to $[\eta, \pi - \eta]$ are bounded above and below by strictly positive constants.

The bounds (3-53) then imply that u_n^* (resp. v_n) are uniformly $\frac{1}{2}$ -Hölder continuous on $[\eta, \pi - \eta]$ for every $\eta \in (0, \pi/2)$.

Thus, by the Arzelà–Ascoli theorem combined with a standard diagonal argument, there exists $u^* : [0, \pi] \rightarrow \mathbb{R}$ (resp. $v : [0, \pi] \rightarrow \mathbb{R}$) and a (nonrelabelled for simplicity) subsequence such that $u_n^* \rightarrow u^*$ (resp. $v_n \rightarrow v$) uniformly on $[\eta, \pi - \eta]$ for every $\eta \in (0, \pi/2)$. It is also easy to check that

$$\int_{[0, \pi]} u_n^* h_n \phi \mathcal{L}^1 \rightarrow \int_{[0, \pi]} u^* h_N \phi \mathcal{L}^1, \quad \int_{[0, \pi]} v_n h_n \phi \mathcal{L}^1 \rightarrow \int_{[0, \pi]} v h_N \phi \mathcal{L}^1 \quad \text{for all } \phi \in C([0, \pi]).$$

Combining the last weak convergence statement with the bounds (3-52), (3-53) and with [Gigli et al. 2015, Theorem 6.3] gives

$$\|u^*\|_{L^2([0, \pi], \mathfrak{m}_N)} = \|v\|_{L^2([0, \pi], \mathfrak{m}_N)} = 1, \quad \left| \int_{[0, \pi]} u^* v \mathfrak{m}_N \right| \leq \frac{1}{2},$$

$$\int_{[0, \pi]} |(u^*)'|^2 \mathfrak{m}_N \leq N, \quad \int_{[0, \pi]} |v'|^2 \mathfrak{m}_N \leq N.$$

Therefore, both u^* and v are first Neumann eigenfunctions on the model space $([0, \pi], d_{\text{eucl}}, \mathfrak{m}_N)$. However the first eigenfunction is unique up to a sign, thus it must hold

$$\left| \int_{[0, \pi]} u^* v \mathfrak{m}_N \right| = 1,$$

a contradiction. □

Corollary 3.9. *For every $N > 1$ there exists $\beta = \beta(N) > 0$ with the following property. Let $(I, d_{\text{eucl}}, \mathfrak{m})$ be a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. with $\mathfrak{m}(I) = 1$ and satisfying*

$$\lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} - N < \beta.$$

Then, for any $u \in W^{1,2}(I, d_{\text{eucl}}, \mathfrak{m})$ with $\|u\|_{L^2(\mathfrak{m})} = 1$ and $|\int_I uu^ \mathfrak{m}| \leq \frac{1}{2}$, where u^* is a first Neumann eigenfunction with $\|u^*\|_{L^2(\mathfrak{m})} = 1$, it holds*

$$\lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} + \beta < \int |u'|^2 \mathfrak{m}.$$

Proof. First choose $\beta > 0$ sufficiently small so that, by Proposition 3.3, the diameter of $(I, d_{\text{eucl}}, \mathfrak{m})$ is bigger than \bar{D} . Then conclude by Lemma 3.8 (and decrease the constant $\beta > 0$ if necessary). \square

Proposition 3.10. *For every $N > 1$ there exists $\beta = \beta(N) > 0$ with the following property. Let $(I, d_{\text{eucl}}, \mathfrak{m})$ be a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. with $\mathfrak{m}(I) = 1$. Assume there exists $v \in W^{1,2}(I, d_{\text{eucl}}, \mathfrak{m})$ with $\|v\|_{L^2(\mathfrak{m})} = 1$ satisfying*

$$\int_I |v'|^2 \mathfrak{m} - N < \beta. \tag{3-54}$$

Then it holds

$$\min\{\|v - u^*\|_{W^{1,2}(\mathfrak{m})}^2, \|v + u^*\|_{W^{1,2}(\mathfrak{m})}^2\} \leq C \left(\int |v'|^2 \mathfrak{m} - \int |(u^*)'|^2 \mathfrak{m} \right), \tag{3-55}$$

where u^ is a first Neumann eigenfunction with $\|u^*\|_{L^2(\mathfrak{m})} = 1$.*

Proof. We begin by rewriting

$$\begin{aligned} \int |v'|^2 \mathfrak{m} - \int |(u^*)'|^2 \mathfrak{m} &= \int |(v - u^*)'|^2 \mathfrak{m} + 2 \int (v - u^*)' (u^*)' \mathfrak{m} \\ &= \int |(v - u^*)'|^2 \mathfrak{m} - 2\lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} \left(1 - \int v u^* \mathfrak{m} \right) \\ &= \int |(v - u^*)'|^2 \mathfrak{m} - \lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} \int (v - u^*)^2 \mathfrak{m}. \end{aligned} \tag{3-56}$$

Now (3-54) implies that $|\int v u^* \mathfrak{m}| > \frac{1}{2}$ by Corollary 3.9. Hence, assuming without loss of generality that $\int v u^* \mathfrak{m} > \frac{1}{2}$, we get $|\int u^* (u^* - v) \mathfrak{m}| < \frac{1}{2}$. Therefore, Corollary 3.9 yields

$$\int |(v - u^*)'|^2 \mathfrak{m} \geq (\lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} + \beta) \|v - u^*\|_2^2.$$

The combination of the last estimate with (3-56) gives

$$\|v - u^*\|_2^2 \leq C \left(\int |v'|^2 \mathfrak{m} - \int |(u^*)'|^2 \mathfrak{m} \right), \tag{3-57}$$

with $C := 1/\beta$. We now improve (3-57) to $W^{1,2}$ -closeness, namely (3-55). In order to do so, it suffices to observe that the estimates we obtained above yield

$$\begin{aligned} \int |(v - u^*)'|^2 \mathfrak{m} &\leq \lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2} \|v - u^*\|_2^2 + \int |v'|^2 \mathfrak{m} - \int |(u^*)'|^2 \mathfrak{m} \\ &\leq C(1 + \lambda_{(I, d_{\text{eucl}}, \mathfrak{m})}^{1,2}) \left(\int |v'|^2 \mathfrak{m} - \int |(u^*)'|^2 \mathfrak{m} \right). \end{aligned} \quad \square$$

Theorem 3.11 (one-dimensional quantitative Obata's theorem on the function). *For every $N > 1$ there exist constants $C = C(N) > 0$ and $\delta_0 = \delta_0(N) > 0$ with the following property. Let $([0, D], d_{\text{eucl}}, \mathfrak{m})$ be a one-dimensional $\text{CD}(N - 1, N)$ m.m.s. and let $u \in \text{Lip}(I)$ satisfy $\int u \, \mathfrak{m} = 0$ and $\int u^2 \, \mathfrak{m} = 1$. If*

$$\delta := \int |u'|^2 \, \mathfrak{m} - N \leq \delta_0,$$

then

$$\min\{\|u - \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}(\mathfrak{m})}, \|u + \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}(\mathfrak{m})}\} \leq C\delta^{\min\{1/2, 1/N\}}. \tag{3-58}$$

Moreover, setting $r = \delta^{\gamma/N}$ for some $\gamma \in (0, 1)$, for any $\eta \in (0, \frac{1}{10}r)$ it holds

$$\min\{\|u - \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}([0,r],\mathfrak{m})}, \|u + \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}([0,r],\mathfrak{m})}\} \leq C(\delta^{1/2} + r^{N/2}\delta^{\min\{1/2, 1/N\}}), \tag{3-59}$$

$$\min\{\|u^* - \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}([r-\eta,r+\eta],\mathfrak{m})}, \|u^* + \sqrt{N + 1} \cos(\cdot)\|_{W^{1,2}([r-\eta,r+\eta],\mathfrak{m})}\} \leq C(N)(\delta^{1/2} + (r^{N-1}\eta)^{1/2}\delta^{\min\{1/2, 1/N\}}). \tag{3-60}$$

Proof. First apply [Proposition 3.10](#) to bound the $W^{1,2}(\mathfrak{m})$ -distance between u and a first eigenfunction of the Neumann Laplacian on $([0, D], d_{\text{eucl}}, \mathfrak{m})$, then apply [Proposition 3.6](#) (respectively [Corollary 3.7](#)) to bound the $W^{1,2}(\mathfrak{m})$ -distance (respectively the $W^{1,2}([0, r], \mathfrak{m})$ or $W^{1,2}([r - \eta, r + \eta], \mathfrak{m})$ distance) between the first eigenfunction and the normalized cosine. The sought estimate follows by the triangle inequality. □

4. Quantitative Obata's theorem on the diameter

Building on top of the one-dimensional results obtained in [Section 3](#), we will derive several quantitative estimates for a general essentially nonbranching m.m.s. (X, d, \mathfrak{m}) satisfying $\text{CD}(K, N)$.

Given an m.m.s. (X, d, \mathfrak{m}) , the perimeter $P(E)$ of a Borel subset $E \subset X$ is defined as

$$P(E) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_X |\nabla u_n| \, \mathfrak{m} : u_n \in \text{Lip}(X), u_n \rightarrow \chi_E \text{ in } L^1_{\text{loc}}(X) \right\}, \tag{4-1}$$

where χ_E is the characteristic function of E . Accordingly $E \subset X$ has finite perimeter in (X, d, \mathfrak{m}) if and only if $P(E) < \infty$.

The isoperimetric profile $\mathcal{I}_{(X,d,\mathfrak{m})} : [0, 1] \rightarrow [0, \infty)$ is given by

$$\mathcal{I}_{(X,d,\mathfrak{m})}(v) := \inf\{P(E) : E \subset X, \mathfrak{m}(E) = v\}. \tag{4-2}$$

Given a smooth Riemannian manifold (M, g) with finite Riemannian volume $\text{vol}_g(M) < \infty$, let us denote by

$$\mathfrak{m}_g := \frac{1}{\text{vol}_g(M)} \text{vol}_g$$

the normalized Riemannian volume measure.

We next recall the improved Lévy–Gromov inequality obtained by Bérard, Besson and Gallot [[Bérard et al. 1985](#), Remark 3.1] for smooth Riemannian N -manifolds with $\text{Ricci} \geq N - 1$ and with upper bound on the diameter (see also [[Milman 2015](#)]).

Theorem 4.1. *Let (M, d, m_g) be the metric measure space associated to a Riemannian manifold (M, g) with dimension $N \in \mathbb{N}$, $N \geq 2$, Ricci bounded from below by $N - 1$ and diameter D (recall that, by the Bonnet–Myers theorem, $D \leq \pi$). Then, for any $v \in (0, 1)$, it holds*

$$\frac{\mathcal{I}_{(X,d,m)}(v)}{\mathcal{I}_N(v)} \geq \left(\frac{\int_0^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{1/N} =: C_{N,D} \geq 1, \tag{4-3}$$

where \mathcal{I}_N , defined in (3-3), for $N \geq 2$, $N \in \mathbb{N}$, is the isoperimetric profile of the normalized round sphere of constant sectional curvature 1 $(\mathbb{S}^N, d_{\mathbb{S}^N}, m_{g_{\mathbb{S}^N}})$.

We extend Theorem 4.1 to the class of essentially nonbranching $CD(N - 1, N)$ metric measure spaces, $N > 1$ any real parameter. In view of [Cavalletti and Mondino 2017b; 2018] the result follows from the one-dimensional improved Lévy–Gromov inequality proved in Lemma 3.1.

Theorem 4.2 (Bérard–Besson–Gallot improved Lévy–Gromov for $CD(N - 1, N)$ e.n.b. spaces). *Let (X, d, m) be an essentially nonbranching $CD(N - 1, N)$ m.m.s. with $\text{diam}(X) \leq D$ for some $N > 1$, $D \in (0, \pi]$. Then, for any $v \in (0, 1)$, it holds*

$$\frac{\mathcal{I}_{(X,d,m)}(v)}{\mathcal{I}_N(v)} \geq \left(\frac{\int_0^{\pi/2} (\cos t)^{N-1} dt}{\int_0^{D/2} (\cos t)^{N-1} dt} \right)^{1/N} =: C_{N,D} \geq 1, \tag{4-4}$$

where \mathcal{I}_N was defined in (3-3).

Proof. One of the main results in [Cavalletti and Mondino 2017b; 2018] is that for (X, d, m) as in the assumptions of the theorem it holds

$$\mathcal{I}_{(X,d,m)}(v) \geq \mathcal{I}_{N,D}(v), \tag{4-5}$$

where $\mathcal{I}_{N,D}$ stands for the model isoperimetric profile defined in (3-3).

The claimed (4-4) follows by combining (4-5) with Lemma 3.1. □

It is also possible to obtain a quantitative spectral gap inequality for Neumann boundary conditions. The analogous result in the case of smooth Riemannian manifolds was established in [Croke 1982, Theorem B] building upon a quantitative improvement of the Lévy–Gromov inequality and on [Bérard and Meyer 1982] (see also [Bérard et al. 1985, Corollary 17]).

Theorem 4.3 (improved spectral gap and quantitative Obata’s theorem for $CD(N - 1, N)$ e.n.b. spaces). *Let (X, d, m) be an essentially nonbranching $CD(N - 1, N)$ m.m.s. with $\text{diam}(X) \leq D$ for some $N > 1$, $D \in (0, \pi]$. Then*

$$\lambda_{(X,d,m)}^{1,2} \geq NC_{N,D}^2, \tag{4-6}$$

where $C_{N,D}$ is given in (4-4). Moreover, there exists $C = C_N > 0$ (more precisely one can choose $C_N = \bar{C}N$ where \bar{C} was defined in Lemma 3.2) such that

$$C_N(\pi - \text{diam}(X))^N \leq \lambda_{(X,d,m)}^{1,2} - N.$$

Proof. Thanks to [Cavalletti and Mondino 2017c, Theorem 4.4] (see also Proposition 3.3) we know that $\lambda_{(X,d,m)}^{1,2} \geq \lambda_{N,D}^{1,2}$, where $\lambda_{N,D}^{1,2}$ was defined in (3-21).

Let us briefly outline the argument since it will be relevant for addressing the quantitative inequality for the first eigenfunction later in the note. By the very definition of $\lambda_{(X,d,m)}^{1,2}$ it suffices to prove that, for any $u \in \text{Lip}(X)$ with $\int u \, m = 0$ and $\int u^2 \, m = 1$, it holds

$$\delta(u) := \int_X |\nabla u|^2 \, m - N \geq C_N(\pi - \text{diam}(X))^N.$$

To this aim, we perform the one-dimensional localization associated to the function u which by assumption has null mean value (this is analogous to the proof of [Cavalletti and Mondino 2017c, Theorem 4.4]; see Section 2D for some basics about one-dimensional localization). We obtain

$$\begin{aligned} \int_X |\nabla u|^2 \, m - N \int_X u^2 \, m &\geq \int_Q \left(\int_{X_q} |u'_q|^2 \, m_q - N \int_{X_q} u_q^2 \, m_q \right) \mathfrak{q}(dq) \\ &\geq \int_Q \left(\lambda_{N, \text{diam}(X_q)}^{1,2} \int_{X_q} u_q^2 \, m_q - N \int_{X_q} u_q^2 \, m_q \right) \mathfrak{q}(dq) \\ &\geq \int_Q (\lambda_{N, \text{diam}(X)}^{1,2} - N) \int_{X_q} u_q^2 \, m_q \, \mathfrak{q}(dq) = \lambda_{N, \text{diam}(X)}^{1,2} - N. \end{aligned}$$

Taking into account Proposition 3.3, we conclude that

$$\delta(u) \geq \lambda_{\text{diam}(X), N}^{1,2} - N \geq C_N(\pi - \text{diam}(X))^N$$

and (4-6) can be obtained in an analogous way. □

Remark 4.4. In [Jiang and Zhang 2016] the authors obtained a quantitative version of the estimate for the gap of the diameters in terms of the deficit in the spectral gap for RCD spaces (see Remark 1.3 therein). Their estimate reads as follows: if (X, d, m) is an $\text{RCD}(N - 1, N)$ space of diameter $D \leq \pi$, then

$$\lambda_{(X,d,m)}^{1,2} \geq \frac{N}{1 - \cos^N(D/2)}.$$

Theorem 4.3 extends such quantitative control to essentially nonbranching $\text{CD}(N - 1, N)$ spaces whose Sobolev space $W^{1,2}$ is a priori non-Hilbert (but just Banach, as for instance on Finsler manifolds).

4A. Volume control. The aim of this brief subsection is to prove that for a $\text{CD}(N - 1, N)$ m.m.s. with diameter close to π we have a quantitative volume control for balls centered at extrema of long rays. The proof is inspired by [Ohta 2007, Lemma 5.1], where the case of maximal diameter π is treated (see also [Cavalletti et al. 2019, Proposition 5.1]).

Proposition 4.5. *Let (X, d, m) be an m.m.s. satisfying $\text{CD}(N - 1, N)$ (actually $\text{MCP}(N - 1, N)$ is enough). Let $P_N, P_S \in X$ be such that $d(P_N, P_S) = \pi - \delta$ for some $\delta \geq 0$. Then, for any $0 < r < \pi - \delta$, it holds*

$$m_N([0, r]) \leq m(B_r(P_N)) \leq m_N([0, r]) + m_N([r, r + \delta]), \tag{4-7}$$

where we recall that $m_N = (1/\omega_N)(\sin t)^{N-1} dt$ is the model measure on the interval $[0, \pi]$.

Proof. First of all, since $d(P_N, P_S) = \pi - \delta$, it holds $B_r(P_N) \cap B_{\pi-r-\delta}(P_S) = \emptyset$.

By the Bishop–Gromov inequality implied by the $CD(N - 1, N)$ condition (actually $MCP(N - 1, N)$ is enough), and using that $m(X) = 1$, we have

$$m(B_r(P_N)) \geq m_N([0, r]), \quad m(B_{\pi-r-\delta}(P_S)) \geq m_N([0, \pi - r - \delta]) = m_N([r + \delta, \pi]), \tag{4-8}$$

where the last equality follows from the symmetries of the density $\sin^{N-1}(\cdot)$. Hence we can compute

$$\begin{aligned} m(B_r(P_N)) &\leq 1 - m(B_{\pi-r-\delta}(P_S)) \leq 1 - m_N([0, \pi - r - \delta]) \\ &= m_N([0, r]) + m_N([r, r + \delta]). \end{aligned}$$

The claimed conclusion (4-7) follows. □

5. Quantitative Obata’s theorem on almost optimal functions

Consider $u \in \text{Lip}(X)$ such that

$$\int_X u \, m = 0, \quad \int_X u^2 \, m = 1;$$

denote its spectral gap deficit by

$$\delta(u) := \int_X |\nabla u|^2 \, m - N. \tag{5-1}$$

Since we are interested in quantitative estimates when the spectral gap deficit is small, it is enough to consider the case $\delta(u) \leq 1$. Recall that N is the first eigenvalue for the Neumann Laplacian for the one-dimensional metric measure space $([0, \pi], |\cdot|, m_N)$, where $m_N := \sin^{N-1}(t) \, dt / \omega_N$ and ω_N is the normalizing constant. In particular

$$N = (N + 1) \int_{(0,\pi)} \sin^2(t) \, m_N(dt),$$

since, as we already observed, $\int_{(0,\pi)} \cos^2(t) \, m_N(dt) = 1/(N + 1)$.

Consider the localization associated to the zero-mean function u (see Section 2D for the background and for the relevant bibliography):

$$m_{\perp \mathcal{T}} = \int_Q m_q \, q(dq),$$

where \mathcal{T} is the transport set associated to the L^1 -optimal transport problem between $u^+ m$ and $u^- m$, the positive and the negative parts of u , respectively. It follows that

$$\int_Q \int_{X_q} |u|^2 \, m_q \, q(dq) = \int_{\mathcal{T}} |u|^2 \, m = \int_X |u|^2 \, m = 1, \quad \int_{X \setminus \mathcal{T}} |\nabla u|^2 \, m = 0. \tag{5-2}$$

Setting $u_q := u|_{X_q}$ and $|c_q| := (\int_{X_q} |u_q|^2 \, m_q)^{1/2}$ (for the sign of c_q , see before (5-13)), observe that (5-2) gives

$$\int_Q c_q^2 \, q(dq) = 1. \tag{5-3}$$

Moreover, the integral constraint $\int_X u \, \mathbf{m} = 0$ localizes to almost every ray:

$$\int_{X_q} u_q \, \mathbf{m}_q = 0. \tag{5-4}$$

Since almost each ray $(X_q, d|_{X_q}, \mathbf{m}_q)$ is a one-dimensional $\text{CD}(N-1, N)$ space, the Lichnerowicz spectral gap gives

$$\int_{X_q} |u'_q|^2 \, \mathbf{m}_q \geq N c_q^2, \tag{5-5}$$

where $|u'_q|(x)$ denotes the local Lipschitz constant of $u_q : (X_q, d|_{X_q}) \rightarrow \mathbb{R}$ at $x \in X_q$. It is clear that, for each $x \in X_q \subset X$, $|u'_q|(x)$ is bounded by the local Lipschitz constant $|\nabla u|(x)$ of $u : (X, d) \rightarrow \mathbb{R}$:

$$|u'_q|(x) \leq |\nabla u|(x) \quad \text{for all } x \in X_q, \text{ q-a.e. } q \in Q. \tag{5-6}$$

With a slight abuse of notation, in order to keep the formulas short, in the following we will often identify q and $q \llcorner_{\{q \in Q : c_q > 0\}}$. Localizing the spectral gap deficit using (5-6) gives

$$\begin{aligned} \delta(u) &= \int_X |\nabla u|^2 \, \mathbf{m} - N \geq \int_Q \left(\int_{X_q} \frac{|u'_q|^2}{c_q^2} \, \mathbf{m}_q \right) c_q^2 \, \mathbf{q}(dq) - N \\ &= \int_Q \left[\int_{X_q} \left(\frac{|u'_q|^2}{c_q^2} - N \right) \, \mathbf{m}_q \right] c_q^2 \, \mathbf{q}(dq) \end{aligned} \tag{5-7}$$

$$= \int_Q \delta(u_q) c_q^2 \, \mathbf{q}(dq), \tag{5-8}$$

where we set

$$\delta(u_q) := \int_{X_q} \left(\frac{|u'_q|^2}{c_q^2} - N \right) \, \mathbf{m}_q,$$

the one-dimensional spectral gap deficit of u_q . From now on, in order to keep notation short, we will write δ for $\delta(u)$. Let $\beta \in (0, 1)$ be a real parameter to be optimized later in the proof and denote the set of “long rays” by

$$Q_\ell := \{q \in Q : \delta(u_q) \leq \delta^\beta \text{ and } c_q > 0\}.$$

It follows from (5-8), Chebyshev’s inequality and (5-3) that

$$\int_{Q \setminus Q_\ell} c_q^2 \, \mathbf{q}(dq) \leq \delta^{1-\beta}, \quad \int_{Q_\ell} c_q^2 \, \mathbf{q}(dq) \geq 1 - \delta^{1-\beta}. \tag{5-9}$$

Hence we can use Proposition 3.3 to deduce that, for all $q \in Q_\ell$,

$$(\pi - |X_q|)^N \leq C_N \delta^\beta, \tag{5-10}$$

where $|X_q|$ denotes the length of the ray X_q . Being the preimage of a measurable function, Q_ℓ is a measurable subset of Q . Adopting the notation $R(E) := \bigcup_{q \in E} X_q$, so that $R(E)$ is the span of the rays corresponding to equivalence classes in E , we claim that

$$\int_{X \setminus R(Q_\ell)} |\nabla u|^2 \, \mathbf{m} \leq (N + 1) \delta^{1-\beta}. \tag{5-11}$$

Indeed (5-6), (5-5) and (5-9) yield

$$\int_{R(Q_\ell)} |\nabla u|^2 \mathfrak{m} \geq \int_{Q_\ell} \int_{X_q} |u'_q|^2 \mathfrak{m}_q \mathfrak{q}(dq) \geq N \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) \geq N(1 - \delta^{1-\beta}).$$

The claim (5-11) follows by combining the last estimate with

$$\int_{X \setminus R(Q_\ell)} |\nabla u|^2 \mathfrak{m} + \int_{R(Q_\ell)} |\nabla u|^2 \mathfrak{m} = \int_X |\nabla u|^2 \mathfrak{m} \leq N + \delta.$$

For each $q \in Q$, we denote by $a(X_q)$ (resp. $b(X_q)$) the initial (resp. final) point of the ray X_q .

Throughout this last section we will often make the identification between the ray X_q and the interval $(0, |X_q|)$.

Proposition 5.1. *There exists a distinguished $\bar{q} \in Q_\ell$ having initial point P_N and final point P_S such that*

$$d(P_N, a(X_q)) \leq C(N)\delta^{\beta/N}, \quad d(P_S, b(X_q)) \leq C(N)\delta^{\beta/N} \quad \text{for all } q \in Q_\ell. \tag{5-12}$$

Proof. Fix any $\bar{q} \in Q_\ell$ and set $P_N := a(X_{\bar{q}})$, $P_S := b(X_{\bar{q}})$. By d-cyclical monotonicity of the transport set \mathcal{T} , for any other $q \in Q_\ell$ it holds

$$2\pi - d(a(X_q), b(X_q)) - d(P_N, P_S) \geq 2\pi - d(a(X_q), P_S) - d(b(X_q), P_N),$$

which we rewrite as

$$\pi - |X_q| + \pi - |X_{\bar{q}}| \geq \pi - d(a(X_q), P_S) + \pi - d(b(X_q), P_N).$$

Combining the last estimate with (5-10) gives

$$2C_N\delta^{\beta/N} \geq \pi - d(a(X_q), P_S) + \pi - d(b(X_q), P_N).$$

Finally by [Cavalletti et al. 2019, Proposition 5.1] we deduce the existence of a constant, depending only on the dimension N , such that

$$d(a(X_q), P_N) \leq C(N)\delta^{\beta/N}, \quad d(b(X_q), P_S) \leq C(N)\delta^{\beta/N},$$

and the claim follows. □

From now on, for every $q \in Q_\ell$ choose the sign of c_q so that

$$\left\| \frac{u_q}{c_q} - \sqrt{N+1} \cos(\cdot) \right\|_{L^2(X_q, \mathfrak{m}_q)} = \min \left\{ \left\| \frac{u_q}{|c_q|} + \sqrt{N+1} \cos(\cdot) \right\|_{L^2(X_q, \mathfrak{m}_q)}, \left\| \frac{u_q}{|c_q|} - \sqrt{N+1} \cos(\cdot) \right\|_{L^2(X_q, \mathfrak{m}_q)} \right\}.$$

From Theorem 3.11 we obtain that for all $q \in Q_\ell$ it holds

$$\left\| \frac{u_q}{c_q} - \sqrt{N+1} \cos(\cdot) \right\|_{L^2(X_q, \mathfrak{m}_q)} \leq C(N)\delta^{\beta \min\{1/2, 1/N\}}. \tag{5-13}$$

The goal of the next section is to globalize estimate (5-13) to the whole space X .

The sought bound will be obtained through two intermediate steps: Firstly, in Proposition 5.2, we control the variance of the map $q \mapsto c_q$ with respect to the measure \mathfrak{q} on the set of long rays Q_ℓ . Then, in Proposition 5.3, we estimate $(1 - \mathfrak{q}(Q_\ell))$ in terms of a power of the deficit.

Below we briefly present the strategy of the proof. In order to fix the ideas, we discuss the heuristics in the rigid case of zero deficit. Actually in the case of zero deficit there is a more streamlined argument (the assumption that u is Lipschitz, combined with the fourth bullet below, gives immediately that $q \mapsto c_q$ is constant); however, the point here is to present a strategy which generalizes to the nonrigid case of nonzero deficit.

In the case where $\delta(u) = 0$, the results of the previous sections give the following conclusions:

- Almost all the transport rays have length π . Moreover, they start from a common point P_N , with $u(P_N) > 0$, and end in a common point P_S , with $u(P_S) < 0$.
- $m(B_r(P_N)) = m_N([0, r])$ for any $r \in [0, \pi]$.
- For q -a.e. $q \in Q$, it holds that $m_q = m_N$ is the model measure for the $CD(N - 1, N)$ condition.
- For q -a.e. $q \in Q$, it holds that $u_q(\cdot) = c_q \cos(d(P_N, \cdot))$.

Our aim is to prove that $q(Q) = 1$ and that $c_q = 1$ for q -a.e. $q \in Q$. The basic idea is to apply the Poincaré inequality to balls centered at P_N and having radii converging to 0.

Observe that we can compute

$$\int_{B_r(P_N)} u \, m = \frac{1}{m_N([0, r])} \int_Q \int_0^r c_q \cos(t) m_N(dt) = \left(\int_Q c_q \, q \right) \int_0^r \cos(t) m_N(dt). \tag{5-14}$$

Moreover, recalling that $u = 0$ m -a.e. outside of the transport set, we have

$$\begin{aligned} & \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \, m \right|^2 m \\ & \stackrel{(5-14)}{=} (1 - q(Q)) \left(\int_{B_r(P_N)} u \, m \right)^2 + \int_Q \int_0^r \left| c_q \cos(t) - \int_Q c_q \, q(dq) \int_0^r \cos(t) m_N(dt) \right|^2 m_N(dt) \, q(dq) \\ & \sim (1 - q(Q)) \left(\int_Q c_q \, q(dq) \right)^2 + \int_Q \left| c_q - \int_Q c_q \, q(dq) \right|^2 q(dq) \quad \text{as } r \rightarrow 0, \end{aligned} \tag{5-15}$$

where in the last step we relied on the asymptotic $\cos(t) = 1 + o(t)$ as $t \rightarrow 0$. Eventually we can compute

$$\int_{B_{2r}(P_N)} |\nabla u|^2 m = \int_Q c_q^2 \, q(dq) \int_0^{2r} \sin^2(t) m_N(dt) = \int_0^{2r} \sin^2(t) m_N(dt) \sim r^2 \quad \text{as } r \rightarrow 0,$$

where in the last step we relied on the asymptotic $\sin(t) = t + o(t)$ as $t \rightarrow 0$.

An application of the Poincaré inequality, in the asymptotic regime $r \downarrow 0$, yields that

$$\int_Q \left| c_q - \int_Q c_q \, q(dq) \right|^2 q(dq) = 0, \tag{5-16}$$

which implies both the conclusions $q(Q) = 1$ and $q \mapsto c_q$ constant q -a.e. Due to the constraint $\int_Q c_q^2 q(dq) = 1$ and the fact that $u(P_N) > 0$, we also get that $c_q = 1$ q -a.e., as we claimed.

A second heuristic motivation of the fact that the oscillation of the map $q \mapsto c_q$ is controlled by (a power of) the deficit is that “the gradient of u is almost aligned along the rays” in a quantitative L^2 -sense, suggesting that u “should not oscillate much in the direction orthogonal to the rays”. Note that in the

current framework of $CD(K, N)$ spaces there is no scalar product and the set Q is far from regular, this is the reason why we cannot directly implement this heuristic strategy. However, let us make precise the fact that “the gradient of u is almost aligned along the rays” in a quantitative L^2 -sense, since this will be used in the arguments below:

$$\begin{aligned} 0 &\stackrel{(5-6)}{\leq} \int_Q \left(\int_{X_q} |\nabla u|^2 - |u'_q|^2 m_q \right) q(dq) = \int_X |\nabla u|^2 m - \int_Q \left(\int_{X_q} |u'_q|^2 m_q \right) q(dq) \quad (\text{by (5-1),(5-5)}) \\ &\leq N + \delta - N \int_Q c_q^2 q(dq) \stackrel{(5-3)}{=} \delta. \end{aligned} \tag{5-17}$$

The proofs of Propositions 5.2 and 5.3 below are based on the idea we just presented, although they are quite technical since one has to handle all the various error terms occurring in the nonrigid case $\delta(u) > 0$.

5A. Control on the variance.

Proposition 5.2. *The following estimate holds:*

$$\int_{Q_\ell} \left| c_q - \int_{Q_\ell} c_q q(dq) \right|^2 q(dq) \leq C(N) (\delta^{4\gamma/N} + \delta^{1-\beta-\gamma+(2\gamma/N)} + \delta^{(\beta-\gamma)\min\{2/N,1\}}) \tag{5-18}$$

for any $0 < \beta < 1$ and for any $0 < \gamma < \min\{\beta, 1 - \beta\}$.

Proof. In order to bound the variance of $q \mapsto c_q$ on Q_ℓ we wish to prove that it can be controlled by an integral depending on the variation of the function u on a small ball $B_r(P_N)$. Next we will appeal to the fact that in the rigid case the L^2 -norm squared of the gradient of u on $B_r(P_N)$ is comparable with r^{N+2} and, at least heuristically, this has to be the case also when dealing with almost rigidity. Some intermediate steps are devoted to reducing to the case where the function u coincides with $c_q \cos(\cdot)$ when restricted to any long ray X_q .

In order to slightly shorten the notation, we will write C in place of $C(N)$ to denote a dimensional constant.

Step 1: We will set $r = \delta^{\gamma/N}$ for a suitable $\gamma \in (0, \beta)$. First of all, notice that the triangle inequality and (5-12) yield

$$[0, r - C\delta^{\beta/N}] \subset X_q \cap B_r(P_N) \subset [0, r + C\delta^{\beta/N}] \tag{5-19}$$

for any $q \in Q_\ell$, where we have identified $[0, r \pm C\delta^{\beta/N}]$ with the set

$$\{z \in X_q : d(z, a(X_q)) \leq r \pm C\delta^{\beta/N}\}.$$

The minimality of the mean combined with the inclusion (5-19) and with the weak local 2-2 Poincaré inequality (2-6) gives

$$\int_{Q_\ell \times [0, r - C\delta^{\beta/N}]} \left| u - \int_{Q_\ell \times [0, r - C\delta^{\beta/N}]} u m \right|^2 m \leq \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \right|^2 m \leq Cr^2 \int_{B_{10r}(P_N)} |\nabla u|^2 m. \tag{5-20}$$

Step 2: Next we will obtain a more explicit expression of $\int_{Q \times [0, r - C\delta^{\beta/N}]} u m$.

Recall that we will often tacitly identify the ray X_q with the interval $(0, |X_q|)$.

Using [Theorem 3.11](#), [Corollary 2.7](#) and that $\delta_q \leq \delta^\beta$ for $q \in Q_\ell$, we estimate

$$\begin{aligned} & \left| \int_{Q_\ell} \int_{[0,r]} u \mathfrak{m}_q \mathfrak{q}(dq) - \sqrt{N+1} \int_{Q_\ell} \int_{[0,r]} c_q \cos(\cdot) \mathfrak{m}_q \mathfrak{q}(dq) \right| \\ & \leq \int_{Q_\ell} |c_q| \int_{[0,r]} \left| \frac{u}{c_q} - \sqrt{N+1} \cos(\cdot) \right| \mathfrak{m}_q \mathfrak{q}(dq) \\ & \leq \int_{Q_\ell} |c_q| \sqrt{\mathfrak{m}_q([0,r])} \left\| \frac{u}{c_q} - \sqrt{N+1} \cos(\cdot) \right\|_{L^2([0,r], \mathfrak{m}_q)} \mathfrak{q}(dq) \\ & \leq C r^{N/2} (r^{N/2} \delta^{\beta \min\{1/2, 1/N\}} + \delta^{\beta/2}) \int_{Q_\ell} |c_q| \mathfrak{q}(dq). \end{aligned} \tag{5-21}$$

Also, using [Corollary 2.8](#), it holds

$$\left| \int_{Q_\ell} \int_{[0,r]} c_q \cos(\cdot) \mathfrak{m}_q \mathfrak{q}(dq) - \int_{Q_\ell} \int_{[0,r]} c_q \cos(\cdot) \mathfrak{m}_N \mathfrak{q}(dq) \right| \leq C \delta^{\beta/N} r^{N-1} \int_{Q_\ell} |c_q| \mathfrak{q}(dq). \tag{5-22}$$

With an analogous estimate involving [Corollary 2.8](#), we also obtain

$$|\mathfrak{m}(Q_\ell \times [0,r]) - \mathfrak{q}(Q_\ell) \mathfrak{m}_N([0,r])| \leq C \mathfrak{q}(Q_\ell) r^{N-1} \delta^{\beta/N}. \tag{5-23}$$

The combination of (5-21), (5-22) and (5-23), setting $\bar{r} := r - C \delta^{\beta/N}$, yields

$$\begin{aligned} & \left| \int_{Q_\ell \times [0,\bar{r}]} u \mathfrak{m} - \frac{\sqrt{N+1} \int_{Q_\ell \times [0,\bar{r}]} c_q \cos(\cdot) \mathfrak{m}_N \mathfrak{q}(dq)}{\mathfrak{q}(Q_\ell) (\mathfrak{m}_N([0,\bar{r}]) - C r^{N-1} \delta^{\beta/N})} \right| \\ & \leq \frac{C (\int_{Q_\ell} |c_q| \mathfrak{q}(dq)) (r^N \delta^{\beta \min\{1/2, 1/N\}} + r^{N/2} \delta^{\beta/2} + r^{N-1} \delta^{\beta/N})}{\mathfrak{q}(Q_\ell) (\mathfrak{m}_N([0,\bar{r}]) - C r^{N-1} \delta^{\beta/N})}. \end{aligned} \tag{5-24}$$

Step 3: In this step we estimate the order in δ of the right-hand side of (5-24) and choose r as

$$r = \delta^{\gamma/N}, \quad \text{with } \gamma \in (0, \beta). \tag{5-25}$$

Approximating the cosine with its first-order Taylor expansion near to the origin in (5-24), we have

$$\int_{Q_\ell \times [0,\bar{r}]} u \mathfrak{m} = \frac{\int_{Q_\ell} c_q \mathfrak{q}(dq) + (\int_{Q_\ell} |c_q| \mathfrak{q}(dq)) O(\delta^{(\beta-\gamma) \min\{1/2, 1/N\}})}{\mathfrak{q}(Q_\ell) / \sqrt{N+1}}.$$

Since by Cauchy–Schwarz inequality and (5-3) it holds

$$\left(\int_{Q_\ell} c_q \mathfrak{q}(dq) \right)^2 \leq \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) \leq \frac{1}{\mathfrak{q}(Q_\ell)},$$

the last estimate can be rewritten as

$$\left| \int_{Q_\ell \times [0,\bar{r}]} u \mathfrak{m} - \sqrt{N+1} \int_{Q_\ell} c_q \mathfrak{q}(dq) \right|^2 \leq \frac{C}{\mathfrak{q}(Q_\ell)} \delta^{(\beta-\gamma) \min\{1, 2/N\}}. \tag{5-26}$$

Step 4: The aim of this step is to eventually gain (5-18). We first need the following intermediate inequality, where we assume that $r \gg \delta^{\beta/N}$ is a free parameter, that we will set later:

$$\begin{aligned}
 & \int_{Q_\ell} \int_{[0,r]} |u - \sqrt{N+1}c_q|^2 m_q \, q(dq) \\
 & \leq 2 \int_{Q_\ell} \int_{[0,r]} |u - \sqrt{N+1}c_q \cos(\cdot)|^2 m_q \, q(dq) + 2 \int_{Q_\ell} \int_{[0,r]} (\sqrt{N+1}|c_q| |\cos(\cdot) - 1|)^2 m_q \, q(dq) \\
 & \leq C\delta^{\beta \min\{1,2/N\}} r^N \int_{Q_\ell} c_q^2 \, q(dq) + C\delta^\beta + Cr^4 \int_{Q_\ell} c_q^2 m_q([0, r]) \, q(dq) \quad (\text{by (3-59)}) \\
 & \leq C\delta^{\beta \min\{1,2/N\}} r^N + C\delta^\beta + Cr^4 \int_{Q_\ell} c_q^2 (m_N([0, r]) + Cr^{N-1}\delta^{\beta/N}) \, q(dq) \quad (\text{by (5-10)+(2-12)}) \\
 & \leq C\delta^{\beta \min\{1,2/N\}} r^N + Cr^4 m_N([0, r]) \int_{Q_\ell} c_q^2 \, q(dq) + C\delta^\beta \leq Cr^N (\delta^{\beta \min\{1,2/N\}} + r^4) + C\delta^\beta. \quad (5-27)
 \end{aligned}$$

In particular, the previous inequality holds true substituting $\bar{r} := r - C\delta^{\beta/N}$ in place of r , and $r = \delta^{\gamma/N}$ is as in the previous Step 3. We deduce

$$\begin{aligned}
 & m_N([0, \bar{r}]) (N+1) \int_{Q_\ell} \left| c_q - \int_{Q_\ell} c_q \, q(dq) \right|^2 \, q(dq) \\
 & \leq (N+1) \int_{Q_\ell} \left| c_q - \int_{Q_\ell} c_q \, q(dq) \right|^2 (m_q([0, \bar{r}]) + Cr^{N-1}\delta^{\beta/N}) \, q(dq) \\
 & \leq C\delta^{\beta/N} r^{N-1} + (N+1) \int_{Q_\ell} \left| c_q - \int_{Q_\ell} c_q \, q(dq) \right|^2 m_q([0, \bar{r}]) \, q(dq) \\
 & \leq C\delta^{\beta/N} r^{N-1} + 2 \int_{Q_\ell} \int_{[0,\bar{r}]} |u - \sqrt{N+1}c_q|^2 m_q \, q(dq) + 2 \int_{Q_\ell} \int_{[0,\bar{r}]} \left| u - \int_{Q_\ell} \sqrt{N+1}c_q \, q(dq) \right|^2 m_q \, q(dq) \\
 & \leq C\delta^{\beta/N} r^{N-1} + 2 \int_{Q_\ell} \int_{[0,\bar{r}]} |u - \sqrt{N+1}c_q|^2 m_q \, q(dq) + 4 \int_{Q_\ell \times [0,\bar{r}]} \left| u - \int_{Q_\ell \times [0,\bar{r}]} u \, m \right|^2 m \\
 & \quad + 4 \int_{Q_\ell} \int_{[0,\bar{r}]} \left| \int_{Q_\ell \times [0,\bar{r}]} u \, m - \sqrt{N+1} \int_{Q_\ell} c_q \, q(dq) \right|^2 m_q \, q(dq).
 \end{aligned}$$

Now use (5-20), (5-26), (5-25), (5-27) to continue the chain of inequalities

$$\leq C\delta^\gamma (\delta^{\beta-\gamma \min\{1,2/N\}} + \delta^{4\gamma/N}) + Cr^2 \int_{B_{10r}(P_N)} |\nabla u|^2 m. \quad (5-28)$$

Next we wish to bound the term $\int_{B_{10r}(P_N)} |\nabla u|^2 m$. To this aim we observe that

$$\begin{aligned}
 & \int_{B_{10r}(P_N)} |\nabla u|^2 m \\
 & \leq \int_{X \setminus R(Q_\ell)} |\nabla u|^2 m + \int_{Q_\ell} \int_0^{10r+C\delta^{\beta/N}} |u'_q|^2 m_q \, q(dq) + \delta \quad (\text{by (5-17)}) \\
 & \leq C(\delta^{1-\beta} + \delta^\beta + r^N \delta^{\beta \min\{2/N,1\}} + \delta^{\beta/N} r^{N-1} r^2) + C \int_0^{10r+C\delta^{\beta/N}} \sin(\cdot)^2 m_N \quad (\text{by (5-11), (3-59), (2-12)}) \\
 & \leq C(\delta^{1-\beta} + \delta^\beta + r^N (\delta^{\beta \min\{2/N,1\}} + r^2)). \quad (5-29)
 \end{aligned}$$

Combine now (5-28) with (5-29) and recall that $r = \delta^{\gamma/N}$ for $0 < \gamma < \min\{\beta, 1 - \beta\}$ to get

$$\delta^\gamma \int_{Q_\ell} \left| c_q - \int_{Q_\ell} c_q \mathfrak{q}(dq) \right|^2 \mathfrak{q}(dq) \leq C\delta^\gamma (\delta^{4\gamma/N} + \delta^{1-\beta-\gamma+(2\gamma/N)} + \delta^{(\beta-\gamma)\min\{1,2/N\}}),$$

which gives the desired estimate (5-18). □

5B. Control of the measure of long rays. Following Proposition 5.2, we set

$$\bar{c} := \int_{Q_\ell} c_q \mathfrak{q}(dq). \tag{5-30}$$

Next we proceed proving that $\mathfrak{q}(Q_\ell)$ is quantitatively close to 1 up to an error of the order of a suitable power of the deficit.

Proposition 5.3. *The following estimate holds:*

$$(1 - \mathfrak{q}(Q_\ell))^2 \leq C(N)(\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N} + \delta^{1-\beta-\gamma}) \tag{5-31}$$

for any $0 < \beta < 1$ and for any $0 < \gamma < \min\{\beta, 1 - \beta\}$.

Proof. In order to slightly shorten the notation, we will write C in place of $C(N)$ to denote constants depending only on N . Moreover, we will continue to tacitly identify the ray X_q with the interval $(0, |X_q|)$. We achieve (5-31) through three intermediate steps.

Step 1: Aim of this first step is to prove that, for $r = \delta^{\gamma/N}$, $\gamma \in (0, \min\{\beta, 1 - \beta\})$, letting $\bar{r} := r - C\delta^{\beta/N}$, it holds

$$\begin{aligned} (N + 1) \int_{Q_\ell} \int_{[0, \bar{r}]} \left| c_q \cos(\cdot) - \bar{c} \mathfrak{q}(Q_\ell) \int_{[0, \bar{r}]} \cos(\cdot) \mathfrak{m}_N \right|^2 \mathfrak{m}_N \mathfrak{q}(dq) \\ \leq \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m} + C(\delta^{\gamma+(\beta-\gamma)/N} + \delta^{1-\beta}). \end{aligned} \tag{5-32}$$

Arguing as in the first steps of the proof of Proposition 5.2, we estimate

$$\begin{aligned} \int_{Q_\ell} \int_0^{\bar{r}} \left| \sqrt{N+1} c_q \cos(\cdot) - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m}_N \mathfrak{q}(dq) \\ \leq \int_{Q_\ell} \int_0^{\bar{r}} \left| \sqrt{N+1} c_q \cos(\cdot) - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m}_q \mathfrak{q}(dq) + C\delta^{\beta/N} r^{N-1} \quad (\text{by (2-12), (5-10)}) \\ \leq 2 \int_{Q_\ell} \int_0^{\bar{r}} |\sqrt{N+1} c_q \cos(\cdot) - u|^2 \mathfrak{m}_q \mathfrak{q}(dq) + C\delta^{\beta/N} r^{N-1} + 2 \int_{Q_\ell} \int_0^{\bar{r}} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m}_q \mathfrak{q}(dq) \\ \leq 2 \int_{Q_\ell} c_q^2 \left\| \frac{u}{c_q} - \sqrt{N+1} \cos(\cdot) \right\|_{L^2([0, \bar{r}], \mathfrak{m}_q)}^2 \mathfrak{q}(dq) + C\delta^{\beta/N} r^{N-1} + 2 \int_{Q_\ell} \int_0^{\bar{r}} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m}_q \mathfrak{q}(dq) \mathfrak{m} \\ \leq 2 \int_{B_r(P_N) \cap R(Q_\ell)} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m} + C\delta^{\beta/N} r^{N-1} \quad (\text{by (3-59), (5-19)}). \end{aligned} \tag{5-33}$$

In order to achieve (5-32), having in mind to argue by triangle inequality, we are left to bound

$$\mathfrak{m}_N([0, \bar{r}]) \mathfrak{q}(Q_\ell) \left| \int_{B_r(P_N)} u \mathfrak{m} - \sqrt{N+1} \bar{c} \mathfrak{q}(Q_\ell) \int_{[0, \bar{r}]} \cos(\cdot) \mathfrak{m}_N \right|^2. \tag{5-34}$$

We start by observing that

$$\begin{aligned} & \left| \int_{B_r(P_N)} u \, m - \sqrt{N+1} \, \bar{c} \, q(Q_\ell) \int_0^r \cos(\cdot) \, m_N \right| \\ & \leq \left| \int_{B_r(P_N) \cap R(Q_\ell)} u \, m - \sqrt{N+1} \, \bar{c} \, q(Q_\ell) \int_0^r \cos(\cdot) \, m_N \right| + \left| \int_{B_r(P_N) \setminus R(Q_\ell)} u \, m \right|. \end{aligned} \tag{5-35}$$

We first treat the second term of the right-hand side.

From (5-9) we know that $\int_{X \setminus R(Q_\ell)} u^2 \, m \leq \delta^{1-\beta}$; an application of Hölder’s inequality and (2-12) yields

$$\int_{B_r(P_N) \setminus R(Q_\ell)} |u| \, m \leq \delta^{(1-\beta)/2} \sqrt{m(B_r(P_N) \setminus R(Q_\ell))} \leq C \delta^{(1-\beta)/2} r^{N/2}. \tag{5-36}$$

We estimate the first term in the right-hand side of (5-35) by reducing to (5-21) in the second step of the proof of Proposition 5.2:

$$\begin{aligned} & \left| \int_{B_r(P_N) \cap R(Q_\ell)} u \, m - \sqrt{N+1} \, \bar{c} \, q(Q_\ell) \int_0^r \cos(\cdot) \, m_N \right| \\ & \leq \left| \int_{B_r(P_N) \cap R(Q_\ell)} u \, m - \int_{Q_\ell} \int_{[0,r]} u \, m_q \, q(dq) \right| \\ & \quad + \left| \int_{Q_\ell} \int_{[0,r]} u \, m_q \, q(dq) - \sqrt{N+1} \int_{Q_\ell} \int_{[0,r]} c_q \cos(\cdot) \, m_q \, q(dq) \right| \\ & \quad + \left| \int_{Q_\ell} \int_{[0,r]} \sqrt{N+1} c_q \cos(\cdot) \, m_q \, q(dq) - \sqrt{N+1} \bar{c} q(Q_\ell) \int_0^r \cos(\cdot) \, m_N \right|. \end{aligned}$$

Using (2-12), (3-59), (5-10), (5-19), (5-21), we continue as follows:

$$\leq \int_{Q_\ell} \int_{r-C\delta^{\beta/N}}^{r+C\delta^{\beta/N}} |u| \, m_q \, q(dq) + C r^{N/2} (\delta^{\beta \min\{1/2, 1/N\}} r^{N/2} + \delta^{\beta/2} + r^{(N/2)-1} \delta^{\beta/N}) \int_{Q_\ell} |c_q| \, q(dq). \tag{5-37}$$

Arguing by triangle inequality bounding first the distance from the normalized cosine (with (3-60)) and then replacing the measures m_q with the model measure m_N (with (2-12)), we estimate the first summand in the right-hand side of (5-37) as

$$\int_{Q_\ell} \int_{r-C\delta^{\beta/N}}^{r+C\delta^{\beta/N}} |u| \, m_q \, q(dq) \leq C (r^{N-1} \delta^{\beta/N} + r^{(N-1)/2} \delta^{\beta(1/2+1/(2N))}) \int_{Q_\ell} |c_q| \, q(dq). \tag{5-38}$$

Combining (5-37), (5-38), and choosing $r = \delta^\gamma$ with $\gamma \in (0, \min\{\beta, 1-\beta\})$ yields

$$\left| \int_{B_r(P_N) \cap R(Q_\ell)} u \, m - \bar{c} \sqrt{N+1} q(Q_\ell) \int_0^{\bar{r}} \cos(\cdot) \, m_N \right| \leq C (r^{N-1} \delta^{\beta/N} + r^N \delta^{\beta \min\{1/2, 1/N\}} + r^{N/2} \delta^{\beta/2}). \tag{5-39}$$

The combination of (5-35) (5-36) and (5-39) gives

$$\begin{aligned} & \left| \int_{B_r(P_N)} u \, m - \bar{c} \sqrt{N+1} q(Q_\ell) \int_0^{\bar{r}} \cos(\cdot) \, m_N \right| \\ & \leq C (r^{N-1} \delta^{\beta/N} + r^N \delta^{\beta \min\{1/2, 1/N\}} + r^{N/2} \delta^{\beta/2} + \delta^{(1-\beta)/2} r^{N/2}). \end{aligned} \tag{5-40}$$

To bound (5-34), approximating the measure of the ball $B_r(P_N)$ and then the function u with the respective model behaviors, we now estimate

$$\begin{aligned}
 & \mathfrak{m}_N([0, \bar{r}])\mathfrak{q}(Q_\ell) \left| \int_{B_r(P_N)} u \mathfrak{m} - \bar{c}\sqrt{N+1} \mathfrak{q}(Q_\ell) \int_0^{\bar{r}} \cos(\cdot) \mathfrak{m}_N \right|^2 \\
 & \leq 2\mathfrak{m}_N([0, \bar{r}])\mathfrak{q}(Q_\ell) \left| \int_{B_r(P_N)} u \mathfrak{m} - \frac{1}{\mathfrak{m}_N([0, \bar{r}])} \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \\
 & \quad + 2\mathfrak{m}_N([0, \bar{r}])\mathfrak{q}(Q_\ell) \left| \frac{1}{\mathfrak{m}_N([0, \bar{r}])} \int_{B_r(P_N)} u \mathfrak{m} - \bar{c} \mathfrak{q}(Q_\ell)\sqrt{N+1} \int_0^{\bar{r}} \cos(\cdot) \mathfrak{m}_N \right|^2 \\
 & \leq 2\mathfrak{m}_N([0, \bar{r}])\mathfrak{q}(Q_\ell) \left(\int_{B_r(P_N)} u \mathfrak{m} \right)^2 \left(\frac{1}{\mathfrak{m}(B_r(P_N))} - \frac{1}{\mathfrak{m}_N([0, \bar{r}])} \right)^2 \\
 & \quad + 2 \frac{1}{\mathfrak{m}_N([0, \bar{r}])} \mathfrak{q}(Q_\ell) \left| \int_{B_r(P_N)} u \mathfrak{m} - \bar{c} \mathfrak{q}(Q_\ell)\sqrt{N+1} \int_0^{\bar{r}} \cos(\cdot) \mathfrak{m}_N \right|^2. \tag{5-41}
 \end{aligned}$$

Estimate the first term by Cauchy–Schwarz and the second term by (5-40):

$$\leq 2\mathfrak{q}(Q_\ell) \left[\frac{(\mathfrak{m}(B_r(P_N)) - \mathfrak{m}_N([0, \bar{r}]))^2}{\mathfrak{m}(B_r(P_N))\mathfrak{m}_N([0, \bar{r}])} \right] \int_{B_r(P_N)} u^2 \mathfrak{m} + C\mathfrak{q}(Q_\ell) [\delta^{\gamma+2(\beta-\gamma)/N} + \delta^{\gamma+\beta \min\{1, 2/N\}} + \delta^\beta + \delta^{1-\beta}].$$

Now use Proposition 4.5 and choose $r = \delta^{\gamma/N}$, $\gamma \in (0, \min\{\beta, 1 - \beta\})$:

$$\begin{aligned}
 & \leq 2 \left(\int_{B_r(P_N)} u^2 \mathfrak{m} \right) \left(\frac{\mathfrak{m}_N([\bar{r}, r + C\delta^{\beta/N}])}{\mathfrak{m}_N([0, \bar{r}])} \right)^2 + C(\delta^{\gamma+2(\beta-\gamma)/N} + \delta^{\gamma+\beta \min\{1, 2/N\}} + \delta^\beta + \delta^{1-\beta}) \\
 & \leq C(\delta^{\gamma+2(\beta-\gamma)/N} + \delta^{\gamma+\beta \min\{1, 2/N\}} + \delta^\beta + \delta^{1-\beta}), \tag{5-42}
 \end{aligned}$$

where the second inequality is obtained by observing that

$$\begin{aligned}
 \int_{B_r(P_N)} u^2 \mathfrak{m} &= \int_{B_r(P_N) \setminus Q_\ell} u^2 \mathfrak{m} + \int_{B_r(P_N) \cap Q_\ell} u^2 \mathfrak{m} \\
 &\leq \delta^{1-\beta} + 2 \int_{Q_\ell} \int_{[0, r+C\delta^{\beta/N}]} (u - c_q\sqrt{N+1} \cos(\cdot))^2 \mathfrak{m}_q \mathfrak{q}(dq) \\
 & \quad + 2 \int_{Q_\ell} \int_{[0, r+C\delta^{\beta/N}]} c_q^2(N+1) \cos^2(\cdot) \mathfrak{m}_q \mathfrak{q}(dq) \\
 &\leq C(\delta^{1-\beta} + \delta^\beta + \delta^{\gamma+\beta \min\{1, 2/N\}} + \delta^\gamma).
 \end{aligned}$$

The claimed estimate (5-32) is eventually obtained via triangle inequality from (5-33) and (5-42)

Step 2: Building upon Proposition 5.2, we shall obtain the bound

$$\begin{aligned}
 & \int_{Q_\ell} \int_{[0, \bar{r}]} (N+1) \left| \bar{c} \cos(\cdot) - \bar{c} \mathfrak{q}(Q_\ell) \int_0^{\bar{r}} \cos(\cdot) \mathfrak{m}_N \right|^2 \mathfrak{m}_N \mathfrak{q}(dq) \\
 & \leq 2 \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m} + C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N}) + C\delta^{1-\beta}. \tag{5-43}
 \end{aligned}$$

Thanks to the triangle inequality, the error we introduce by replacing $c_q \cos(\cdot)$ with $\bar{c} \cos(\cdot)$ can be controlled by

$$\int_{Q_\ell} \int_{[0, \bar{r}]} |c_q - \bar{c}|^2 \cos^2(t) \mathfrak{m}_N(dt) \mathfrak{q}(dq) \leq \mathfrak{m}_N([0, \bar{r}]) \int_{Q_\ell} |c_q - \bar{c}|^2 \mathfrak{q}(dq) \leq C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma) \min\{1, 2/N\}}) + C\delta^{1-\beta+2\gamma/N}, \tag{5-44}$$

where the last inequality is a consequence of (5-18) and the fact that $\bar{r} \leq r = \delta^{\gamma/N}$, $\gamma \in (0, \min\{\beta, 1 - \beta\})$. The claimed (5-43) follows from (5-44) and (5-32) via triangle inequality.

Step 3: Using the Taylor expansion $\cos(t) = 1 + O(t^2)$ in the left-hand side of (5-43), we obtain

$$\int_{Q_\ell} \int_0^{\bar{r}} (N + 1) |\bar{c} - \bar{c}\mathfrak{q}(Q_\ell)|^2 \mathfrak{m}_N \mathfrak{q}(dq) \leq \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m} + C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N}) + C\delta^{1-\beta},$$

giving

$$\mathfrak{m}_N([0, \bar{r}]) (N + 1) \bar{c}^2 (1 - \mathfrak{q}(Q_\ell))^2 \mathfrak{q}(Q_\ell) \leq 2 \int_{B_r(P_N)} \left| u - \int_{B_r(P_N)} u \mathfrak{m} \right|^2 \mathfrak{m} + C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N}) + C\delta^{1-\beta}.$$

Using the 2-2 Poincaré inequality (2-6) (combined with Bishop–Gromov volume comparison), we obtain

$$\mathfrak{m}_N([0, \bar{r}]) (N + 1) \bar{c}^2 (1 - \mathfrak{q}(Q_\ell))^2 \mathfrak{q}(Q_\ell) \leq Cr^2 \int_{B_{10r}(P_N)} |\nabla u|^2 \mathfrak{m} + C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N}) + C\delta^{1-\beta} \leq C\delta^\gamma (\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N}) + C\delta^{1-\beta}, \tag{5-45}$$

where in the last estimate we used (5-29) (recall that $r = \delta^{\gamma/N}$).

Using again that $\int_{Q_\ell} |c_q - \bar{c}|^2 \mathfrak{q}(dq) \leq C\delta^{\alpha(N)}$ from (5-18) for some $\alpha(N) > 0$, observing that

$$\int_{Q_\ell} (c_q^2 - \bar{c}^2) \mathfrak{q}(dq) = \int_{Q_\ell} |c_q - \bar{c}|^2 \mathfrak{q}(dq), \tag{5-46}$$

and recalling (5-9), we get

$$\begin{aligned} \bar{c}^2 \mathfrak{q}(Q_\ell) &= \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) + \int_{Q_\ell} (\bar{c}^2 - c_q^2) \mathfrak{q}(dq) \geq 1 - \delta^{1-\beta} - \int_{Q_\ell} |c_q - \bar{c}|^2 \mathfrak{q}(dq) \\ &\geq 1 - \delta^{1-\beta} - C\delta^{\alpha(N)} > \frac{1}{C(N)} > 0. \end{aligned} \tag{5-47}$$

Plugging (5-47) into (5-45) yields

$$(1 - \mathfrak{q}(Q_\ell))^2 \leq C(\delta^{4\gamma/N} + \delta^{(\beta-\gamma)/N} + \delta^{1-\beta-\gamma}), \tag{5-48}$$

completing the proof. □

Remark 5.4. Observe that a direct consequence of Proposition 5.3 above is an estimate of the measure of the region of the space which is not covered by transport rays, that is, $\{u = 0\}$.

Indeed (5-31) implies in particular that

$$\mathfrak{m}(X \setminus \mathcal{T}) \leq 1 - \mathfrak{q}(Q_\ell) \leq C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}). \tag{5-49}$$

5C. Proof of the main theorem. We are now ready to prove the main result putting together the estimates we proved so far. First we reduce to the set spanned by long rays using [Proposition 5.3](#); then, building upon [Proposition 5.2](#) and on [Theorem 3.11](#), we prove that on the set of long rays the function is close to a fixed multiple of the cosine of the distance from the endpoint. Eventually we change the distance from the endpoint of the ray into the distance from a pole thanks to [\(5-12\)](#).

Theorem 5.5. *For any $N \in (1, \infty)$ there exist $C(N) > 0$ and $\delta_0 = \delta_0(N) > 0$ with the following properties. Let (X, d, \mathfrak{m}) be an essentially nonbranching $CD(N - 1, N)$ m.m.s. Then, for any $u \in \text{Lip}(X)$ with $\int_X u \, \mathfrak{m} = 0, \int_X u^2 \, \mathfrak{m} = 1$ and*

$$\delta := \int_X |\nabla u|^2 \, \mathfrak{m} - N \leq \delta_0, \tag{5-50}$$

there exists a distinguished point $P \in X$ such that

$$\|u - \sqrt{N + 1} \cos(d(P, \cdot))\|_{L^2(\mathfrak{m})} \leq C(N)\delta^{1/(6N+4)}. \tag{5-51}$$

Proof. Step 1: Let us begin observing that [Proposition 5.2](#) combined with [\(5-30\)](#) and [\(5-46\)](#) gives

$$\left| \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) - \bar{c}^2 \mathfrak{q}(Q_\ell) \right| \leq C(N)(\delta^{4\gamma/N} + \delta^{1-\beta-\gamma+(2\gamma/N)} + \delta^{(\beta-\gamma)\min\{2/N, 1\}}). \tag{5-52}$$

Since from [\(5-9\)](#) we know that

$$1 - \delta^{1-\beta} \leq \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) \leq 1,$$

and in [Proposition 5.3](#) we proved that

$$\mathfrak{q}(Q_\ell) \geq 1 - C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}), \tag{5-53}$$

from [\(5-52\)](#) we infer that

$$|1 - \bar{c}^2| \leq C(N)(\delta^{2\gamma/N} + \delta^{(1-\beta-\gamma)/2} + \delta^{(\beta-\gamma)/2N}). \tag{5-54}$$

Notice that [\(5-54\)](#) implies (see for instance the proof of [\(3-48\)](#))

$$\min\{|1 - \bar{c}|, |1 + \bar{c}|\} \leq C(N)(\delta^{2\gamma/N} + \delta^{(1-\beta-\gamma)/2} + \delta^{(\beta-\gamma)/2N}). \tag{5-55}$$

Without loss of generality (up to switching the sign of u) we can assume that

$$|1 - \bar{c}| = \min\{|1 - \bar{c}|, |1 + \bar{c}|\}.$$

The combination of [Proposition 5.2](#) and [\(5-55\)](#) gives

$$\begin{aligned} \int_{Q_\ell} |c_q - 1|^2 \mathfrak{q}(dq) &\leq 2 \int_{Q_\ell} |c_q - \bar{c}|^2 \mathfrak{q}(dq) + 2|\bar{c} - 1|^2 \mathfrak{q}(Q_\ell) \\ &\leq C(N)(\delta^{4\gamma/N} + \delta^{1-\beta-\gamma} + \delta^{(\beta-\gamma)/N}). \end{aligned} \tag{5-56}$$

Step 2: Next we let P be equal to P_N given in [Proposition 5.1](#). We get

$$\begin{aligned} & \|u - \sqrt{N+1} \cos(d(P, \cdot))\|_{L^2(\mathfrak{m})}^2 \\ &= \int_Q \int_{X_q} |u - \sqrt{N+1} \cos(d(P, \cdot))|^2 \mathfrak{m}_q \mathfrak{q}(dq) + \int_{X \setminus \mathcal{T}} (N+1) \cos(d(P, \cdot))^2 \mathfrak{m} \\ &\leq \int_{Q_\ell} \int_{X_q} |u - \sqrt{N+1} \cos(d(P, \cdot))|^2 \mathfrak{m}_q \mathfrak{q}(dq) \\ &\qquad\qquad\qquad + 2 \int_{X \setminus R(Q_\ell)} u^2 \mathfrak{m} + 2(N+1) \mathfrak{q}(Q \setminus Q_\ell) + (N+1) \mathfrak{m}(X \setminus \mathcal{T}). \end{aligned}$$

Using [\(5-9\)](#), [\(5-53\)](#) and [Remark 5.4](#), recalling that we are tacitly identifying the ray X_q with the interval $(0, |X_q|)$,

$$\begin{aligned} &\leq \int_{Q_\ell} \int_{X_q} |u - \sqrt{N+1} \cos(d(P, \cdot))|^2 \mathfrak{m}_q \mathfrak{q}(dq) + C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}) \\ &\leq 2 \int_{Q_\ell} \int_{X_q} |u - \sqrt{N+1} \cos(\cdot)|^2 \mathfrak{m}_q \mathfrak{q}(dq) \\ &\qquad\qquad\qquad + 2 \int_{Q_\ell} \int_{X_q} |\cos(\cdot) - \cos(d(P, \cdot))|^2 \mathfrak{m}_q \mathfrak{q}(dq) + C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}). \end{aligned}$$

Using triangle inequality to estimate the first term and [\(5-12\)](#) for the second,

$$\begin{aligned} &\leq 4 \int_{Q_\ell} \int_{X_q} |u - \sqrt{N+1} c_q \cos(\cdot)|^2 \mathfrak{m}_q \mathfrak{q}(dq) + C(N) \int_{Q_\ell} |c_q - 1|^2 \mathfrak{q}(dq) \\ &\qquad\qquad\qquad + C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}). \end{aligned}$$

By [\(5-13\)](#) and [\(5-56\)](#),

$$\begin{aligned} &\leq C(N) \delta^{\beta \min\{1, 2/N\}} \int_{Q_\ell} c_q^2 \mathfrak{q}(dq) + C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}) \\ &\leq C(N)(\delta^{2\gamma/N} + \delta^{(\beta-\gamma)/2N} + \delta^{(1-\beta-\gamma)/2}). \end{aligned} \tag{5-57}$$

The optimal choice of parameters in [\(5-57\)](#) is $\beta = 5N/(6N+4)$ and $\gamma = N/(6N+4)$ giving the claim [\(5-51\)](#). □

Acknowledgements

Part of the work was developed when Semola was visiting Cavalletti at SISSA and Mondino at the Mathematics Institute of the University of Warwick; he wishes to thank both institutes for the excellent working conditions and the stimulating atmosphere.

Mondino is supported by the EPSRC First Grant EP/R004730/1 ‘‘Optimal transport and geometric analysis’’ and by the ERC Starting Grant 802689 ‘‘CURVATURE’’.

The authors are grateful to the anonymous reviewers, for their suggestions that helped to improve a previous version of the paper.

References

- [Ambrosio 2018] L. Ambrosio, “Calculus, heat flow and curvature-dimension bounds in metric measure spaces”, pp. 301–340 in *Proceedings of the International Congress of Mathematicians, I: Invited lectures*, edited by B. Sirakov et al., World Sci., Hackensack, NJ, 2018. [MR](#) [Zbl](#)
- [Ambrosio et al. 2014] L. Ambrosio, N. Gigli, and G. Savaré, “Metric measure spaces with Riemannian Ricci curvature bounded from below”, *Duke Math. J.* **163**:7 (2014), 1405–1490. [MR](#) [Zbl](#)
- [Ambrosio et al. 2015] L. Ambrosio, N. Gigli, A. Mondino, and T. Rajala, “Riemannian Ricci curvature lower bounds in metric measure spaces with σ -finite measure”, *Trans. Amer. Math. Soc.* **367**:7 (2015), 4661–4701. [MR](#) [Zbl](#)
- [Ambrosio et al. 2019] L. Ambrosio, A. Mondino, and G. Savaré, *Nonlinear diffusion equations and curvature conditions in metric measure spaces*, Mem. Amer. Math. Soc. **1270**, Amer. Math. Soc., Providence, RI, 2019. [MR](#) [Zbl](#)
- [Bacher and Sturm 2010] K. Bacher and K.-T. Sturm, “Localization and tensorization properties of the curvature-dimension condition for metric measure spaces”, *J. Funct. Anal.* **259**:1 (2010), 28–56. [MR](#) [Zbl](#)
- [Bakry and Qian 2000] D. Bakry and Z. Qian, “Some new results on eigenvectors via dimension, diameter, and Ricci curvature”, *Adv. Math.* **155**:1 (2000), 98–153. [MR](#) [Zbl](#)
- [Bérard and Meyer 1982] P. Bérard and D. Meyer, “Inégalités isopérimétriques et applications”, *Ann. Sci. École Norm. Sup. (4)* **15**:3 (1982), 513–541. [MR](#) [Zbl](#)
- [Bérard et al. 1985] P. Bérard, G. Besson, and S. Gallot, “Sur une inégalité isopérimétrique qui généralise celle de Paul Lévy–Gromov”, *Invent. Math.* **80**:2 (1985), 295–308. [MR](#) [Zbl](#)
- [Bertrand 2007] J. Bertrand, “Pincement spectral en courbure de Ricci positive”, *Comment. Math. Helv.* **82**:2 (2007), 323–352. [MR](#) [Zbl](#)
- [Bianchini and Caravenna 2009] S. Bianchini and L. Caravenna, “On the extremality, uniqueness and optimality of transference plans”, *Bull. Inst. Math. Acad. Sin. (N.S.)* **4**:4 (2009), 353–454. [MR](#) [Zbl](#)
- [Brasco and Pratelli 2012] L. Brasco and A. Pratelli, “Sharp stability of some spectral inequalities”, *Geom. Funct. Anal.* **22**:1 (2012), 107–135. [MR](#) [Zbl](#)
- [Brasco et al. 2015] L. Brasco, G. De Philippis, and B. Velichkov, “Faber–Krahn inequalities in sharp quantitative form”, *Duke Math. J.* **164**:9 (2015), 1777–1831. [MR](#) [Zbl](#)
- [Cavalletti and Milman 2021] F. Cavalletti and E. Milman, “The globalization theorem for the curvature-dimension condition”, *Invent. Math.* **226**:1 (2021), 1–137. [MR](#) [Zbl](#)
- [Cavalletti and Mondino 2017a] F. Cavalletti and A. Mondino, “Optimal maps in essentially non-branching spaces”, *Commun. Contemp. Math.* **19**:6 (2017), art. id. 1750007. [MR](#) [Zbl](#)
- [Cavalletti and Mondino 2017b] F. Cavalletti and A. Mondino, “Sharp and rigid isoperimetric inequalities in metric-measure spaces with lower Ricci curvature bounds”, *Invent. Math.* **208**:3 (2017), 803–849. [MR](#) [Zbl](#)
- [Cavalletti and Mondino 2017c] F. Cavalletti and A. Mondino, “Sharp geometric and functional inequalities in metric measure spaces with lower Ricci curvature bounds”, *Geom. Topol.* **21**:1 (2017), 603–645. [MR](#) [Zbl](#)
- [Cavalletti and Mondino 2018] F. Cavalletti and A. Mondino, “Isoperimetric inequalities for finite perimeter sets under lower Ricci curvature bounds”, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **29**:3 (2018), 413–430. [MR](#) [Zbl](#)
- [Cavalletti et al. 2019] F. Cavalletti, F. Maggi, and A. Mondino, “Quantitative isoperimetry à la Levy–Gromov”, *Comm. Pure Appl. Math.* **72**:8 (2019), 1631–1677. [MR](#) [Zbl](#)
- [Cheng 1975] S. Y. Cheng, “Eigenvalue comparison theorems and its geometric applications”, *Math. Z.* **143**:3 (1975), 289–297. [MR](#) [Zbl](#)
- [Chodosh et al. 2023] O. Chodosh, M. Engelstein, and L. Spolaor, “The Riemannian quantitative isoperimetric inequality”, *J. Eur. Math. Soc. (JEMS)* **25**:5 (2023), 1711–1741. [MR](#) [Zbl](#)
- [Cicalese and Leonardi 2012] M. Cicalese and G. P. Leonardi, “A selection principle for the sharp quantitative isoperimetric inequality”, *Arch. Ration. Mech. Anal.* **206**:2 (2012), 617–643. [MR](#) [Zbl](#)
- [Croke 1982] C. B. Croke, “An eigenvalue pinching theorem”, *Invent. Math.* **68**:2 (1982), 253–256. [MR](#) [Zbl](#)

- [Erbar et al. 2015] M. Erbar, K. Kuwada, and K.-T. Sturm, “On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces”, *Invent. Math.* **201**:3 (2015), 993–1071. [MR](#) [Zbl](#)
- [Faber 1923] G. Faber, “Beweis, daß unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt”, *Sitz. Bayer Acad. Wiss.* **1923**:8 (1923), 169–172. [Zbl](#)
- [Figalli et al. 2010] A. Figalli, F. Maggi, and A. Pratelli, “A mass transportation approach to quantitative isoperimetric inequalities”, *Invent. Math.* **182**:1 (2010), 167–211. [MR](#) [Zbl](#)
- [Fremlin 2006] D. H. Fremlin, *Measure theory, IV: Topological measure spaces*, Torres Fremlin, Colchester, 2006. [MR](#) [Zbl](#)
- [Fusco et al. 2008] N. Fusco, F. Maggi, and A. Pratelli, “The sharp quantitative isoperimetric inequality”, *Ann. of Math. (2)* **168**:3 (2008), 941–980. [MR](#) [Zbl](#)
- [Fusco et al. 2009] N. Fusco, F. Maggi, and A. Pratelli, “Stability estimates for certain Faber–Krahn, isocapacitary and Cheeger inequalities”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **8**:1 (2009), 51–71. [MR](#) [Zbl](#)
- [Gigli 2015] N. Gigli, *On the differential structure of metric measure spaces and applications*, Mem. Amer. Math. Soc. **1113**, Amer. Math. Soc., Providence, RI, 2015. [MR](#) [Zbl](#)
- [Gigli et al. 2015] N. Gigli, A. Mondino, and G. Savaré, “Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows”, *Proc. Lond. Math. Soc. (3)* **111**:5 (2015), 1071–1129. [MR](#) [Zbl](#)
- [Gromov 1999] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Progr. Math. **152**, Birkhäuser, Boston, 1999. [MR](#) [Zbl](#)
- [Gromov and Milman 1987] M. Gromov and V. D. Milman, “Generalization of the spherical isoperimetric inequality to uniformly convex Banach spaces”, *Compos. Math.* **62**:3 (1987), 263–282. [MR](#) [Zbl](#)
- [Hajlasz and Koskela 2000] P. Hajlasz and P. Koskela, *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **688**, Amer. Math. Soc., Providence, RI, 2000. [MR](#) [Zbl](#)
- [Hansen and Nadirashvili 1994] W. Hansen and N. Nadirashvili, “Isoperimetric inequalities in potential theory”, *Potential Anal.* **3**:1 (1994), 1–14. [MR](#) [Zbl](#)
- [Jiang and Zhang 2016] Y. Jiang and H.-C. Zhang, “Sharp spectral gaps on metric measure spaces”, *Calc. Var. Partial Differential Equations* **55**:1 (2016), art. id. 14. [MR](#) [Zbl](#)
- [Kannan et al. 1995] R. Kannan, L. Lovász, and M. Simonovits, “Isoperimetric problems for convex bodies and a localization lemma”, *Discrete Comput. Geom.* **13**:3-4 (1995), 541–559. [MR](#) [Zbl](#)
- [Ketterer 2015] C. Ketterer, “Obata’s rigidity theorem for metric measure spaces”, *Anal. Geom. Metr. Spaces* **3**:1 (2015), 278–295. [MR](#) [Zbl](#)
- [Klartag 2017] B. Klartag, *Needle decompositions in Riemannian geometry*, Mem. Amer. Math. Soc. **1180**, Amer. Math. Soc., Providence, RI, 2017. [MR](#) [Zbl](#)
- [Krahn 1925] E. Krahn, “Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises”, *Math. Ann.* **94**:1 (1925), 97–100. [MR](#) [Zbl](#)
- [Li and Yau 1980] P. Li and S. T. Yau, “Estimates of eigenvalues of a compact Riemannian manifold”, pp. 205–239 in *Geometry of the Laplace operator* (Honolulu, 1979), edited by R. Osserman and A. Weinstein, Proc. Sympos. Pure Math. **36**, Amer. Math. Soc., Providence, RI, 1980. [MR](#) [Zbl](#)
- [Lichnerowicz 1958] A. Lichnerowicz, *Géométrie des groupes de transformations*, Travaux et Rech. Math. **3**, Dunod, Paris, 1958. [MR](#) [Zbl](#)
- [Lott and Villani 2009] J. Lott and C. Villani, “Ricci curvature for metric-measure spaces via optimal transport”, *Ann. of Math. (2)* **169**:3 (2009), 903–991. [MR](#) [Zbl](#)
- [Lovász and Simonovits 1993] L. Lovász and M. Simonovits, “Random walks in a convex body and an improved volume algorithm”, *Random Structures Algorithms* **4**:4 (1993), 359–412. [MR](#) [Zbl](#)
- [Matei 2000] A.-M. Matei, “First eigenvalue for the p -Laplace operator”, *Nonlinear Anal.* **39**:8 (2000), 1051–1068. [MR](#) [Zbl](#)
- [Melas 1992] A. D. Melas, “The stability of some eigenvalue estimates”, *J. Differential Geom.* **36**:1 (1992), 19–33. [MR](#) [Zbl](#)
- [Milman 2015] E. Milman, “Sharp isoperimetric inequalities and model spaces for the curvature-dimension-diameter condition”, *J. Eur. Math. Soc.* **17**:5 (2015), 1041–1078. [MR](#) [Zbl](#)

- [Mondino and Semola 2020] A. Mondino and D. Semola, “Polya–Szegő inequality and Dirichlet p -spectral gap for non-smooth spaces with Ricci curvature bounded below”, *J. Math. Pures Appl.* (9) **137** (2020), 238–274. [MR](#) [Zbl](#)
- [Obata 1962] M. Obata, “Certain conditions for a Riemannian manifold to be isometric with a sphere”, *J. Math. Soc. Japan* **14** (1962), 333–340. [MR](#) [Zbl](#)
- [Ohta 2007] S.-I. Ohta, “Products, cones, and suspensions of spaces with the measure contraction property”, *J. Lond. Math. Soc.* (2) **76**:1 (2007), 225–236. [MR](#) [Zbl](#)
- [Ohta 2009] S.-i. Ohta, “Finsler interpolation inequalities”, *Calc. Var. Partial Differential Equations* **36**:2 (2009), 211–249. [MR](#) [Zbl](#)
- [Payne and Weinberger 1960] L. E. Payne and H. F. Weinberger, “An optimal Poincaré inequality for convex domains”, *Arch. Ration. Mech. Anal.* **5** (1960), 286–292. [MR](#) [Zbl](#)
- [Petersen 1999] P. Petersen, “On eigenvalue pinching in positive Ricci curvature”, *Invent. Math.* **138**:1 (1999), 1–21. [MR](#) [Zbl](#)
- [Petrunin 2011] A. Petrunin, “Alexandrov meets Lott–Villani–Sturm”, *Münster J. Math.* **4** (2011), 53–64. [MR](#) [Zbl](#)
- [Pólya and Szegő 1951] G. Pólya and G. Szegő, *Isoperimetric inequalities in mathematical physics*, Ann. of Math. Stud. **27**, Princeton Univ. Press, 1951. [MR](#) [Zbl](#)
- [Rajala 2012] T. Rajala, “Local Poincaré inequalities from stable curvature conditions on metric spaces”, *Calc. Var. Partial Differential Equations* **44**:3-4 (2012), 477–494. [MR](#) [Zbl](#)
- [Rajala and Sturm 2014] T. Rajala and K.-T. Sturm, “Non-branching geodesics and optimal maps in strong $CD(K, \infty)$ -spaces”, *Calc. Var. Partial Differential Equations* **50**:3-4 (2014), 831–846. [MR](#) [Zbl](#)
- [Rayleigh 1877] J. W. Strutt, 3rd Baron Rayleigh, *The theory of sound*, Macmillan, London, 1877.
- [Sturm 2006a] K.-T. Sturm, “On the geometry of metric measure spaces, I”, *Acta Math.* **196**:1 (2006), 65–131. [MR](#) [Zbl](#)
- [Sturm 2006b] K.-T. Sturm, “On the geometry of metric measure spaces, II”, *Acta Math.* **196**:1 (2006), 133–177. [MR](#) [Zbl](#)
- [Villani 2019] C. Villani, “Inégalités isopérimétriques dans les espaces métriques mesurés [d’après F. Cavalletti and A. Mondino]”, exposé 1127, pp. 213–265 in *Séminaire Bourbaki*, 2016/2017, Astérisque **407**, Soc. Math. France, Paris, 2019. [MR](#) [Zbl](#)
- [Zhang and Zhu 2010] H.-C. Zhang and X.-P. Zhu, “Ricci curvature on Alexandrov spaces and rigidity theorems”, *Comm. Anal. Geom.* **18**:3 (2010), 503–553. [MR](#) [Zbl](#)

Received 24 Jul 2021. Revised 20 Dec 2021. Accepted 31 Jan 2022.

FABIO CAVALLETTI: cavallet@sissa.it

Scuola Internazionale Superiore di Studi Avanzati, Trieste, Italy

ANDREA MONDINO: andrea.mondino@maths.ox.ac.uk

Mathematical Institute, University of Oxford, Oxford, United Kingdom

DANIELE SEMOLA: daniele.semola@sns.it

Scuola Normale Superiore, Pisa, Italy

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard [Universit  Paris Sud XI, France](mailto:patrick.gerard@universite-paris-saclay.fr)
patrick.gerard@universite-paris-saclay.fr

Cl ment Mouhot [Cambridge University, UK](mailto:c.mouhot@dpmms.cam.ac.uk)
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

| | | | |
|--------------------|--|-----------------------|--|
| Massimiliano Berti | Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it | William Minicozzi II | Johns Hopkins University, USA minicozz@math.jhu.edu |
| Zbigniew Blocki | Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl | Werner Müller | Universit t Bonn, Germany mueller@math.uni-bonn.de |
| Charles Fefferman | Princeton University, USA cf@math.princeton.edu | Igor Rodnianski | Princeton University, USA irod@math.princeton.edu |
| David G rard-Varet | Universit  de Paris, France david.gerard-varet@imj-prg.fr | Yum-Tong Siu | Harvard University, USA siu@math.harvard.edu |
| Colin Guillarmou | Universit  Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr | Terence Tao | University of California, Los Angeles, USA tao@math.ucla.edu |
| Ursula Hamenstaedt | Universit t Bonn, Germany ursula@math.uni-bonn.de | Michael E. Taylor | Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu |
| Vadim Kaloshin | University of Maryland, USA vadim.kaloshin@gmail.com | Gunther Uhlmann | University of Washington, USA gunther@math.washington.edu |
| Izabella Laba | University of British Columbia, Canada ilaba@math.ubc.ca | Andr s Vasy | Stanford University, USA andras@math.stanford.edu |
| Anna L. Mazzucato | Penn State University, USA alm24@psu.edu | Dan Virgil Voiculescu | University of California, Berkeley, USA dvv@math.berkeley.edu |
| Richard B. Melrose | Massachussets Inst. of Tech., USA rbm@math.mit.edu | Jim Wright | University of Edinburgh, UK j.r.wright@ed.ac.uk |
| Frank Merle | Universit  de Cergy-Pontoise, France merle@ihes.fr | Maciej Zworski | University of California, Berkeley, USA zworski@math.berkeley.edu |

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

  2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 6 2023

| | |
|--|------|
| The Cauchy problem for the infinitesimal model in the regime of small variance FLORIAN PATOUT | 1289 |
| The “good” Boussinesq equation: long-time asymptotics CHRISTOPHE CHARLIER, JONATAN LENELLS and DENG-SHAN WANG | 1351 |
| Quantitative Obata’s theorem FABIO CAVALLETTI, ANDREA MONDINO and DANIELE SEMOLA | 1389 |
| Couniversality and controlled maps on product systems over right LCM semigroups EVGENIOS T.A. KAKARIADIS, ELIAS G. KATSOULIS, MARCELO LACA and XIN LI | 1433 |