

DE VRIES POWERS AND PROXIMITY SPECKER ALGEBRAS

G. BEZHANISHVILI, L. CARAI, P. J. MORANDI, AND B. OLBERDING

ABSTRACT. By de Vries duality [9], the category \mathbf{KHaus} of compact Hausdorff spaces is dually equivalent to the category \mathbf{DeV} of de Vries algebras. In [5] an alternate duality for \mathbf{KHaus} was developed, where de Vries algebras were replaced by proximity Baer-Specker algebras. The functor associating with each compact Hausdorff space a proximity Baer-Specker algebra was described by generalizing the notion of a boolean power of a totally ordered domain to that of a de Vries power. It follows that \mathbf{DeV} is equivalent to the category \mathbf{PBSp} of proximity Baer-Specker algebras. The equivalence is obtained by passing through \mathbf{KHaus} , and hence is not choice-free. In this paper we give a direct algebraic proof of this equivalence, which is choice-free. To do so, we give an alternate choice-free description of de Vries powers of a totally ordered domain.

2020 Mathematics subject classification. 06F25; 13G05; 54E05.

Keywords. Proximity, de Vries algebra, Specker algebra, Baer ring, integral domain, boolean power.

1. INTRODUCTION

By the celebrated Stone duality, the category \mathbf{BA} of boolean algebras is dually equivalent to the category \mathbf{Stone} of Stone spaces (zero-dimensional compact Hausdorff spaces). The functor from \mathbf{Stone} to \mathbf{BA} associates with each Stone space X , the boolean algebra $\mathbf{Clop}(X)$ of clopen subsets of X . Thinking of clopen subsets of X as continuous characteristic functions, we can identify $\mathbf{Clop}(X)$ with the idempotents of the \mathbb{R} -algebra $C(X)$ of all continuous real-valued functions on X . The \mathbb{R} -subalgebra of $C(X)$ generated by the idempotents of $C(X)$ is then the \mathbb{R} -algebra of finitely-valued continuous real-valued functions on X . The reals can be replaced by an arbitrary domain D with the discrete topology, thus yielding the notion of a Specker D -algebra, which, as the D -algebra of (finitely-valued) continuous D -valued functions on a Stone space, is nothing more than a boolean power of D (see [6]). Let \mathbf{Sp}_D be the category of Specker D -algebras. Then \mathbf{Sp}_D is dually equivalent to \mathbf{Stone} [6, Cor. 3.9], hence \mathbf{Sp}_D is equivalent to \mathbf{BA} , and we arrive at the following commutative diagram:

$$\begin{array}{ccc} \mathbf{Sp}_D & \xleftrightarrow{\quad} & \mathbf{BA} \\ & \searrow & \swarrow \\ & \mathbf{Stone} & \end{array}$$

Stone duality was generalized to compact Hausdorff spaces by de Vries [9]. In de Vries duality, with each compact Hausdorff space X , we associate the complete boolean algebra $\mathbf{RO}(X)$ of regular open subsets of X equipped with the proximity relation $<$ given by $U < V$ iff $\text{cl}(U) \subseteq V$. The resulting structures are known as de Vries algebras [3]. Stone duality

then lifts to a dual equivalence between the categories \mathbf{KHaus} of compact Hausdorff spaces and \mathbf{DeV} of de Vries algebras.

The same way $\mathbf{Clop}(X)$ can be identified with the idempotents of the \mathbb{R} -algebra $C(X)$ of continuous real-valued functions, $\mathcal{RO}(X)$ can be identified with the idempotents of the \mathbb{R} -algebra $N(X)$ of normal real-valued functions (see, e.g., [4, Lem. 6.5]). The notion of a normal function originates in the work of Dilworth [10], where the MacNeille completion of the lattice $C(X)$ was characterized as the lattice $N(X)$.

Dilworth's notion of a normal real-valued function requires working with the underlying order of \mathbb{R} . In [5], the notion of a finitely-valued normal function was generalized to an arbitrary totally ordered algebra D . This paved a way to generalize boolean powers of D , which are defined over a Stone space, to the more general notion of de Vries powers of D , which are defined over a compact Hausdorff space. More precisely, if $(B, <)$ is a de Vries algebra and X is its dual compact Hausdorff space, then the de Vries power of D by $(B, <)$ is the algebra $FN(X)$ of finitely-valued normal functions $f : X \rightarrow D$. The de Vries algebra $(B, <)$ can then be identified with the idempotents of $FN(X)$, and the proximity $<$ can be lifted to a proximity \triangleleft on $FN(X)$. Thus, the de Vries power of D by $(B, <)$ is the proximity algebra $(FN(X), \triangleleft)$.

As was shown in [5], in the special case when D is a totally ordered domain, de Vries powers can be characterized using the notion of Baer-Specker D -algebras. These algebras have a long history. We refer to [7] for details. In [5] de Vries proximities on boolean algebras were generalized to proximity relations on Specker D -algebras which has resulted in the category \mathbf{PBSp}_D of proximity Baer-Specker D -algebras. One of the main results of [5] establishes that this category is dually equivalent to \mathbf{KHaus} , and hence is equivalent to \mathbf{DeV} , thus yielding the following commutative diagram, which lifts the diagram given above for Stone duality via boolean powers to de Vries duality via de Vries powers.

$$\begin{array}{ccc}
 \mathbf{PBSp}_D & \xleftrightarrow{\quad} & \mathbf{DeV} \\
 & \swarrow \quad \searrow & \\
 & \mathbf{KHaus} &
 \end{array}$$

The equivalence between \mathbf{DeV} and \mathbf{PBSp}_D is obtained by passing through \mathbf{KHaus} , hence is not choice-free. In this article we give a purely algebraic proof of this equivalence, which is choice-free. This we do by going back to the original definition of boolean powers by Foster [11, 12, 13]. Using this approach, we can see that Specker D -algebras, defined as idempotent-generated torsion-free D -algebras, are boolean powers by utilizing orthogonal decompositions of elements of these algebras (see Section 2). However, orthogonal decompositions are ill suited for working with a proximity on a Specker D -algebra, and so we introduce a different decomposition for the elements in the algebra, a decreasing decomposition that is reminiscent of Mundici's good sequences ([17, p. 28]). These decreasing decompositions are the key ingredient in lifting de Vries proximities to proximities on Baer-Specker algebras in a choice-free manner.

The paper is organized as follows. In Section 2 we use the original definition of Foster to give a choice-free proof that boolean powers of a domain D are exactly the Specker D -algebras. Starting from Section 3, we assume that D is a totally ordered domain. In Section 3 we give a choice-free proof that there is a unique partial ordering on a Specker D -algebra S making it a torsion-free f -algebra over D . In Section 4 we recall the notion of a proximity on a Specker D -algebra and give an alternate description of a boolean power of D using decreasing decompositions. In Section 5 we use decreasing decompositions to lift a de Vries proximity from the boolean algebra $\text{Id}(S)$ of idempotents to the Specker D -algebra S . This allows us to give a choice-free definition of a de Vries power of D and prove that de Vries powers of D are exactly the Baer-Specker D -algebras. Finally, in Section 6 we give a direct choice-free proof that the category DeV of de Vries algebras is equivalent to the category PBSp_D of proximity Baer-Specker D -algebras.

2. SPECKER ALGEBRAS AND BOOLEAN POWERS

In this section D is an arbitrary fixed integral domain. For a commutative unital D -algebra S , we let $\text{Id}(S)$ be the boolean algebra of its idempotents.

Definition 2.1. Let S be a commutative unital D -algebra.

- (1) We call S *idempotent-generated* if S is generated as a D -algebra by $\text{Id}(S)$.
- (2) We call S a *Specker D -algebra* if S is idempotent-generated and torsion-free as a D -module.
- (3) We denote by \mathbf{Sp}_D the category of Specker D -algebras and unital D -algebra homomorphisms.

Various characterizations of Specker D -algebras can be found in [6]. Some of these we collect in the next theorem. Recall that the boolean power of D by a boolean algebra B is the D -algebra $C(X, D)$ of continuous functions $f : X \rightarrow D$, where X is the Stone space of B and D is given the discrete topology.

Theorem 2.2. *For a commutative unital D -algebra S , the following are equivalent:*

- (1) S is a Specker D -algebra.
- (2) S is isomorphic to an idempotent-generated subalgebra of a power of D .
- (3) S is isomorphic to a boolean power of D .
- (4) S is idempotent-generated and a free D -module.

The definition of a boolean power given before the theorem is due to Jónsson (see [1, p. 5]). Because the definition involves the Stone space of B , it is not choice-free. Since this definition was used in [6], the proof of Theorem 2.2 is also not choice-free. To avoid this reliance on the axiom of choice, we revert to the original definition of a boolean power given by Foster (see, e.g., [13, p. 31]):

Definition 2.3. The (bounded) *boolean power of D by B* is the D -algebra $D[B]^*$ of finitely-valued functions $f : D \rightarrow B$ such that $f(a) \wedge f(b) = 0$ for all $a \neq b$ in D and $\bigvee \text{Im} f = 1$. The algebra operations on $D[B]^*$ are defined as follows, where $a, b \in D$ and $f, g \in D[B]^*$:

- $(f + g)(a) = \bigvee \{f(b) \wedge g(c) : b + c = a\}$.

- $(fg)(a) = \bigvee \{f(b) \wedge g(c) : bc = a\}$.
- $(bf)(a) = \bigvee \{f(c) : bc = a\}$.

Using Definition 2.3, a choice-free proof that S is a Specker D -algebra iff S is isomorphic to a boolean power was outlined in [6, Rem. 2.9]. Since this observation is important to our point of view in the present article, we give the details below. For this we recall orthogonal decompositions of elements of Specker D -algebras.

Definition 2.4. Let S be a Specker D -algebra and $B = \text{Id}(S)$. An *orthogonal decomposition* of $s \in S$ is a representation $s = \sum_{i=0}^n a_i e_i$ with $a_i \in D$ (not necessarily distinct) and $e_i \in B$ are pairwise orthogonal (that is, $e_i \wedge e_j = 0$ for each $i \neq j$). If, in addition, $e_0 \vee \dots \vee e_n = 1$, we call this a *full orthogonal decomposition*.

By [6, Lem. 2.1], each $s \in S$ has a unique full orthogonal decomposition with distinct coefficients. To connect orthogonal decompositions with the boolean power of D by B , let $s = \sum_{i=0}^n a_i e_i$ be a full orthogonal decomposition of $s \in S$ with the a_i distinct. Define $s^\perp : D \rightarrow B$ by

$$s^\perp(a) = \begin{cases} e_i & \text{if } a = a_i \text{ for some } i, \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to see that $s^\perp \in D[B]^*$ and $s = \sum_{a \in D} a s^\perp(a)$. Conversely, if $f \in D[B]^*$, then $s = \sum_{a \in D} a f(a)$ is a full orthogonal decomposition of s with distinct coefficients such that $s^\perp = f$. We show that this correspondence is an isomorphism.

Theorem 2.5. *Let S be a Specker D -algebra and $B = \text{Id}(S)$. Then the map $(-)^{\perp} : S \rightarrow D[B]^*$ is a D -algebra isomorphism. Moreover, the restriction of $(-)^{\perp}$ to B is a boolean isomorphism from B to $\text{Id}(D[B]^*)$.*

Proof. We just saw that $(-)^{\perp}$ is a bijection. It remains to show that $(-)^{\perp}$ is a D -algebra homomorphism. Let $s, t \in S$ and let $s = \sum_i a_i e_i$ and $t = \sum_j b_j f_j$ be full orthogonal decompositions. Then $s = \sum_{i,j} a_i (e_i \wedge f_j)$ and $t = \sum_{i,j} b_j (e_i \wedge f_j)$. Therefore, s, t have full orthogonal decompositions with the same set of idempotents (but not necessarily distinct coefficients). Thus, without loss of generality we may assume that s and t have orthogonal decompositions $s = \sum_i a_i e_i$ and $t = \sum_i b_i e_i$. Then $s^\perp(a) = \bigvee \{e_i : a = a_i\}$, and a similar description holds for t^\perp . Applying the definition of $s^\perp + t^\perp$ and the fact that the e_i are pairwise orthogonal, we obtain

$$\begin{aligned} (s^\perp + t^\perp)(a) &= \bigvee \{s^\perp(b) \wedge t^\perp(c) : b + c = a\} \\ &= \bigvee \{\bigvee \{e_i : b = a_i\} \wedge \bigvee \{e_j : c = b_j\} : b + c = a\} \\ &= \bigvee \{\bigvee \{e_i \wedge e_j : b = a_i \text{ and } c = b_j\} : b + c = a\} \\ &= \bigvee \{\bigvee \{e_i : b = a_i \text{ and } c = b_i\} : b + c = a\} \\ &= \bigvee \{e_i : a_i + b_i = a\} \\ &= (s + t)^\perp(a), \end{aligned}$$

where the last equality holds since $s + t = \sum_i (a_i + b_i) e_i$. Therefore, $(s + t)^\perp = s^\perp + t^\perp$. The proofs that $(st)^\perp = s^\perp t^\perp$ and $(bs)^\perp = b s^\perp$ for $b \in D$ are similar. Thus, $(-)^{\perp}$ is a D -algebra isomorphism. Finally, since $(-)^{\perp}$ is a ring isomorphism from S to $D[B]^*$, it restricts to a boolean isomorphism between $B = \text{Id}(S)$ and $\text{Id}(D[B]^*)$. \square

It follows from Theorem 2.5 that if S is a Specker D -algebra, then S is isomorphic to the boolean power of D by $\text{Id}(S)$. To prove that every boolean power of D is a Specker D -algebra, we require the following construction that has its roots in the work of Bergman [2] and Rota [18].

Definition 2.6. [6, Def. 2.4] For a boolean algebra B , let $D[B]$ be the quotient ring of the polynomial ring $D[\{x_e : e \in B\}]$ over D in variables indexed by the elements of B modulo the ideal I_B generated by the following elements, as e, f range over B :

$$x_{e \wedge f} - x_e x_f, \quad x_{e \vee f} - (x_e + x_f - x_e x_f), \quad x_{\neg e} - (1 - x_e), \quad x_0.$$

The following result follows from the definition of $D[B]$ and [6, Lem. 3.2(4)].

Theorem 2.7. $D[B]$ is a Specker D -algebra and B is isomorphic to $\text{Id}(D[B])$.

We thus are ready for a choice-free proof that Specker D -algebras are boolean powers of D .

Corollary 2.8. Boolean powers of a domain D are, up to isomorphism in the category of D -algebras, precisely the Specker D -algebras.

Proof. If S is a Specker D -algebra, then Theorem 2.5 yields that S is isomorphic to the boolean power $D[\text{Id}(S)]^*$. Conversely, by Theorem 2.7, $D[B]$ is a Specker D -algebra and B is isomorphic to $\text{Id}(D[B])$. Thus, by Theorem 2.5, the boolean power of D by B is isomorphic to the Specker D -algebra $D[B]$. \square

Remark 2.9. Let $B \in \mathbf{BA}$ and let $i_B : B \rightarrow \text{Id}(D[B])$ be the boolean isomorphism of Theorem 2.7. Composing i_B with the boolean isomorphism $(-)^{\perp} : \text{Id}(D[B]) \rightarrow \text{Id}(D[B]^*)$ of Theorem 2.5 yields a boolean isomorphism which sends $e \in B$ to $e^{\perp} \in \text{Id}(D[B]^*)$, given by

$$e^{\perp}(a) = \begin{cases} e & \text{if } a = 1, \\ \neg e & \text{if } a = 0, \\ 0 & \text{if } a \neq 0, 1. \end{cases}$$

Remark 2.10. More generally, Specker algebras can be defined over an arbitrary commutative ring R with 1, but the definition is more subtle when zero divisors are present. This was done in [6], where the notion of a faithful generating algebra of idempotents was introduced. If we replace $\text{Id}(S)$ by such a generating algebra, then the proofs of Theorem 2.5 and Corollary 2.8 generalize, thus yielding a choice-free proof of [6, Thm. 2.7] that boolean powers of R are precisely the Specker R -algebras. The reason we restrict to domains will become clear when we introduce proximities on Specker algebras; see Remark 4.2(4).

3. SPECKER ALGEBRAS OVER TOTALLY ORDERED DOMAINS

From now on we assume that D is a totally ordered domain. It was shown in [6, Thm. 5.1] that there is a unique ordering on a Specker D -algebra S that makes S into a torsion-free f -algebra over D . But the proof is not choice-free. In Theorem 3.2 we give a choice-free proof of this result, and also show that the isomorphism of Theorem 2.5 is an order isomorphism.

We start by recalling some basic definitions of ordered rings (see, e.g., [8, Ch. XVII.5]). A ring R with a partial ordering \leq is an ℓ -ring (lattice-ordered ring) if

- (i) (R, \leq) is a lattice;
- (ii) $s \leq t$ implies $s + r \leq t + r$ for each r ;
- (iii) $0 \leq s, t$ implies $0 \leq st$.

An ℓ -ring R is an f -ring if for each $r, s, t \in R$ with $s \wedge t = 0$ and $r \geq 0$, we have $rs \wedge t = 0$.

Definition 3.1. Let (S, \leq) be a partially ordered D -algebra.

- (1) We call S an ℓ -algebra over D if S is both an ℓ -ring and a D -algebra such that whenever $0 \leq s \in S$ and $0 \leq a \in D$, then $as \geq 0$.
- (2) We call S an f -algebra over D if S is both an ℓ -algebra over D and an f -ring.

Theorem 3.2. *Let S be a Specker D -algebra. Then there is a unique partial ordering \leq on S for which (S, \leq) is an f -algebra over D , given by $s \leq t$ if $t - s$ has an orthogonal decomposition whose coefficients are nonnegative. Moreover, \leq restricts to the usual order on $\text{Id}(S)$.*

Proof. Let P be the set of elements in S that have an orthogonal decomposition whose coefficients are nonnegative. We prove that $P \cap -P = \{0\}$ and P is closed under addition, multiplication, and multiplication by positive scalars. Let $s, t \in P$ and let $s = \sum_i a_i e_i$ and $t = \sum_j b_j f_j$ be orthogonal decompositions with $0 \leq a_i, b_j$ for each i, j . As in the proof of Theorem 2.5, we may write $s = \sum_{i,j} a_i (e_i \wedge f_j)$ and $t = \sum_{i,j} b_j (e_i \wedge f_j)$. Therefore, $s + t = \sum_{i,j} (a_i + b_j) (e_i \wedge f_j)$ and $st = \sum_{i,j} a_i b_j (e_i \wedge f_j)$, so $s + t, st \in P$. Moreover, it is clear that $as \in P$ for each $0 \leq a \in D$. To see that $P \cap -P = \{0\}$, suppose that $s = \sum_i a_i e_i = \sum_j -b_j f_j$ are orthogonal decompositions with each $a_i, b_j \geq 0$. Then $s = \sum_{i,j} a_i (e_i \wedge f_j) = \sum_{i,j} -b_j (e_i \wedge f_j)$, so $0 = \sum_{i,j} (a_i + b_j) (e_i \wedge f_j)$. Multiplying by $e_i \wedge f_j$ yields $(a_i + b_j) (e_i \wedge f_j) = 0$. Since S is a torsion free D -module, $a_i + b_j = 0$ or $e_i \wedge f_j = 0$. If $e_i \wedge f_j = 0$, then $a_i (e_i \wedge f_j) = 0$. Otherwise $a_i = 0 = b_j$ since both are nonnegative. In either case, $a_i (e_i \wedge f_j) = 0$ for each i, j , and so $s = 0$. Thus, if we set $s \leq t$ whenever $t - s \in P$, then \leq is a partial ordering on S and (S, \leq) satisfies conditions (ii) and (iii) of the definition of an ℓ -ring (see [14, Thm. VI.1.1]).

To see that (S, \leq) also satisfies (i), let $s, t \in S$ and let $s = \sum_i a_i e_i$ and $t = \sum_i b_i e_i$ be orthogonal decompositions of s and t with the same set of idempotents. The join and meet of s, t exist and are given by:

Claim 3.3. $s \vee t = \sum_i \max(a_i, b_i) e_i$ and $s \wedge t = \sum_i \min(a_i, b_i) e_i$.

Proof of Claim: The proofs of the two parts of the claim are similar, so we only prove the second. Set $r = \sum_i \min(a_i, b_i) e_i$. The definition of P shows that $r \leq s, t$. Next, let $q \in S$ be a lower bound of s, t . By refining the decompositions and eliminating zero idempotents if necessary, we may assume that $q = \sum_i d_i e_i$ for some $d_i \in D$ and that all $e_i \neq 0$. Since $q \leq s$, the e_i are pairwise orthogonal, and $e_i \geq 0$, we have that $d_i e_i = q e_i \leq s e_i = a_i e_i$, so $(a_i - d_i) e_i \in P$. If $a_i < d_i$, then $(d_i - a_i) e_i \in P$. Because $P \cap -P = \{0\}$, this forces $(a_i - d_i) e_i = 0$, so $a_i = d_i$ since S is torsion-free over D . This contradiction shows that $d_i \leq a_i$. Similarly, $d_i \leq b_i$, so $d_i \leq \min(a_i, b_i)$. Therefore, $q \leq r$. Thus, r is the greatest lower bound of s, t , and so $s \wedge t$ exists in S and is equal to $\sum_i \min(a_i, b_i) e_i$. \square

Consequently, S is an ℓ -ring. That $0 \leq s \in S$ and $0 \leq a \in D$ imply $as \geq 0$ is easy to see. Thus, S is an ℓ -algebra over D . To see that S is an f -algebra, let $s \wedge t = 0$ and $r \in S$ with $r \geq 0$. As above, s, t , and r have orthogonal decompositions $s = \sum_i a_i e_i$, $t = \sum_i b_i e_i$, and

$r = \sum_i c_i e_i$ with the same set of idempotents and $0 \leq a_i, b_i, c_i$, and we may assume without loss of generality that each $e_i \neq 0$. By the claim, $s \wedge t = \sum_i \min(a_i, b_i) e_i$. Since $s \wedge t = 0$, for each i , either $a_i = 0$ or $b_i = 0$. Because $sr \wedge t = \sum_i \min(a_i c_i, b_i) e_i$, we see that $sr \wedge t = 0$. Consequently, S is an f -algebra over D . This in particular implies that the order on S restricts to the usual order on $\text{Id}(S)$ (see [5, Lem 4.9(2)]). Finally, the proof of uniqueness of \leq is a direct adaptation of that given in [6, Thm. 5.1]. \square

It was proved in [6, Cor. 5.3] that each unital D -algebra homomorphism between Specker D -algebras is an ℓ -algebra homomorphism. The proof used [6, Thm. 5.1] and hence was not choice-free. By using the original argument from [6] but substituting the choice-free Theorem 3.2 for that of choice-dependent Theorem 5.1 from [6], we therefore obtain a choice-free proof of this result.

Theorem 3.4. *Each unital D -algebra homomorphism between Specker D -algebras is an ℓ -algebra homomorphism.*

As an immediate consequence of Theorem 3.4 we obtain:

Corollary 3.5. *Let S be a Specker D -algebra and $B = \text{Id}(S)$. The map $(-)^{\perp} : S \rightarrow D[B]^*$ of Theorem 2.5 is an ℓ -algebra isomorphism.*

Remark 3.6. Let D be a totally ordered domain and B a boolean algebra. Clearly D is a lattice, where $a \wedge b = \min(a, b)$ and $a \vee b = \max(a, b)$ for each $a, b \in D$.

(1) The positive cone P of $D[B]^*$ for the partial order \leq defined in Theorem 3.2 can be described by

$$f \in P \text{ iff } f(a) = 0 \text{ for each } a < 0.$$

To see this, if $f \in D[B]^*$, then by the comments before Theorem 2.5 applied to the map $(-)^{\perp} : D[B] \rightarrow D[B]^*$, we have $f = (\sum_{a \in D} a f(a))^{\perp}$. If $f(a) = 0$ for each $a < 0$, then the description of P in the proof of Theorem 3.2 shows that $f \in P$. Conversely, if $f \in P$, then there are $e_i \in B$ and $0 \leq a_i \in D$ with $f = \sum_{i=1}^n a_i e_i^{\perp}$. Let $a < 0$. Then

$$f(a) = \bigvee \{a_1 e_1^{\perp}(b_1) \wedge \cdots \wedge a_n e_n^{\perp}(b_n) : a_1 b_1 + \cdots + a_n b_n = a\}.$$

Because $a < 0$ and all $a_i \geq 0$, if $\sum_i a_i b_i = a$, then some $b_i < 0$, and so $e_i^{\perp}(b_i) = 0$. This shows that $f(a) = 0$. From this we see that

$$f \leq g \text{ iff } (g - f)(a) = 0 \text{ for each } a < 0.$$

(2) The meet and join in $D[B]^*$ are calculated by

$$\begin{aligned} (f \wedge g)(a) &= \bigvee \{f(b) \wedge g(c) : \min(b, c) = a\}, \\ (f \vee g)(a) &= \bigvee \{f(b) \wedge g(c) : \max(b, c) = a\}. \end{aligned}$$

We only prove the first equality as the second is proved similarly. Define $h : D \rightarrow B$ by $h(a) = \bigvee \{f(b) \wedge g(c) : \min(b, c) = a\}$. We show that $h = f \wedge g$. It is easy to see

that $h \in D[B]^*$. Therefore, $-h(c) = h(-c)$ for each $c \in D$ by the definition of scalar multiplication in $D[B]^*$. To see that $h \leq f, g$, let $a < 0$. Then

$$\begin{aligned} (f - h)(a) &= \bigvee \{f(b) \wedge (-h)(c) : b + c = a\} = \bigvee \{f(b) \wedge h(-c) : b + c = a\} \\ &= \bigvee \{f(b) \wedge \bigvee \{f(d) \wedge g(e) : \min(d, e) = -c\} : b + c = a\} \\ &= \bigvee \{f(b) \wedge f(d) \wedge g(e) : \min(d, e) = -c, b + c = a\}. \end{aligned}$$

If $b + c = a$, then $b < -c$ since $a < 0$. Therefore, if $\min(d, e) = -c$, then $b < d$. This implies that $f(b) \wedge f(d) = 0$ and hence $(f - h)(a) = 0$. Thus, $h \leq f$. Similarly, $h \leq g$, which gives $h \leq f \wedge g$.

To see the reverse inequality, suppose that $k \in D[B]^*$ with $k \leq f, g$. We show that $k \leq h$. It follows from (1) that $k \leq f$ implies $(f - k)(a) = 0$ for all $a < 0$. As we saw above,

$$(f - k)(a) = \bigvee \{f(b) \wedge k(-c) : b + c = a\}.$$

Therefore, $f(b) \wedge k(-c) = 0$ whenever $b + c < 0$. Similarly, $g(b) \wedge k(-c) = 0$ whenever $b + c < 0$. We have

$$\begin{aligned} (h - k)(a) &= \bigvee \{h(b) \wedge k(-c) : b + c = a\} \\ &= \bigvee \{\bigvee \{f(b_1) \wedge g(b_2) : \min(b_1, b_2) = b\} \wedge k(-c) : b + c = a\} \\ &= \bigvee \{f(b_1) \wedge g(b_2) \wedge k(-c) : \min(b_1, b_2) = b, b + c = a\}. \end{aligned}$$

If $\min(b_1, b_2) = b$ and $b + c = a < 0$, then either $b_1 + c < 0$ or $b_2 + c < 0$. Therefore, either $f(b_1) \wedge k(-c) = 0$ or $f(b_2) \wedge k(-c) = 0$. Consequently, if $\min(b_1, b_2) + c = a$, then $f(b_1) \wedge g(b_2) \wedge k(-c) = 0$. From this it follows that $(h - k)(a) = 0$ if $a < 0$, so $k \leq h$. In particular, $f \wedge g \leq h$. Thus, $h = f \wedge g$.

Remark 3.7. Let S be a Specker D -algebra and $s, t \in S$. It is not true in general that $s \leq t$ iff $s^\perp(a) \leq t^\perp(a)$ for all $a \in D$. For example, while $0 \leq 1$, we have $1 = 0^\perp(0) \not\leq 1^\perp(0) = 0$ because the full orthogonal decompositions of $0, 1$ are $0 = 0 \cdot 1$ and $1 = 1 \cdot 1$, respectively. This drawback will be corrected in Lemma 5.1(2) using a different way of viewing boolean powers of D , which we turn to next.

4. PROXIMITIES ON SPECKER ALGEBRAS AND DECREASING DECOMPOSITIONS

As we pointed out in the introduction, for a compact Hausdorff space X , there is a standard notion of proximity on the boolean algebra $\mathcal{RO}(X)$ of regular open subsets of X given by $U < V$ iff $\text{cl}(U) \subseteq V$. De Vries [9] axiomatized proximity relations on arbitrary boolean algebras. This has resulted in the notion of a de Vries proximity $<$ on a boolean algebra B . A *de Vries algebra* is a pair $(B, <)$, where B is a complete boolean algebra and $<$ is a de Vries proximity on B (see [3]).

In [5] de Vries proximities on boolean algebras were generalized to proximities on arbitrary torsion-free f -algebras over a totally ordered domain D .

Definition 4.1. [5, Def. 4.2] Let S be a torsion-free f -algebra over D . We call a binary relation \triangleleft on S a *proximity* if the following axioms are satisfied:

$$(P1) \quad 0 \triangleleft 0 \text{ and } 1 \triangleleft 1.$$

- (P2) $s \triangleleft t$ implies $s \leq t$.
- (P3) $s \leq t \triangleleft r \leq u$ implies $s \triangleleft u$.
- (P4) $s \triangleleft t, r$ implies $s \triangleleft t \wedge r$.
- (P5) $s \triangleleft t$ implies $-t \triangleleft -s$.
- (P6) $s \triangleleft t$ and $r \triangleleft u$ imply $s + r \triangleleft t + u$.
- (P7) $s \triangleleft t$ implies $as \triangleleft at$ for each $0 < a \in D$, and $as \triangleleft at$ for some $0 < a \in D$ implies $s \triangleleft t$.
- (P8) $s, t, r, u \geq 0$ with $s \triangleleft t$ and $r \triangleleft u$ imply $sr \triangleleft tu$.
- (P9) $s \triangleleft t$ implies there is $r \in S$ with $s \triangleleft r \triangleleft t$.
- (P10) $s > 0$ implies there is $0 < t \in S$ with $t \triangleleft s$.

We call a pair (S, \triangleleft) a *proximity D -algebra* if S is a torsion-free f -algebra over D and \triangleleft is a proximity on S . If S is a Specker D -algebra, then we call (S, \triangleleft) a *proximity Specker D -algebra*.

Remark 4.2.

- (1) The axioms (P1)–(P5) and (P9)–(P10) are direct analogues of the corresponding de Vries axioms, while the axioms (P6)–(P8) govern the interaction between the algebra operations and proximity on S .
- (2) Since every Specker D -algebra S is torsion-free, if S is nonzero, we always identify D with a subalgebra of S by sending $a \in D$ to $a \cdot 1 \in S$.
- (3) It is an easy consequence of the axioms that $s \triangleleft t$ and $r \triangleleft u$ imply $s \wedge r \triangleleft t \wedge u$ and $s \vee r \triangleleft t \vee u$, and that $s \triangleleft t$ iff $as \triangleleft bt$ for $0 < a \leq b \in D$. Hence, it follows from (P1), (P7), and (P5) that for each $a \in D$, we have $a \triangleleft a$.
- (4) The right-to-left implication in (P7) plays an important role in our considerations (see the proofs of Propositions 5.5 and 6.4). This implication is problematic if D is not a domain, so in Definition 4.1 it is essential that D is a domain.

Let S be a Specker D -algebra and $B = \text{Id}(S)$ the boolean algebra of idempotents of S . If \triangleleft is a proximity on S , we can consider its restriction to B . It was shown in [5] using orthogonal decompositions that the restriction of \triangleleft is a de Vries proximity on B . The proof in [5] is choice-free.

Proposition 4.3. [5, Prop. 5.1] *Let \triangleleft be a proximity on a Specker D -algebra S . Then \triangleleft restricts to a de Vries proximity on $\text{Id}(S)$.*

Our next goal is to prove the converse of Proposition 4.3, that a de Vries proximity on $\text{Id}(S)$ has a unique extension to a proximity on S . For this we need to work with decreasing decompositions instead of orthogonal decompositions. Decreasing decompositions, which are similar to Mundici's good sequences [17, p. 28], were studied for Specker algebras in [5, Sec. 5], where it was shown how to go back and forth between orthogonal and decreasing decompositions.

Remark 4.4. We will briefly describe how to go from an orthogonal decomposition to a decreasing decomposition since this will be used in Proposition 4.11. Let S be a Specker D -algebra, $s \in S$, and $s = \sum_{i=0}^n a_i f_i$ be an orthogonal decomposition of s with the $a_i \in D$ distinct and nonzero. Without loss of generality we may assume that $a_0 < \dots < a_n$. We can

then write

$$s = a_0(f_0 + \cdots + f_n) + (a_1 - a_0)(f_1 + \cdots + f_n) + \cdots + (a_n - a_{n-1})f_n.$$

Therefore, $s = \sum_{i=0}^n b_i e_i$, where $b_0 = a_0$, $b_i = a_i - a_{i-1}$ for $i \geq 1$, and $e_i = \sum_{j=0}^i f_j = \bigvee_{j=0}^i f_j$, where the second equality follows from [8, Eqn. XIII.3(14)]. This exhibits s as a linear combination of a sequence of strictly decreasing idempotents. Moreover, all the coefficients are nonzero and all of them except possibly b_0 are positive. Furthermore, if $s = \sum_{i=0}^n a_i f_i$ is a full orthogonal decomposition of s , then $e_0 = 1$. In this case we will write the corresponding decreasing decomposition as $s = a_0 + \sum_{i=1}^n b_i e_i$.

Definition 4.5. Let S be a Specker D -algebra and let $s \in S$.

- (1) We say that s is in *decreasing form* if $s = \sum_{i=0}^n b_i e_i$ with $e_0 \geq \cdots \geq e_n$, $b_0 \neq 0$ and $b_i > 0$ for $i \geq 1$.
- (2) We say that s is in *full decreasing form* if $s = a_0 + \sum_{i=1}^n b_i e_i$ is in decreasing form (with $e_0 = 1$).

Remark 4.6.

- (1) Because each element of S has a full orthogonal decomposition, each element has a full decreasing decomposition. Moreover, since a full orthogonal decomposition with distinct nonzero coefficients is unique, each $s \in S$ has a unique representation as $s = a_0 + \sum_{i=1}^n b_i e_i$ with each $b_i > 0$ and $1 = e_0 > e_1 > \cdots > e_n$.
- (2) As we saw in Section 2, to write two elements in compatible orthogonal form, we cannot assume coefficients are distinct. Similarly, we will see in Lemma 5.4(2) that two elements have a compatible decreasing decomposition, but we cannot assume that the idempotents are strictly decreasing. It is for this reason that the idempotents in Definition 4.5(1) are not assumed to be strictly decreasing.

Using decreasing decompositions, we give an alternative view of boolean powers of D .

Definition 4.7. Let B be a boolean algebra. We define $D[B]^b$ to be the set of all decreasing functions $f : D \rightarrow B$ for which there exist $1 = e_0 > e_1 > \cdots > e_n > 0$ in B and $a_0 < a_1 < \cdots < a_n$ in D such that

$$f(a) = \begin{cases} 1 & \text{if } a \leq a_0, \\ e_i & \text{if } a_{i-1} < a \leq a_i, \\ 0 & \text{if } a_n < a. \end{cases}$$

Let S be a Specker D -algebra and $B = \text{Id}(S)$. The following proposition illustrates that $D[B]^b$ encodes decreasing decompositions of elements of S into an algebra of functions from D to B .

Proposition 4.8. *Let S be a Specker D -algebra and $B = \text{Id}(S)$.*

- (1) *Let $s \in S$ be in full decreasing form $s = a_0 + \sum_{i=1}^n b_i e_i$ and set $a_i = a_0 + b_1 + \cdots + b_i$ for $1 \leq i \leq n$. Define $s^b : D \rightarrow B$ by*

$$s^b(a) = \begin{cases} 1 & \text{if } a \leq a_0, \\ e_i & \text{if } a_{i-1} < a \leq a_i, \\ 0 & \text{if } a_n < a. \end{cases}$$

Then $s^b \in D[B]^b$.

- (2) Conversely, for $f \in D[B]^b$, let the image of f in B be $\{1 = e_0 > e_1 > \dots > e_n > 0\}$, and for each $i \leq n$, let a_i be the largest element of $f^{-1}(e_i)$. Then $s = a_0 + \sum_{i=1}^n (a_i - a_{i-1})f(a_i)$ is an element of S in full decreasing form and $s^b = f$.

Proof. Straightforward. \square

Let S be a Specker D -algebra and $B = \text{Id}(S)$. The reader probably already anticipates that $D[B]^b$ is a D -algebra and that $(-)^b : S \rightarrow D[B]^b$ is a D -algebra isomorphism. This in particular implies that $D[B]^*$ and $D[B]^b$ provide two alternative representations of S , one that encodes orthogonal decompositions and the other that encodes decreasing decompositions. Consequently, $D[B]^b$ also provides an alternative way to view boolean powers of a totally ordered domain D .

We first prove that $D[B]^*$ is in bijective correspondence with $D[B]^b$, and describe the D -algebra structure of $D[B]^b$ induced by this bijection. From this we will then derive that $(-)^b : S \rightarrow D[B]^b$ is a D -algebra isomorphism.

Theorem 4.9. *For a boolean algebra B , there is a bijection between $D[B]^*$ and $D[B]^b$ that induces on $D[B]^b$ the structure of a Specker D -algebra whose operations satisfy, for all $f, g \in D[B]^b$ and $a, b \in D$,*

- (1) $(f + g)(a) = \bigvee \{f(b_1) \wedge g(b_2) : b_1 + b_2 \geq a\}$.
- (2) If $b > 0$, then $(bf)(a) = \bigvee \{f(c) : bc \geq a\}$.
- (3) If $f, g \geq 0$, then $(fg)(a) = \bigvee \{f(b_1) \wedge g(b_2) : b_1, b_2 \geq 0, b_1 b_2 \geq a\}$.

Proof. Define $\alpha : D[B]^* \rightarrow D[B]^b$ by $\alpha(f)(a) = \bigvee \{f(b) : b \geq a\}$ for each $f \in D[B]^*$ and $a \in D$. To see that α is well defined, let $f \in D[B]^*$, and let $\{a \in D : f(a) \neq 0\} = \{a_0 < \dots < a_n\}$. Set $e_i = f(a_i) \vee \dots \vee f(a_n)$ for $0 \leq i \leq n$. Then $1 = e_0 > e_1 > \dots > e_n > 0$, $\alpha(f)^{-1}(1) = (-\infty, a_0]$, and $\alpha(f)^{-1}(0) = (a_n, \infty)$. Moreover, if $a_{i-1} < a \leq a_i$, then $\alpha(f)(a) = f(a_i) \vee \dots \vee f(a_n) = e_i$, so that $\alpha(f)^{-1}(e_i) = (a_{i-1}, a_i]$. Thus, $\alpha(f) \in D[B]^b$, and α is well defined.

To see that α is onto, let $g \in D[B]^b$, and let $\{1 = e_0 > e_1 > \dots > e_n > 0\}$ be the image of g in B . For each i , let a_i be the largest element of $g^{-1}(e_i)$. Define $f : D \rightarrow B$ by $f(a_0) = 1$, $f(a_n) = e_n$, $f(a_i) = e_i \wedge \neg e_{i+1}$ for each $1 \leq i \leq n-1$, and $f(a) = 0$ for all $a \in D \setminus \{a_0, \dots, a_n\}$. Then $f \in D[B]^*$. We show that $\alpha(f) = g$. If $a \leq a_0$, then $\alpha(f)(a) = 1$ as it is the join of the $f(b)$ over all $b \geq a$, so $\alpha(f)(a) = g(a)$. If $a_0 < a \leq a_1$, then $\alpha(f)(a)$ is the join of $e_1 \wedge \neg e_2, e_2 \wedge \neg e_3, \dots, e_{n-1} \wedge \neg e_n, e_n$, which is $e_1 = g(a)$. Similarly, if $a_{i-1} < a \leq a_i$, then $\alpha(f)(a) = e_i = g(a)$, and if $a_n < a$, then $\alpha(f)(a) = 0 = g(a)$. Thus, $\alpha(f) = g$.

To see that α is 1-1, let $f, g \in D[B]^*$ with $\alpha(f) = \alpha(g)$. For $a \in D$ we have

$$\alpha(f)(a) = f(a) \vee \bigvee \{f(b) : b > a\}.$$

Since the values of f are pairwise orthogonal, $f(a) = \alpha(f)(a) \wedge \neg \bigvee \{f(b) : b > a\}$. However, $\bigvee \{f(b) : b > a\} = \bigvee \{\alpha(f)(b) : b > a\}$, so $f(a) = \alpha(f)(a) \wedge \neg \bigvee \{\alpha(f)(b) : b > a\}$. Similarly, $g(a) = \alpha(g)(a) \wedge \neg \bigvee \{\alpha(g)(b) : b > a\}$. Since $\alpha(f) = \alpha(g)$, we see that $f(a) = g(a)$. Thus, $f = g$.

Now since $\alpha : D[B]^* \rightarrow D[B]^b$ is a bijection, $D[B]^b$ inherits the structure of a Specker D -algebra from $D[B]^*$. Therefore, what remains to verify is that the algebraic structure

that $D[B]^b$ inherits from $D[B]^*$ satisfies (1)–(3) of the theorem. In light of the bijection with $D[B]^*$, it suffices to show that if $f, g \in D[B]^*$, then $\alpha(f), \alpha(g)$ both behave as stated in (1)–(3). Thus, we assume that $f, g \in D[B]^*$ and $a, b \in D$. Since the proofs are similar, we only prove (1).

Using the definition of $f + g$ in $D[B]^*$, we have:

$$\begin{aligned} (\alpha(f) + \alpha(g))(a) &= \bigvee_{b_1+b_2 \geq a} \alpha(f)(b_1) \wedge \alpha(g)(b_2) \\ &= \bigvee_{b_1+b_2 \geq a} \left(\bigvee_{c_1 \geq b_1} f(c_1) \right) \wedge \left(\bigvee_{c_2 \geq b_2} g(c_2) \right) = \bigvee_{b_1+b_2 \geq a} \left(\bigvee_{c_i \geq b_i} f(c_1) \wedge g(c_2) \right) \\ &= \bigvee_{c \geq a} \left(\bigvee_{c_1+c_2=c} f(c_1) \wedge g(c_2) \right) = \bigvee_{c \geq a} (f + g)(c) = \alpha(f + g)(a). \end{aligned}$$

□

Remark 4.10. By Remark 2.9, there is an isomorphism $B \rightarrow \text{Id}(D[B]^*)$ sending e to e^\perp for each $e \in B$. Since $\alpha : D[B]^* \rightarrow D[B]^b$ restricts to an isomorphism from $\text{Id}(D[B]^*)$ to $\text{Id}(D[B]^b)$, the composition τ_B is an isomorphism from B to $\text{Id}(D[B]^b)$. If $e^b = \tau_B(e)$, then it follows from the definition of α and the description of e^\perp that

$$e^b(a) = \begin{cases} 1 & \text{if } a \leq 0, \\ e & \text{if } 0 < a \leq 1, \\ 0 & \text{if } 1 < a. \end{cases}$$

In particular, $0^b(a) = 1$ if $a \leq 0$ and $0^b(a) = 0$ if $0 < a$. Similarly, $1^b(a) = 1$ if $a \leq 1$ and $1^b(a) = 0$ if $1 < a$. We note that 0^b and 1^b are then the 0 and 1 of $D[B]^b$, respectively.

Proposition 4.11. *Let S be a Specker D -algebra and $B = \text{Id}(S)$. The following diagram commutes.*

$$\begin{array}{ccc} & & D[B]^b \\ & \nearrow^{(-)^b} & \uparrow \alpha \\ S & & \\ & \searrow_{(-)^\perp} & D[B]^* \end{array}$$

Consequently, $(-)^b$ is an ℓ -algebra isomorphism.

Proof. We first show that $\alpha \circ (-)^\perp = (-)^b$. Let $s \in S$ and write $s = \sum_{i=0}^n a_i e_i$ in full orthogonal form with $a_0 < \dots < a_n$. Then the full decreasing form of s is $a_0 + (a_1 - a_0)f_1 + \dots + (a_n - a_{n-1})f_n$, where $f_i = e_i \vee \dots \vee e_n$ by Remark 4.4. Therefore, $s^b(a) = 1$ if $a \leq a_1$, $s^b(a) = f_i$ if $a_{i-1} < a \leq a_i$, and $s^b(a) = 0$ if $a_n < a$. On the other hand, $\alpha(s^\perp)(a) = \bigvee \{e_i : a \geq a_i\}$. Thus, $\alpha(s^\perp)(a) = 1$ if $a \leq a_0$, $\alpha(s^\perp)(a) = e_i \vee \dots \vee e_n = f_i$ if $a_{i-1} < a \leq a_i$, and $\alpha(s^\perp)(a) = 0$ if $a_n < a$. Consequently, $\alpha(s^\perp) = s^b$, and hence the diagram commutes.

To conclude the proof, it follows from Theorem 2.5 that $(-)^\perp : S \rightarrow D[B]^*$ is a D -algebra isomorphism, and it follows from Theorem 4.9 that $\alpha : D[B]^* \rightarrow D[B]^b$ is a D -algebra isomorphism. Therefore, $(-)^b$ is a D -algebra isomorphism, thus an ℓ -algebra isomorphism by Theorem 3.4. □

5. DE VRIES POWERS

In this section we use decreasing decompositions to lift de Vries proximities on boolean algebras to proximities on Specker D -algebras. In the particular case in which $(B, <)$ is a de Vries algebra, we lift $<$ to a proximity $<^b$ on $D[B]^b$ to obtain that $(D[B]^b, <^b)$ is a proximity Specker D -algebra, which in addition is a Baer ring (defined below). Following [5], we term such algebras proximity Baer-Specker D -algebras. The pair $(D[B]^b, <^b)$ provides a choice-free description of the de Vries power of D by $(B, <)$ that in [5, Def. 3.3] was defined in a choice-dependent way via the dual compact Hausdorff space of $(B, <)$.

We start by showing that the order on $D[B]^b$ is pointwise.

Lemma 5.1. *Let B be a boolean algebra. For $f, g \in D[B]^b$ we have:*

- (1) $(f \wedge g)(a) = f(a) \wedge g(a)$ for each $a \in D$.
- (2) $f \leq g$ iff $f(a) \leq g(a)$ for each $a \in D$.

Proof. (1) Since $\alpha : D[B]^* \rightarrow D[B]^b$ is a bijection, there are $s, t \in D[B]^*$ with $f = \alpha(s)$ and $g = \alpha(t)$. By Remark 3.6, $(s \wedge t)(a) = \bigvee \{s(b) \wedge t(c) : \min(b, c) = a\}$. Therefore,

$$\begin{aligned} f(a) \wedge g(a) &= \alpha(s)(a) \wedge \alpha(t)(a) = \left(\bigvee_{b_1 \geq a} s(b_1) \right) \wedge \left(\bigvee_{b_2 \geq a} t(b_2) \right) \\ &= \bigvee_{b_1, b_2 \geq a} s(b_1) \wedge t(b_2) = \bigvee_{b \geq a} \left(\bigvee_{\min(b_1, b_2) = b} s(b_1) \wedge t(b_2) \right) \\ &= \bigvee_{b \geq a} (s \wedge t)(b) = \alpha(s \wedge t)(a) = (f \wedge g)(a). \end{aligned}$$

- (2) We have $f \leq g$ iff $f = f \wedge g$. Therefore, (2) follows from (1). □

Let S be a Specker D -algebra and $B = \text{Id}(S)$. Since $(-)^b : S \rightarrow D[B]^b$ is an ℓ -algebra isomorphism by Proposition 4.11, the following is an immediate consequence of Lemma 5.1.

Lemma 5.2. *Let S be a Specker D -algebra. For $s, t \in S$ we have:*

- (1) $(s \wedge t)^b(a) = s^b(a) \wedge t^b(a)$ for each $a \in D$.
- (2) $s \leq t$ iff $s^b(a) \leq t^b(a)$ for each $a \in D$.

Remark 5.3. In contrast to Lemma 5.2(2), as we observed in Remark 3.6, it is not the case that $s \leq t$ iff $s^\perp(a) \leq t^\perp(a)$ for all $a \in D$.

The next technical lemma is needed in Proposition 5.5.

Lemma 5.4. *Let S be a Specker D -algebra.*

- (1) *Let $s \in S$ and let $a, b \in D$ with $a < b$. If $s^\perp(c) = 0$ for all c with $a < c < b$, then*

$$(s \wedge b) - (s \wedge a) = [(s - a) \wedge (b - a)] \vee 0 = (b - a)s^b(b).$$

- (2) *Let $s, t \in S$. Then there exist $a_0 < \dots < a_n$ in D with $a_0 \leq s, t \leq a_n$ such that s and t have compatible decreasing decompositions $s = a_0 + \sum_{i=1}^n (a_i - a_{i-1})s^b(a_i)$ and $t = a_0 + \sum_{i=1}^n (a_i - a_{i-1})t^b(a_i)$. Moreover, if $s, t \geq 0$, then we may assume $a_0 = 0$.*

Proof. (1) The proof that $(s \wedge b) - (s \wedge a) = [(s - a) \wedge (b - a)] \vee 0$ is given in [5, Claim 6.8]. We show that $(s \wedge b) - (s \wedge a) = (b - a)s^b(b)$. As discussed in Section 2, we may write $s = \sum_{b \in D} bs^\perp(b)$. By assumption, $\sum_{a < c < b} cs^\perp(c) = 0$, so

$$s = \sum_{c \leq a} cs^\perp(c) + \sum_{b \leq c} cs^\perp(c).$$

Because $\{s^\perp(c) : c \in D\}$ is a set of orthogonal idempotents whose join is 1 and $a, b \in D$, we have $a = \sum_{c \in D} as^\perp(c)$ and $b = \sum_{c \in D} bs^\perp(c)$. Therefore, by Claim 3.3,

$$s \wedge b = \sum_{c \leq a} \min(b, c)s^\perp(c) + \sum_{b \leq c} \min(b, c)s^\perp(c) = \sum_{c \leq a} cs^\perp(c) + \sum_{b \leq c} bs^\perp(c),$$

while

$$s \wedge a = \sum_{c \leq a} \min(a, c)s^\perp(c) + \sum_{b \leq c} \min(a, c)s^\perp(c) = \sum_{c \leq a} cs^\perp(c) + \sum_{b \leq c} as^\perp(c).$$

Thus, as the $s^\perp(c)$ are orthogonal,

$$(s \wedge b) - (s \wedge a) = (b - a) \sum_{b \leq c} s^\perp(c) = (b - a)s^b(b).$$

(2) We first show that for each $s \in S$ there is $0 \leq b \in D$ with $-b \leq s \leq b$. Write $s = \sum_{i=0}^n c_i e_i$ with $a_i \in D$ and $e_i \in \text{Id}(S)$. Let $b_i = \max(c_i, -c_i)$ and $b = \sum_{i=0}^n b_i$. Since $0 \leq e_i \leq 1$ for each i , we have $-b_i \leq c_i e_i \leq b_i$, so $-b \leq s \leq b$. Therefore, there are $a_0, a_n \in D$ with $a_0 \leq s, t \leq a_n$.

Since s^\perp, t^\perp have finitely many nonzero values, there are a_1, \dots, a_{n-1} in D such that $a_0 < a_1 < \dots < a_{n-1} < a_n$ and for each $a \notin \{a_0, \dots, a_n\}$, we have $s^\perp(a) = 0 = t^\perp(a)$. From $a_0 \leq s \leq a_n$ we get $(s \wedge a_n) - (s \wedge a_0) = s - a_0$. Thus, by (1),

$$s - a_0 = \sum_{i=1}^n ((s \wedge a_i) - (s \wedge a_{i-1})) = \sum_{i=1}^n (a_i - a_{i-1})s^b(a_i).$$

So $s = a_0 + \sum_{i=1}^n (a_i - a_{i-1})s^b(a_i)$, and a similar argument gives $t = a_0 + \sum_{i=1}^n (a_i - a_{i-1})t^b(a_i)$. Finally, if $s, t \geq 0$, then we may choose a_0 above to be 0. \square

As we showed in Lemma 5.2(2), if s and t are elements of a Specker D -algebra, then $s \leq t$ iff $s^b(a) \leq t^b(a)$ for all $a \in D$. We strengthen this in the next proposition and show that the analogous property holds for the proximity relation on a proximity Specker D -algebra. The desire to have such a simple functional interpretation of the proximity relation motivates our use of the $(-)^b$ representation of a Specker D -algebra in place of the $(-)^\perp$ representation.

Proposition 5.5. *Let (S, \triangleleft) be a proximity Specker D -algebra and let $s, t \in S$. Then $s \triangleleft t$ iff $s^b(b) \triangleleft t^b(b)$ for all $b \in D$.*

Proof. Let $s, t \in S$. We first show that $s \triangleleft t$ iff $[(s - a) \wedge b] \vee 0 \triangleleft [(t - a) \wedge b] \vee 0$ for all $a, b \in D$. First suppose that $s \triangleleft t$. By Remark 4.2(3), $a \triangleleft a$. Therefore, $s - a \triangleleft t - a$. Applying Remark 4.2(3) again, we first get $(s - a) \wedge b \triangleleft (t - a) \wedge b$, and then that $[(s - a) \wedge b] \vee 0 \triangleleft [(t - a) \wedge b] \vee 0$. Conversely, as S is bounded, there exist $a, b \in D$ with $a \leq s, t \leq a + b$. Therefore, $[(s - a) \wedge b] \vee 0 = s - a$ and $[(t - a) \wedge b] \vee 0 = t - a$. Thus, $s - a \triangleleft t - a$. Since $a \triangleleft a$, we conclude that $s \triangleleft t$.

Next, let $s \triangleleft t$ and $b \in D$. Choose $a < b$ so that if $a < c < b$, then $s^\perp(c) = 0 = t^\perp(c)$. By Lemma 5.4(1), $[(s - a) \wedge (b - a)] \vee 0 = (b - a)s^b(b)$ and $[(t - a) \wedge (b - a)] \vee 0 = (b - a)t^b(b)$.

Consequently, by the previous paragraph, we have $(b-a)s^b(b) \triangleleft (b-a)t^b(b)$. Since $b-a > 0$, it follows from (P7) that $s^b(b) \triangleleft t^b(b)$.

Conversely, suppose that $s^b(b) \triangleleft t^b(b)$ for all $b \in D$. By Lemma 5.4(2), we may write $s = a_0 + \sum_{i=1}^n (a_i - a_{i-1})s^b(a_i)$ and $t = a_0 + \sum_{i=1}^n (a_i - a_{i-1})t^b(a_i)$ for appropriate $a_0 < \dots < a_n$ in D . Since \triangleleft preserves addition and scalar multiplication by nonnegative scalars, from these representations we conclude that $s \triangleleft t$. \square

Let S be a Specker D -algebra and $<$ be a de Vries proximity on $\text{Id}(S)$. Proposition 5.5 suggests a way to lift $<$ to a proximity \triangleleft on S . We will show in Corollary 5.8 that the relation in the following definition is a proximity on S and that it is the unique proximity extending $<$.

Definition 5.6.

- (1) Let $<$ be a de Vries proximity on a boolean algebra B . Define $<^b$ on $D[B]^b$ by $f <^b g$ if $f(b) < g(b)$ for each $b \in D$.
- (2) Let S be a Specker D -algebra and let $<$ be a de Vries proximity on $\text{Id}(S)$. Define \triangleleft on S by $s \triangleleft t$ if $s^b <^b t^b$.

Theorem 5.7. *Let $<$ be a de Vries proximity on a boolean algebra B . Then $<^b$ is a proximity on $D[B]^b$, and is the unique proximity on $D[B]^b$ such that $e < f$ iff $e^b <^b f^b$ for each $e, f \in B$.*

Proof. The proofs of (P1)–(P4) are straightforward. Among the axioms (P5)–(P8), we only verify (P6) since the other axioms follow along similar lines.

(P6) Suppose that $s, t, r, u \in D[B]^b$ with $s <^b t$ and $r <^b u$. Let $a \in D$. By Theorem 4.9(1),

$$\begin{aligned} (s+r)(a) &= \bigvee_{b_1+b_2 \geq a} s(b_1) \wedge r(b_2) \\ (t+u)(a) &= \bigvee_{b_1+b_2 \geq a} t(b_1) \wedge u(b_2) \end{aligned}$$

Because $<$ preserves finite meets and joins (see Remark 4.2(3)), it follows that $s+r <^b t+u$.

(P9) Let $s <^b t$. By Lemma 5.4(2), there are $a_0 < \dots < a_n$ in D and decreasing $e_i, f_i \in B$ with $s(a) = e_i$ and $t(a) = f_i$ for $a_{i-1} < a \leq a_i$. From $s <^b t$ it follows that $e_i < f_i$ for each i . Since $<$ is a de Vries proximity, there is $g_i \in B$ with $e_i < g_i < f_i$ for each i . As the e_i and f_i decrease and $<$ preserves finite meets, without loss of generality we may assume that the g_i decrease. Define $r \in D[B]^b$ by $r(a) = g_i$ when $a_{i-1} < a \leq a_i$. Then $s(a) < r(a) < t(a)$ for each $a \in D$. Thus, $s <^b r <^b t$.

(P10) Let $0 < s$. By Lemma 5.4(2), there are $a_0 < \dots < a_n$ in D and $1 = f_0 > f_1 > \dots > f_n > 0$ in B with $s(a) = f_i$ for $a_{i-1} < a \leq a_i$. Since $s > 0$ we may assume that $a_0 = 0$. Since $<$ is a de Vries proximity, there is $0 < e \in B$ with $e < f_n$. Define $t \in D[B]^b$ by $t(a) = 1$ if $a \leq 0$, $t(a) = e$ if $0 < a \leq a_n$, and $t(a) = 0$ if $a_n < a$. Then $t(a) < s(a)$ for each $a \in D$, so $t <^b s$. Also, by Remark 4.10 and Lemma 5.2(2), $0 < t$.

Finally, for $e, f \in B$, it follows from Remark 4.10 and Proposition 5.5 that $e < f$ iff $e^b <^b f^b$, and that $<^b$ is the unique proximity on $D[B]^b$ satisfying this property. \square

Corollary 5.8. *Let S be a Specker D -algebra and let $<$ be a de Vries proximity on $\text{Id}(S)$. If \triangleleft is the extension of $<$ to S given in Definition 5.6(2), then \triangleleft is a proximity on S .*

Furthermore, \triangleleft is the unique extension of $<$ to a proximity on S . Consequently, there is a 1-1 correspondence between proximities on S and de Vries proximities on $\text{Id}(S)$.

Proof. Let S be a Specker D -algebra and let $<$ be a de Vries proximity on $\text{Id}(S)$. By Proposition 4.11, $(-)^b : S \rightarrow D[\text{Id}(S)]^b$ is an ℓ -algebra isomorphism. Moreover, for each $s, t \in S$, we have $s \triangleleft t$ iff $s^b <^b t^b$. Therefore, by Theorem 5.7, \triangleleft is the unique proximity on S extending $<$. \square

We recall (see, e.g., [15, Def. 7.45]) that a commutative ring R is a *Baer ring* if for each ideal I of R the annihilator $\{r \in R : rs = 0 \ \forall s \in I\}$ of I is a principal ideal generated by an idempotent. By [6, Cor. 4.4], a Specker D -algebra S is a Baer ring iff $\text{Id}(S)$ is a complete boolean algebra. Thus, if $(B, <)$ is a de Vries algebra, then $(D[B]^b, <^b)$ is a proximity Specker D -algebra and $D[B]^b$ is a Baer ring.

Definition 5.9. Let S be a Specker D -algebra. If S is a Baer ring, then we call S a *Baer-Specker D -algebra*. If in addition \triangleleft is a proximity on S , then we call (S, \triangleleft) a *proximity Baer-Specker D -algebra*.

We are ready to give a choice-free definition of de Vries powers of D .

Definition 5.10. Let D be a totally ordered domain and $(B, <)$ a de Vries algebra. The *de Vries power of D by $(B, <)$* is the proximity D -algebra $(D[B]^b, <^b)$.

The next theorem shows that de Vries powers of D are exactly the proximity Baer-Specker D -algebras. It was first proved in [5, Thm. 4.10, Cor. 5.6] using choice. Our proof here is choice-free.

Theorem 5.11.

- (1) If $(B, <)$ is a de Vries algebra, then $(D[B]^b, <^b)$ is a proximity Baer-Specker D -algebra.
- (2) If (S, \triangleleft) is a proximity Baer-Specker D -algebra and $B = \text{Id}(S)$, then $(-)^b : S \rightarrow D[B]^b$ is an ℓ -algebra isomorphism such that $s \triangleleft t$ iff $s^b <^b t^b$.

Proof. (1) Let $(B, <)$ be a de Vries algebra. By Theorem 4.9, $D[B]^b$ is a Specker D -algebra, and $B \cong \text{Id}(D[B]^b)$ by Remark 4.10. Since B is complete, $D[B]^b$ is a Baer-Specker D -algebra. Thus, the de Vries power $(D[B]^b, <^b)$ is a proximity Baer-Specker D -algebra by Theorem 5.7.

(2) Let (S, \triangleleft) be a proximity Baer-Specker algebra and $B = \text{Id}(S)$. The restriction $<$ of \triangleleft to B is a de Vries proximity by Proposition 4.3. Since S is Baer, B is a complete boolean algebra. Therefore, $(B, <)$ is a de Vries algebra, and so $(D[B]^b, <^b)$ is a de Vries power and $(-)^b : S \rightarrow D[B]^b$ is an ℓ -algebra isomorphism by Proposition 4.11. Moreover, $s \triangleleft t$ iff $s^b <^b t^b$ by Proposition 5.5. \square

6. DE VRIES ALGEBRAS AND PROXIMITY BAER-SPECKER ALGEBRAS

In this final section we extend the correspondence of Section 5 between de Vries proximities on boolean algebras and proximities on Specker D -algebras to a categorical equivalence between the category DeV of de Vries algebras and the category PbSp_D of proximity Baer Specker D -algebras. As we pointed out in the introduction, this equivalence follows from

de Vries duality between DeV and KHaus and the duality of [5] between PBSp_D and KHaus . However, the proof requires going through KHaus and hence is not choice-free. We give a purely algebraic choice-free proof of this equivalence.

Definition 6.1. [5, Def. 6.4] Let (S, \triangleleft) and (T, \triangleleft) be proximity Baer-Specker D -algebras. A *proximity morphism* is a map $\alpha : S \rightarrow T$ satisfying

- (M1) $\alpha(0) = 0$.
- (M2) $\alpha(s \wedge t) = \alpha(s) \wedge \alpha(t)$.
- (M3) $s \triangleleft t$ implies $-\alpha(-s) \triangleleft \alpha(t)$.
- (M4) $\alpha(t) = \bigvee \{\alpha(s) : s \triangleleft t\}$.
- (M5) $s \in S$ and $a \in D$ imply $\alpha(s + a) = \alpha(s) + a$.
- (M6) $s \in S$ and $0 \leq a \in D$ imply $\alpha(as) = a\alpha(s)$.
- (M7) $s \in S$ and $a \in D$ imply $\alpha(s \vee a) = \alpha(s) \vee a$.

Remark 6.2.

- (1) It is immediate from (M1) and (M5) that $\alpha(a) = a$ for each $a \in D$.
- (2) The reading of axiom (M4) should be that the least upper bound of $\{\alpha(s) : s \triangleleft t\}$ exists and is equal to $\alpha(t)$.
- (3) The axioms (M1)–(M4) are direct analogues of the corresponding axioms for de Vries morphisms, while the axioms (M5)–(M7) govern the behavior of proximity morphisms with respect to addition, multiplication, and join by a scalar.

It was proved in [5, Prop. 6.6] in a choice-dependent way that a proximity morphism between proximity Baer-Specker D -algebras restricts to a de Vries morphism between the de Vries algebras of idempotents. To give a choice-free proof, we require the following lemma, which gives a strictly order-theoretic characterization of idempotents in an f -ring.

Lemma 6.3. *Let A be an f -ring and $e \in A$. Then $e \in \text{Id}(A)$ iff $e = 2e \wedge 1$.*

Proof. Let $e \in A$. We have

$$e = 2e \wedge 1 \text{ iff } (2e \wedge 1) - e = 0 \text{ iff } e \wedge (1 - e) = 0.$$

Therefore, if $e = 2e \wedge 1$, then $0 \leq e, 1 - e \leq 1$, and hence $e(1 - e) = 0$ because $e(1 - e) \leq e \wedge (1 - e)$. Thus, $e^2 = e$. Conversely, let $e \in \text{Id}(A)$. Since A is an f -ring, the proof of [5, Lem. 4.9(2)] shows that the order on $\text{Id}(A)$ is the restriction of the order on A . Therefore,

$$(1 \wedge 2e) - e = (1 - e) \wedge e = 0.$$

Thus, $e = 2e \wedge 1$. □

To prove that each de Vries morphism lifts to a proximity morphism, we need the following lemma. A choice-dependent proof of Item (1) was given in [5, Prop. 6.6]. We give a choice-free proof of (1) which together with [5, Thm. 6.7] then yields a choice-free proof of Item (2).

Lemma 6.4. *Let (S, \triangleleft) and (T, \triangleleft) be proximity Baer-Specker D -algebras and let $\alpha : S \rightarrow T$ be a proximity morphism.*

- (1) $\alpha(\text{Id}(S)) \subseteq \text{Id}(T)$ and $\alpha|_{\text{Id}(S)}$ is a de Vries morphism from $\text{Id}(S)$ to $\text{Id}(T)$.
- (2) Let $s \in S$ and write $s = a_0 + \sum_{i=1}^n b_i e_i$ in decreasing form. Then $\alpha(s) = a_0 + \sum_{i=1}^n b_i \alpha(e_i)$.

Proof. (1) Let $e \in \text{Id}(S)$. By Lemma 6.3, $e = 2e \wedge 1$, so

$$\alpha(e) = \alpha(2e \wedge 1) = \alpha(2e) \wedge \alpha(1) = 2\alpha(e) \wedge 1.$$

Therefore, $\alpha(e) \in \text{Id}(T)$.

It follows that $\alpha|_{\text{Id}(S)} : \text{Id}(S) \rightarrow \text{Id}(T)$ is well defined. It is also clear that $\alpha|_{\text{Id}(S)}$ satisfies (M1) and (M2). Suppose that $e, f \in \text{Id}(S)$ with $e \triangleleft f$. Since $\neg e = 1 - e$, we have

$$\neg\alpha(\neg e) = 1 - \alpha(1 - e) = 1 - [1 + \alpha(\neg e)] = -\alpha(\neg e).$$

Because $\neg\alpha(\neg e) \triangleleft \alpha(f)$, we conclude that $\neg\alpha(\neg e) \triangleleft \alpha(f)$. Therefore, $e \triangleleft f$ implies that $\neg\alpha(\neg e) \triangleleft \alpha(f)$, which is the de Vries analogue of (M3). Let $f \in \text{Id}(S)$. Then $\alpha(f) = \bigvee\{\alpha(s) : s \in S, s \triangleleft f\}$. Suppose that $0 \leq s \triangleleft f$. As above, write $s = \sum_{i=1}^n b_i e_i$ in orthogonal form with each $b_i > 0$. Then $b_i e_i \leq s \triangleleft f$, so $b_i e_i \triangleleft f$ by (P3). It follows from the proof of [5, Prop. 5.1], which uses (P7), that $b_i \leq 1$ and $e_i \triangleleft f$ for each i . Consequently, $s \leq e_1 \vee \dots \vee e_n \triangleleft f$. Since $\alpha(s) \leq \alpha(e_1 \vee \dots \vee e_n)$ and $e_1 \vee \dots \vee e_n \in \text{Id}(S)$, we see that $\alpha(f) = \bigvee\{\alpha(e) : e \in \text{Id}(S), e \triangleleft f\}$. Thus, $\alpha|_{\text{Id}(S)}$ satisfies (M4).

(2) In the proof of [5, Thm. 6.7] substitute the choice-dependent proof of (1) with the choice-free proof above. \square

We recall the isomorphism $\tau_B : B \rightarrow \text{Id}(D[B]^b)$ from Remark 4.10, given by $\tau_B(e) = e^b$.

Theorem 6.5. *Let (A, \triangleleft) and (B, \triangleleft) be de Vries algebras and let $\sigma : A \rightarrow B$ be a de Vries morphism. Then there is a unique proximity morphism $\sigma^b : D[A]^b \rightarrow D[B]^b$ such that $\sigma^b \circ \tau_A = \tau_B \circ \sigma$.*

$$\begin{array}{ccc} A & \xrightarrow{\sigma} & B \\ \tau_A \downarrow & & \downarrow \tau_B \\ D[A]^b & \xrightarrow{\sigma^b} & D[B]^b \end{array}$$

Proof. Define $\sigma^b : D[A]^b \rightarrow D[B]^b$ by $\sigma^b(f) = \sigma \circ f$. It is easy to see that σ^b is well defined. Let $e \in A$ and consider the corresponding idempotent $\tau_A(e) = e^b \in D[A]^b$. It follows from Remark 4.10 that $\sigma \circ e^b = \sigma(e)^b$. Thus, $\sigma^b \circ \tau_A = \tau_B \circ \sigma$.

We now show that σ^b is a proximity morphism. Verifying (M1) and (M2) is straightforward, so we begin with (M3).

(M3) We first show that

$$(-f)(a) = \bigwedge\{\neg f(b) : b > -a\}. \quad (*)$$

for each $f \in D[A]^b$ and $a \in D$. There is $s \in D[A]^*$ with $f = \alpha(s)$. Since α is a D -algebra homomorphism, $\neg f = \alpha(\neg s)$. Therefore,

$$(-f)(a) = \bigvee\{(-s)(b) : b \geq a\} = \bigvee\{s(-b) : b \geq a\} = \bigvee\{s(c) : c \leq -a\}.$$

Since the values of s are pairwise orthogonal and $\bigvee\{s(c) : c \in D\} = 1$, we have

$$(-f)(a) = \neg \bigvee\{s(c) : c > -a\} = \bigwedge\{\neg s(c) : c > -a\}.$$

In addition,

$$\begin{aligned} \bigwedge\{\neg f(b) : b > -a\} &= \bigwedge\{\neg \bigvee\{s(c) : c \geq b\} : b > -a\} \\ &= \bigwedge\{\neg s(c) : c \geq b > -a\} = \bigwedge\{\neg s(c) : c > -a\}. \end{aligned}$$

This verifies Equation (*).

Now let $f, g \in D[A]^b$ with $f <^b g$ and let $a \in D$. By Equation (*) applied twice,

$$(-\sigma^b(-f))(a) = \bigwedge_{b>a} -\sigma^b(-f)(b) = \bigwedge_{b>a} -\sigma((-f)(b)) = \bigwedge_{b>a} -\sigma\left(\bigwedge_{c>b} \neg f(c)\right).$$

Thus, as σ preserves finite meets,

$$\begin{aligned} (-\sigma^b(-f))(a) &= \bigwedge_{b>a} -\sigma\left(\bigwedge_{c>b} \neg f(c)\right) \\ &= \bigwedge_{b>a} \neg \bigwedge_{c>b} \sigma(\neg f(c)) = \bigwedge_{b>a} \bigvee_{c>b} -\sigma(\neg f(c)) \\ &= \bigwedge_{a>d} \bigvee_{c>d} -\sigma(\neg f(c)). \end{aligned}$$

Since $f <^b g$, we have $f(c) < g(c)$ for each $c \in D$, and so $-\sigma(\neg f(c)) < \sigma(g(c))$. Hence, $\bigwedge_{a>d} \bigvee_{c>d} -\sigma(\neg f(c)) < \bigwedge_{a>d} \bigvee_{c>d} \sigma(g(c))$. We show that $\bigwedge_{a>d} \bigvee_{c>d} \sigma(g(c)) = \sigma(g(a))$. There are $a_0 < \dots < a_n$ in D and $1 = e_0 > \dots > e_n > 0$ in A with $g(a) = e_i$ if $a_{i-1} < a \leq a_i$, $g(a) = 1$ if $a \leq a_0$, and $g(a) = 0$ if $a > a_n$. Let $a \in D$. There is i with $a_{i-1} < a \leq a_i$. Take $d < a$. If $a_{i-1} \leq d$, then $\bigvee_{c>d} \sigma(g(c)) = \sigma(e_i) = \sigma(g(a))$ since e_i is the largest element of $\{g(c) : c > d\}$. On the other hand, if $d < a_{i-1}$, then $\bigvee_{c>d} \sigma(g(c)) = \sigma(e_{i-1}) \geq \sigma(e_i)$. Thus, $\bigwedge_{a>d} \bigvee_{c>d} \sigma(g(c)) = \sigma(e_i) = \sigma(g(a))$, as desired. Consequently, we have $(-\sigma^b(-f))(a) < \sigma^b(g)(a)$ for each $a \in D$, which yields $(-\sigma^b(-f)) <^b \sigma^b(g)$.

(M4) Let $g \in D[A]^b$. Clearly $\sigma^b(g)$ is an upper bound of $\{\sigma^b(f) : f <^b g\}$. To see that $\sigma^b(g)$ is the least upper bound, it is sufficient to show that

$$\sigma(g(a)) = \bigvee \{\sigma(f(a)) : f <^b g\} \text{ for each } a \in D. \quad (**)$$

Indeed, suppose Equation (**) holds and $h \in D[B]^b$ is an upper bound of $\{\sigma^b(f) : f <^b g\}$. Then, by Lemma 5.1(2), $h(a) \geq \sigma^b(f)(a) = \sigma(f(a))$ for each $a \in D$. Therefore, $h(a) \geq \sigma(g(a)) = \sigma^b(g)(a)$ for each $a \in D$, and so $h \geq \sigma^b(g)$, again by Lemma 5.1(2).

To prove Equation (**), there are $a_0 < \dots < a_n$ in D and $1 = f_0 > \dots > f_n > 0$ in A with $g(a) = f_i$ if $a_{i-1} < a \leq a_i$, $g(a) = 1$ if $a \leq a_0$, and $g(a) = 0$ if $a > a_n$. If $1 = e_0 \geq e_1 \geq \dots \geq e_n$ are elements of A , then there is $f \in D[A]^b$ satisfying $f(a) = 1$ if $a \leq a_0$, $f(a) = e_i$ if $a_{i-1} < a \leq a_i$ for each i , and $f(a) = 0$ if $a > a_n$. We call f the function associated to $e_0 \geq \dots \geq e_n$ (fixing $a_0 < \dots < a_n$). Note that $f <^b g$ iff $e_i < f_i$ for each i by the definition of $<^b$.

Let $a \in D$. If $a \leq a_0$, then $\sigma(g(a)) = \sigma(1) = 1$. Let $f \in D[A]^b$ be associated to $1 > 0 \geq 0 \geq \dots \geq 0$. Then $f <^b g$ and $f(a) = 1$, so $\sigma(f(a)) = 1$. Therefore, Equation (**) holds for $a \leq a_0$. Next, suppose that $a > a_n$. Then $\sigma(g(a)) = \sigma(0) = 0$. With the same f , we have $\sigma(f(a)) = 0$, and so Equation (**) also holds for $a > a_n$. Finally, let $a_{i-1} < a \leq a_i$. Since σ is a de Vries morphism, $\sigma(g(a)) = \bigvee \{e : e < g(a)\}$. Let $e \in A$ with $e < g(a)$ and let f be the function associated to $1 \geq \dots \geq 1 \geq e \geq 0 \geq \dots \geq 0$, where e is the i -th term in the sequence. Then $f <^b g$ and $f(a) = e$. Since we can produce $f <^b g$ for each e with $e < g(a)$, this shows that Equation (**) holds for a . Therefore, Equation (**) holds for all $a \in D$.

The proofs of (M5)–(M7) are similar to each other, so we only give the proof of (M6).

(M6) Let $f \in D[A]^b$ and $0 < b \in D$. Let $\{1 = e_0 > \dots > e_n > 0\}$ be the image of f in A . For each i , let a_i be the largest element of $f^{-1}(e_i)$. We claim that $(bf)(a) = e_i$ for

$ba_{i-1} < a \leq ba_i$. For suppose that $ba_{i-1} < a \leq ba_i$. By Theorem 4.9(2), $(bf)(a) = \bigvee_{bc \geq a} f(c)$. Since $bc \geq a > ba_{i-1}$, we have $b(c - a_{i-1}) = bc - ba_{i-1} > 0$. If $c \leq a_{i-1}$, then $c - a_{i-1} \leq 0$, so $b(c - a_{i-1}) \leq 0$, a contradiction. Therefore, $c > a_{i-1}$, and hence $f(c) \leq e_i$. Thus, by choosing $c = a_i$, we see that the join is $f(a_i) = e_i$, so that $(bf)(a) = e_i$ for $ba_{i-1} < a \leq ba_i$, as claimed. We see then that $\sigma^b(bf)(a) = \sigma((bf)(a)) = \sigma(e_i)$ if $ba_{i-1} < a \leq ba_i$. By the same reasoning, since $\{1 = \sigma(e_0) > \dots > \sigma(e_n) > 0\}$ is the image of $\sigma^b(f)$, we have $(b\sigma^b(f))(a) = \sigma(e_i)$ if $ba_{i-1} < a \leq ba_i$. Therefore, $\sigma^b(bf) = b\sigma^b(f)$.

Finally, to prove uniqueness, suppose $\gamma : D[A]^b \rightarrow D[B]^b$ is another proximity morphism with $\gamma \circ \tau_A = \tau_B \circ \sigma$. Let $f \in D[A]^b$. Then there are $a_0 < \dots < a_n$ in D and $1 = e_0 > \dots > e_n > 0$ in A with $f(a) = e_i$ when $a_{i-1} < a \leq a_i$. By Proposition 4.8(2), if $b_i = a_i - a_{i-1}$ for $1 \leq i \leq n$, then $f = a_0 + \sum_i b_i e_i^b$. By Lemma 6.4(2), $\gamma(f) = a_0 + \sum_i b_i \sigma(e_i)^b$. Therefore, $\gamma(f)(a) = \sigma(e_i)$ when $a_{i-1} < a \leq a_i$. Thus, $\gamma(f) = \sigma \circ f = \sigma^b(f)$. \square

For a Specker D -algebra S let $\eta_S : S \rightarrow D[\text{Id}(S)]^b$ be given by $\eta_S(s) = s^b$. By Theorem 4.11 η_S is an ℓ -algebra isomorphism.

Corollary 6.6. *If (S, \triangleleft) and (T, \triangleleft) are proximity Baer-Specker D -algebras and $\sigma : \text{Id}(S) \rightarrow \text{Id}(T)$ is a de Vries morphism, then there is a unique proximity morphism $\alpha : S \rightarrow T$ such that α extends σ and $\sigma^b \circ \eta_S = \eta_T \circ \sigma$.*

$$\begin{array}{ccc} S & \xrightarrow{\alpha} & T \\ \eta_S \downarrow & & \downarrow \eta_T \\ D[\text{Id}(S)]^b & \xrightarrow{\sigma^b} & D[\text{Id}(T)]^b \end{array}$$

Proof. By Theorem 6.5, there is a unique proximity morphism $\sigma^b : D[\text{Id}(S)]^b \rightarrow D[\text{Id}(T)]^b$ such that $\sigma^b \circ \tau_{\text{Id}(S)} = \tau_{\text{Id}(T)} \circ \sigma$.

$$\begin{array}{ccc} \text{Id}(S) & \xrightarrow{\sigma} & \text{Id}(T) \\ \left(\begin{array}{ccc} \downarrow & & \downarrow \\ S & \xrightarrow{\alpha} & T \\ \downarrow \eta_S & & \downarrow \eta_T \end{array} \right) & & \tau_{\text{Id}(T)} \\ D[\text{Id}(S)]^b & \xrightarrow{\sigma^b} & D[\text{Id}(T)]^b \end{array}$$

Set $\alpha = \eta_T^{-1} \circ \sigma^b \circ \eta_S$. Then α is a proximity morphism and $\sigma^b \circ \eta_S = \eta_T \circ \sigma$. Let $e \in \text{Id}(S)$. We have $\alpha(e) = \eta_T^{-1} \sigma^b \eta_S(e) = \eta_T^{-1} \eta_T \sigma(e) = \sigma(e)$, hence α extends σ . Finally, since σ^b is unique, so is α . \square

Let $\sigma_1 : (B_1, <) \rightarrow (B_2, <)$ and $\sigma_2 : (B_2, <) \rightarrow (B_3, <)$ be de Vries morphisms. We recall that the composition $\sigma_2 \star \sigma_1$ in DeV is defined by

$$(\sigma_2 \star \sigma_1)(e) = \bigvee \{ \sigma_2 \sigma_1(f) : f < e \}.$$

Theorem 6.7. *Proximity Baer-Specker D -algebras and proximity morphisms between them form a category PBSp_D , where the composition $\alpha_2 \star \alpha_1$ of two proximity morphisms $\alpha_1 : S_1 \rightarrow$*

S_2 and $\alpha_2 : S_2 \rightarrow S_3$ is the unique proximity morphism extending the de Vries morphism $\alpha_2|_{\text{Id}(S_2)} \star \alpha_1|_{\text{Id}(S_1)}$. It is given by

$$(\alpha_2 \star \alpha_1)(s) = \bigvee \{ \alpha_2 \alpha_1(t) : t \triangleleft s \}.$$

Proof. Let $\alpha_1 : S_1 \rightarrow S_2$ and $\alpha_2 : S_2 \rightarrow S_3$ be proximity morphisms. By Proposition 6.4, their restrictions to the idempotents are de Vries morphisms. Therefore, $\alpha_2|_{\text{Id}(S_2)} \star \alpha_1|_{\text{Id}(S_1)}$ is a de Vries morphism. By Corollary 6.6, $\alpha_2 \star \alpha_1$ is the unique proximity morphism extending $\alpha_2|_{\text{Id}(S_2)} \star \alpha_1|_{\text{Id}(S_1)}$. That \star is associative follows from Corollary 6.6 and the fact that de Vries composition is associative. Since identity morphisms are identity functions, it is then clear that PBSp_D forms a category.

It is left to show that $(\alpha_2 \star \alpha_1)(s) = \bigvee \{ \alpha_2 \alpha_1(t) : t \triangleleft s \}$. First, suppose that $t \triangleleft s$. By Lemma 5.4(2), write $s = a_0 + \sum_{i=1}^n b_i s^b(a_i)$ and $t = a_0 + \sum_{i=1}^n b_i t^b(b_i)$. Set $e_i = s^b(a_i)$ and $f_i = t^b(b_i)$. Then $f_i \triangleleft e_i$ for each i by Theorem 5.5. We have $\alpha_2 \alpha_1(t) = a_0 + \sum_{i=1}^n b_i \alpha_2 \alpha_1(f_i)$. Also, since $\alpha_2 \star \alpha_1$ is a proximity morphism, $(\alpha_2 \star \alpha_1)(s) = a_0 + \sum_i b_i (\alpha_2 \star \alpha_1)(e_i)$ by Lemma 6.4(2). As $\alpha_2|_{\text{Id}(S_2)} \star \alpha_1|_{\text{Id}(S_1)}$ is a de Vries morphism, $(\alpha_2 \star \alpha_1)(e_i) = \bigvee \{ \alpha_2 \alpha_1(e) : e \in \text{Id}(S_1), e \triangleleft e_i \}$, so $\alpha_2 \alpha_1(t) \leq (\alpha_2 \star \alpha_1)(s)$. Therefore, $(\alpha_2 \star \alpha_1)(s)$ is an upper bound of $\{ \alpha_2 \alpha_1(t) : t \triangleleft s \}$. To see that it is the least upper bound, let r be an upper bound of $\{ \alpha_2 \alpha_1(t) : t \triangleleft s \}$. Let $E_i = \{ e : e \triangleleft e_i \}$. By [8, Eqn. XIII.3(8)] and Lemma 6.4(2),

$$\begin{aligned} (\alpha_2 \star \alpha_1)(s) &= a_0 + \sum_{i=1}^n b_i (\alpha_2 \star \alpha_1)(e_i) = a_0 + \sum_{i=1}^n b_i \bigvee \{ \alpha_2 \alpha_1(e) : e \in E_i \} \\ &= \bigvee \{ a_0 + \sum_{i=1}^n b_i \alpha_2 \alpha_1(k_i) : k_i \in E_i, 1 \leq i \leq n \} = \bigvee \{ \alpha_2 \alpha_1(a_0 + \sum_{i=1}^n b_i k_i) : k_i \in E_i \}. \end{aligned}$$

Since $k_i \triangleleft e_i$ for each i , we have $a_0 + \sum_{i=1}^n b_i k_i \triangleleft s$ by (P6) and (P7). Therefore, $(\alpha_2 \star \alpha_1)(s) = \bigvee \{ \alpha_2 \alpha_1(a_0 + \sum_{i=1}^n b_i k_i) : k_i \in E_i \} \leq r$. Thus, $(\alpha_2 \star \alpha_1)(s) = \bigvee \{ \alpha_2 \alpha_1(t) : t \triangleleft s \}$. \square

Although proximity morphisms are not in general D -algebra homomorphisms, as was shown in [5, Lem. 8.3] with a choice-free proof, proximity isomorphisms are D -algebra isomorphisms that preserve and reflect proximity. This is similar to what happens in DeV [9, Prop. I.5.5].

Lemma 6.8. [5, Lem. 8.3] *Let $(S, \triangleleft), (T, \triangleleft) \in \text{PBSp}_D$ and let $\alpha : S \rightarrow T$ be a proximity morphism. Then α is an isomorphism in PBSp_D iff α is a D -algebra isomorphism such that $s \triangleleft t$ in (S, \triangleleft) iff $\alpha(s) \triangleleft \alpha(t)$ in (T, \triangleleft) .*

We are finally ready to give a choice-free proof that PBSp_D is equivalent to DeV .

Theorem 6.9 (Main Theorem). *The category PBSp_D of proximity Baer-Specker D -algebras is equivalent to the category DeV of de Vries algebras.*

Proof. Define a covariant functor $\text{Id} : \text{PBSp}_D \rightarrow \text{DeV}$ by sending $(S, \triangleleft) \in \text{PBSp}_D$ to the de Vries algebra $(\text{Id}(S), \triangleleft|_{\text{Id}(S)})$ and a proximity morphism $\alpha : S \rightarrow T$ to the de Vries morphism $\alpha|_{\text{Id}(S)}$. It follows from Proposition 4.3, Lemma 6.4, and the definition of compositions in PBSp_D and DeV that Id is well defined.

Define a covariant functor $\text{Sp} : \text{DeV} \rightarrow \text{PBSp}_D$ by sending $(B, \triangleleft) \in \text{DeV}$ to the de Vries power D -algebra $(D[B]^b, \triangleleft^b)$ and a de Vries morphism $\sigma : A \rightarrow B$ to the proximity morphism

σ^b . It follows from Theorem 5.11, Theorem 6.5, and the definition of compositions in \mathbf{PBSp}_D and \mathbf{DeV} that \mathbf{Sp} is well defined.

To show that \mathbf{Id} and \mathbf{Sp} form an equivalence, we first show that $\eta : 1_{\mathbf{PBSp}_D} \rightarrow \mathbf{Sp} \circ \mathbf{Id}$ is a natural isomorphism. Let $\alpha : (S_1, \triangleleft) \rightarrow (S_2, \triangleleft)$ be a proximity morphism. Set $B_i = \mathbf{Id}(S_i)$ and $\sigma = \alpha|_{B_1}$. Then $\sigma^b = \mathbf{Sp}(\mathbf{Id}(\alpha))$, and we have the following diagram,

$$\begin{array}{ccc} S_1 & \xrightarrow{\alpha} & S_2 \\ \eta_{S_1} \downarrow & & \downarrow \eta_{S_2} \\ D[B_1]^b & \xrightarrow{\mathbf{Sp}(\mathbf{Id}(\alpha))} & D[B_2]^b \end{array}$$

which commutes by Corollary 6.6. Thus, η is a natural transformation, and it is then a natural isomorphism by Proposition 4.11.

We next show that $\tau : 1_{\mathbf{DeV}} \rightarrow \mathbf{Id} \circ \mathbf{Sp}$ is a natural isomorphism. Let $\sigma : (B_1, <) \rightarrow (B_2, <)$ be a de Vries morphism. We have the following diagram,

$$\begin{array}{ccc} B_1 & \xrightarrow{\sigma} & B_2 \\ \tau_{B_1} \downarrow & & \downarrow \tau_{B_2} \\ \mathbf{Id}(D[B_1]^b) & \xrightarrow{\mathbf{Id}(\mathbf{Sp}(\sigma))} & \mathbf{Id}(D[B_2]^b) \end{array}$$

which commutes by Theorem 6.5. Thus, τ is a natural transformation, and it is then a natural isomorphism by Remark 4.10. Consequently, \mathbf{Sp} and \mathbf{Id} establish an equivalence of \mathbf{PBSp}_D and \mathbf{DeV} . \square

Remark 6.10. Let A be an algebra of a fixed type. Generalizing [14, p. 5], we say that a binary relation R on A is *compatible* with the operations of A if for each n -ary operation λ on A there is a subalgebra B of A such that from $a_1 R b_1, \dots, a_n R b_n$ it follows that $\lambda(a_1, \dots, a_n) R \lambda(b_1, \dots, b_n)$ or $\lambda(b_1, \dots, b_n) R \lambda(a_1, \dots, a_n)$ for each $a_1, \dots, a_n, b_1, \dots, b_n \in B$. Such a pair (A, R) is a particular case of an algebraic system of Malcev [16]. Let B be a boolean algebra and let r be a binary relation on B . We let $A[B]^*$ be the boolean power of A , as defined by Foster and discussed in Section 2. Define a relation \mathcal{R} on $A[B]^*$ by

$$f \mathcal{R} g \text{ iff } \bigvee \{f(b) : a R b\} r \bigvee \{g(b) : a R b\} \text{ for all } a \in A.$$

Then \mathcal{R} lifts r and R to the boolean power $A[B]^*$. If A is a totally ordered domain, R is \leq and r is $<$, then this generalization of a boolean power is exactly our de Vries power. It would be interesting to study in more detail this generalization of boolean powers when additional relations are also at play. Of course, the binary relations R and r can further be generalized to arbitrary relations. A particular case of such a generalization, when R is present but r is not, is briefly discussed by Banaschewski and Nelson [1, Concluding Remarks].

Conflict of interest. The authors declare that they have no conflict of interest.

REFERENCES

1. B. Banaschewski and E. Nelson, *Boolean powers as algebras of continuous functions*, Dissertationes Math. (Rozprawy Mat.) **179** (1980), 51.
2. G. M. Bergman, *Boolean rings of projection maps*, J. London Math. Soc. **4** (1972), 593–598.

3. G. Bezhanishvili, *Stone duality and Gleason covers through de Vries duality*, *Topology Appl.* **157** (2010), no. 6, 1064–1080.
4. G. Bezhanishvili, L. Carai, and P. J. Morandi, *A point-free approach to canonical extensions of boolean algebras and bounded archimedean ℓ -algebras*, submitted, 2021.
5. G. Bezhanishvili, V. Marra, P. J. Morandi, and B. Olberding, *De Vries powers: a generalization of Boolean powers for compact Hausdorff spaces*, *J. Pure Appl. Algebra* **219** (2015), no. 9, 3958–3991.
6. ———, *Idempotent generated algebras and Boolean powers of commutative rings*, *Algebra Universalis* **73** (2015), 183–204.
7. G. Bezhanishvili, P. J. Morandi, and B. Olberding, *Specker algebras: A survey*, *Algebraic Techniques and Their Use in Describing and Processing Uncertainty. To the Memory of Professor Elbert A. Walker. Hung Nguyen and Vladik Kreinovich, editors, Springer, 2020*, pp. 1–19.
8. G. Birkhoff, *Lattice theory*, third ed., American Mathematical Society Colloquium Publications, vol. 25, American Mathematical Society, Providence, R.I., 1979.
9. H. de Vries, *Compact spaces and compactifications. An algebraic approach*, Ph.D. thesis, University of Amsterdam, 1962.
10. R. P. Dilworth, *The normal completion of the lattice of continuous functions*, *Trans. Amer. Math. Soc.* **68** (1950), 427–438.
11. A. L. Foster, *Generalized “Boolean” theory of universal algebras. I. Subdirect sums and normal representation theorem*, *Math. Z.* **58** (1953), 306–336.
12. ———, *Generalized “Boolean” theory of universal algebras. II. Identities and subdirect sums of functionally complete algebras*, *Math. Z.* **59** (1953), 191–199.
13. ———, *Functional completeness in the small. Algebraic structure theorems and identities*, *Math. Ann.* **143** (1961), 29–58.
14. L. Fuchs, *Partially ordered algebraic systems*, Pergamon Press, Oxford, 1963.
15. T. Y. Lam, *Lectures on modules and rings*, Graduate Texts in Mathematics, vol. 189, Springer-Verlag, New York, 1999.
16. A. I. Malcev, *Algebraic systems*, Springer-Verlag, New York, 1973, Posthumous edition, edited by D. Smirnov and M. Tačlin, Translated from the Russian by B. D. Seckler and A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 192.
17. D. Mundici, *Interpretation of AF C^* -algebras in Łukasiewicz sentential calculus*, *J. Funct. Anal.* **65** (1986), no. 1, 15–63.
18. G.-C. Rota, *The valuation ring of a distributive lattice*, Proceedings of the University of Houston Lattice Theory Conference (Houston, Tex., 1973), Dept. Math., Univ. Houston, Houston, Tex., 1973, pp. 574–628.

NEW MEXICO STATE UNIVERSITY

Email address: guram@nmsu.edu

UNIVERSITÀ DEGLI STUDI DI SALERNO

Email address: lcarai@unisa.it

NEW MEXICO STATE UNIVERSITY

Email address: pmorandi@nmsu.edu

NEW MEXICO STATE UNIVERSITY

Email address: bruce@nmsu.edu