

1 **Polystyrene microplastics exposure modulated the content and**
2 **the profile of fatty acids in the Cladoceran *Daphnia magna***

3

4 Marco Parolini^{a*}, Beatrice De Felice^a, André Gois^b, Marisa Faria^{b,c}, Nereida
5 Cordeiro^{b,c}, Natacha Nogueira^{c,d}

6

7 ^a Department of Environmental Science and Policy, University of Milan, via Celoria 26, I-20133
8 Milan, Italy

9 ^b LB3, Faculty of Exact Science and Engineering, University of Madeira, 9020-105 Funchal, Portugal

10 ^c CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto,
11 4450-208 Porto, Portugal

12 ^d Mariculture Center of Calheta (CMC), Regional Directorate for the Sea, Av. D. Manuel I 7, 9370-
13 135 Calheta, Madeira (Portugal)

14

15

16

17 * Correspondence should be addressed to Marco Parolini (marco.parolini@unimi.it)

18

19 **Abstract**

20 A growing number of studies has shown that the exposure to microplastics (MPs) of different
21 polymeric compositions can induce diverse adverse effects towards several aquatic species. The vast
22 majority of such studies has been focused on the effects induced by the administration of MPs made
23 by polystyrene (PS; hereafter PS-MPs). However, despite the increase in the knowledge on the
24 potential toxicity of PS-MPs, there is a dearth of information concerning their role in affecting energy
25 resources and/or their allocation. The present study aimed at exploring the impact of 21-days exposure
26 to three concentrations (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of PS-MPs of different sizes (1 and 10 μm) on
27 fatty acids (FAs) profile of the freshwater Cladoceran *Daphnia magna*. The exposure to the highest
28 tested concentration of PS-MPs induced an overall decrease in *D. magna* total FAs content,
29 independently of the particle size. Moreover, a change in the accumulation of essential FAs by the
30 diet was noted, with an enhanced synthesis of monounsaturated FAs-rich storage lipids. However, a
31 sort of adaptation to counteract the adverse effects and to re-establish the FAs homeostasis was
32 observed in individuals treated with high PS-MPs concentration, independently of their size. These
33 results indicate that the exposure to PS-MPs could alter the allocation or induce changes in FAs
34 composition in *D. magna*, with potential long-term consequences on life-history traits of this
35 zooplanktonic species.

36

37 **Keywords:** essential fatty acids; freshwaters; microplastics; polystyrene

38

39 **1. Introduction**

40 Plastic contamination represents one of the main environmental issues that our society must
41 tackle. Despite the undeniable benefits of plastics (Andrady and Neal, 2009), the improper disposal,
42 and management, coupled with the persistence of plastic waste result in the accumulation of plastics
43 in terrestrial and aquatic ecosystems. Once in the environment, plastics undergo weathering processes
44 due to mechanical erosion, physical abrasion, solar radiation and biological degradation (De Sá et al.,
45 2018) leading to their breakage and fragmentation in small-sized items. Microplastics (MPs) have
46 been recently categorised as any plastic item in the 1 to < 1,000 µm size range (Hartmann et al.,
47 2019). They are considered as a hot topic in environmental studies because of their presence in every
48 ecosystem worldwide, as well as their potential toxicity towards organisms. A growing number of
49 monitoring surveys has detected MPs in the atmosphere (Can-Güven, 2021), marine (Horton and
50 Barnes, 2020), freshwater (Bellasi et al., 2020) and terrestrial (Rillig and Lehmann, 2020) ecosystems,
51 as well as in the so-called remote areas, such as the deep sea, Arctic and Antarctica (Shahul Hamid
52 et al., 2018; González-Pleiter et al., 2021), and high-mountain areas (Ambrosini et al., 2019;
53 Bergmann et al., 2019; Parolini et al., 2021; Crosta et al., 2022).

54 Freshwaters ecosystems have been identified as the primary source of plastic contamination for
55 seas and oceans (Wagner et al., 2014; Eerkes-Medrano et al., 2015). In recent years there has been
56 growing interest in the assessment of MPs contamination in rivers and lakes, and their impact on
57 freshwater organisms (Wagner and Lambert, 2018). The MPs contamination has been estimated as
58 high as 0.001 - 0.1 items/m² in lake and 0.1 - 1 items/m² in river water, while 10 - 10,000 items/m²
59 and 1 - 1,000 items/m² range has been estimated for lake and river sediments, respectively (Dris et
60 al., 2015). These estimates have been confirmed by recent field studies (Li et al., 2018; 2020).
61 Depending on their physical and chemical features, a wide array of MPs with different environmental
62 fate contributes to freshwater contamination. The pattern is dominated by polyethylene (PE),

63 polypropylene (PP), polystyrene (PS) and polyethylene terephthalate (PET), according to the demand
64 and use of plastics (PlasticsEurope, 2020).

65 In addition to surveys on the presence and the abundance of MPs in freshwaters, a growing
66 number of investigations has focused on their impact on organisms. All the studies agreed that MPs
67 of different size, shape and polymeric composition are efficiently ingested by many aquatic organisms
68 at different trophic levels and stages of development (*e.g.*, Cole et al., 2013; Imhof and Laforsch,
69 2016; Scherer et al., 2017; Canniff and Hoang, 2018; Al-Jaibachi et al., 2018; Al-Jaibachi and
70 Callaghan, 2018; An et al., 2021). However, the effects induced by the ingestion of diverse MPs by
71 freshwater and marine invertebrates resulted as contrasting and depended on different factors,
72 including the size, the shape, the polymer and the administered concentrations of MPs. For instance,
73 some experimental studies has demonstrated that the exposure to MPs reduced food uptake and
74 energy-supply related changes as a consequence of a false sense of satiation in the marine copepod
75 *Calanus helgolandicus* (75 microplastics mL⁻¹ of 20 µm PS beads; Cole et al. 2015), in the shore crab
76 *Carcinus maenas* (9.4 × 10⁵ microspheres L⁻¹ of 10 µm PS microspheres; Watts et al. 2014) and in
77 the freshwater amphipod *Gammarus fossarum* (range 100 - 13,380 fibres cm⁻² base area of glass
78 beakers of 500 × 20 µm polyamide fibres; Blarer and Burkhardt-Holm, 2016), the onset of oxidative
79 stress in in larvae of the sea urchin *Pseudechinus huttoni* (range 10 - 10,000 microspheres mL⁻¹ of 1–
80 5 µm of unknown polymer; Richardson et al., 2021) and the decrease in growth and reproduction rate
81 in the oyster *Crassostrea gigas* (0.023 mg L⁻¹ of 2 and 6 µm PS microspheres; Sussarellu et al., 2016)
82 and growth and development of veliger of the bivalve *Crepidula onyx* (6 × 10⁴ particles mL⁻¹ and 1.4
83 × 10⁵ particles mL⁻¹ of 2.0–2.4 µm PS microspheres; Lo and Chan, 2018). In contrast, other
84 investigations have reported slight or null effects on growth and survival of larvae of the sea urchin
85 *Tripneustes gratilla* (1 - 300 spheres mL⁻¹ range of 10-45 µm of polyethylene (PE) microspheres;
86 Kaposi et al., 2014), survival, morphological traits and reproductive parameters in the freshwater
87 cladoceran *Daphnia magna* (290 - 580 particles mL⁻¹ of two mixtures of ~40 µm PA, polycarbonate,
88 polyethylene terephthalate (PET) and polyvinylchloride and acrylonitrile-butadiene-styrene

89 terpolymer, plasticized polyvinyl chloride, polyoxymethylene homopolymer and styrene-
90 acrylonitrile copolymer MPs; Imhof et al., 2017), and on the survival, development, metabolism and
91 feeding activity of the freshwater amphipod *Gammarus pulex* (0.8 – 4,000 particles mL⁻¹ of 0 – 150
92 µm PET items; Weber et al. 2018).

93 Zooplanktonic filter-feeder species indiscriminately ingest MPs during swimming and feeding
94 activity (Gorokhova, 2015) resulted as particularly prone to the effects of these contaminants. Several
95 studies have demonstrated that the presence of MPs in the digestive tract of different zooplanktonic
96 species can result in a series of negative effects, including changes in physiological (e.g., movement,
97 growth, feeding, survival, gene expression), systemic (i.e., digestive, reproductive and
98 neuromodulation systems) and reproductive (i.e., amount and size of offspring) endpoints (see He et
99 al., 2021 and references therein). Focusing on freshwater zooplanktonic species, studies of *Daphnia*
100 *magna* have demonstrated that the ingestion of MPs can negatively affect food availability (Al-
101 Jaibachi and Callaghan, 2018), oxidative stress response (Liu et al., 2022), growth, swimming activity
102 and reproduction (Jemec et al., 2016; Ogonowski et al., 2016), with different outcomes depending on
103 the polymer composition, size and shape (Rosenkranz et al., 2009; Jemec et al., 2016; Frydkjær et al.,
104 2017; Na et al., 2021; Song et al., 2021a,b). However, our previous study has demonstrated that the
105 21-days exposure to increasing concentrations (0.125, 1.25 and 12.5 µg mL⁻¹) of polystyrene
106 microplastics (PS-MPs) of different size (1 and 10 µm in diameter) induced ‘putatively positive’
107 effects (De Felice et al., 2019). Indeed, independently of MPs size, individuals exposed to the highest
108 tested concentration grew more, swam longer distances and faster, and generated more offspring than
109 conspecifics from the control group, suggesting that PS-MPs caused a modulation of the energy
110 reserves of cladocerans (De Felice et al., 2019).

111 To check for this hypothesis, the present study aimed at replicating the experimental design
112 used by De Felice et al. (2019) to explore if the 21-days exposure to three concentrations (0.125, 1.25
113 and 12.5 µg mL⁻¹) of differently sized PS-MPs (1 and 10 µm in diameter) could affect the energy
114 reserves of *D. magna* individuals in terms of modulation of the fatty acids (FAs) profile and content.

115 In fact, FAs are critical for the permeability and generation of the cell membrane, function as essential
116 nutrients and energy reserves in the metabolic systems at all trophic levels (Lee et al. 2018; Neves et
117 al. 2015) and are crucial for growth and reproduction (Arts and Kohler, 2009). Moreover, FAs are
118 considered as markers of environmental stressors because when stressors emerge, FAs metabolism
119 can be regulated through cellular physiological and biochemical responses (Gonçalves et al., 2016;
120 Yang et al., 2021).

121 **2. Materials and Methods**

122 ***2.1 Exposure to polystyrene microplastics (PS-MPs)***

123 *Daphnia magna* individuals were reared in the facility located at the University of Milan,
124 (Italy) according to the procedure described by Parolini et al. (2018) and De Felice et al. (2019).
125 Individuals were reared in a commercial mineral water (San Benedetto®). Forty individuals/L were
126 maintained in glass beakers at 20.0 ± 0.5 °C under a natural photoperiod (16 h light: 8 h dark) to
127 ensure continuative parthenogenic reproduction. In the husbandry individuals were fed *ad libitum*
128 with a suspension of the unicellular green alga *Raphidocelis subcapitata* (8×10^6 cells individuals⁻¹
129 day⁻¹ until they were 8-days old, then 16×10^6 cells individuals⁻¹ day⁻¹ up to the end of the experiment)
130 and the yeast *Saccharomyces cerevisiae* (15×10^6 cells mL⁻¹). The culture medium was renewed
131 every second day.

132 In the present study, the experimental design performed by De Felice et al. (2019) was faithfully
133 replicated. Red PS-MPs with two different sizes (1 and 10 µm of diameter) were purchased from
134 Sigma-Aldrich (product number 89904 and 72986 for particles of 1 and 10 µm of diameter,
135 respectively; Milan, Italy). Chemical-physical properties of PS-MPs were provided by the supplier
136 (1 µm nominal diameter - calibrated particle diameter = 1.07 ± 0.03 µm, density 1.51 g cm⁻³; 10 µm
137 nominal diameter - calibrated particle diameter = 9.86 ± 0.13 µm, density = 1.51 g cm⁻³). The size
138 and polymeric composition of MPs used in the experiments were confirmed through Scanning

139 Electron Microscopy and Fourier Transformed Infrared Spectroscopy (FT-IR) analyses, respectively,
140 according to De Felice et al. (2019). A stock solution for each MPs type was prepared through a
141 1:1,000 (v/v) dilution of the commercial standard in the mineral water used to prepare the culture
142 medium used. Briefly, *D. magna* individuals were independently exposed to three increasing,
143 unrealistic concentrations of 1 μm and 10 μm PS-MPs (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$). The
144 concentrations were the same tested in our previous study and in other experiments performed on
145 marine and freshwater organisms (e.g., Messinetti et al., 2018; De Felice et al., 2018). Three replicates
146 per every single experimental condition were performed, for a total of 60 individuals per treatment.
147 In each replicate, twenty daphnids (< 24 hours old) were exposed in 500 mL beakers filled with
148 commercial mineral water (San Benedetto[®]) and maintained at 20.0 ± 0.5 °C under 16 h light: 8 h
149 dark photoperiod. Experiments were performed under semi-static conditions for 21 days, similarly to
150 the *Daphnia magna* reproduction test (OECD test N° 211), with a daily renewal of the exposure
151 medium (*i.e.*, water, feed and PS-MPs). During the duration of the experiments, the individuals were
152 fed *ad libitum* as described for husbandry. At the end of the experiments, all live individuals were
153 collected from each beaker, quickly frozen in liquid nitrogen and stored at -80 °C. *D. magna*
154 individuals were then lyophilised in a freeze-dryer (Edwards Pirani 1001) for 36 h and stored in a
155 desiccator until the FAs analysis. Three pools of twenty individuals each were collected per
156 experimental group from both 1 and 10 μm PS-MPs experiments, including controls (without
157 exposure to PS-MPs). Residual moisture was determined (Gibertini Eurotherm dry weight balance)
158 to correct to dry weight (DW).

159 **2.2 Fatty acids analysis**

160 The fatty acids (FAs) composition of dried *D. magna* individuals was determined and
161 expressed as FA methyl esters (FAMES), as described by Nogueira et al. (2020). Briefly, FAs were
162 converted to FAMES by adding a mixture of ethyl acetate-methanol (1:19 v/v) to dry biomass that
163 was then left at 80 °C for 1 h and then further extracted with heptane. FAMES were analyzed by gas

164 chromatography (Agilent HP 6890) equipped with a mass selective detector (Agilent 5973) and a
165 capillary column DB-225 J&W (30 m × 0.25 mm inner diameter, 0.15 μm film thickness). The
166 chromatographic conditions were as follows: initial temperature of the oven was 35 °C for 0.5 min;
167 was increased by 25 °C min⁻¹ to 195 °C; followed by 3 °C min⁻¹ to 205 °C; and 8 °C min⁻¹ until
168 reaching the final temperature of 230 °C for 3 min. The temperature of the injector was 250 °C, that
169 of the transfer line, 280 °C; and the split ratio was 1:100. Helium was used as the carrier gas, with a
170 flow rate of 2.6 mL min⁻¹. At least two replicates were performed for each gas chromatography
171 analysis.

172 FAMES were identified based on comparing their mass spectra with the equipment mass
173 spectral library (Wiley-Nist) and comparing the retention times and mass spectra fragmentation to
174 known standards (Supelco 37 component FAME Mix; Sigma-Aldrich CRM47885). Heneicosanoic
175 acid (C21:0) was used as an internal standard. For quantitative analysis, GC–MS was calibrated with
176 pure reference compounds (Supelco 37 component FAME Mix; Sigma-Aldrich CRM47885) relative
177 to the internal standard. The respective response factors were calculated as an average of six GC–MS
178 runs. The results presented are the mean values ± standard deviation (SD) of FAMES expressed in
179 the percentage of total FAs detected.

180 **2.3 Statistical analysis**

181 The effects of PS-MPs exposure on the levels of FAs in *D. magna* individuals were
182 investigated by a one-way analysis of variance (ANOVA) followed by Dunnett's multiple
183 comparisons test. Normality and homogeneity of variance was previously checked by means of
184 Shapir-Wilk's and Bartlett's tests, respectively. Pearson' correlation analysis was performed to check
185 for the relationships and changes in the relationships among different fatty acids after the exposure to
186 increasing concentrations of differently shaped PS-MPs. *p*-values < 0.05 were considered as
187 statistically significant. Statistical analyses were performed using GraphPad Prism version 9.3.0 for
188 Windows, GraphPad Software (San Diego, California, USA).


189 3. Results and discussion

190 A previous study performed under these same experimental conditions demonstrated that a
191 21-days exposure to 1 and 10 μm PS-MPs resulted in a quick PS-MPs ingestion and accumulation
192 into the digestive tract of *D. magna* individuals without causing mortality (De Felice et al., 2019).
193 Similarly, a quick and efficient ingestion (data not shown) and no mortality was observed over the
194 whole duration of the present experiment, in agreement with the findings reported by other studies
195 on *Daphnia magna* individuals exposed to different PE (17 – 34 μm ; An et al., 2021) and PS (100 nm
196 and 2 μm ; Rist et al., 2017) microbeads in a similar range of concentrations.

197 3.1. Effect of PS-MPs exposure on the total FAs content

198 The 21-days exposure to three increasing concentrations of 1 and 10 μm PS-MPs induced the
199 modulation of fatty acids (FAs) profile in *D. magna* individuals. The exposure to the highest tested
200 concentration of PS-MPs induced an overall decrease in total FAs content independently of the
201 particle size (Fig. 1). FAs can be used to monitor environmentally-induced stress (Filimonova et al.,
202 2016; Gonçalves et al., 2016; Guschina et al., 2020; Yang et al., 2021), which can affect energy
203 reserves (Lee et al., 2018; Neves et al. 2015). The significant decrease in the total content of FAs
204 observed in *D. magna* exposed to 12.5 $\mu\text{g mL}^{-1}$ of both PS-MPs sizes suggests that treated individuals
205 might suffer an high stressful situation, forcing them to activate biochemical or behavioural energy-
206 demanding defense responses. Indeed, the exposure to the highest concentration of PS-MPs might
207 induce a generic or oxidative stress condition in treated organisms, which could activate physiological
208 or behavioral processes to prevent or to counteract negative consequences towards organism health
209 status. Previous studies on aquatic organisms showed an increase in antioxidant defenses to tackle
210 the overproduction of reactive oxygen species (ROS) due to PS-MPs (Zhang et al., 2019; Liang et al.,
211 2021; Umamaheswari et al., 2021). For instance, a recent study by Tang and coauthors (2019),
212 showed that the 10-days exposure to 1.25 μm PS-MPs (concentration range 2 - 8 mg L^{-1}) altered the
213 expression of genes involved in oxidative stress defense, specifically the expression of thioredoxin

214 reductase (TRxR), an enzyme involved in the thioredoxin system that plays a crucial role in cellular
215 antioxidant defense eliminating the excess of ROS (Lushchak, 2011), with subsequent deregulation
216 of genes related to energy production. Moreover, our previous study demonstrated that the exposure
217 to 12.5 $\mu\text{g mL}^{-1}$ of PS-MPs induced an increase in swimming activity of *D. magna* individuals,
218 suggesting an avoidance behavior fulfilled by the organism to swim away from a highly contaminated
219 environment or an attempt made by the cladocerans to get rid of the particles on their body and
220 appendices (De Felice et al., 2019). Thus, the implementation of physiological or behavioral defense
221 in a stressful situation requires a massive amount of extra energy (Tang et al., 2019). As PS-MPs
222 retention in the digestive tract of *D. magna* can result in a false sense of satiation, reducing food
223 ingestion and energy reserves (Wright et al., 2013), the decrease in FAs content might be due to their
224 use or re-allocation to support defense mechanisms of treated individuals. The allocation of food-
225 derived resources towards growth and reproduction depends on their availability (Mariash et al.,
226 2017; Nogueira et al., 2004). Previous studies have shown the importance of storage lipids for the
227 fitness of *D. magna* individuals (Garreta et al., 2016; Mariash et al., 2017). The life-history traits of
228 *D. magna* are strongly influenced by food concentration. Under food limitation, individuals mature
229 later, present smaller body and clutch sizes compared to conspecifics experiencing high food
230 availability (Klintworth and Von Elert, 2020). Although *D. magna* individuals were fed *ad libitum*
231 during the whole duration of the experiments, the ingestion of PS-MPs and the filling of the digestive
232 tract might cause a sort of fasting condition in treated organisms, reducing the uptake of food and
233 forcing individuals to the mobilisation, and use of FAs as an energetic source to support organism
234 functions. When organisms experience a stressful situation such as fasting, storage lipids can be
235 metabolised, and FAs become available as a metabolic fuel (Tessier et al., 1983, Raclot, 2003). Under
236 a potential limitation of food, due to the presence of PS-MPs in the digestive tract, individuals might
237 consume internal lipid reserves to sustain the basic metabolic processes of the somatic body, causing
238 the decrease of FAs content, as observed in both neonates and juveniles of *D. magna* experiencing
239 short-term starvation (Yang et al., 2021).



240


241 **Fig. 1.** Total fatty acids content of *Daphnia magna* after 21-days exposure to PS-MPs, with different sizes (1
 242 and 10 μm) and concentrations (0.125, 1.25 and 12.5 μg mL⁻¹). Statistical significance: ** $p < 0.01$, **** $p <$
 243 0.0001 after ANOVA followed by Dunnett's multiple comparisons test. Data are reported as mean ± SD (n=3).

244 The decrease of FAs content could be also related to the allocation of storage lipids from lipid
 245 droplets to eggs, a process that could explain the increase in offspring number observed in De Felice
 246 et al. (2019). The authors showed an increase in the reproductive effort of *D. magna* individuals
 247 exposed to the same high concentration of PS-MPs. This result was explained as an effort
 248 accomplished by cladocerans in a highly contaminated environment, where adults preferred to invest
 249 energy in their reproductive fitness rather than survival. The significant reduction of the total FAs
 250 content, especially after the exposure to 12.5 μg mL⁻¹ of PS-MPs of both sizes might promote the
 251 allocation of FAs to the eggs in order to provide offspring with sufficient energetic resources to tackle
 252 stressful situations caused by the exposure to a high concentration of MPs, as well as to outgrow the
 253 most vulnerable instars faster. Indeed, the allocation of FAs to offspring represents a part of life-
 254 history changes in response to environmental stressors, as demonstrated in *D. magna* adults exposed
 255 to predator-borne kairomones (Stibor and Müller-Navarra, 2000; Hahn et al., 2019; Klintworth and
 256 Von Elert, 2020). This hypothesis might be further confirmed through the comparison of the FAs
 257 content and profile of *D. magna* adults treated with PS-MPs and their offspring.

258


259 **3.2. Size-related effect of PS-MPs on FAs profile**

260 The analysis of the relative abundance of FAs allowed to elucidate changes in FAs accumulation
 261 and to identify potentially enhanced metabolic pathways (Fig. **Errore. L'origine riferimento non è**
 262 **stata trovata.**2 and 3**Errore. L'origine riferimento non è stata trovata.**). In the presence of 1 μm
 263 PS-MPs, the relative abundance of monounsaturated fatty acids (MUFAs) was significantly
 264 increased, while polyunsaturated fatty acids (PUFAs) decreased according to the increase of the
 265 exposure concentrations. Although a similar trend was observed at the end of the exposure to 10 μm
 266 PS-MPs, it was only verified at the lower tested concentrations, as the exposure to the highest PS-
 267 MPs concentration led to a re-establishment of the homeostasis regarding to MUFAs/PUFAs
 268 abundance.



269
 270 **Fig. 2.** Relative abundance of fatty acid groups from *Daphnia magna* grown 21-days in the presence of
 271 different concentrations (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of 1 μm PS-MPs. (Control: without exposure to PS-
 272 MPs; SFAs: saturated fatty acids, MUFAs: monounsaturated fatty acids, PUFAs: polyunsaturated fatty acids.

273 Statistical significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ after ANOVA followed by
 274 Dunnett's multiple comparisons test. Data are reported as mean \pm SD (n=3)).



275
 276 **Fig. 3.** Relative abundance of fatty acid groups from *Daphnia magna* grown 21-days in the presence of
 277 different concentrations (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of 10 μm PS-MPs concentrations. (Control: without
 278 exposure to PS-MPs; SFAs: saturated fatty acids, MUFAs: monounsaturated fatty acids, PUFAs:
 279 polyunsaturated fatty acids. Statistical significance: **** $p < 0.0001$ after ANOVA followed by Dunnett's
 280 multiple comparisons test. Data are reported as mean \pm SD (n=3)).

281 The analysis of FAs profile of *D. magna* exposed to 1 μm PS-MPs (Table 1 **Errore. L'origine**
 282 **riferimento non è stata trovata.**) showed a significant increase in the abundance of MUFAs, except
 283 C18:1 ω 9 (oleic acid). A similar modulation of the FAs profile was also observed in a common and
 284 widespread algal species, *Chlorella sorokiniana*, exposed to PS-MPs (< 70 μm in size), whereby the
 285 reduction in the content of essential FAs C18:2 ω 6 (linoleic acid) and C18:3 ω 3 (α -linolenic acid) was
 286 accompanied by an increase in C18:1 ω 9 content (Guschina et al., 2020).


287 **Table 1.** Relative abundance (mean \pm standard deviation) of fatty acids from *Daphnia magna* grown 21-days in the presence of different concentrations (0.125,
 288 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of 10 μm PS-MPs concentrations.

Fatty acid (%)	Control		0.125 $\mu\text{g mL}^{-1}$		1.25 $\mu\text{g mL}^{-1}$		12.5 $\mu\text{g mL}^{-1}$	
	1 μm	10 μm	1 μm	10 μm	1 μm	10 μm	1 μm	10 μm
C14:1 ω9c	0.49 \pm 0.01	0.47 \pm 0.03	0.44 \pm 0.02	0.60 \pm 0.02	0.52 \pm 0.02	0.55 \pm 0.02	0.60 \pm 0.03	0.50 \pm 0.04
C14:1 ω9t	1.50 \pm 0.02	1.69 \pm 0.04	1.54 \pm 0.03	1.97 \pm 0.05	1.59 \pm 0.02	1.90 \pm 0.07	1.78 \pm 0.04**	1.78 \pm 0.07
C14:0	1.77 \pm 0.03	1.48 \pm 0.01	1.84 \pm 0.03	1.55 \pm 0.04	1.89 \pm 0.03	1.46 \pm 0.05	1.68 \pm 0.05	1.46 \pm 0.07
C15:1 ω5c	0.00 \pm 0.00	0.16 \pm 0.01	0.18 \pm 0.02	0.00 \pm 0.00	0.21 \pm 0.01*	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00
C15:0	1.26 \pm 0.02	1.23 \pm 0.02	1.33 \pm 0.03	1.39 \pm 0.04	1.39 \pm 0.01	1.30 \pm 0.05	1.32 \pm 0.02	1.27 \pm 0.03
C16:1 ω9c	8.30 \pm 0.05	8.30 \pm 0.06	8.12 \pm 0.11	9.18 \pm 0.14**	8.32 \pm 0.03	8.77 \pm 0.67	9.24 \pm 0.06****	9.02 \pm 0.09*
C16:1 ω7c	30.98 \pm 0.26	33.91 \pm 0.21	32.04 \pm 0.48****	35.77 \pm 0.40****	32.25 \pm 0.11****	33.85 \pm 0.39	33.70 \pm 0.07****	33.36 \pm 0.34
C16:1 ω9t	0.23 \pm 0.01	0.22 \pm 0.01	0.23 \pm 0.01	0.27 \pm 0.01	0.25 \pm 0.01	0.00 \pm 0.00	0.24 \pm 0.01	0.25 \pm 0.00
C16:1 ω7t	0.33 \pm 0.01	0.25 \pm 0.01	0.30 \pm 0.02	0.27 \pm 0.01	0.34 \pm 0.01	0.24 \pm 0.02	0.18 \pm 0.01	0.27 \pm 0.01
C16:0	10.67 \pm 0.08	9.22 \pm 0.04	10.91 \pm 0.14*	9.01 \pm 0.07	10.78 \pm 0.03	8.79 \pm 0.11	9.34 \pm 0.00	9.07 \pm 0.11
C18:3 ω6c	0.16 \pm 0.02	0.16 \pm 0.01	0.14 \pm 0.01	0.18 \pm 0.01	0.18 \pm 0.00	0.17 \pm 0.02	0.07 \pm 0.00	0.19 \pm 0.01
C18:2 ω6c	3.09 \pm 0.02	3.09 \pm 0.02	2.94 \pm 0.15	0.00 \pm 0.00****	2.82 \pm 0.01**	3.02 \pm 0.14	2.46 \pm 0.01****	3.26 \pm 0.06
C18:3 ω3c	7.11 \pm 0.11	9.98 \pm 0.11	6.02 \pm 0.20****	6.81 \pm 0.67****	5.85 \pm 0.05****	9.49 \pm 0.35	6.67 \pm 0.28****	9.55 \pm 1.75
C18:1 ω9c	28.28 \pm 0.30	24.13 \pm 0.23	27.78 \pm 0.51****	27.14 \pm 0.49****	27.94 \pm 0.19***	24.78 \pm 1.45*	27.55 \pm 0.01****	23.70 \pm 1.06
C18:1 ω9t	2.47 \pm 0.02	2.73 \pm 0.01	2.74 \pm 0.01**	2.74 \pm 0.14	2.57 \pm 0.02	2.73 \pm 0.05	2.37 \pm 0.01	3.04 \pm 0.03
C18:0	2.67 \pm 0.02	2.39 \pm 0.02	2.68 \pm 0.03	2.42 \pm 0.05	2.64 \pm 0.01	2.44 \pm 0.03	2.36 \pm 0.00**	2.54 \pm 0.04
C20:4 ω6c	0.32 \pm 0.01	0.26 \pm 0.02	0.34 \pm 0.03	0.30 \pm 0.03	0.00 \pm 0.00***	0.17 \pm 0.02	0.16 \pm 0.01	0.44 \pm 0.01
C20:5 ω3c	0.19 \pm 0.02	0.17 \pm 0.02	0.25 \pm 0.03	0.21 \pm 0.01	0.03 \pm 0.00	0.14 \pm 0.02	0.18 \pm 0.00	0.14 \pm 0.00
C20:1 ω9c	0.07 \pm 0.01	0.06 \pm 0.01	0.07 \pm 0.01*	0.08 \pm 0.01	0.31 \pm 0.02	0.08 \pm 0.01	0.00 \pm 0.00	0.07 \pm 0.01
C22:0	0.08 \pm 0.01	0.07 \pm 0.01	0.07 \pm 0.01	0.07 \pm 0.00	0.07 \pm 0.01	0.07 \pm 0.01	0.05 \pm 0.01	0.06 \pm 0.01
C24:0	0.05 \pm 0.01	0.04 \pm 0.00	0.05 \pm 0.00	0.04 \pm 0.01	0.05 \pm 0.00	0.03 \pm 0.01	0.03 \pm 0.00	0.04 \pm 0.01

289 Statistical significance: without* - no statistical significance; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$ after ANOVA followed by Dunnett's multiple comparisons

290 test. Data are reported as mean \pm SD. (n=3)).

291 The correlation analysis on the abundance of FAs (Fig. 4) allowed the identification of
 292 relationships occurring among different lipid species and to hypothesize biosynthetic events translating
 293 these variations. The synthesis of C14:1 and C16:1 was performed at the expense of saturated fatty
 294 acids (SFAs), while the increase of C16:1 and other MUFAs resulted as inversely correlated with
 295 C18:3 and C18:2 content (Fig. 4; correlations highlighted in box named with the letter A).
 296 Additionally, C18:1ω9 was also consumed to synthesize C14:1 and C16:1. Considering these
 297 findings, we might speculate that under the exposure to high concentrations of PS-MPs, treated
 298 individuals redirected the synthesis of FAs towards shorter and less unsaturated molecules, increasing
 299 C14 and C16 MUFAs. The variation of SFAs (Fig. 4; correlations highlighted in box named with the
 300 letter **Errore. L'origine riferimento non è stata trovata.**B) was also simultaneous, suggesting a
 301 whole regulation, as supported by the positive correlation indexes. As PUFAs are accumulated and
 302 transferred upwards in the food web, *D. magna* individuals accumulate these FAs according to their
 303 availability in the diet (Masclaux et al., 2012). Therefore, it is also possible that the decrease in the
 304 PUFAs/MUFAs ratio is related to less PUFAs being obtained.



305


306 **Fig. 4.** Correlation matrix of fatty acids from *Daphnia magna* grown 21 days in the presence of different
307 concentrations (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of 1 μm PS-MPs. Values are presented as linear correlation
308 index between variables; the colour gradient represent the strength of the correlations.

309 Regarding the individuals grown in the presence of 10 μm PS-MPs, an increase in MUFAs,
310 particularly C16:1, was observed (Table 1). However, while the C18:1 ω 9 content increased, a
311 decrease in C18:3 occurred, suggesting an interconversion of these two FAs. In the 1 μm PS-MPs
312 experiments, C18:1 ω 9 content increased, suggesting a different impact of this size of PS-MPs on FAs
313 metabolism. All the effects caused by 10 μm PS-MPs were only observed at the lowest tested
314 concentration, while homeostasis seems to be re-established after the exposure to the higher
315 concentrations. These results might suggest i) a sort of adaptation in *D. magna* to the increasing
316 concentrations of PS-MPs and ii) that also low concentrations of PS-MPs are sufficient to induce
317 significant modulation of FAs metabolism and accumulation in this cladoceran species. Thus, smaller
318 sized PS-MPs exposure should lead to a decrease in the accumulation of essential FAs by the diet,
319 promoting the synthesis of monounsaturated FAs-rich storage lipids, as well as allocating resources
320 towards their offspring. Larger MPs caused similar events on fatty acids management. However, at
321 higher concentrations, the individuals look like to experience a sort of adaptation that counters the
322 environmental stress and could promotes mobility, reproduction and food procurement, reestablishing
323 fatty acids homeostasis and consuming storage lipids in the process.

324 The analysis of the correlation among FAs content measured in individuals exposed to 10 μm
325 PS-MPs (Fig. 5), confirmed that C18 PUFAs were consumed when an increase in the abundance of
326 C14, C16 and C18 MUFAs occurred (Fig. 5; correlations highlighted in box named with the letter
327 A). Moreover, the positive correlation between all MUFAs (Fig. 5; correlations highlighted in box
328 named with the letter B) suggested a synchronism in this regularory mechanism, as the synthesis of
329 these FAs seems to be modulated collectively. Overall, these results showed that the presence of PS-
330 MPs in the environment might force *D. magna* individuals to accumulate higher amounts of MUFAs

331 rather than PUFAs. These changes might be induced by a limited food availability, leading to a
 332 rewiring of the FAs biosynthesis that promotes the conversion of longer and more unsaturated fatty
 333 acids into shorter and more saturated ones. The increase in MUFAs content could benefit individuals
 334 experiencing stressful conditions through the increase of the amount of storage lipids. Indeed,
 335 MUFAs and their derived fatty alcohols are important components of esters (Brett et al., 2009), the
 336 main form of storage lipid in daphnids (Lee et al., 2006).

337



338

339 **Fig. 5.** Correlation matrix of fatty acids from *Daphnia magna* grown 21 days in presence of different
 340 concentrations (0.125, 1.25 and 12.5 $\mu\text{g mL}^{-1}$) of 10 μm PS-MPs concentrations. Values are presented as linear
 341 correlation index between variables; the colour gradient represent the strength of the correlations.

342 **3.3. Effect of PS-MPs exposure on the biosynthetic $\omega 3$, $\omega 6$ and $\omega 9$ pathways**

343 A significant decrease in the abundance of $\omega 3$ and $\omega 6$ fatty acids, but an increase in $\omega 9$, was
 344 observed at the end of the exposure to 1 μm PS-MPs. Interestingly, at 10 μm PS-MPs exposure, such
 345 effects were observed only at the lowest tested concentrations (0.125 $\mu\text{g mL}^{-1}$), suggesting that the
 346 effect on $\omega 3$, $\omega 6$ and $\omega 9$ FAs are related to the mechanisms previously discussed regarding

347 PUFAs/MUFAs abundance. For instance, the most abundant ω 3 and ω 6 FAs are C18:2 and C18:3,
348 which were also hallmarks of the MUFAs decrease (Table 1).

349 The correlation analysis highlighted a collective variation in ω 6 FAs, with positive correlations
350 among FAs from this biosynthetic pathway (Fig. 4; correlations highlighted in box named with the
351 letter D). Accordingly, the synthesis of the ω 6 ARA, an essential precursor of signalling biomolecules
352 and a key element in the reproduction in *D. magna* (Ginjupalli et al., 2015), is accumulated together
353 with C18:3 ω 6 (Fig. 5; correlations highlighted in box named with the letter D).

354 These results also suggest that the ω 9 pathway is regulated at the expense of C16:0, leading to the
355 synthesis of C20:1 ω 9. When this synthesis is enhanced, there is an increment of C16, C18 and C20
356 MUFAs (Fig. 5; correlations highlighted in box named with the letter E). Overall, these results
357 suggest the allocation of resources towards ω 9 FAs, in addition to the MUFAs accumulation. Finally,
358 a sharp increase in the ω 3/ ω 6 ratio was observed in individuals that experienced the exposure to the
359 lowest concentration of 10 μ m and the highest one of 1 μ m PS-MPs. This situation could likely occur
360 as a consequence of a competition in the intake of essential ω 3 FAs, which are not accumulated, as
361 the synthesis of ω 6 and ω 9 storage lipids increases. These findings confirmed that FAs are involved
362 in the response to PS-MPs exposure and their modulation might be considered as a valuable marker
363 of MPs contamination in natural ecosystems, as daphnids seem to accumulate shorter FAs in the
364 absence of ω 3-rich food.

365 **4. Conclusions**

366 The present study showed that the exposure to increasing concentration of 1 and 10 μ m PS-MPs,
367 resulted in the modulation of fatty acids profile and content of *D. magna* individuals. Differently sized
368 MPs affected fatty acids composition and profiles in distinct manners. In fact, higher concentrations
369 of smaller MPs and lower concentrations of larger MPs exerted the most significant changes in the
370 FAs profile. Although the concentrations of PS-MPs we tested in the present study can be considered
371 as unrealistic, considering the amount of large-sized plastics in the environment and their potential

372 fragmentation, the levels of MPs could increase, returning effects similar to those observed in this
373 experiments. These results enlarged the knowledge of the sub-lethal effects of MPs in zooplanktonic
374 species. They returned helpful information to understand the mechanisms of action underlying the
375 impact of these contaminants on different life-history traits, confirming that the ingestion of MPs can
376 influence the energy budget and allocation, with potential consequences on growth, reproduction, and
377 survival. In addition, further studies need to explore the adverse effects induced by mixture of MPs
378 of different size, shape and polymeric composition towards aquatic organisms. This approach should
379 increase the ecological realism of the experiments and allow to shed light on the effects induced by
380 different interacting substances, returning useful, although not conclusive, information to assess the
381 real risk of MPs exposure in natural ecosystems.

382

383 **5. References**

- 384 Al-Jaibachi, R., & Callaghan, A. (2018). Impact of polystyrene microplastics on *Daphnia magna*
385 mortality and reproduction in relation to food availability. PeerJ, 6, e4601.
- 386 Al-Jaibachi, R., Cuthbert, R. N., & Callaghan, A. (2018). Up and away: ontogenic transference as a
387 pathway for aerial dispersal of microplastics. Biology Letters, 14(9), 20180479.
- 388 Ambrosini, R., Azzoni, R. S., Pittino, F., Diolaiuti, G., Franzetti, A., & Parolini, M. (2019). First
389 evidence of microplastic contamination in the supraglacial debris of an alpine glacier.
390 Environmental Pollution, 253, 297-301.
- 391 An, D., Na, J., Song, J., Jung, J., 2021. Size-dependent chronic toxicity of fragmented polyethylene
392 microplastics to *Daphnia magna*. Chemosphere 271, 129591.
- 393 Andrady, A.L., & Neal, M.A. (2009). Applications and societal benefits of plastics. Phil. Trans. R.
394 Soc. B. 364, 1977-1984.

- 395 Arts, M. T., & Kohler, C. C. (2009). Health and condition in fish: the influence of lipids on membrane
396 competency and immune response. *Lipids in aquatic ecosystems*, 237-256.
- 397 Barboza, L. G. A., Vethaak, A. D., Lavorante, B. R., Lundebye, A. K., & Guilhermino, L. (2018).
398 Marine microplastic debris: An emerging issue for food security, food safety and human health.
399 *Marine Pollution Bulletin*, 133, 336-348.
- 400 Bellasi, A., Binda, G., Pozzi, A., Galafassi, S., Volta, P., & Bettinetti, R. (2020). Microplastic
401 contamination in freshwater environments: A review, focusing on interactions with sediments and
402 benthic organisms. *Environments*, 7(4), 30.
- 403 Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J., & Gerdt, G. (2019). White and
404 wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Science Advances*, 5(8),
405 eaax1157.
- 406 Blarer, P., & Burkhardt-Holm, P. (2016). Microplastics affect assimilation efficiency in the
407 freshwater amphipod *Gammarus fossarum*. *Environmental Science and Pollution Research*,
408 23(23), 23522-23532.
- 409 Brett, M. T., Müller-Navarra, D. C., & Persson, J. (2009). Crustacean zooplankton fatty acid
410 composition. *Lipids in Aquatic Ecosystems*, 9780387893662, 115–146.
- 411 Can-Güven, E. (2021). Microplastics as emerging atmospheric pollutants: a review and bibliometric
412 analysis. *Air Quality, Atmosphere & Health*, 14(2), 203-215.
- 413 Canniff, P. M., & Hoang, T. C. (2018). Microplastic ingestion by *Daphnia magna* and its
414 enhancement on algal growth. *Science of the Total Environment*. 633, 500-507.
- 415 Cole, M., Lindeque, P., Fileman, E., Halsband, C., & Galloway, T.S. (2015). The impact of
416 polystyrene microplastics on feeding, function and fecundity in the marine copepod *Calanus*
417 *helgolandicus*. *Environmental Science & Technology* 49, 1130–1137.

418 Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., & Galloway, T.S. (2013).
419 Microplastic ingestion by zooplankton. *Environmental Science & Technology* 47, 6646–6655.

420 Crosta, A., De Felice, B., Antonioli, D., Chiarcos, R., Perin, E., Ortenzi, M. A., et al. (2022).
421 Microplastic contamination of supraglacial debris differs among glaciers with different anthropic
422 pressures. *Science of The Total Environment*, 158301.

423 De Felice, B., Bacchetta, R., Santo, N., Tremolada, P., & Parolini, M. (2018). Polystyrene
424 microplastics did not affect body growth and swimming activity in *Xenopus laevis* tadpoles.
425 *Environmental Science and Pollution Research*, 25(34), 34644-34651.

426 De Felice, B., Sabatini, V., Antenucci, S., Gattoni, G., Santo, N., Bacchetta, R., & Parolini, M. (2019).
427 Polystyrene microplastics ingestion induced behavioral effects to the cladoceran *Daphnia magna*.
428 *Chemosphere*, 231, 423-431.

429 De Sá, L. C., Oliveira, M., Ribeiro, F., Rocha, T. L., & Futter, M. N. (2018). Studies of the effects of
430 microplastics on aquatic organisms: what do we know and where should we focus our efforts in
431 the future? *Science of the Total Environment*, 645, 1029-1039.

432 Dris, R., Imhof, H., Sanchez, W., Gasperi, J., Galgani, F., Tassin, B., & Laforsch, C. (2015). Beyond
433 the ocean: contamination of freshwater ecosystems with (micro-)plastic particles. *Environmental*
434 *Chemistry*. 12, 539–550.

435 Eerkes-Medrano, D., Thompson R.C., & Aldridge, D.C. (2015). Microplastics in freshwater systems:
436 A review of the emerging threats, identification of knowledge gaps and prioritisation of research
437 needs. *Water Research*. 75, 63-82.

438 Filimonova, V., Gonçalves, F., Marques, J. C., De Troch, M., & Gonçalves, A. M. (2016). Fatty acid
439 profiling as bioindicator of chemical stress in marine organisms: a review. *Ecological Indicators*,
440 67, 657-672.

441 Frydkjær, C., Iversen, N., & Roslev, P. (2017). Ingestion and egestion of microplastics by the
442 Cladoceran *Daphnia magna*: effects of regular and irregular shaped plastic and Sorbed
443 Phenanthrene. *Bulletin of Environmental Contamination and Toxicology*, 99, 655–661.

444 Ginjupalli, G. K., Gerard, P. D., & Baldwin, W. S. (2015). Arachidonic acid enhances reproduction
445 in *Daphnia magna* and mitigates changes in sex ratios induced by pyriproxyfen. *Environmental*
446 *Toxicology and Chemistry*, 34(3), 527–535.

447 Gonçalves, A. M. M., Mesquita, A. F., Verdelhos, T., Coutinho, J. A. P., Marques, J. C., & Gonçalves,
448 F. (2016). Fatty acids' profiles as indicators of stress induced by of a common herbicide on two
449 marine bivalves species: *Cerastoderma edule* (Linnaeus, 1758) and *Scrobicularia plana* (da Costa,
450 1778). *Ecological Indicators*, 63, 209-218.

451 González-Pleiter, M., Lacerot, G., Edo, C., Pablo Lozoya, J., Leganés, F., Fernández-Piñas, F., ... &
452 Teixeira-de-Mello, F. (2021). A pilot study about microplastics and mesoplastics in an Antarctic
453 glacier. *The Cryosphere*, 15(6), 2531-2539.

454 Gorokhova E.,2015. Screening for microplastic particles in plankton samples: How to integrate
455 marine litter assessment into existing monitoring programs? *Marine Pollution Bulletin*. 99, 271-
456 275.

457 Hahn, M. A., Effertz, C., Bigler, L., & von Elert, E. (2019). 5 α -cyprinol sulfate, a bile salt from fish,
458 induces diel vertical migration in *Daphnia*. *Elife*, 8, e44791.

459 Hartmann, N. B., Hüffer, T., Thompson, R. C., Hassellöv, M., Verschoor, A., Daugaard, A. E., ..., &
460 Wagner, M. (2019). Are We Speaking the Same Language? Recommendations for a Definition
461 and Categorization Framework for Plastic Debris. *Environmental Science and Technology*, 53(3),
462 1039–1047.

- 463 He, M., Yan, M., Chen, X., Wang, X., Gong, H., Wang, W., & Wang, J. (2021). Bioavailability and
464 toxicity of microplastics to zooplankton. *Gondwana Research*, 108, 120-126.
- 465 Horton, A. A., & Barnes, D. K. (2020). Microplastic pollution in a rapidly changing world:
466 implications for remote and vulnerable marine ecosystems. *Science of The Total Environment*,
467 738, 140349.
- 468 Imhof, H. K., & Laforsch, C. (2016). Hazardous or not–Are adult and juvenile individuals of
469 *Potamopyrgus antipodarum* affected by non-buoyant microplastic particles? *Environmental*
470 *Pollution*, 218, 383-391.
- 471 Imhof, H.K., Rusek, J., Thiel, M., Wolinska, J., & Laforsch, C. (2017). Do microplastic particles
472 affect *Daphnia magna* at the morphological, life history and molecular level? *PlosOne* DOI
473 12:e0187590.
- 474 Jemec, A., Horvat, P., Kunej, U., Bele, M., & Kržan, A. (2016). Uptake and effects of microplastic
475 textile fibers on freshwater crustacean *Daphnia magna*. *Environmental Pollution*, 219, 201-209.
- 476 Kaposi, K.L., Mos, B., Kelaher, B.P., & Dworjanyn, S.A. (2014). Ingestion of Microplastic has
477 limited impact on a marine larva. *Environmental Science & Technology*. 48, 1638-1645.
- 478 Klintworth, S., & Von Elert, E. (2020). Risk of predation alters resource allocation in *Daphnia* under
479 food limitation. *Journal of Plankton Research*, 42(1), 45-56.
- 480 Lee, M.C., Park, J.C., & Lee, J.S. (2018). Effects of environmental stressors on lipid metabolism in
481 aquatic invertebrates. *Aquatic Toxicology*, 200, 83-92.
- 482 Lee, R. F., Hagen, W., & Kattner, G. (2006). Lipid storage in marine zooplankton. *Marine Ecology*
483 *Progress Series*, 307, 273–306.

- 484 Li, C., Busquets, R., & Campos, L. C. (2020). Assessment of microplastics in freshwater systems: A
485 review. *Science of the Total Environment*, 707, 135578.
- 486 Li, J., Liu, H., & Chen, J.P. (2018). Microplastics in freshwater systems: A review on occurrence,
487 environmental effects, and methods for microplastics detection. *Water research*, 137, 362-374.
- 488 Liang, Y., Yang, X., Wang, Y., Liu, R., Gu, H., & Mao, L. (2021). Influence of polystyrene
489 microplastics on rotifer (*Brachionus calyciflorus*) growth, reproduction, and antioxidant
490 responses. *Aquatic Ecology*, 55(3), 1097-1111.
- 491 Liu, Y., Zhang, J., Zhao, H., Cai, J., Sultan, Y., Fang, H., ... & Ma, J. (2022). Effects of polyvinyl
492 chloride microplastics on reproduction, oxidative stress and reproduction and detoxification-
493 related genes in *Daphnia magna*. *Comparative Biochemistry and Physiology Part C: Toxicology*
494 & *Pharmacology*, 109269.
- 495 Lo, H.K.A., & Chan, K.Y.K. (2018). Negative effects of microplastic exposure on growth and
496 development of *Crepidula onyx*. *Environmental Pollution*. 233, 588–595.
- 497 Lu, Y., Zhang, Y., Deng, Y., Jiang W., Zhao, Y., Geng, J., Ding, L., & Ren, H. (2016). Uptake and
498 Accumulation of Polystyrene Microplastics in Zebrafish (*Danio rerio*) and Toxic Effects in Liver.
499 *Environmental Science & Technology*. 50, 4054-4060.
- 500 Lushchak, V. I. (2011). Environmentally induced oxidative stress in aquatic animals. *Aquatic*
501 *toxicology*, 101(1), 13-30.
- 502 Mariash, H. L., Cusson, M., & Rautio, M. (2017). Fall Composition of Storage Lipids is Associated
503 with the Overwintering Strategy of *Daphnia*. *Lipids*, 52(1), 83–91.
- 504 Masclaux, H., Bec, A., Kainz, M. J., Perrière, F., Desvillettes, C., & Bourdier, G. (2012).
505 Accumulation of polyunsaturated fatty acids by cladocerans: effects of taxonomy, temperature and
506 food. *Freshwater Biology*, 57(4), 696–703.

507 Messinetti, S., Mercurio, S., Parolini, M., Sugni, M., & Pennati, R. (2018). Effects of polystyrene
508 microplastics on early stages of two marine invertebrates with different feeding strategies.
509 *Environmental Pollution*, 237, 1080-1087.

510 Na, J., Song, J., Achar, J.C., Jung, J., 2021. Synergistic effect of microplastic fragments and
511 benzophenone-3 additives on lethal and sublethal *Daphnia magna* toxicity. *Journal of Hazardous*
512 *Materials*, 402, 123845.

513 Neves, M. F. J. V., Castro, B. B., Vidal, T., Vieira, R. H. S. D. F., Marques, J. C., Coutinho, J. A. P.,
514 & Gonçalves, A. M. M. (2015). Biochemical and populational responses of an aquatic bioindicator
515 species, *Daphnia longispina*, to a commercial formulation of a herbicide (Primextra® Gold TZ)
516 and its active ingredient (S-metolachlor). *Ecological Indicators*, 53, 220-230.

517 Nogueira, A. J. A., Baird, D. J., & Soares, A. M. V. M. (2004). Testing physiologically-based resource
518 allocation rules in laboratory experiments with *Daphnia magna* Straus. *Annales de Limnologie -*
519 *International Journal of Limnology*, 40(4), 257–267.

520 Nogueira, N., Nascimento, F., Cunha, C., & Cordeiro, N. (2020). *Nannochloropsis gaditana* grown
521 outdoors in annular photobioreactors: Operation strategies. *Algal Research*, 48, 101913.

522 Ogonowski, M., Schür, C., Jarsén, Å., & Gorokhova, E. (2016). The effects of natural and
523 anthropogenic microparticles on individual fitness in *Daphnia magna*. *PloS one*, 11(5), e0155063.

524 Parolini, M., Antonioli, D., Borgogno, F., Gibellino, M. C., Fresta, J., Albonico, C., ... & Cavallo, R.
525 (2021). Microplastic Contamination in Snow from Western Italian Alps. *International Journal of*
526 *Environmental Research and Public Health*, 18(2), 768.

527 Parolini, M., De Felice, B., Ferrario, C., Salgueiro-González, N., Castiglioni, S., Finizio, A., &
528 Tremolada, P. (2018). Benzoylecgonine exposure induced oxidative stress and altered swimming
529 behavior and reproduction in *Daphnia magna*. *Environmental Pollution*, 232, 236-244.

530 PlasticsEurope, 2020. <https://plasticseurope.org/it/knowledge-hub/plastics-the-facts-2020/>

531 Raclot, T. (2003). Selective mobilisation of fatty acids from adipose tissue triacylglycerols. Progress
532 in Lipid Research, 42(4), 257-288.

533 Richardson, C. R., Burritt, D. J., Allan, B. J., & Lamare, M. D. (2021). Microplastic ingestion induces
534 asymmetry and oxidative stress in larvae of the sea urchin *Pseudechinus huttoni*. Marine Pollution
535 Bulletin, 168, 112369.

536 Rillig, M. C., & Lehmann, A. (2020). Microplastic in terrestrial ecosystems. Science, 368(6498),
537 1430-1431.

538 Rist, S., Baun, A., & Hartmann, N. (2017). Ingestion of micro- and nanoplastics in *Daphnia magna*-
539 quantification of body burdens and assessment of feeding rates and reproduction. Environmental
540 Pollution. 228, 398–407.

541 Rosenkranz, P., Chaudhry, Q., Stone, V., & Fernandes, T.F. (2009). A comparison of nanoparticle
542 and fine particle uptake by *Daphnia magna*. Environmental Toxicology and Chemistry. 28,10.

543 Scherer, C., Brennholt, N., Reifferscheid, G., & Wagner, M. (2017). Feeding type and development
544 drive the ingestion of microplastics by freshwater invertebrates. Scientific Reports. 7,17006.

545 Shahul Hamid, F., Bhatti, M. S., Anuar, N., Anuar, N., Mohan, P., & Periathamby, A. (2018).
546 Worldwide distribution and abundance of microplastic: how dire is the situation?. Waste
547 Management & Research, 36(10), 873-897.

548 Sighicelli, M., Pietrelli, L., Lecce, F., Iannilli, V., Falconieri, M., Coscia, L., & Zampetti, G. (2018).
549 Microplastic pollution in the surface waters of Italian Subalpine Lakes. Environmental Pollution,
550 236, 645-651.

551 Song, J., Na, J., An, D., Jung, J., 2021a. Role of benzophenone-3 additive in chronic toxicity of
552 polyethylene microplastic fragments to *Daphnia magna*. *Science of the Total Environment*, 800,
553 149638.

554 Song, J., Kim, C., Na, J., Sivri, N., Samanta, P., Jung, J., 2021b. Transgenerational effects of
555 polyethylene microplastic fragments containing benzophenone-3 additive in *Daphnia magna*.
556 *Journal of Hazardous Materials*, 436, 129225.

557 Stibor, H., & Navarra, D. M. (2000). Constraints on the plasticity of *Daphnia magna* influenced by
558 fish-kairomones. *Functional Ecology*, 455-459.

559 Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M.E.J., Le Goïc, N.,
560 Quillien, V., Mingant, C., Epelboin, Y., Corporeau, C., Guyomarch, J., Robbens, J., Paul-Pont, I.,
561 Soudant, P., & Huvet, A. (2016). Oyster reproduction is affected by exposure to polystyrene
562 microplastics. *Proceedings of the National Academy of Sciences. USA* 113:2430–2435.

563 Tang, J., Wang, X., Yin, J., Han, Y., Yang, J., Lu, X., ... & Yang, Z. (2019). Molecular
564 characterization of thioredoxin reductase in waterflea *Daphnia magna* and its expression
565 regulation by polystyrene microplastics. *Aquatic Toxicology*, 208, 90-97.

566 Tessier, A.J., Henry, L.L., Goulden, C.E., & Durand, M.W. (1983). Starvation in *Daphnia*: Energy
567 reserves and reproductive allocation 1. *Limnology and Oceanography*, 28(4), 667-676.

568 Umamaheswari, S., Priyadarshinee, S., Kadirvelu, K., & Ramesh, M. (2021). Polystyrene
569 microplastics induce apoptosis via ROS-mediated p53 signaling pathway in zebrafish. *Chemico-
570 Biological Interactions*, 345, 109550.

571 Wagner, M., & Lambert, S. (2018). Freshwater microplastics: emerging environmental
572 contaminants? (p. 303). Springer Nature.

573 Wagner, M., Scherer, C., Alvarez-Muñoz, D., Brennholt, N., Bourrain, X., Buchinger, S., Fries, E.,
574 Grosbois, C., Klasmeier, J., Marti, T., Rodriguez-Mozaz, S., Urbatzka, R., Vethaak, A.D.,
575 Winther-Nielsen, M., & Reifferscheid, G. (2014). Microplastics in freshwater ecosystems: what
576 we know and what we need to know. *Environmental Sciences Europe*. 24,12.

577 Watts, A.J.R., Lewis, C., Goodhead, R.M., Beckett, S.J., Moger, J., Tyler, C.R., & Gallowat, T.S.
578 (2014). Uptake and retention of microplastics by the shore crab *Carcinus maenas*. *Environmental*
579 *Science Technology*. 48,8823–8830.

580 Weber, A., Scherer, C., Brennholt, N., Reifferscheid, G., & Wagner, M. (2018). PET microplastics
581 do not negatively affect the survival, development, metabolism and feeding activity of the
582 freshwater invertebrate *Gammarus pulex*. *Environmental Pollution*. 234,181–189.

583 Wright, S.L., Thompson, R.C., & Galloway, T.S. (2013). The physical impacts of microplastics on
584 marine organisms: a review. *Environmental Pollution* 178, 483–492

585 Yang, M., Wei, J., Wang, Y., Shen, C., & Xie, X. (2021). Short-term starvation affects fatty acid
586 metabolism of *Daphnia magna* neonates and juveniles. *Aquatic Sciences*, 83(1), 1-11.

587 Zhang, P., Yan, Z., Lu, G., & Ji, Y. (2019). Single and combined effects of microplastics and
588 roxithromycin on *Daphnia magna*. *Environmental Science and Pollution Research*, 26(17), 17010-
589 17020.

590