

Amphiphilic Pt(II) complex for in vivo imaging applications
Tania Pecoraro^{a,b}, Giorgio Facchetti^b, Isabella Rimoldi^b, Claudia Tortiglione^c,
Giuseppina Tommasini^c, Alessandro Aliprandi^d, Luisa De Cola^{a,b}

^a Department of Molecular Biochemistry and Pharmacology, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, 2056 Milano

^bDISFARM, Università degli Studi di Milano, 20133 Milano

^c Istituto di Scienze Applicate e Sistemi Intelligenti "E. Caianiello", CNR, Pozzuoli

^d Université de Strasbourg, ISIS, & CNRS UMR 7006, Strasbourg, France

Transition metal complexes have attracted much attention because of their rich and peculiar chemical and spectroscopic properties that have found applications in different fields and even as therapeutics.^[1] In this field, Pt(II) complexes have been extensively studied especially for cancer treatment thanks to their high cytotoxicity. Platinum complexes can be very luminescent and their emission has been investigated for optoelectronics, while their use as luminescent probes, for cellular imaging, has been much less explored. One of the main drawbacks of phosphorescent compounds as optical labels is the efficient quenching of dioxygen in solution. This problem can be overcome by exploiting the high tendency of such square planar compounds, containing conjugated ligands, to self-assembly in supramolecular structures. This phenomenon can significantly enhance the emissive properties of Pt(II) compounds, because of the formation of new excited states (metal-metal ligand charge transfer, MMLCT) and an increasing rigidity due to the packing of the units, and as a consequence, also a slower or negligible diffusion of dioxygen in the assembled structures. The assemblies therefore can become better probes for imaging application due to their enhanced emission and reduced reactivity.^[2] In this regard, our research group has already demonstrated the aggregation induced emission for Pt(II) complexes^[3,4] in different media. In this contribution we describe the synthesis and characterization of a luminescent amphiphilic platinum compound, soluble in water, containing a N⁺N⁺N⁺ pyridil-triazolate and a positively charged functionalized pyridine and its behaviour in vivo. The complex aggregates in water giving a very bright orange luminescence. As an in vivo model we have employed an invertebrate, transparent, freshwater polyp, *Hydra vulgaris*. It was treated with the phosphorescent complexes at only 20 μM concentration.,

The compound self-assembles in vivo and in particular accumulates in the tentacles of the animal. Interestingly, the results obtained suggest not only the use of this compound for bioimaging and cell tracking but also as an enhancer of cell proliferation.

References:

[1] K.J. Franz et al. *Chem. Rev.* **2019**, 119, 2, 727–729

[2] S. Sinn et al. *J. Am. Chem. Soc.* **2018**, 140, 2355–2362

[3] A. Aliprandi et al. *Nature Chemistry*, **2016**, 10-15

[4] A. Aliprandi, et al. *Isr. J. Chem.*, **2019**, 59, 892–897