

JOURNAL OF AVIAN BIOLOGY

Viewpoint

Insufficient considerations of seasonality, data selection and validation lead to biased species–climate relationships in mountain birds

Mattia Brambilla, Chiara Bettega, Maria M. Delgado, Miguel De Gabriel-Hernando, Martin Päckert, Raphaël Arlettaz, Sebastian. Dirren, Philippe Fontanilles, Juan Antonio Gil, Mylene Herrmann, Sabine Hille, Fränzi Korner-Nievergelt, Paolo Pedrini, Jaime Resano-Mayor, Christian Schano and Davide Scridel

M. Brambilla (<https://orcid.org/0000-0002-7643-4652>) ✉ (mattia.brambilla@unimi.it), Univ. degli Studi di Milano Statale, Dip.to Scienze Politiche e Ambientali, Milano, Italy. – C. Bettega (<https://orcid.org/0000-0002-0814-0046>) ✉ (chiara.bettega@gmail.com) and P. Pedrini, MUSE-Museo delle Scienze, Corso del Lavoro e della Scienza 3, Trento, Italy. – M. M. Delgado (<https://orcid.org/0000-0002-3009-738X>), Biodiversity Research Inst. (IMIB, UO-CSIC-PA), Campus Mieres, Mieres, Spain. – M. De Gabriel-Hernando (<https://orcid.org/0000-0002-8722-6146>), Dept of Conservation Biology, Estación Biológica de Doñana-CSIC, Sevilla, Spain. – M. Päckert (<https://orcid.org/0000-0001-5045-0139>), Senckenberg Natural History Collections Dresden, Dresden, Germany. – R. Arlettaz, Division of Conservation Biology, Inst. of Ecology and Evolution, Univ. of Bern, Bern, Switzerland. – S. Dirren, F. Korner-Nievergelt (<https://orcid.org/0000-0001-9081-3563>) and C. Schano, Swiss Ornithological Inst., Sempach, Switzerland. – P. Fontanilles, Parc National des Pyrénées, Villa Fould, Tarbes, France. – J. A. Gil, Fundación para la Conservación del Quebrantahuesos, Zaragoza, Spain and Estación Zaragoza-Delicias, Grupo Aragón de Anillamiento Científico de Aves, Zaragoza, Spain. – M. Herrmann, Parc National de la Vanoise, Val d'Isère, France. – S. Hille, Univ. für Bodenkultur Wien, Wien, Austria. – J. Resano-Mayor, Grup d'anellament PARUS, Lliça d'Amunt, Barcelona, Spain. CS also at: Univ. of Zurich, Dept of Evolutionary Biology and Environmental Studies, Zurich, Switzerland. – D. Scridel, CNR-IRSA National Research Council – Water Research Institute, via del mulino 19, Brugherio (MB), Italy

Journal of Avian Biology

2022: e03015

doi: 10.1111/jav.03015

Subject Editor: Theresa M. Burg

Editor-in-Chief: Staffan Bensch

Accepted 7 June 2022



www.avianbiology.org

Keywords: distribution, ecological niche, *Montifringilla nivalis*, mountains, seasonality, snowfinch

Linking organism distribution to climate is key to understanding factors determining species occurrence and evaluating the potential impacts of ongoing climate change. A common analytical tool to assess the link between species and climate is represented by ecological niche modelling and by the tightly related species distribution models (SDMs). Those approaches have been widely used to explain key features related to species occurrence in relation to current climatic and other environmental variables, or as a response to past colonization and extinction events (Pons et al. 2021), as well as to predict the future distribution of target species (Scridel et al. 2021). Moreover, the link between climatic changes and niche evolution, i.e. niche changes (or conservatism) over time and especially across phylogenies remains complex and fascinating.

In the context of change, the rise of community science ('citizen science') data provides an invaluable opportunity to expand our knowledge on species distribution. While much attention has been given to aspects related to the statistical approaches and algorithm settings adopted in distribution or niche modelling (Engler et al. 2017, Ortega-Huerta and Rivera 2017, Kozma et al. 2018), less attention has been paid to validate very fundamental features which can greatly impact models performance, such

© 2022 The Authors. Journal of Avian Biology published by John Wiley & Sons Ltd on behalf of Nordic Society Oikos

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

as the quality of data for a given species, its spatial and temporal congruence with the climatic and environmental data used, and the risks associated with the inclusion of records of occasional vagrant individuals observed outside of their usual environment (but see Engler et al. 2017, Eyres et al 2017, Andrew and Fox 2020, Taheri et al. 2020).

Mountain birds and their ecological niche

Modelling the ecological niche of mountain bird species is particularly challenging. As they depend on dynamic habitats subject to large variation in local climate over small spatio-temporal scales (Scridel et al. 2018), they often show strong seasonal preferences during their annual cycle (Engler et al. 2014, Peña-Peniche et al. 2018). Moreover, they use a high diversity of movement strategies (e.g. elevational migration, nomadic or erratic movements), which are related to the very unpredictable seasonality of alpine environments (e.g. climate conditions and resource availability; Borrás et al. 2010, Henry 2011, Boyle 2018, Novoa et al. 2020, Resano-Mayor et al. 2020, Barras et al. 2021). Even for species thought to be largely resident or to move over short distances, there is some evidence of relatively long movements of individuals during the non-breeding season (Resano-Mayor et al. 2020), which might increase the chances of finding some transient individuals outside what could be regarded as the species' niche. Our knowledge of mountain bird movement strategies during the non-breeding period is still very limited and highlights the need to be particularly careful when assessing a species' ecological niche across the annual cycle. For these reasons, separate modelling of breeding versus non-breeding distribution should be required for mountain birds. Otherwise, model outputs may lead to misleading evaluations of species' ecological niches.

A worked example: the thermal niche of the white-winged snowfinch *Montifringilla nivalis*

The white-winged snowfinch *Montifringilla nivalis*, a well-known inhabitant of high alpine zones in Eurasia, is one of the species breeding at the highest elevations in the Western Palearctic (Cramp and Perrins 1994). In a recent paper, Cobos and colleagues (2021) implemented a framework to evaluate niche evolution and related range expansion in the so-called 'snowfinches' (Passeridae), a group of species specialized to high-elevations. They suggested that, while snowfinch species exhibit a cold thermal niche (i.e. average annual temperature from -8 to 7°C), the white-winged snowfinch, the only species that also occurs in Europe, exhibits a much broader thermal niche (i.e. average annual temperature from -9 to 20°C). According to the authors, such a wider niche should be related to a 'dramatic niche evolution event', leading to a niche expansion dated perhaps 2.6 mya ago, which allowed this species to occupy warmer areas in the western Palearctic,

such as southern Europe. However, we are convinced that the reason for such a finding is not an effective niche expansion in the species, but a question of methods – the inclusion of non-representative occurrence records in the analyses of the species' thermal niche –, neglecting crucial facts about the species' ecology.

There is striking evidence that white-winged snowfinches are strictly associated with cold alpine environments, in terms of their distribution (Brambilla et al. 2017a, 2020, Brambilla and Delgado 2020, BirdLife International 2021, de Gabriel Hernando et al. 2021), foraging areas (Antor et al. 1995, Brambilla et al. 2017b, 2018), social behaviour (Delgado et al. 2021), survival rates (Strinella et al. 2020) and habitat and nest-site selection (Heiniger 1991a, b, Brambilla et al. 2017b, Bettega et al. 2020, Niffenegger 2021). The dependence of this species on cold environments is particularly strong during the breeding season (Brambilla et al. 2019, 2022, Resano-Mayor et al. 2019, Schano et al. 2021, Alessandrini et al. 2022, de Gabriel Hernando et al. 2022): the breeding distribution of the species in European mountains is associated with low values of annual average temperature (Brambilla et al. 2017a, 2020, 2022, de Gabriel Hernando et al. 2021), suggesting a thermal niche much narrower than that suggested by Cobos et al. (2021). Similarly, during winter the species appears almost invariably tied to mountain sites above the treeline (Heiniger 1991a, Bettega et al. 2020, Resano-Mayor et al. 2020, Delgado et al. 2021, de Gabriel Hernando et al. 2021; see also the wintering distribution according to BirdLife International 2021), with season-specific distribution models showing how suitable areas during the cold season extend only partly to lower mountain ranges (de Gabriel Hernando et al. 2021). In fact, unlike other typical alpine species like wheatears, water pipits and alpine choughs, observations in warmer areas at low elevation are extremely rare despite the potential suitability of habitat (Knaus et al. 2018) and are more likely to occur in the case of prolonged, exceptionally cold or snowy periods and mostly linked to food shortage at higher elevation (Cramp and Perrins 1994, Glutz von Blotzheim and Bauer 1997). Such evidence contrasts with Cobos et al. (2021) conclusions about the thermal niche of white-winged snowfinches.

In the work of Cobos et al. (2021), seasonality has not been taken into account, and there is a lack of phenological concordance between records and climate data. The authors have pooled community science records related to different time of the year, and used climate data over the period 1979–2013 without considering inter-seasonal climatic variability, but showing results based only on annual average temperature. For mobile species inhabiting highly seasonal environments, such as the white-winged snowfinch in European mountains, this could be strongly misleading (Engler et al. 2017, Andrew and Fox 2020, de Gabriel Hernando et al. 2021). The same set of annual climatic variables could be used to model species distribution in different seasons, but records need to be split as the relationship between species occurrence and the same climatic predictor would be different when considering different periods of the annual cycle

(de Gabriel Hernando et al. 2021). For example, by including occasional observations in warm location, most likely to occur in winter, the annual mean temperature would be overestimated. This likely happened for snowfinch records in Europe and Western Asia (Supporting information) used by Cobos et al. (2021). In this line, we noticed that the authors included a relatively large number of unusual records located in lower and warmer areas (approx. 10% of records were below 1000 m), which we believe are most likely attributable to vagrant birds occasionally visiting areas outside the species' range, if not to incorrect identification/reporting (e.g. records in central and southwestern Iberian Peninsula and southern Italy). We acknowledge that community science data are an emergent, increasingly important tool contributing to ecological research and conservation, and they have been frequently used (also by many of us) for snowfinches as well as for other mountain species (Delgado et al. 2021, de Gabriel Hernando et al. 2021, Schano et al. 2021, Scridel et al. 2021, Brambilla et al. 2022). However, an accurate filtering of observations to remove potentially biased records, based not only on spatial locations and species identification but also on the basic ecology of the species, is crucial to improve model performance and outcome reliability (Steen et al. 2019, Johnston et al. 2021, Scridel et al. 2021, Brambilla et al. 2022).

Although we recognize the important aims of Cobos et al. (2021), we find their conclusions about the broad thermal niche of white-winged snowfinches inconsistent with the species' ecology and distribution. This overestimation of the species' thermal niche and distribution is potentially detrimental for a proper appreciation of the conservation status of the species and its sensitivity to global warming. The white-winged snowfinch is currently classified as a species of 'least concern' in the European and Global IUCN Red Lists and only Spain and Switzerland have recently changed its status to 'nearly threatened' (Knaus et al. 2021, Laiolo et al. 2021). Such an underrated status for a species highly threatened by climate change is likely partly due to the poorly known population trends. In fact, the ongoing range contraction (Scridel et al. 2017, Patrinat 2019, Brambilla and Delgado 2020), the evidence for regional declines (Knaus et al. 2018), the sensitivity to warming (Strinella et al. 2020) and the habitat changes it induces (Brambilla et al. 2018), as well as the forecast dramatic decline that could take place in the next decades (de Gabriel Hernando et al. 2021), suggest this to be one of the species most threatened by climate change in Europe (Brambilla et al. 2018, Knaus et al. 2021, Laiolo et al. 2021, Schano et al. 2021).

Conclusion

Modelling the ecological niche of organisms with seasonal distributions and/or high mobility requires integrating complex ecological aspects, and for many bird species a season-specific approach (Martínez-Meyer et al. 2004, Engler et al. 2014, Eyres et al. 2017). Thanks to their extraordinary mobility, birds

can use different habitats to track suitable climate or resources across seasons (Engler et al. 2017, and references therein), thus occasionally occur outside their usual environments. Species–climate relationships may hence vary across time and space, or according to the climate parameters considered (e.g. annual versus seasonal temperature): neglecting these potential dynamics could lead to wrong conclusions regarding distribution and niche characteristics, as well as to overpredicted ranges (Reside et al. 2010). The careful selection of reliable, spatially accurate records, and the use of accurate information about climate and life history stages is key to model the actual species–climate relationship, while properly considering its seasonality, especially in mountain regions, where abrupt changes may occur over limited spatial and temporal spans.

Acknowledgements – We thank the European Snowfinch Group for all the efforts in monitoring and studying the snowfinch.

Funding – The authors have no funding to declare.

Author contributions

Mattia Brambilla, Chiara Bettega, Maria Delgado, Davide Scridel: Conceptualization (lead); Writing – original draft (lead); Writing – review and editing (lead). **M. De Gabriel-Hernando, M. Päckert, S. Hille, F. Korner-Nievergelt, C. Schano, R. Arlettaz, S. Dirren, P. Fontanilles, J. A. Gil, M. Herrmann, P. Pedrini, J. Resano-Mayor:** Writing – review and editing (supporting).

Transparent peer review

The peer review history for this article is available at <<https://publons.com/publon/10.1111/jav.03015>>.

Data availability statement

This paper contain no original data.

Supporting information

The Supporting information associated with this article is available with the online version.

References

- Alessandrini, C., Scridel, D., Boitani, L., Pedrini, P. and Brambilla, M. 2022. Remotely sensed variables explain microhabitat selection and reveal buffering behaviours against warming in a climate-sensitive bird species. – *Remote Sens. Ecol. Conserv.* doi: 10.1002/rse2.265.
- Andrew, M. E. and Fox, E. 2020. Modelling species distribution in dynamic landscapes: the importance of the temporal dimension. – *J. Biogeogr.* 47: 1510–1529.
- Antor, R. J. 1995. The importance of arthropod fallout on snow patches for the foraging of high-alpine birds. – *J. Avian Biol.* 26: 81–85.

- Barras, A. G., Liechti, F. and Arlettaz, R. 2021. Seasonal and daily movement patterns of an Alpine passerine suggest high flexibility in relation to environmental conditions. – *J. Avian Biol.* 52: e02860.
- Bettega, C., Fernandez-Gonzalez, A., Ramon Obeso, J. and Delgado, M. D. M. 2020. Circannual variation in habitat use of the white-winged snowfinch *Montifringilla nivalis nivalis*. – *Ibis* 162: 1251–1261.
- BirdLife International 2021. Species factsheet: *Montifringilla nivalis*. –
- Borras, A., Senar, J. C., Alba-Sánchez, F., López-Sáez, J. A., Cabrera, J., Colomer, X. and Cabrera, T. 2010. Citril finches during the winter: patterns of distribution, the role of pines and implications for the conservation of the species. – *Anim. Biodiv. Conserv.* 33: 89–115.
- Boyle, A. W. 2018. Altitudinal bird migration in North America. – *Auk* 134: 443–465.
- Brambilla, M. and Delgado, M. M. 2020. *Montifringilla nivalis* white-winged snowfinch. – In: Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanesi, P., Martí, D., Anton, M., Klvaňová, A., Kalyakin, M. V., Bauer, H.-G. and Foppen, R. P. B. (eds), *European Breeding Bird Atlas 2: distribution, abundance and change*. European Bird Census Council & Lynx Edicions, pp. 794–795.
- Brambilla, M., Caprio, E., Assandri, G., Scridel, D., Bassi, E., Bionda, R., Celada, C., Falco, R., Bogliani, G., Pedrini, P., Rolando, A. and Chamberlain, D. 2017a. A spatially explicit definition of conservation priorities according to population resistance and resilience, species importance and level of threat in a changing climate. – *Divers. Distrib.* 23: 727–738.
- Brambilla, M., Cortesi, M., Capelli, F., Chamberlain, D., Pedrini, P. and Rubolini, D. 2017b. Foraging habitat selection by Alpine white-winged snowfinches *Montifringilla nivalis* during the nestling rearing period. – *J. Ornithol.* 158: 277–286.
- Brambilla, M., Resano-Mayor, J., Arlettaz, R., Bettega, C., Binggeli, A., Bogliani, G., Braunisch, V., Celada, C., Chamberlain, D., Chiffard Carricaburu, J., Delgado, M. M., Fontanilles, P., Kmecl, P., Korner, F., Lindner, R., Pedrini, P., Pöhacker, J., Rubinic, B., Schano, C., Scridel, D., Strinella, E., Teufelbauer, N. and De Gabriel Hernando, M. 2020. Potential distribution of a climate sensitive species, the white-winged snowfinch *Montifringilla nivalis* in Europe. – *Bird Conserv. Int.* 30: 522–532.
- Brambilla, M., Resano-Mayor, J., Scridel, D., Anderle, M., Bogliani, G., Braunisch, V., Capelli, F., Cortesi, M., Horrenberger, N., Pedrini, P., Sangalli, B., Chamberlain, D., Arlettaz, R. and Rubolini, D. 2018. Past and future impact of climate change on foraging habitat suitability in a high-alpine bird species: management options to buffer against global warming effects. – *Biol. Conserv.* 221: 209–218.
- Brambilla, M., Rubolini, D., Appukuttan, O., Calvi, G., Karger, D. N., Kmecl, P., Mihelič, T., Sattler, T., Seaman, B., Teufelbauer, N., Wahl, J. and Celada, C., , 2022. Identifying climate refugia for high-elevation Alpine birds under current climate warming predictions. – *Global Change Biol.* 28: 4276–4291.
- Brambilla, M., Scridel, D., Sangalli, B., Capelli, F., Pedrini, P., Bogliani, G. and Rubolini, D. 2019. Ecological factors affecting foraging behaviour during nestling rearing in a high-elevation species, the white-winged snowfinch *Montifringilla nivalis*. – *Ornis Fenn.* 96: 142–151.
- Cobos, M. E., Cheng, Y., Song, G., Lei, F. and Peterson, T. A. 2021. New distributional opportunities with niche innovation in Eurasian snowfinches. – *J. Avian Biol.* 52: e02868.
- Cramp, S. and Perrins, C. M. 1994. *Handbook of the birds of Europe the Middle East and North Africa. The birds of the Western Palearctic. Volume VIII. Crows to finches.* – Oxford Univ. Press.
- de Gabriel Hernando, M., Fernández-Gil, J., Roa, I., Juan, J., Ortega, F., de la Calzada, F. and Revilla, E. 2021. Warming threatens habitat suitability and breeding occupancy of rear-edge alpine bird specialists. – *Ecography* 44: 1–14.
- de Gabriel Hernando, M., Roa, I., Fernández-Gil, J., Juan, J., Fuertes, B., Reguera, B. and Revilla, E. 2022. Trends in weather conditions favor generalist over specialist species in rear-edge alpine bird communities. – *Ecosphere* 13: e3953.
- Delgado, M. M., Arlettaz, R., Bettega, C., Brambilla, M., de Gabriel Hernando, M., España, A., Fernández-González, Á., Fernández-Martín, Á., Gil, J. A., Hernández-Gómez, S., Laiolo, P., Resano-Mayor, J., Obeso, J. R., Pedrini, P., Roa-Álvarez, I., Schano, C., Scridel, D., Strinella, E., Toranzo, I. and Korner-Nievergelt, F. 2021. Spatio-temporal variation in the wintering associations of an alpine bird. – *Proc. R. Soc. B* 288: 20210690.
- Engler, J., Rödder, D., Stiels, D. and Förschler, M. I. 2014. Suitable, reachable but not colonised: seasonal niche duality in an endemic mountainous songbird. – *J. Ornithol.* 155: 657–669.
- Engler, J. O., Stiels, D., Schidelko, K., Strubbe, D., Quillfeldt, P. and Brambilla, M. 2017. Avian SDMs: current state, challenges and opportunities. – *J. Avian Biol.* 48: 1483–1504.
- Eyres, A., Böhning-Gaese, K. and Fritz, S. A. 2017. Quantification of climatic niches in birds: adding the temporal dimension. – *J. Avian Biol.* 48: 1517–1531.
- Glutz von Blotzheim, U. N. and Bauer, K. M. 1997. *Handbuch der Vögel Mitteleuropas. Band 14/I. Passeriformes (5. teil).* – Aula Verlag.
- Heiniger, P. 1991a. Adaptations of the snow finch *Montifringilla nivalis* to his cold-dominated alpine environment. – *Der Ornithol. Beobachter* 88: 193–207.
- Heiniger, P. 1991b. Ecology of the snowfinch *Montifringilla nivalis* – use of home range in winter and summer with special reference to the winter roosting sites. – *Revue Suisse Zool.* 98: 897–924.
- Henry, P.-Y. 2011. Differential migration in the polygynandrous Alpine accentor *Prunella collaris*. – *Bird Study* 58: 160–170.
- Johnston, A., Hochachka, W. M., Strimas-Mackey, M. E., Gutierrez, V. R., Robinson, O. J., Miller, E. T., Auer, T., Kelling, S. T. and Fink, D. 2021. Analytical guidelines to increase the value of community science data: an example using eBird data to estimate species distributions. – *Divers. Distrib.* 27: 1265–1277.
- Knaus, P., Antoniazza, S., Keller, V., Sattler, T., Schmid, H., Guelat, J. and Strebel, N. 2021. Rote Liste 2021 der Brutvögel: Grundlagen, Hintergründe der Einstufungen und Dokumentation der Arten. – Schweizerische Vogelwarte.
- Knaus, P., Antoniazza, S., Wechsler, S., Guelat, J., Kery, M., Strebel, N. and Sattler, T. 2018. *Swiss Breeding Bird Atlas 2013–2016. Distribution and population trends of birds in Switzerland and Liechtenstein.* – Swiss Ornithol. Inst.
- Kozma, R., Lillie, M., Benito, B. M., Svenning, J. C. and Höglund, J. 2018. Past and potential future population dynamics of three grouse species using ecological and whole genome coalescent modeling. – *Ecol. Evol.* 8: 6671–6681.
- Laiolo, P., Delgado, M. M., López-Bao, J. V. and Obeso, J. R. 2021. Gorrión alpino, *Montifringilla nivalis*. – In: López-Jiménez, N. (ed.), *Libro Rojo de las Aves de España*. SEO/BirdLife, pp. 819–820.

- Martínez-Meyer, E., Peterson, A. T. and Navarro-Sigüenza, A. 2004. Evolution of seasonal ecological niches in the Passerina buntings (Aves: Cardinalidae). – Proc. R. Soc. B 271: 1151–1157.
- Niffenegger, C. 2021. Nest site selection of the white-winged snowfinch *Montifringilla nivalis* in the Swiss Alps. – MS thesis, Swiss Ornithol. Inst., Sempach and Division of Conservation Biology, Univ. of Bern.
- Novoa, C., Resseguier, J., Muffat-Joly, B., Blanch Casadesus, J., Arvin-Bérod, M., Gracia Moya, J. and Desmet, J.-F. 2020. Natal dispersal and survival of juvenile rock ptarmigan *Lagopus muta* in the French Alps and Pyrenees. – Ardeola 68: 123–141.
- Ortega-Huerta, M. A. and Vega-Rivera, J. H. 2017. Validating distribution models for twelve endemic bird species of tropical dry forest in western Mexico. – Ecol. Evol. 7: 7672–7686.
- Peña-Peniche, A., Ruvalcaba-Ortega, I. and Rojas-Soto, O. 2018. Climate complexity in the migratory cycle of *Ammodramus bairdii*. – PLoS One 13: e0202678.
- Pons, J.-M., Campián, D., Chiozzi, G., Ettwein, A., Grangé, J.-L., Kajtoch, Ł., Mazgajski, T. D., Rakovic, M. Winkler, H. and Fuchs, J. 2021. Phylogeography of a widespread Palaearctic forest bird species: the white-backed woodpecker (Aves, Picidae). – Zool. Scr. 50: 155–172.
- Resano-Mayor, J., Bettega, C., Delgado, M. M., Fernández-Martín, A., Hernández-Gómez, S., Toranzo, I., España, A., de Gabriel, M., Roa-Alvarez, I., Gil, J. A., Strinella, E., Hobson, K. A. and Arlettaz, R. 2020. Partial migration of white-winged snowfinches is correlated with winter weather conditions. – Global Ecol. Conserv. 24: e01346.
- Resano-Mayor, J., Korner-Nievergelt, F., Vignali, S., Horrenberger, N., Barras, A. G., Braunisch, V., Pernollet, C. A. and Arlettaz, R. 2019. Snow cover phenology is the main driver of foraging habitat selection for a high-alpine passerine during breeding: implications for species persistence in the face of climate change. – Biodivers. Conserv. 28: 2669e2685.
- Reside, A. E., VanDerWal, J. J., Kutt, A. S. and Perkins, G. C. 2010. Weather, not climate, defines distributions of vagile bird species. – PLoS One 5: e13569.
- Schano, C., Niffenegger, C., Jonas, T. and Korner-Nievergelt, F. 2021. Hatching phenology is lagging behind an advancing snowmelt pattern in a high-alpine bird. – Sci. Rep. 11: 22191.
- Scridel, D., Bogliani, G., Pedrini, P., Iemma, A., von Hardenberg, A. and Brambilla, M. 2017. Thermal niche predicts recent changes in range size for bird species. – Clim. Res. 73: 207–216.
- Scridel, D., Brambilla, M., de Zwaan, D., Froese, N., Wilson, S., Pedrini, P. and Martin, K. 2021. A genus at risk: predicted current and future distribution of all three *Lagopus* species reveal sensitivity to climate change and efficacy of protected areas. – Divers. Distrib. 27: 1759–1774.
- Scridel, D., Brambilla, M., Martin, K., Lehikoinen, A., Iemma, A., Matteo, A., Jähnig, S., Caprio, E., Bogliani, G., Pedrini, P., Rolando, A., Arlettaz, R. and Chamberlain, D. 2018. A review and meta-analysis of the effects of climate change on Holarctic mountain and upland bird populations. – Ibis 160: 489–515.
- Steen, V. A., Elphick, C. S. and Tingley, M. W. 2019. An evaluation of stringent filtering to improve species distribution models from citizen science data. – Divers. Distrib. 25: 1857–1869.
- Strinella, E., Scridel, D., Brambilla, M., Schano, C. and Korner-Nievergelt, F. 2020. Potential sex-dependent effects of weather on apparent survival of a high elevation specialist. – Sci. Rep. 10: 8386.
- Taheri, S., García-Callejas, D. and Araújo, M. B. 2020. Discriminating climate, land-cover and random effects on species range dynamics. – Global Change Biol. 27: 1309–1317.
- UMS Patrinat (coord.) 2019. Résultats synthétiques de l'évaluation des statuts et tendances des espèces d'oiseaux sauvages en France, période 2013–2018. – Rapportage article 12 envoyé à la Commission européenne.