

- JJ, Biemond BJ. Sick cell patients are characterized by a reduced glycocalyx volume. *Haematologica* 2008;93:307-8.
6. Kemna EH, Tjalsma H, Podust VN, Swinkels DW. Mass spectrometry-based hepcidin measurements in serum and urine: analytical aspects and clinical implications. *Clin Chem* 2007;53:620-8.
 7. Swinkels DW, Girelli D, Laarakkers C, Kroot J, Camprostrini N, Kemna EH, et al. Advances in quantitative hepcidin measurements by time-of-flight mass spectrometry. *PLoS ONE* 2008;3:e2706.
 8. Kemna EH, Kartikasari AE, van Tits LJ, Pickkers P, Tjalsma H, Swinkels DW. Regulation of hepcidin: insights from biochemical analyses on human serum samples. *Blood Cells Mol Dis* 2008;40:339-46.
 9. Swinkels DW, Drenth JP. Hepcidin in the management of patients with mild non-hemochromatotic iron overload: Fact or fiction? *J Hepatol* 2008;49:680-5.
 10. Bekri S, Gual P, Anty R, Luciani N, Dahman M, Ramesh B, et al. Increased adipose tissue expression of hepcidin in severe obesity is independent from diabetes and NASH. *Gastroenterology* 2006;131:788-96.

Predictive value of $\beta 2$ -microglobulin ($\beta 2$ -m) levels in chronic lymphocytic leukemia since Binet A stages

We read with interest the study by Rossi and co-workers, reporting CD49d expression as risk factor of treatment free survival (TFS) in Binet A CLL patients.¹ In this paper, a close association between CD49d and CD38, LDH and $\beta 2$ -m is also described. We would like to add further information about the prognostic power of $\beta 2$ -m. It is generally believed that $\beta 2$ -m is released constitutively by CLL cells and that its level approximately correlates with tumor mass.² Based on these premises the predictive value of $\beta 2$ -m serum concentration could vary in the course of the disease and be relatively low in the early disease stages, when tumor mass is low, irrespective of the subsequent clinical outcome. Therefore, $\beta 2$ -m determination could exhibit a lower predictive power particularly at the early disease stages compared to the newer biological markers, such as IgVH gene status, ZAP-70 and CD38, which represent intrinsic cell features that can be determined since the earliest disease stages and never (IgVH) or rarely (ZAP-70 and CD38) change in the course of the disease.³

In order to explore this issue, we have measured $\beta 2$ -m value in 222 Binet stage A patients at diagnosis. IgVH gene status and CD38 expression were also determined in all cases studied. Unlike $\beta 2$ -m, which was measured at diagnosis, these markers were determined in the course of the disease when marker determinations became available. This approach, although irrelevant for the IgVH gene status, may introduce some, albeit minor, biases for CD38 for the reasons alluded to above. The median $\beta 2$ -m value was 2 mg/dL (range 0.4-19). ROC analysis determined that the cut-off value capable of discriminating between patients whose disease progressed and required treatment from those with stable disease was 2.4 mg/dL (AUC:0.67, $p=0.005$). Accordingly 149/222 patients (67%) were $\beta 2$ -m^{neg} and 73/222 (33%) as $\beta 2$ -m^{pos}. Overall, the results did not substantially change when arbitrary cut offs used by other authors⁴⁻⁷ were employed.

The patients' features are summarized in Table 1. $\beta 2$ -m levels overlap with CD38 expression in 128/219 cases (63%) [$\beta 2$ -m^{pos}/CD38 $\geq 30\%$ cases: 23/55 (41.8%), $\beta 2$ -m^{neg}/CD38<30% cases: 115/164 (70.1%)], while β -m

levels overlap with IgVH status in 125/195 cases (64.1%) [$\beta 2$ -m^{pos}/IgVHunmutated cases: 29/62 (46.8%), $\beta 2$ -m^{neg}/IgVHmutated: 96/133 (72.2%)]. Finally, the concordance between CD38 expression and IgVH mutational status was 77.6% (149/192 cases) [IgVHunmutated/CD38 $\geq 30\%$ cases: 35/52 (67.3%), IgVHmutated/CD38<30% cases: 114/140 (81.4%)].

After a median follow-up of 3.5 years, 55 of 222 Binet stage A (25%) required treatment. $\beta 2$ -m^{neg} cases showed a significantly longer TFS than $\beta 2$ -m^{pos} cases; in particular the projected median TFS was 5.3 years for $\beta 2$ -m^{pos} versus not reached for $\beta 2$ -m^{neg} (Figure 1A). TFS represented a reliable measure of disease progression since all centers agreed to follow NCI guidelines for treatment start.

In order to ascertain whether $\beta 2$ -m identifies a patient subset of those with good prognostic markers, we calculated TFS of both CD38<30% and IgVHmutated CLL cases grouped according to the $\beta 2$ -m expression. $\beta 2$ -m^{pos} CD38<30% cases exhibited a TFS which was significantly lower than that of $\beta 2$ -m^{neg} CD38<30% cases (3.5-years TFS probability: $\beta 2$ -m^{neg} vs. $\beta 2$ -m^{pos} 91% vs. 83%; $p=0.05$). However, these differences were not seen in the IgVHmutated cases (3.5-years TFS probability: $\beta 2$ -m^{neg} vs. $\beta 2$ -m^{pos} 89% vs. 84%; $p=ns$).

At Cox univariate analysis, $\beta 2$ -m^{pos} (HR:2.3, $p=0.003$), CD38>30% (HR:3.9, $p<0.0001$) and IgVHunmutated (HR:3.2, $p<0.0001$) showed a statistically significant impact on TFS. At Cox multivariate analysis, all the three markers maintained an independent prognostic impact ($\beta 2$ -m^{pos}, HR:1.8, $p=0.047$; CD38>30%, HR:2.0, $p=0.03$; IgVHunmutated, HR:2.7, $p=0.022$). When a scoring system in which one point was assigned to each unfavorable prognostic marker was utilized, the risk of an early treatment was highest (Figure 1B) in patients presenting all the three adverse prognostic markers. Cases with two, one or none of the unfavorable prognostic factors showed lower risk for an early treatment (Figure 1C).

Collectively, this study shows that $\beta 2$ -m levels represent valuable predictors in early CLL stages, when the neoplastic cell burden is low. This finding raises a number of questions regarding the mechanisms governing the $\beta 2$ -m levels. This molecule is constantly shedded⁸

Table 1. Comparisons of clinical and laboratory features among chronic lymphocytic leukemia patients devised according to $\beta 2$ -m expression.

	All patients	$\beta 2$ -m <2.4 mg/dL	$\beta 2$ -m ≥ 2.4 mg/dL	p
N. of patients	222	149 (67)	73 (33)	
Age (years)				
≤65	124 (56)	94 (63)	30 (41)	0.002
>65	98 (44)	55 (37)	43 (59)	
Gender				
Female	82 (37)	60 (40)	22 (30)	0.14
Male	140 (63)	99 (60)	51 (70)	
IgVH mutational status (n=195)				
Mutated	133 (68)	96 (74)	37 (56)	0.014
Germline	62 (32)	33 (26)	29 (44)	
CD38 expression (n=219)				
<30%	164 (75)	115 (78)	42 (58)	0.02
$\geq 30\%$	55 (25)	32 (22)	30 (42)	
Therapy				
no	167 (75)	123 (83)	44 (60)	<0.0001
yes	55 (25)	26 (17)	29 (40)	




Figure 1. (A) Kaplan-Meier estimates of treatment-free survival (TFS). Comparison between $\beta 2\text{-m}^{\text{neg}}$ patients versus $\beta 2\text{-m}^{\text{pos}}$ patients. (B) Cox univariate analysis for the predictive value of marker combinations for the risk categories. (C) Cox derived estimated TTT curves according to the combination of the three prognostic factors.

by lymphocytes and it is expected that its levels steadily increase together with the progressive expansion of the leukemic clone suggesting a close correlation between stage (which measures tumor burden) and $\beta 2\text{-m}$ levels. Although a correlation with disease stage likely exists, there was a substantial proportion of patients with high $\beta 2\text{-m}$ levels already at Binet A stage (low tumor burden). Possibly, CLL cells from these patients are more activated *in vivo* and shed more abundant $\beta 2\text{-m}$. Taken all the above into consideration, the data indicate that the role of $\beta 2\text{-m}$ as a prognostic tool should be re-evaluated possibly in prospective studies involving large patient cohorts.

Massimo Gentile,¹ Giovanna Cutrona,² Antonino Neri,³ Stefano Molica,⁴ Manlio Ferrarini,^{2,5} and Fortunato Morabito¹

¹Unità Operativa Complessa di Ematologia, Azienda Ospedaliera di Cosenza, Cosenza; ²Divisione di Oncologia Medica C, Istituto Nazionale per la Ricerca sul Cancro, IST, Genova; ³Centro per lo Studio delle Leucemie, Dipartimento di Scienze Mediche, Università di Milano, Unità Operativa di Ematologia 1, Fondazione IRCCS Policlinico, Milano; ⁴Dipartimento di Oncologia/Ematologia, Azienda Ospedaliera

Pugliese-Ciaccio, Catanzaro; ⁵Dipartimento di Oncologia, Biologia e Genetica Università degli Studi di Genova, Italy.

Funding: supported from Associazione Italiana Ricerca sul Cancro (AIRC) (to FM and MF) and Fondazione 'Amelia Scorza' Onlus, Cosenza, Italy.

Acknowledgments: we would like to acknowledge Dr. Vincenzo Callea, Prof Luca Baldini, Dr Ugo Consoli and Dr Serena Matis for their contribution and useful suggestions. We thank Laura Veroni and Brigida Gulino for precious secretarial assistance.

Key words: $\beta 2$ -microglobulin, CD38, IgVH mutational status, CLL, prognosis.

Correspondence: Fortunato Morabito, Unità Operativa Complessa di Ematologia, Dipartimento di Medicina Interna, Azienda Ospedaliera di Cosenza, Viale della Repubblica, 87100 Cosenza, Italy. Phone: international +39.0984.681329. Fax: international +39.0984.791751. E-mail: fortunato_morabito@tin.it

Citation: Gentile M, Cutrona G, Neri A, Molica S, Ferrarini M, and Morabito F. Predictive value of B2-microglobulin ($\beta 2\text{-m}$) levels in chronic lymphocytic leukemia since Binet A stages. *Haematologica* 2009; 94:887-888. doi:10.3324/haematol.2009.005561

References

- Rossi D, Zucchetto A, Rossi FM, Capello D, Cerri M, Deambrogi C, et al. CD49d expression is an independent risk factor of progressive disease in early stage chronic lymphocytic leukemia. *Haematologica* 2008;93:1575-9.
- Simonsson B, Wibell L, Nilsson K. $\beta 2$ -microglobulin in chronic lymphocytic leukaemia. *Scand J Haematol* 1980; 24:174-80.
- Gentile M, Mauro FR, Guarini A, Foà R. New developments in the diagnosis, prognosis and treatment of chronic lymphocytic leukemia. *Curr Opin Oncol* 2005;17:597-604.
- Keating MJ, Lerner S, Kantarjian H, Freireich EJ, O'Brien S. The serum $\beta 2$ -microglobulin ($\beta 2\text{m}$) level is more powerful than stage in predicting response and survival in chronic lymphocytic leukemia (CLL). *Blood* 1995;86 (Suppl 1):606a[Abstract].
- Wierda WG, O'Brien S, Wang X, Faderl S, Ferrajoli A, Do KA, et al. Prognostic nomogram and index for overall survival in previously untreated patients with chronic lymphocytic leukemia. *Blood* 2007;109:4679-85.
- Del Poeta G, Maurillo L, Venditti A, Buccisano F, Episcopo AM, Capelli G, et al. Clinical significance of CD38 expression in chronic lymphocytic leukemia. *Blood* 2001; 98:2633-9.
- Desablens B, Garidi R, Lamour C. Prognostic value of LDH and $\beta 2$ microglobuline in B CLL. Analysis of 275 patients. *Hematol Cell Ther* 1997;39:S93.
- Vilpo J, Vilpo L, Hurme M, Vuorinen P. Induction of $\beta 2$ -microglobulin release in vitro by chronic lymphocytic leukaemia cells: relation to total protein synthesis. *Leuk Res* 1999;23:913-20.

Detection of continuous erythropoietin receptor activator in blood and urine in anti-doping control

Anti-doping control of erythropoietin (Epo) relies on the differentiation by isoelectric profile of natural endogenous hormone from the recombinant hormone used for doping. The first and second generations of recombinant Epo were detectable in urine.^{1,2} The third generation, Continuous Erythropoietin Receptor Activator (CERA), was obtained by linking a methoxy polyethylene glycol to epoetin β , a first generation rHuEPO, resulting in significantly greater stability in blood. CERA, approved in Europe in July 2007, was