

Sensitivity of the electric dipole polarizability to the neutron skin thickness in 208Pb

X. Roca-Maza, B. K. Agrawal, G. Colò, W. Nazarewicz, N. Paar et al.

Citation: *AIP Conf. Proc.* **1491**, 204 (2012); doi: 10.1063/1.4764239

View online: <http://dx.doi.org/10.1063/1.4764239>

View Table of Contents: <http://proceedings.aip.org/dbt/dbt.jsp?KEY=APCPCS&Volume=1491&Issue=1>

Published by the [American Institute of Physics](#).

Related Articles

Fermion realization of the nuclear Sp(6,R) model

J. Math. Phys. **39**, 5123 (1998)

Application of a distributed nucleus approximation in grid based minimization of the Kohn–Sham energy functional

J. Chem. Phys. **103**, 227 (1995)

Additional information on AIP Conf. Proc.

Journal Homepage: <http://proceedings.aip.org/>

Journal Information: http://proceedings.aip.org/about/about_the_proceedings

Top downloads: http://proceedings.aip.org/dbt/most_downloaded.jsp?KEY=APCPCS

Information for Authors: http://proceedings.aip.org/authors/information_for_authors

ADVERTISEMENT

**AIPAdvances**

Submit Now

**Explore AIP's new
open-access journal**

- **Article-level metrics
now available**
- **Join the conversation!
Rate & comment on articles**

Sensitivity of the Electric Dipole Polarizability to the Neutron Skin Thickness in ^{208}Pb

X. Roca-Maza*, B. K. Agrawal†, G. Colò**,‡, W. Nazarewicz§,||, N. Paar††, J. Piekarewicz‡‡, P.-G. Reinhard§§ and D. Vretenar††

*INFN, sezione di Milano, via Celoria 16, I-20133 Milano, Italy

†Saha Institute of Nuclear Physics, Kolkata 700064, India

**Dipartimento di Fisica, Università degli Studi di Milano

‡INFN, Sezione di Milano, 20133 Milano, Italy

§Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA

¶Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

||Institute of Theoretical Physics, University of Warsaw, Hoza 69, PL-00-681 Warsaw, Poland

††Physics Department, Faculty of Science, University of Zagreb, Zagreb, Croatia

‡‡Department of Physics, Florida State University, Tallahassee, FL 32306, USA

§§Institut für Theoretische Physik II, Universität Erlangen-Nürnberg, Staudtstrasse 7, D-91058 Erlangen, Germany

Abstract. The static dipole polarizability, α_D , in ^{208}Pb has been recently measured with high-resolution via proton inelastic scattering at the Research Center for Nuclear Physics (RCNP) [1]. This observable is thought to be intimately connected with the neutron skin thickness, r_{skin} , of the same nucleus and, more fundamentally, it is believed to be associated with the density dependence of the nuclear symmetry energy. The impact of r_{skin} on α_D in ^{208}Pb is investigated and discussed on the basis of a large and representative set of relativistic and non-relativistic nuclear energy density functionals (EDF) [2].

Keywords: electric dipole polarizability, neutron nuclear symmetry energy, energy density functionals

PACS: 21.60.Jz, 21.65.Cd, 21.65.Mn

INTRODUCTION

The Lead Radius Experiment (PREX) [3, 4] has recently measured r_{skin} , defined as the difference between the neutron and proton root mean square radii, of ^{208}Pb [5]. This experiment is performed via parity-violating electron scattering [6] and provides the first purely electroweak measurement of the neutron distribution of a heavy nucleus. The neutron skin is strongly dependent on the isovector properties of nuclei and impacts on a variety of areas such as nuclear structure [7, 8, 9, 10, 11], atomic parity violation [12], and neutron-star structure [13, 14]. By measuring the neutron form factor of ^{208}Pb at $q \approx 0.475 \text{ fm}^{-1}$, PREX was able to determine $r_{\text{skin}} = 0.33_{-0.18}^{+0.16} \text{ fm}$ [5].

Alternatively, although the estimation of the neutron distribution in nuclei based on measurements using hadronic probes are model dependent and display large theoretical uncertainties [15, 16], the use of these probes for the direct or indirect determination of such an observable is nowadays growing due to the necessity of improving our knowledge in the isovector channel of the nuclear effective interaction [17, 18, 19, 20, 21]. The analysis from recent experiments have led to values for $r_{\text{skin}} = 0.16 \pm$

$(0.02)_{\text{stat}} \pm (0.04)_{\text{syst}}$ fm [18] and $r_{\text{skin}} = 0.211^{+0.054}_{-0.063}$ fm [21] in ^{208}Pb .

The electric dipole polarizability, α_{D} , is another observable sensitive to the isovector properties of the nuclear effective interaction. This quantity is obtainable from the linear response of the system to an external dipolar field of the form, $F_{\text{D}} = (Z/A) \sum_i^N r_n Y_{1M}(\hat{r}_n) - (N/A) \sum_i^Z r_p Y_{1M}(\hat{r}_p)$, being N , Z , A , r_j with $j = n$ or p and $Y(\hat{r}_j)$ the neutron, proton and mass numbers, the radial position of the j -th nucleon and the spherical harmonic, respectively. If $|0\rangle$ is the ground state and $|v\rangle$ is a complete set of excited states, the polarizability can be written as follows, $\alpha_{\text{D}} = (8\pi/9)e^2 m_{-1} = (8\pi/9)e^2 \sum_v (|\langle v|F_{\text{D}}|0\rangle|^2 / \omega_v)$ where m_{-1} is the inverse energy weighted sum rule or inverse energy moment of the strength function, $R_{E1}(\omega) = \sum_v |\langle v|F_{\text{D}}|0\rangle|^2 \delta(\omega - \omega_v)$ which evaluates the dipole response. For stable medium and heavy nuclei, the dipole response is largely concentrated in the giant dipole resonance (GDR) [22]. In this isovector mode, commonly viewed as an oscillation of neutrons against protons, the symmetry energy at some sub-saturation density acts as the restoring force [23]. In addition, it is important to notice that the possible presence of a low-lying dipole strength may have a non negligible effect on α_{D} [24, 25, 26].

Actually, r_{skin} is expected to be linearly correlated with α_{D} based on both macroscopic arguments [27, 28] and microscopic calculations [10, 24]. The high-resolution measurement at RCNP of the $E1$ strength distribution $R_{E1}(\omega)$, where ω is the excitation energy, in ^{208}Pb [1] has allowed to deduce the experimental value of $\alpha_{\text{D}} = 20.1 \pm 0.6$ fm³. Actually, Tamii *et al.* [1], relying on the predictions of one single EDF [29] deduced a value of $r_{\text{skin}} = 0.156^{+0.025}_{-0.021}$ fm for ^{208}Pb . However, systematic errors were not estimated. Motivated by the interesting physics behind this observable, we present an exhaustive analysis of the correlation between α_{D} and r_{skin} in ^{208}Pb within a large set of EDFs (see Ref. [2] for more details).

RESULTS

In this contribution, we present our recent results on the correlation between r_{skin} and α_{D} in ^{208}Pb using a representative set of both relativistic and non-relativistic EDFs [2]. In all cases, these self-consistent models have been calibrated to selected global properties of finite nuclei and infinite nuclear matter. These models have been used without any further adjustment to compute R_{E1} using the consistent random-phase approximation (RPA).

We have chosen 48 EDFs and show their predictions in Fig. 1 (for details and original references see Ref. [2]). The up triangles correspond to Skyrme EDFs that have been widely used in the literature and fitted using very different protocols. In addition, we found interesting to analyze the predictions of three different families of relativistic (squares) and Skyrme EDFs (circles and down triangles) in which the value of the symmetry energy have been systematically varied around an optimal model (depicted by a star in Fig. 1). All the models within a family remain still accurate although their departure from the optimal model. This is basically due to the fact that the isovector channel of the nuclear interaction is not tightly constrained by available data.

Although some scatter is shown in Fig. 1, the approximate linear relation between α_{D} and r_{skin} discussed above is roughly confirmed by the different models. Specifically, the




FIGURE 1. α_D and r_{skin} of ^{208}Pb as predicted by 48 nuclear EDFs (see [2] for more details). Constrains on r_{skin} from PREX [5] and on α_D from RCNP [1] are also shown.

predicted linear correlation coefficient is around 0.8 when all the 48 models are taken into account while it is close to one when specific families of interactions are considered separately. The exception to this rule comes from the subset of Skyrme interactions that do not belong to a family of systematically varied interactions (black circles). This points to the fact that other quantities that do not appreciably vary in the different families of interactions are, indeed, affecting the value of α_D . It might be the case of some (isoscalar) properties that are almost constant within each one of the families and change when one looks at different families. This should be further investigated.

In Fig. 1, the constrains on r_{skin} from PREX [5] and on α_D from RCNP [1] are also shown. By looking at the models that are compatible with the RCNP measurement, we perform an average of our theoretical results and obtain a $r_{\text{skin}} = 0.168 \pm 0.022$ fm. Almost all theoretical predictions that agree with α_D (within the experimental error bars) are consistent with the value of r_{skin} measured by PREX.

SUMMARY AND CONCLUSIONS

We have analyzed the correlation between α_D and r_{skin} in ^{208}Pb using a large set of representative EDFs. Macroscopic analyses suggest that these two observables should be correlated. We have seen that in our study, that within families of accurately calibrated models a strong correlation between r_{skin} and α_D in ^{208}Pb arise. When these models are combined and more differences appear between them, the correlation weakens. We have estimated the model or theoretical systematic error of r_{skin} compatible with the measurement at RCNP and found that the obtained value is compatible but still far from

the central value obtained by PREX.

ACKNOWLEDGMENTS

Work partially supported by the Italian Research Project “Many-Body Theory of Nuclear Systems and Implications on the Physics of Neutron Stars” (PRIN 2008), by the Office of Nuclear Physics, U.S. Department of Energy under Contract Nos. DE-FG05-92ER40750 (FSU), DEFG02-96ER40963 (UTK); and by the BMBF under Contract 06ER9063.

REFERENCES

1. A. Tamii *et al.*, Phys. Rev. Lett. **107**, 062502 (2011).
2. J. Piekarewicz, B. K. Agrawal, G. Colò, W. Nazarewicz, N. Paar, P.-G. Reinhard, X. Roca-Maza, and D. Vretenar Phys. Rev. C **85**, 041302(R) (2012).
3. C. J. Horowitz, S. J. Pollock, P. A. Souder, and R. Michaels, Phys. Rev. C **63**, 025501 (2001).
4. R. Michaels *et al.*, Lead Radius Experiment proposal 2005, [<http://hallaweb.jlab.org/parity/prex/>].
5. S. Abrahamyan *et al.*, Phys. Rev. Lett. **108**, 112502 (2012).
6. T. Donnelly, J. Dubach, and I. Sick, Nucl. Phys. A **503**, 589 (1989).
7. B. A. Brown, Phys. Rev. Lett. **85**, 5296 (2000); S. Typel and B. A. Brown, Phys. Rev. C **64**, 027302 (2001).
8. R. J. Furnstahl, Nucl. Phys. A **706**, 85 (2002).
9. M. Centelles, X. Roca-Maza, X. Viñas, and M. Warda, Phys. Rev. Lett. **102**, 122502 (2009); Phys. Rev. C **82**, 054314 (2010).
10. P.-G. Reinhard and W. Nazarewicz, Phys. Rev. C **81**, 051303(R) (2010).
11. X. Roca-Maza, M. Centelles, X. Viñas, and M. Warda, Phys. Rev. Lett. **106**, 252501 (2011).
12. S. J. Pollock, E. N. Fortson, and L. Willets, Phys. Rev. C **46**, 2587 (1992). T. Sil, M. Centelles, X. Viñas, and J. Piekarewicz, *ibid.* **71**, 045502 (2005).
13. C. J. Horowitz and J. Piekarewicz, Phys. Rev. Lett. **86**, 5647 (2001); Phys. Rev. C **66**, 055803 (2002).
14. A. W. Steiner *et al.*, Phys. Rep. **411**, 325 (2005). B.-A. Li and A. W. Steiner, Phys. Lett. B **642**, 436 (2006); F. J. Fattoyev and J. Piekarewicz, Phys. Rev. C **82**, 025810 (2010).
15. L. Ray and G. W. Hoffmann, Phys. Rev. C **31**, 538 (1985).
16. L. Ray, G. W. Hoffmann, and W. R. Coker, Phys. Rep. **212**, 223 (1992).
17. A. Trzcińska *et al.*, Phys. Rev. Lett. **87**, 082501 (2001).
18. B. Kłos *et al.*, Phys. Rev. C **76**, 014311 (2007).
19. B. A. Brown, G. Shen, G. C. Hillhouse, J. Meng, and A. Trzcińska, Phys. Rev. C **76**, 034305 (2007).
20. B. C. Clark, L. J. Kerr, and S. Hama, Phys. Rev. C **67**, 054605 (2003).
21. J. Zenihiro *et al.*, Phys. Rev. C **82**, 044611 (2010).
22. M. N. Harakeh and A. van der Woude, Giant Resonances—Fundamental High-Frequency Modes of Nuclear Excitation (Clarendon, Oxford, UK, 2001).
23. L. Trippa, G. Colò, and E. Vigezzi, Phys. Rev. C **77**, 061304(R) (2008).
24. J. Piekarewicz, Phys. Rev. C **83**, 034319 (2011).
25. X. Roca-Maza, G. Pozzi, M. Brenna, K. Mizuyama, and G. Colò, Phys. Rev. C **85**, 024601 (2011).
26. D. Vretenar, Y. F. Niu, N. Paar, and J. Meng, Phys. Rev. C **85**, 044317 (2012).
27. E. Lipparini and S. Stringari, Phys. Lett. B **112**, 421 (1982); S. Stringari and E. Lipparini, *ibid.* **117**, 141 (1982).
28. W. Satuła, R. A. Wyss, and M. Rafalski, Phys. Rev. C **74**, 011301(R) (2006).
29. P. Klüpfel *et al.*, P.-G. Reinhard, T. J. Burvenich, and J. A. Maruhn, Phys. Rev. C **79**, 034310 (2009).