

## SCIENTIFIC OPINION

### Scientific Opinion on Dietary Reference Values for energy<sup>1</sup>

#### EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA)<sup>2,3</sup>

European Food Safety Authority (EFSA), Parma, Italy

#### ABSTRACT

Following a request from the European Commission, the Panel on Dietetic Products, Nutrition and Allergies (NDA) derived dietary reference values for energy, which are provided as average requirements (ARs) of specified age and sex groups. For children and adults, total energy expenditure (TEE) was determined factorially from estimates of resting energy expenditure (REE) plus the energy needed for various levels of physical activity (PAL) associated with sustainable lifestyles in healthy individuals. To account for uncertainties inherent in the prediction of energy expenditure, ranges of the AR for energy were calculated with several equations for predicting REE in children (1-17 years) and adults. For practical reasons, only the REE estimated by the equations of Henry (2005) was used in the setting of the AR and multiplied with PAL values of 1.4, 1.6, 1.8 and 2.0, which approximately reflect low active (sedentary), moderately active, active and very active lifestyles. For estimating REE in adults, body heights measured in representative national surveys in 13 EU Member States and body masses calculated from heights assuming a body mass index of 22 kg/m<sup>2</sup> were used. For children, median body masses and heights from the WHO Growth Standards or from harmonised growth curves of children in the EU were used. Energy expenditure for growth was accounted for by a 1 % increase of PAL values for each age group. For infants (7-11 months), the AR was derived from TEE estimated by regression equation based on doubly labelled water (DLW) data, plus the energy needs for growth. For pregnant and lactating women, the additional energy for the deposition of newly formed tissue, and for milk output, was derived from data obtained by the DLW method and from factorial estimates, respectively. The proposed ARs for energy may need to be adapted depending on specific objectives and target populations.

© European Food Safety Authority, 2013

#### KEY WORDS

Energy, resting energy expenditure, prediction equation, physical activity level, total energy expenditure, factorial method, average requirement

<sup>1</sup> On request from the European Commission, Question No EFSA-Q-2008-465, adopted on 29 November 2012.

<sup>2</sup> Panel members: Carlo Agostoni, Roberto Berni Canani, Susan Fairweather-Tait, Marina Heinonen, Hannu Korhonen, Sébastien La Vieille, Rosangela Marchelli, Ambroise Martin, Androniki Naska, Monika Neuhäuser-Berthold, Grażyna Nowicka, Yolanda Sanz, Alfonso Siani, Anders Sjödin, Martin Stern, Sean (J.J.) Strain, Inge Tetens, Daniel Tomé, Dominique Turck and Hans Verhagen. Correspondence: [nda@efsa.europa.eu](mailto:nda@efsa.europa.eu)

<sup>3</sup> Acknowledgement: The Panel wishes to thank the members of the Working Group on Population Reference Intakes: Carlo Agostoni, Jean-Louis Bresson, Patrick Even, Susan Fairweather-Tait, Albert Flynn, Ambroise Martin, Monika Neuhäuser-Berthold, Hildegard Przyrembel, Sean (J.J.) Strain, Inge Tetens and Daniel Tomé for the preparatory work on this scientific opinion and EFSA staff: Anja Brönstrup for the support provided to this scientific opinion.

## SUMMARY

Following a request from the European Commission, the Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a scientific opinion on dietary reference values (DRVs) for the European population, including energy.

The DRVs for food energy provide a best estimate of the food energy needs of population groups within Europe. They are given as average requirements (ARs) of specified age and sex groups and are of limited use for individuals. The reference values need to be adapted to specific objectives, such as dietary assessment, dietary planning, labelling dietary reference values, or development of food-based dietary guidelines. In addition, there is a need to define and characterise the target population.

In this Opinion, total energy expenditure (TEE) in the steady state of a healthy body mass was chosen as the criterion on which to base the AR for energy. In practice, the adequacy of usual energy intakes is best monitored by measuring body mass. In terms of regulation of body mass, the overall energy balance over a prolonged period of time needs to be considered. TEE expended over 24 hours is the sum of basal energy expenditure, the energy expenditure of physical activity and the thermic effect of food. In this Opinion, resting energy expenditure (REE) was used as a proxy for the slightly lower basal energy expenditure, as most studies measured REE. TEE is best measured with the doubly labelled water (DLW) method, which provides energy expenditure data over biologically meaningful periods of time and under normal living conditions.

One approach to determine the AR for energy is to use regression equations which describe how TEE measured with the DLW method varies as a function of anthropometric variables (such as body mass and height) for defined population groups and for an activity constant that accounts for the level of physical activity. However, this approach has been criticised because of the inability of such TEE prediction models to account for the variations in energy expenditure of physical activity in a transparent way. In addition, limited TEE data generated with the DLW method are available, and they may not be representative for the European population; moreover, some age groups are under-represented. Another approach for estimating TEE is by the factorial method in which the energy spent in various activities is added to measured or predicted REE. This is achieved by using the physical activity level (PAL), which is defined as the ratio of TEE to REE per 24 hours and reflects the part of TEE that is due to physical activity. Accordingly, TEE is predicted as  $PAL \times REE$ . During growth, pregnancy and lactation, additional energy is needed for the synthesis and deposition of new tissues, and for milk production.

In this Opinion, TEE of children and adults was estimated factorially to account for the diversity in body size, body composition and habitual physical activity among children and adult populations with different geographic, cultural and economic backgrounds.

To estimate REE, predictive equations were used, derived from regression analysis of measured REE, body masses and heights from groups of subjects. Body mass is the most important determinant of REE and all predictive equations use this parameter. In addition to body mass, height, sex, age and ethnicity can affect REE significantly and numerous equations have been developed to take into account one or several of these parameters. Based on the accuracy of various equations in specified population groups, five widely used equations (Harris and Benedict, 1919; Henry, 2005; Mifflin et al., 1990; Müller et al., 2004; Schofield et al., 1985) were considered as equally valid for estimating the REE of healthy adults in Europe. For healthy children and adolescents, the equations of Schofield et al. (1985) and Henry (2005) derived from large datasets and covering wide age groups were considered to be the most suitable.

PAL can be estimated either from time-allocated lists of daily activities expressed as physical activity ratio values or from the ratio of TEE (measured by the DLW method) to REE (either measured or estimated). However, the same limitations apply to the derivation of PAL values from DLW data as to the estimates of TEE with this method. Within the general population, PALs associated with

sustainable lifestyles have been observed to range between 1.35 and 2.5, and to decrease only marginally with age. When assigning PAL values to descriptions of activities/lifestyles (such as light, moderate or heavy activity), the range of PAL values in each lifestyle category is large. Thus, the allocation of lifestyles to defined PAL values can only be considered a rough indication of PAL, but may be useful for decisions about which PAL values to apply in various circumstances and applications.

In the absence of arguments for the selection of one predictive equation best fitted to adults in the European Union (EU), REE was calculated with five widely applied predictive equations using individual data of measured body heights of adults obtained in 13 representative national surveys in EU Member States, with corresponding body masses calculated for a body mass index (BMI) of  $22 \text{ kg/m}^2$ , i.e. the midpoint of the range of healthy BMI of adults as defined by the WHO. This yielded a range of ARs calculated for PAL values from 1.4 to 2.4 in steps of 0.2, and demonstrated the magnitude of uncertainty inherent in these values. However, for practical reasons, only one AR is proposed for a defined age and sex group with a healthy BMI of 22, and for PAL values selected to approximate corresponding lifestyles. The predictive equations of Henry (2005) were used to estimate REE because, at present, the underlying database is the most comprehensive as regards number of subjects, their nationalities and age groups. To derive TEE as  $\text{REE} \times \text{PAL}$ , PAL values of 1.4, 1.6, 1.8 and 2.0 were chosen to approximately reflect low active (sedentary), moderately active, active and very active lifestyles. Because of a lack of anthropometric data from EU countries for age groups from 80 years onwards, average requirements were not calculated for adults  $\geq 80$  years.

For infants from birth to six months of age, energy requirements were considered to be equal to the energy supply from human milk, and no DRV is proposed. For infants aged 7-11 months, the ARs were estimated from equations for TEE, adding the energy needs for growth. TEE was based on measurements using the DLW method in healthy, full-term infants, exclusively breast-fed for the first four months of life and with adequate body mass. Body masses from the WHO Growth Standards were used to derive ARs for infants growing along the trajectory of this standard. Estimates of the energy requirement for growth were based on protein and fat gains reported in the literature.

The ARs of children from one year upwards are based on predicted REE and adjusted PAL for growth. REE was calculated using the predictive equations of Henry (2005) and Schofield et al. (1985) and median body masses and heights taken from the WHO Growth Standards (for children up to two years) or from harmonised growth curves of EU children (for children from 3 to 17 years). For the same reasons as outlined for adults, and because the results obtained with these two equations were very similar, only the predictive equations of Henry (2005) were applied for the estimation of REE values. PAL values of 1.4, 1.6, 1.8 and 2.0 were used for three age groups (1-3 years,  $>3$ - $<10$  years, and 10-18 years). Energy expenditure for growth was accounted for by a 1 % increase in PAL values for each age group.

For pregnant women, a mean gestational increase in body mass of 12 kg was considered to be associated with optimal maternal and fetal health outcomes. The additional amount of energy required during pregnancy to support this increase in body mass was estimated using the cumulative increment in TEE estimated with the DLW technique plus the energy deposited as protein and fat. Based on these data, the average additional energy requirement for pregnancy is 320 MJ (76,530 kcal) which equates to approximately 0.29 MJ/day (70 kcal/day), 1.1 MJ/day (260 kcal/day) and 2.1 MJ/day (500 kcal/day) during the first, second and third trimesters, respectively.

For women exclusively breastfeeding during the first six months after birth, the additional energy requirement during lactation was estimated factorially as 2.1 MJ/day (500 kcal/day) over pre-pregnancy requirements, taking into account a requirement of 2.8 MJ/day (670 kcal/day) for milk production and an energy mobilisation from maternal tissues of 0.72 MJ/day (170 kcal/day). No additional energy requirement is proposed for women lactating beyond the sixth month because volumes of milk produced during this period are highly variable and depend on the infant's energy intake from complementary foods.

## TABLE OF CONTENTS

Abstract .....	1
Summary .....	2
Table of contents .....	4
Background as provided by the European Commission.....	7
Terms of reference as provided by the European Commission.....	7
Preamble.....	9
Assessment .....	9
1. Introduction .....	9
1.1. Definition of energy requirement.....	10
1.2. Concept of dietary reference values (DRVs) for energy .....	10
1.3. Approach.....	10
2. Definition/Category .....	10
2.1. Components of total energy expenditure (TEE) .....	10
2.1.1. Basal energy expenditure (BEE) .....	10
2.1.2. Resting energy expenditure (REE) .....	10
2.1.3. Sleeping energy expenditure .....	11
2.1.4. Cold-induced thermogenesis .....	11
2.1.5. Thermic effect of food (TEF) .....	11
2.1.6. Energy expenditure of physical activity (EEPA).....	11
2.1.7. Adaptive thermogenesis .....	11
2.2. Methods of assessing energy expenditure and its components .....	12
2.2.1. General principles.....	12
2.2.1.1. Direct calorimetry .....	12
2.2.1.2. Indirect calorimetry.....	12
2.2.1.3. Doubly labelled water (DLW) method .....	12
2.2.1.4. Heart rate (HR) monitoring.....	13
2.2.2. Basal and resting energy expenditure (BEE and REE).....	13
2.2.3. Thermic effect of food (TEF) .....	13
2.2.4. Energy expenditure of physical activity (EEPA).....	13
2.2.5. Total energy expenditure (TEE) .....	14
2.2.6. Energy expenditure for growth.....	14
2.2.7. Energy expenditure of pregnancy .....	14
2.2.8. Energy expenditure of lactation.....	14
2.3. Determinants of energy expenditure .....	14
2.3.1. Body mass and body composition .....	14
2.3.2. Physical activity.....	15
2.3.3. Growth.....	15
2.3.4. Pregnancy .....	15
2.3.5. Lactation .....	16
2.3.6. Endocrinological factors.....	16
2.3.7. Ageing .....	16
2.3.8. Diet .....	16
2.3.9. Sex .....	17
2.3.10. Ethnicity .....	17
2.3.11. Environmental factors.....	17
2.4. Equations to predict resting energy expenditure (REE).....	17
2.4.1. Predictive equations for adults.....	17
2.4.2. Predictive equations for children .....	19
3. Dietary sources of energy and intake data .....	19
3.1. Dietary sources of energy .....	19
3.2. Dietary intake data .....	20
4. Overview of dietary reference values and recommendations .....	21
4.1. Adults.....	21

4.2.	Infants and children.....	22
4.3.	Pregnancy.....	23
4.4.	Lactation .....	25
5.	Criteria and approaches for deriving the Average Requirement (AR) for energy.....	28
5.1.	Criteria .....	28
5.1.1.	Energy balance .....	28
5.1.2.	Body mass, body mass index (BMI) and body composition .....	28
5.1.3.	Body mass gain in pregnancy.....	29
5.1.4.	Physical activity.....	30
5.2.	Approaches .....	30
5.3.	Derivation of energy requirements of various population groups.....	32
5.3.1.	Adults .....	32
5.3.1.1.	Calculation of resting energy expenditure (REE).....	32
5.3.1.2.	Selection of physical activity level (PAL) values.....	34
5.3.1.3.	Ranges of Average Requirement (AR) for energy for adults .....	35
5.3.2.	Infants .....	36
5.3.2.1.	Total energy expenditure (TEE).....	36
5.3.2.2.	Energy deposition in new tissue .....	37
5.3.3.	Children .....	37
5.3.3.1.	Calculation of resting energy expenditure (REE).....	38
5.3.3.2.	Selection of physical activity levels (PALs).....	38
5.3.3.3.	Energy expenditure of children and adolescents for growth.....	39
5.3.3.4.	Ranges of Average Requirement (AR) for energy for children and adolescents.....	39
5.3.4.	Pregnancy .....	40
5.3.4.1.	Energy requirement for the increase in tissue mass during pregnancy.....	41
5.3.4.2.	Calculation of additional AR for energy for tissue deposition in pregnancy.....	41
5.3.5.	Lactation.....	42
6.	Key data on which to base dietary reference values (DRVs).....	42
6.1.	Adults.....	43
6.1.1.	Calculation of resting energy expenditure (REE).....	43
6.1.2.	Selection of physical activity level (PAL) values.....	43
6.2.	Infants .....	43
6.3.	Children and adolescents .....	44
6.3.1.	Calculation of resting energy expenditure (REE).....	44
6.3.2.	Selection of physical activity level (PAL) values.....	44
6.4.	Pregnancy.....	44
6.5.	Lactation .....	45
	Conclusions .....	46
	Recommendations for Research.....	48
	References .....	49
	Appendices.....	68
	Appendix 1: Energy content of human milk from healthy mothers of term infants in the EU.....	68
	Appendix 2: Predictive equations for REE in adults .....	69
	Appendix 3a: Population, methods and period of dietary assessment in children and adolescents in European countries.....	73
	Appendix 3b: Energy intake of children aged ~1-3 years in European countries.....	76
	Appendix 3c: Energy intake of children aged ~4-6 years in European countries.....	77
	Appendix 3d: Energy intake of children aged ~7-9 years in European countries.....	78
	Appendix 3e: Energy intake of children aged ~10-14 years in European countries.....	79
	Appendix 3f: Energy intake of adolescents aged ~15-18 years in European countries .....	80
	Appendix 4a: Population, methods and period of dietary assessment in adults in European countries.....	81
	Appendix 4b: Energy intake of adults aged ~19-65 years in European countries .....	84
	Appendix 4c: Energy intake of adults aged ~19-34 years in European countries .....	85
	Appendix 4d: Energy intake of adults aged ~35-64 years in European countries .....	86

Appendix 4e: Energy intake of adults aged ~65 years and over in European countries.....	88
Appendix 5: Overview of the approaches used by selected authorities for the estimation of average requirements (ARs) for energy for adults .....	89
Appendix 6: Overview of the approaches to estimate average requirements (ARs) for energy for infants and young children of selected countries and authorities other than FAO/WHO/UNU and IoM.....	93
Appendix 7: Overview of the approaches of FAO/WHO/UNU (2004) and IoM (2005) to estimate average requirements (ARs) for energy for infants, children and adolescents .....	94
Appendix 8: Overview of the approaches to estimate daily average requirements (ARs) for energy for children and adolescents of selected countries and authorities other than FAO/WHO/UNU and IoM.....	95
Appendix 9: REE calculated with five most used predictive equations using measured heights from surveys in 13 EU Member States and body masses to yield a BMI of 22.....	97
Appendix 10: Comparison of measured REE of GISELA subjects (last available measurements) with REE calculated with various predictive equations.....	99
Appendix 11: Reference body heights and body masses for infants, children and adults .....	100
Appendix 12a: Examples of relationships reported between lifestyle, activity and physical activity level (PAL).....	101
Appendix 12b: Contribution of various activities to physical activity levels (PALs) .....	101
Appendix 13: Selected predictive equations for REE in children and adolescents .....	103
Appendix 14a: Ranges of average Requirement (AR) for energy for adults based on the factorial method and predicting REE with five most used equations.....	104
Appendix 14b: Ranges of Average Requirement (AR) for energy for children and adolescents based on the factorial method and predicting REE with two predictive equations.....	105
Appendix 15: Derivation of the Average Requirement (AR) for energy for infants aged 7-11 months.....	107
Appendix 16: Summary of Average Requirement (AR) for energy expressed in kcal/day.....	108
Glossary and Abbreviations .....	110



## BACKGROUND AS PROVIDED BY THE EUROPEAN COMMISSION

The scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and if necessary to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on nutrient and energy intakes for the European Community<sup>4</sup>. The report provided reference intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU Reference Intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context the EFSA is requested to consider the existing Population Reference Intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a Population Reference Intake for dietary fibre.

For communication of nutrition and healthy eating messages to the public it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context the EFSA is asked to provide assistance on the translation of nutrient based recommendations for a healthy diet into food based recommendations intended for the population as a whole.

## TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

In accordance with Article 29 (1)(a) and Article 31 of Regulation (EC) No. 178/2002, the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on population reference intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance the EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically advice is requested on the following dietary components:

- Carbohydrates, including sugars;
- Fats, including saturated fatty acids, polyunsaturated fatty acids and monounsaturated fatty acids, *trans* fatty acids;
- Protein;

---

<sup>4</sup> Scientific Committee for Food, Nutrient and energy intakes for the European Community, Reports of the Scientific Committee for Food 31<sup>st</sup> series, Office for Official Publication of the European Communities, Luxembourg, 1993.

- Dietary fibre.

Following on from the first part of the task, the EFSA is asked to advise on population reference intakes of micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, the EFSA is asked to provide guidance on the translation of nutrient based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).



## PREAMBLE

In the Opinion on General Principles (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010), the Panel distinguishes between reference values and recommendations: “Dietary Reference Values are scientific references based on health criteria, taking into account dietary requirements and health outcomes. [...] They represent one of the bases for establishing nutrient recommendations and food based dietary guidelines. [...] Nutrient goals and recommendations may differ between countries depending on health needs, nutritional status and known patterns of intake of foods and nutrients in specific populations and the actual composition of available foods”. In this Opinion, the Panel proposes reference values that need to be adapted to specific objectives and target populations.

In this Opinion on Dietary Reference Values (DRVs) for energy, the Panel has decided to use the scientifically correct term “body mass” instead of “body weight”. In accordance with the International System of Units, the FAO/WHO/UNU consensus (1971) and the European regulations<sup>5</sup>, the AR for energy will be expressed in joules (J). However, because of the continuing use of thermochemical energy units (calories, cal), equivalents<sup>6</sup> will be given in brackets in the text or in separate tables in the Appendices.

## ASSESSMENT

### 1. Introduction

Human beings need energy to perform and regulate all biochemical processes that maintain body structures and functions, and to perform physical activities.

Energy is provided in the diet by carbohydrates, fats, protein and alcohol, and the individual contribution of these sources is variable. Thus, DRVs for energy are not specified as defined amounts of a single nutrient but are expressed in units of energy.

DRVs for energy differ from those for nutrients in that (a) there is a wide inter-individual variation in the behavioural, physiological and metabolic components of energy needs, and the energy requirement of a defined group cannot be applied to other groups or individuals who differ from the defined group in sex, age, body mass, activity level and possibly other factors; and (b) there are differences between the energy supply needed to maintain current body mass and level of actual physical activity and the energy supply needed to maintain desirable body mass and a level of physical activity consistent with good health.

The proposed DRVs for food energy provide a best estimate of the food energy needs of population groups within Europe, and present criteria against which to judge the adequacy of their food energy intakes. They constitute the basis for policy-makers and authorities to make recommendations for populations which can be used for the development and monitoring of nutrition programmes, and for planning agricultural production, food supplies and, if required, the mobilisation and distribution of emergency food aid.

---

<sup>5</sup> Council Directive 90/496/EEC of 24 September 1990 on nutrition labelling for foodstuffs. OJ L 276, 6.10.1990, p. 40–44/ Regulation (EU) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No 1924/2006 and (EC) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/200. OJ L 304, 22.11.2011, p. 18–63.

<sup>6</sup> 1 joule (J) is the amount of mechanical energy required to displace a mass of 1 kg through a distance of 1 m with an acceleration of 1 m per second ( $1 \text{ J} = 1 \text{ kg} \times 1 \text{ m}^2 \times 1 \text{ sec}^{-2}$ ). Multiples of 1,000 (kilojoules, kJ) or 1 million (megajoules, MJ) are used in human nutrition. The conversion factors between joules and calories are:  $1 \text{ kcal} = 4.184 \text{ kJ}$ , or conversely,  $1 \text{ kJ} = 0.239 \text{ kcal}$ .

## 1.1. Definition of energy requirement

Energy requirement is the amount of food energy needed to balance energy expenditure in order to maintain body mass, body composition, and a level of physical activity consistent with long-term good health. This includes the energy needed for the optimal growth and development of children, for the deposition of tissues during pregnancy, and for the secretion of milk during lactation, consistent with the good health of both mother and child (FAO/WHO/UNU, 1985, 2004; IoM, 2005; SCF, 1993).

## 1.2. Concept of dietary reference values (DRVs) for energy

Following the definition of energy requirement, dietary reference values are based on estimates of the requirements of healthy individuals, representative for a particular population group. As a result of biological variability, there is a distribution of energy requirements within each group. Whereas DRVs for protein and various micronutrients are given as population reference intakes (PRI)<sup>7</sup>, DRVs for energy are provided as average requirements (ARs) of specified groups. Due to the very large variation coefficients (CV) induced by large differences in physical activity levels (PAL) and anthropometric parameters, the definition of a PRI would be inappropriate, since it implies an intake above the requirement for nearly all subjects which would lead to a positive energy balance and promote an unfavourable increase in body mass and the development of obesity in the long term. The AR for energy as a reference value exceeds the requirement of half of the individuals of any specified group. The AR for energy relates to groups of healthy people and is of limited use for individuals.

The AR for energy is expressed on a daily basis but represents an average of energy needs over a minimum of one week.

## 1.3. Approach

The AR for energy can be established by two approaches: measurements of energy intake or expenditure of healthy reference populations. Because the day-to-day variation in energy intake is considerably larger than the day-to-day variation in total energy expenditure (TEE) in a steady state of body mass, measurements or estimates of TEE were chosen by experts from FAO/WHO/UNU (1985, 2004) and the US Institute of Medicine (IoM, 2005) as the criterion on which to base the AR for energy. The Panel agrees with this approach.

## 2. Definition/Category

### 2.1. Components of total energy expenditure (TEE)

Total energy expenditure (TEE) expended over 24 hours is the sum of basal energy expenditure (BEE), the energy expenditure of physical activity (EEPA), the thermic effect of food (TEF) and in less frequent situations cold-induced thermogenesis.

#### 2.1.1. Basal energy expenditure (BEE)

Basal energy expenditure (BEE) is the energy used to maintain the basic physiological functions of the body at rest under strictly defined conditions: after an overnight fast corresponding to 12-14 hours of food deprivation, awake, supine, resting comfortably, motionless, no strenuous exercise in the preceding day (or eight hours of physical rest), being in a state of “mental relaxation” and in a thermoneutral environment. BEE is the main component (45-70 %) of TEE (FAO/WHO/UNU, 2004).

#### 2.1.2. Resting energy expenditure (REE)

By definition, resting energy expenditure (REE) is the energy expended when the body is at rest, which is when no extra energy is used for muscular effort. In many studies, for practical reasons since

---

<sup>7</sup> The PRI is defined as the level of intake that is adequate for virtually all people in a population group, which is determined as the average requirement (AR) of the population group plus two standard deviations (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010. Scientific Opinion on principles for deriving and applying Dietary Reference Values. EFSA Journal, 8(3):1458, 30 pp.)

conditions for measuring BEE are more stringent, REE instead of BEE is measured. Changes in REE are used to measure the expenditure of many processes such as thermoregulation, eating and excess post-exercise oxygen consumption. Practically, REE is measured in conditions less stringent than the ones that prevail for measurement of BEE, so that REE is usually slightly higher than BEE (up to 10 %). In this Opinion, REE is used as a proxy for BEE, as most studies measure REE.

### **2.1.3. Sleeping energy expenditure**

Sleeping energy expenditure can be measured instead of BEE or REE to estimate daily energy requirements. It is usually considered to be lower than REE depending on the sleeping phase (Wouters-Adriaens and Westerterp, 2006). Sleeping energy expenditure can be considered as a practical means to approach BEE particularly in infants for whom the criteria related to measurements of BEE would be impractical.

### **2.1.4. Cold-induced thermogenesis**

Cold-induced thermogenesis is the production of heat in response to environmental temperatures below thermoneutrality. Cold-induced thermogenesis can be divided into two types: shivering thermogenesis and non-shivering thermogenesis. The thermoneutral zone (or the critical temperature) is the environmental temperature at which oxygen consumption and metabolic rate are lowest (IoM, 2005). The relative contribution of cold-induced thermogenesis to TEE has decreased in recent decades due to the increase in time spent in enclosed and heated environments.

### **2.1.5. Thermic effect of food (TEF)**

Eating requires energy for the digestion, absorption, transport, interconversion and, where appropriate, deposition/storage of nutrients. These metabolic processes increase REE, and their energy expenditure is known as the thermic effect of food (TEF). It should be noted that the muscular work required for eating is not part of TEF.

### **2.1.6. Energy expenditure of physical activity (EEPA)**

Physical activity can be defined as any body movement produced by skeletal muscles which results in energy expenditure. In practice, physical activity in daily life can be categorised into obligatory and discretionary activity. The term “obligatory” is more appropriate than the term “occupational” that was used in the 1985 report (FAO/WHO/UNU, 1985) because, in addition to occupational work, obligatory activities include a range of daily activities, for example children going to school, adults tending to the home and family, and other demands made on children and adults by their economic, social and cultural environment (FAO/WHO/UNU, 2004). Levine (2004b) has divided the energy expended during physical activity into exercise activity thermogenesis and non-exercise activity thermogenesis. Exercise activity thermogenesis is the energy expended during voluntary exercise (discretionary) which is a type of physical activity that is planned, structured and repetitive. Non-exercise activity thermogenesis is the energy expenditure of all physical activities other than sleeping, eating or sports-like exercise. It includes the energy expended during daily activities such as working, walking, housework and gardening, as well as fidgeting, which corresponds to small unconscious muscle movements (Levine, 2004b).

The physical activity level (PAL) is defined as the ratio of TEE to REE over 24 hours. It reflects the part of TEE that is due to physical activity. The physical activity ratio (PAR) is used to express the increase in energy expenditure per unit of time induced by a given activity, and can also be expressed as a multiple of REE.

### **2.1.7. Adaptive thermogenesis**

Adaptive thermogenesis is defined as the heat that can be added or not to the normal thermogenic response to food and/or cold in order to best adjust energy expenditure to the requirements of energy balance (Wijers et al., 2009). Several studies conducted in recent years suggest that mitochondrial uncoupling protein in brown adipose tissue (Nedergaard et al., 2007) and skeletal muscle tissue in

adult humans (Wijers et al., 2009) can be the main effectors of adaptive thermogenesis. Other mechanisms such as futile calcium cycling, protein turnover and substrate cycling may be involved (Harper et al., 2008). Under normal circumstances in healthy individuals, adaptive thermogenesis does not account for a significant component of TEE.

## 2.2. Methods of assessing energy expenditure and its components

### 2.2.1. General principles

#### 2.2.1.1. Direct calorimetry

As body temperature is kept constant, the energy expended by the body is dissipated as heat and potential external work. Direct calorimetry measures the heat released by the subject through conduction, convection and evaporation. Direct calorimetry has been used in the past to validate the principle of indirect calorimetry, but is less often used currently because of its cost and complexity (Seale et al., 1991; Walsberg and Hoffman, 2005).

#### 2.2.1.2. Indirect calorimetry

Indirect calorimetry is based on the principle that energy production by substrate oxidation in the body is coupled to oxygen consumption ( $\text{VO}_2$ ) and carbon dioxide production ( $\text{VCO}_2$ ), and has become the reference method for measuring energy expenditure. Many equations have been derived to provide an exact measure of energy expenditure from  $\text{VO}_2$  and  $\text{VCO}_2$  (Brouwer, 1957; Elia, 1992; Lusk, 1928; Weir, 1949; Zuntz, 1897). The most widely used is the Weir formula, and the other formulas give results that all lie within  $\pm 1$  % of the results by Weir (1949).

Closed-circuit indirect calorimetry: At a time when no accurate automated gas analysers were available, the closed-circuit system allowed a volumetric measurement of  $\text{VO}_2$  to be performed. In the closed-circuit design,  $\text{VCO}_2$  is absorbed within the system, and  $\text{VO}_2$  is measured either from the decrease in the volume of gas in the system, or by the amount of oxygen required to maintain the pressure in the chamber. Closed-circuit systems are no longer used for measurement of REE in humans.

Open-circuit indirect calorimetry: The principle of the open-circuit device is that the respired gases of the subject are collected in a device ventilated at a known flow-rate, and  $\text{VO}_2$  and  $\text{VCO}_2$  are computed by multiplying the changes in %  $\text{O}_2$  and %  $\text{CO}_2$  in the container by the air flow. Various open-circuit systems have been designed based on this principle. Ventilated open-circuit systems such as ventilated hood, canopy, and whole room calorimeters are the most used for assessing BEE, REE, TEF and TEE. Expiratory collection systems are systems where the subject inspires from the atmosphere and expires via a non-return valve into a measurement unit. They are mostly used for exercise and field measurements via portable systems. Open-circuit indirect calorimeters are reliable with an error of 0.5–2 % (Compher et al., 2006; Schoeller, 2007; Wahrlich et al., 2006).

Since both  $\text{VCO}_2$  and  $\text{VO}_2$  are measured, a main advantage of the open-circuit devices is the possibility to compute  $\text{VCO}_2$  over  $\text{VO}_2$ , which is defined as the respiratory quotient. Values for the respiratory quotient vary depending on the substrate mixture oxidised (0.7 for lipids, 0.82 for proteins and 1.0 for glucose). A precise computation of the respective levels of glucose, lipid and protein oxidation thus requires that protein oxidation be measured. This is usually done by measuring urinary nitrogen excretion, assuming that, on average, nitrogen excreted multiplied by 6.25 is equivalent to the amount of protein oxidised (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2012).

#### 2.2.1.3. Doubly labelled water (DLW) method

The doubly labelled water (DLW) method is used for determining TEE in free-living subjects. It is based on the disappearance rates in body fluids (usually urine sampled at three or more intervals) of two orally administered stable isotopes of water ( $\text{H}_2^{18}\text{O}$  and  $^2\text{H}_2\text{O}$ ) during the 15 following days (which corresponds to about two biological half-lives of the isotopes) (Schoeller, 1988).  $\text{VCO}_2$  is

calculated from the difference between the disappearance rates of  $^{18}\text{O}$  and  $^2\text{H}$ .  $\text{VO}_2$  is calculated from  $\text{VCO}_2$  by estimating the respiratory quotient from the food quotient (sum of the respiratory quotient of individual foods/energy contribution in 24 hours) based on either the reported macronutrient intake of the subject or on average data from population surveys. TEE can then be calculated from the energy equivalent of  $\text{VCO}_2$  for the given diet (Elia, 1991) or from  $\text{VCO}_2$  measured and  $\text{VO}_2$  calculated with the use of the food quotient using the same standard equations as for indirect calorimetry. Computation of TEE with DLW relies on a series of assumptions including the constancy of the water pool throughout the measurement period, the rate of  $\text{H}_2\text{O}$  and  $\text{CO}_2$  fluxes, the isotopic fractionation, and no label-re-entering the body (IDECG, 1990). The reproducibility and accuracy of the DLW technique may vary markedly among analytical centres, and estimates of  $\pm 8.5\%$  for the reliability of TEE have been reported (Goran et al., 1994a). The main advantages of DLW versus calorimetry are that (i) it provides energy expenditure estimations over biologically meaningful periods of time, (ii) it captures energy expenditure of all kinds of activities including spontaneous movements and fidgeting, and (iii) being non-invasive, measurements can be made in subjects leading their usual daily lives.

#### 2.2.1.4. Heart rate (HR) monitoring

Heart rate (HR) monitoring can be used to estimate TEE but individual calibrations of the relationship between HR and oxygen consumption must be performed because the relationship between HR and TEE varies between subjects (Bitar et al., 1996; Ceesay et al., 1989).

### 2.2.2. Basal and resting energy expenditure (BEE and REE)

BEE and REE as a proxy for BEE are best determined by indirect calorimetry measurements under standardised laboratory conditions (Compher et al., 2006; Harris and Benedict, 1919).

### 2.2.3. Thermic effect of food (TEF)

TEF is best measured in laboratory conditions from changes in REE induced by ingestion of a standardised meal of known composition and of 1,700 kJ (~400 kcal) or greater. In practice, first REE is measured (see Section 2.1.2.), then the meal is ingested, and the meal-induced increase in REE versus the pre-meal value is measured.

### 2.2.4. Energy expenditure of physical activity (EEPA)

As is the case for TEF, energy expenditure of physical activity (EEPA) should be measured relative to REE or relative to the energy expenditure of the reference activity (e.g. expenditure of office work relative to energy expenditure while seated, expenditure of walking or running relative to energy expenditure while standing).

The measurement of average daily TEE by the DLW method combined with a measurement of REE permits the calculation of the energy expenditure for the average physical activity of an individual (Westerterp and Goran, 1997) but does not provide information on the expenditure and time spent in the various activities. The energy expenditure of fidgeting has been assessed with indirect calorimetry measurements as the difference between energy expenditure at rest and at various levels of activities with and without fidgeting (Levine et al., 2000). At a population level, systematic data on the amount and expenditure of non-exercise activity thermogenesis are scarce.

Tables have been developed which ascribe to each type of activity a PAR that defines the energy expended while performing this activity relative to REE (e.g. FAO/WHO/UNU, 2005). Such tables are of limited value because of inconsistencies in the way the data were collected and presented, and because of differences in the description of activities, in the computation/prediction of REE, and in the conditions of measurements (Vaz et al., 2005). As a result, PAR values of a given activity can vary greatly between studies.



### **2.2.5. Total energy expenditure (TEE)**

TEE in normal living conditions is best estimated with the DLW method (Coward and Cole, 1991) which, while expensive, allows long-term measurements and preserves normal behaviour better than recording in room calorimeters. Therefore, TEE in a population group is generally estimated by factorial methods in which the energy spent in various activities is added to measured or calculated REE.

The factorial method thus requires (i) accurate recording of daily activities, which is tedious (especially for children), (ii) accurate data on the energy expenditure of most individual daily activities and (iii) a precise value for REE, either measured or calculated from either body mass, body mass and body height, or body composition. The difficulty of complying with all three requirements is a source of large potential errors at the individual level, but the factorial method can be applied to estimate TEE in groups of people.

### **2.2.6. Energy expenditure for growth**

The increase in energy expenditure induced by growth results from the expenditure for protein and lipid synthesis and their deposition in newly-formed tissue. It is significant only in rapidly growing infants and children. It cannot be measured by indirect calorimetry or the DLW method because there is no possibility of access to a reference “growth-free” REE. However, it can be evaluated from changes in body composition measured in groups of healthy growing infants (Torun, 2005). A factorial method which consists of measuring changes in body composition and estimating the energy requirements from the estimated energetic efficiencies of the biochemical pathways involved in protein and lipid synthesis can also be used (Butte, 2005).

### **2.2.7. Energy expenditure of pregnancy**

Energy expenditure related to pregnancy is calculated using two different approaches. Both require that measurements be started before conception, which raises recruitment difficulties. The first approach is based on serial measurements of REE assuming that EEPA and TEF are not affected by pregnancy (Prentice et al., 1996a). In the second approach, calculations can be based on serial measurements of TEE using the DLW method. This method not only includes the energy expenditure for tissue deposition but also any changes in TEF and EEPA.

### **2.2.8. Energy expenditure of lactation**

Energy expenditure for lactation can be computed from the amount of milk produced, the energy content of the milk, and the energetic efficiency of milk synthesis. The efficiency of converting dietary energy into human milk has been estimated from theoretical biochemical efficiencies of synthesising the constituents in milk, and from metabolic balance studies (Prentice and Prentice, 1988). Biochemical efficiency can be calculated from the stoichiometric equations and the obligatory heat losses associated with the synthesis of lactose, protein and fat. When the expenditure for digestion, absorption, inter-conversion and transport is taken into account, the estimate of efficiency of milk synthesis yields a figure of 80–85 % (Butte and King, 2005).

## **2.3. Determinants of energy expenditure**

### **2.3.1. Body mass and body composition**

The relationship of body mass and body composition to energy expenditure is not appropriately reflected in a simple regression of REE and body mass, as this does not pass through the zero intercept, and is not linear because body composition does not evolve linearly with body mass (Müller et al., 2002). The various tissues and organs of the body have very different mass-specific metabolic rates, with very low or null values for plasma, collagen, tendons, fluids and bones, for example, low values for adipose tissues, average values for muscles, and high values for brain, heart, liver and kidneys (Elia, 1992; Müller et al., 2002; Wang et al., 2010). Thus, the contribution of fat mass (FM) to energy expenditure is low in lean subjects, but cannot be neglected in overweight and obese subjects



(Müller et al., 2004; Prentice et al., 1996b; Schulz and Schoeller, 1994). In addition, it has been demonstrated that fat distribution is a key determinant for the contribution of body fat to REE. For example, abdominal fat has a greater metabolic activity than peripheral fat (Lührmann et al., 2001). Prediction of REE can be improved by using multicomponent body composition models based on various techniques. This may be particularly useful in populations for which the current equations may not properly predict REE (Wilms et al., 2010), and for reassessing the validity of ethnic and sex differences.

### 2.3.2. Physical activity

EEPA is the most variable component of TEE, both within and between subjects, ranging from 15 % of TEE in very sedentary individuals to 50 % or more of TEE in highly active individuals. The energy expended with exercise is often negligible or zero in individuals, but even in those who exercise regularly, the energy expended with non-exercise activity thermogenesis is far larger than the energy expended with exercise. Thus, energy requirements related to physical activity mainly arise from non-exercise activity thermogenesis. The latter can vary between two people of similar size by more than 8 MJ/day (1,900 kcal/day) because of different occupations, leisure-time activities and fidgeting. Fidgeting can increase daily energy expenditure above REE levels by 20–40 % (Levine et al., 2000), and has been related to long-term control of body mass (Levine and Kotz, 2005).

The energy expended with physical activity also depends on the energetic efficiency with which activities are performed, and these also vary between individuals. In general, the energy expenditure of body mass-bearing activities (walking, running) increases with body mass (Bray et al., 1977; Levine, 2004a), but, when expressed on a per kilogram basis, the energy expended to walk a fixed distance or at a given speed can be as much as two to three times greater for smaller than for larger individuals (Weyand et al., 2010).

### 2.3.3. Growth

Growth increases energy expenditure through synthesis of new tissues. However, except for the first months of life, the energy requirement for growth relative to the total energy requirement is small; it decreases from about 40 % at age one month to about 3 % at the age of 12 months (Butte, 2005).

### 2.3.4. Pregnancy

The effect of pregnancy on energy expenditure varies during the course of pregnancy and differs considerably between individual women. Pregnancy increases REE due to the metabolic contribution of the uterus and fetus to the expenditure of tissue deposition, and to the increased work of the heart and lungs (Forsum and Löf, 2007; Hytten and Chamberlain, 1980). Pregnancy can also affect EEPA.

Energy expenditure due to pregnancy is primarily related to the increased energy needed for tissue maintenance of the increased tissue mass. In the FAO/WHO/UNU report (2004), for REE, an average cumulative increment of 147.8 MJ (35,330 kcal) for a gain in body mass of 12 kg was estimated from studies of well-nourished women who gave birth to infants with adequate body masses (Cikrikci et al., 1999; de Groot et al., 1994; Durnin et al., 1987; Forsum et al., 1988; Goldberg et al., 1993; Kopp-Hoolihan et al., 1999; Piers et al., 1995; Spaaij et al., 1994b; van Raaij et al., 1987). Corresponding cumulative average increases in REE have been observed to be around 5 %, 10 % and 25 % in the first, second and third trimesters, respectively. However, even within populations of well-nourished women, large variations in the effect of pregnancy on REE are observed (Prentice et al., 1989).

Reviews of numerous studies in a variety of countries provide little evidence that women are less active during pregnancy (IoM, 1992; Prentice et al., 1996a), although these studies do not give information about changes in the intensity of the effort associated with habitual tasks. Compared with non-pregnant values, the energy expended for EEPA in the third trimester of pregnancy ranged from a decrease of 22 % to an increase of 17 %, but on average did not differ significantly from non-pregnant women (Butte and King, 2002). However, when expressed per unit of body mass, there was a tendency towards lower EEPA/kg per day. Three recent studies in healthy well-nourished women

reviewed by Forsum and L f (2007) also concluded that EEPA is not significantly increased during pregnancy. TEF, when expressed as a proportion of energy intake, is generally assumed to remain unchanged during pregnancy (Butte and King, 2005; Forsum and L f, 2007; Kopp-Hoolihan et al., 1999; Prentice et al., 1996a), but considerable intra-individual variations occur.

### 2.3.5. Lactation

The main factors that influence the impact of lactation on energy expenditure are the intensity (exclusive or partial) and duration of breastfeeding; this may vary widely between individuals and populations. In exclusively breastfeeding women, the mean amount of milk produced daily was reported to be from 562 to 854 g/day during the first six months *post partum* (Butte et al., 2002; FAO/WHO/UNU, 2004) with an average gross energy content of 2.8 kJ/g (0.67 kcal/g) (Butte and King, 2002; FAO/WHO/UNU, 2004; Garza and Butte, 1986; Goldberg et al., 1991; IoM, 1991; Panter-Brick, 1993; Prentice and Prentice, 1988; WHO, 1985). The energy content of milk sampled at various stages of lactation from healthy mothers of term infants in Europe is shown in Appendix 1.

Increases in REE of 4 to 5 % have been observed in lactating women (Butte et al., 1999; Forsum et al., 1992; IoM, 2005; Sadurskis et al., 1988; Spaaij et al., 1994a) which is consistent with the additional energy cost of milk synthesis (IoM, 2005). However, others have reported similar REE in lactating women compared to the non-lactating state (Frigerio et al., 1991; Goldberg et al., 1991; Illingworth et al., 1986; Motil et al., 1990; Piers et al., 1995; van Raaij et al., 1991). Thus, during lactation there seem to be no significant changes in REE compared with non-pregnant, non-lactating women; furthermore, there also seem to be no significant changes in the efficiency of work performance or TEE (FAO/WHO/UNU, 2004; IoM, 2005).

### 2.3.6. Endocrinological factors

Several hormones, such as the thyroid hormone (al-Adsani et al., 1997; Danforth and Burger, 1984; Silva, 2006), glucagon or epinephrine (Heppner et al., 2010), glucocorticoids (Silva, 2006), insulin, leptin (Belgardt and Bruning, 2010), estrogens and progesterone (Bisdee et al., 1989; Webb, 1986) are implicated in the regulation of energy expenditure, but their impact on the energy expenditure of healthy subjects is generally considered to be minute.

### 2.3.7. Ageing

There is no clear evidence for a decrease in organ metabolic rate, i.e. per gram of tissue mass, in healthy ageing (Gallagher et al., 1996; Gallagher et al., 2000; Krems et al., 2005). There is also no consistent evidence that TEF changes with age. If differences exist they are assumed to be too small to significantly affect energy requirements (Roberts and Dallal, 2005; Roberts and Rosenberg, 2006). Thus, assuming that REE corrected for body composition does not change in older adults, but that sarcopenia and increased adiposity decrease the metabolically active mass, and considering the fact that EEPA decreases with age (Vaughan et al., 1991), the energy requirement in older adults is generally lower (see Section 5.1.4.). For instance, in a longitudinal study in a well-functioning elderly population aged 67 years at baseline the observed decreases per decade were 6 % and 7.5 % for TEE, 3 % and 5 % for REE, and 12.6 % and 10.7 % for EEPA in women and men, respectively (L hrmann et al., 2009).

### 2.3.8. Diet

It has been hypothesised that when long-term energy intake surpasses energy expenditure a facultative component generated by stimulation of the sympathetic nervous system and heat dissipation in the brown adipose tissue can add to the obligatory TEF (see Section 2.2.3) to increase TEE. This phenomenon, termed "Luxuskonsumption" or diet-induced thermogenesis, was first identified in laboratory rodents (Stock and Rothwell, 1981). The discovery that significant depots of brown fat exist in humans has reactivated the hypothesis that diet-induced thermogenesis exists in humans, and thus that excess dietary intake can increase TEE in humans (Schutz et al., 1984; Wijers et al., 2009). However, the relevance of diet-induced thermogenesis and the role of the brown adipose tissue as an

effector of energy balance was challenged from the very beginning in rodents (Hervey and Tobin, 1983), and is now contested in humans (Kozak, 2010).

### **2.3.9. Sex**

In general, absolute REE, and in consequence TEE, is higher in men than in women; these sex-specific differences are mainly due to differences in body mass and body composition (Buchholz et al., 2001; Klausen et al., 1997). There seem to be no significant differences in PAL values between men and women (Roberts and Dallal, 2005).

### **2.3.10. Ethnicity**

Differences in REE have been reported between groups of different ethnic background (e.g. Africans, Asians and Caucasians) and, more recently, specific predictive equations for REE have been developed to take such differences into account (Vander Weg et al., 2004) (see also Section 2.4.). However, these differences in REE in relation to ethnicity are more the consequences of differences in body mass and composition rather than being related to specific ethnic differences in metabolism (Gallagher et al., 2006; Hunter et al., 2000; Wang et al., 2010).

### **2.3.11. Environmental factors**

Temperature is the main environmental factor that can affect energy expenditure. Humans regulate their body temperature within narrow limits (Danforth and Burger, 1984). This process of thermoregulation can elicit increases in energy expenditure when ambient temperature decreases below the zone of thermoneutrality (Valencia et al., 1992). However, because most people adjust their clothing and environment to maintain comfort, and thus thermoneutrality, the additional energy expenditure of thermoregulation rarely affects TEE to an appreciable extent.

## **2.4. Equations to predict resting energy expenditure (REE)**

In practice, predictive equations are used to calculate an individual's REE instead of directly measuring it. Multiplication of REE with a predetermined factor for physical activity will give TEE and energy needs. An accurate prediction of REE is a prerequisite for obtaining an accurate prediction of TEE.

### **2.4.1. Predictive equations for adults**

Equations for predicting REE are historically based on easily measurable parameters such as body mass, height, sex, age and also ethnicity (see Appendix 2). These equations are derived by regression analysis of the data from a group of subjects whose REE is measured by direct or indirect calorimetry. The accuracy of an equation is usually estimated as the percentage of subjects that have a REE predicted by the equation within 10 % of the measured REE (Frankenfield et al., 2005). The mean percentage difference between the predicted and measured REE is considered a measure of accuracy at a population group level.

The majority of predictive equations use body mass as the most important determinant of REE. In addition to body mass, body height, body composition, sex, age and ethnicity can affect REE significantly. Numerous equations have been devised, and are still under development, to take into account one or several of these parameters. The first set of equations was proposed as early as 1919 (Harris and Benedict, 1919) and has been one of the most used set of equations (Daly et al., 1985; FAO/WHO/UNU, 1985, 2004; Schofield et al., 1985). Many new equations have been proposed since then (see Appendix 2). Among these, the equations developed by Owen et al. (1986; 1987), Mifflin et al. (1990), Schofield et al. (1985), Müller et al. (2004), and Henry (2005) are the most widely used. The multitude of new equations, their growing complexity, the fact that many equations have been developed for specific categories of people, in particular overweight and obese subjects (Weijs, 2008), and the continuous use of the historical Harris-Benedict equation illustrate a persistent problem: none of these equations is really satisfactory in the sense that when applied to a group other than the one from which it was derived, significant differences between measured and predicted values can be

observed. Thus, the predictive value of equations can vary substantially according to sex, BMI (which reflects body composition), age and ethnicity of the subjects (Hasson et al., 2011).

In 1985, the two most frequently used equations were the Harris-Benedict and the Schofield equations, but they are suspected to overestimate REE. The Harris-Benedict database included a relatively small number of subjects, with no children or adolescents below the age of 15 years, and a significant number of measurements were obtained by the use of closed-circuit indirect calorimetry, whilst the Schofield database included a large number (~40 %) of physically very active (Italian) subjects.

The main studies that reassessed the historical equations and generated new equations are as follows: Owen et al. (1986; 1987) reported that the Harris-Benedict equation overestimated REE by 12.8 % in women and by 6.4 % in men, and proposed a new set of equations. Mifflin et al. (1990) observed that the Harris-Benedict equation overestimated REE by 5 % in a group of 498 healthy men and women, and developed new predictive equations that are now considered to be among the most relevant and extensively used equations. They also observed that the Owen equations predicted values very close to the REE measured in their study (-4 % in women and 0.1 % in men). Müller et al. (2004) investigated the application of the FAO/WHO/UNU equations (1985) and concluded that the prediction of REE by FAO/WHO/UNU formulas systematically overestimated REE at low REE and underestimated REE at high REE, and proposed alternative equations, some of which include the use of the BMI. Finally, Henry (2005) also developed a new database including 5,794 males and 4,702 females from 166 studies (the Oxford database) that excluded the very active (Italian) subjects of the Schofield database and included more individuals from the tropics (n=4,018). In general, the equations proposed by Henry (2005) (Oxford equations) predict lower REE values than the current FAO/WHO/UNU equations in 18–30 and 30–60 year-old men, and in all women over 18 years of age.

Despite the development of numerous new equations intended to improve the predictive power of the Harris-Benedict and Schofield equations, the FAO/WHO/UNU consultation in 2001 (FAO/WHO/UNU, 2004), after re-analysis of the data and attempts to define new equations (Cole, 2002; Henry, 2001), decided to keep the equations proposed by Schofield and colleagues in 1985 that formed the basis for the equations used by FAO/WHO/UNU in 1985. Analysis of the literature published between 2005 and 2011 in which the Harris-Benedict, Schofield or FAO/WHO/UNU equations were tested and compared (among others) to the more recent Owen, Mifflin, Müller and/or Henry equations shows that the conclusions can be very different between studies and suggest that the more recent equations do not provide a better prediction than the 1919 Harris-Benedict or the present FAO/WHO/UNU equations (Amirkalali et al., 2008; Boullata et al., 2007; Frankenfield et al., 2005; Hasson et al., 2011; Khalaj-Hedayati et al., 2009; Melzer et al., 2007; Weijs et al., 2008; Weijs, 2008; Weijs and Vansant, 2010). Considering the discrepancies in the results of the various publications, there is no reason to favour one set of predictive equation over another, and the Panel concludes that the equations by Harris-Benedict (1919), Schofield et al. (1985), Mifflin et al. (1990), Müller et al. (2004) and Henry (2005) can be considered as equally valid, whereas the equations by Owen et al. (1986; 1987) are not appropriate for this Opinion because of the large BMI range and the very low number of subjects on which they were based.

Overweight and obese subjects: Recently, Weijs (2008) compared the predictive power of 27 published equations in relation to the country of origin (USA versus the Netherlands) and the BMI of the subjects. Using three validation criteria, Weijs reported that the Mifflin equation predicted best for overweight (BMI 25-30) and class I and class II (BMI 30-40) obese US adults, but not for the taller Dutch subjects for which there was no single accurate equation.

Ethnicity/environment: Lower levels of REE in African-American compared to European-American women have been reported (Gannon et al., 2000; Sharp et al., 2002; Weyer et al., 1999). Equations that fail to consider ethnicity may result in inappropriate reference values. In women of African and European origin, Vander Weg et al. (2004) showed that the Owen equation predicted REE best in African-American women but underestimated it in European-American women, whereas the Mifflin equation predicted best in European-American women. They proposed a new equation including an

ethnicity correction factor. As suggested by Müller et al. (2004), Henry (2005) and Frankenfield et al. (2005), the Harris-Benedict and Schofield equations over-predict REE, and more so in women of African rather than European origin. Recently, Yang (2010) showed that the Harris-Benedict, Schofield and Henry equations overpredict REE in healthy Chinese adults, and Nhung (2005) showed that the FAO/WHO/UNU equations overpredicted in Vietnamese adults. Studies on other racial or ethnic groups also demonstrated differences in REE (Benedict, 1932; Henry and Rees, 1991).

#### 2.4.2. Predictive equations for children

For children and adolescents, several equations based on age, body mass, height and sex are available to predict REE, among them those of Schofield et al. (1985), Maffeis et al. (1993), Molnar et al. (1995), Müller et al. (2004) and Henry (2005). Predictive equations solely derived from overweight/obese cohorts of children are not considered here. The equations of Schofield were the mostly used in the past and have been cross-validated in various settings. While some studies have suggested that the Schofield equations provide inadequate estimates for infants (Duro et al., 2002; Thomson et al., 1995) and obese adolescents (Hofsteenge et al., 2010), they showed the best agreement with actual measurements in other studies which compared predicted to actual measurements (Firouzbakhsh et al., 1993; Rodriguez et al., 2002). Both the Schofield and the Henry equations were derived from large datasets covering the age groups from 0 to 18 years, whereas the equations of Maffeis et al. (1993), Molnar et al. (1995) and Müller et al. (2004) were developed from smaller samples not including all age groups (see Table 1). The Panel concludes that the equations of Schofield and Henry are equally valid for predicting REE in children with a wide age range.

**Table 1:** Number of male (m) and female (f) children in the datasets from which the prediction equations for children and adolescents were derived

Age (years)	Harris and Benedict (1919)	Schofield et al. (1985)	Maffeis et al. (1993)	Molnar et al. (1995)	Müller et al. (2004)	Henry (2005)
0-3		162 m 137 f				277 m 215 f
3-10		338 m 413 f	6-10 y: 62 m 68 f		5-11 y: 99 m 89 f	289 m 403 f
10-18	Only few subjects aged 15 y and older	734 m 575 f		10-16 y: 193 m 178 f	12-17 y: 28 m 27 f	863 m 1,063 f

y, years

### 3. Dietary sources of energy and intake data

#### 3.1. Dietary sources of energy

The energy available for metabolism, namely physiologically available energy, is primarily determined by the chemical energy of the food. This is measured in the laboratory as the heat produced when its organic molecules are fully oxidised. The energy content of food as measured by complete combustion is termed gross energy (GE) or ingested energy (IE). Not all chemical energy in foods is available to humans, and the chemical energy value must therefore be corrected for losses due to incomplete digestion and absorption and, with protein, for incomplete oxidation and losses as urea (FAO, 2003). The term metabolisable energy (ME) encompasses the energy available after accounting for losses of the ingested energy in faeces, urine, gases from fermentation in the large intestine, and waste products lost from surface areas. Not all ME is available for the production of ATP. When energy losses such as the heat of microbial fermentation and obligatory thermogenesis are subtracted from ME, the remainder is the energy content of food that will be available to the body for ATP



production, which is referred to as net metabolisable energy (FAO, 2003). In EU legislation<sup>8</sup>, the energy conversion factors for nutrients for labelling purposes have been calculated as ME.

The energy providers in food are carbohydrate, fat, protein and alcohol. The digestibility and absorption of these, and also the heat of combustion, differ depending on the composition and on the foods in which they are found. Correspondingly, energy conversion factors may vary considerably. Specific factors for calculating the energy content of certain foodstuffs have been presented (FAO, 2003; Livesey et al., 1995).

**Carbohydrate and fibre:** The energy conversion factor for carbohydrate presented in food composition tables is in many cases determined by the ‘difference method’, which defines total carbohydrate as the difference between the total dry matter and the sum of protein, fat and ash, and has a general value of 17 kJ/day (4 kcal/g). The energy conversion factor can also be expressed as monosaccharide equivalents (FAO, 2003). The GE for carbohydrate depends on their composition and number of glycosidic linkages, and ranges from 15.6 to 18 kJ/g (Elia and Cummings, 2007). The energy conversion factor ranges from 16 kJ/g (3.75 kcal/g) to 17 kJ/g (4.0 kcal/g) for available mono- and disaccharides (glucose, galactose, fructose, sucrose) and starch and glycogen, respectively (FAO, 2003). The GE of fibre that reaches the colon does not differ substantially from that of starch and glycogen, but due to large differences in the fermentability of dietary fibre in the colon the energy contribution from fibre is less than for other carbohydrates. Assuming that an average of 70 % of the fibre reaching the colon is fermented, the energy conversion factor for fibre is 8 kJ/g (2 kcal/g) (Elia and Cummings, 2007; FAO, 2003).

**Protein:** Protein is not fully oxidised in the body. The physiologically available energy from protein is therefore reduced due to both incomplete digestibility and urea losses in the urine. The digestibility of protein is lowest in legumes (78 % of GE) and highest in animal products (97 % of GE). Protein in food may be measured as the sum of individual amino acid residues. When these values for protein are not available, determination of protein content based on total nitrogen by Kjeldahl (or a comparable method) multiplied by a factor is the generally accepted approach. Based on the different amino acid composition of various proteins, the nitrogen content of protein varies from around 13 to 19 %. This would equate to nitrogen conversion factors ranging from 5.26 to 7.69. As the average nitrogen content of protein is about 16 %, a general factor of 6.25 to convert nitrogen content to (crude) protein content is used. When protein content is determined in this way, the general energy conversion factor of 17 kJ/g (4 kcal/g) should be applied (FAO, 2003; Merrill and Watt, 1973).

**Fat:** The GE of fat depends on the fatty acid composition of the triglycerides and the content of other lipids in the diet. On average, the ME from fat is calculated as 95 % of GE in most foodstuffs. Fats may be analysed as fatty acids and expressed as triglycerides. For dietary fats, a general energy conversion factor of 37 kJ/g (9 kcal/g) is used (FAO, 2003; Merrill and Watt, 1973).

**Alcohol:** Although consumption of alcohol can contribute to the hepatic *de novo* lipogenesis pathway, about 80 % of the energy liberated contributes to ATP production (Prentice, 1995; Raben et al., 2003). Ethanol is promptly oxidised after ingestion and reduces the oxidation of other substrates used for ATP synthesis. The energy conversion factor for alcohol (ethanol) is 29 kJ/g (7 kcal/g).

### 3.2. Dietary intake data

Estimated energy intakes for children and adolescents in 21 countries and for adults in 24 countries in Europe are presented in Appendix 3 and Appendix 4, respectively. The data refer to food consumption

<sup>8</sup> Council Directive 90/496/EEC of 24 September 1990 on nutrition labelling for foodstuffs. OJ L 276, 6.10.1990, p. 40–44. Regulation (EU) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No 1924/2006 and (EC) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/200. OJ L 304, 22.11.2011, p. 18–63.



surveys conducted from 1989 onwards. Most studies comprise representative national population samples.

As shown in Appendices 3A and 4A, there is a large diversity in the methodology used to assess the individual energy intakes of children, adolescents and adults. These differences in dietary assessment methods make direct comparisons difficult. Age classifications may not be uniform and comparability is also hindered by differences in food composition tables used for the conversion of food consumption data to nutrient intake data (Deharveng et al., 1999). Dietary intake data are prone to reporting errors and there is a varying degree of under-reporting in different surveys (Merten et al., 2011).

Although the differences in methodology have an impact on the accuracy of between-country comparisons, the data presented give an overview of the energy intake in a number of European countries. Most studies reported mean intakes and standard deviations (SD) or mean intakes and intake distributions.

Available data show that average energy intakes in children aged two to six years vary between 4.5 MJ/day (1,077 kcal/day) and 7.9 MJ/day (1,890 kcal/day). Boys usually have somewhat higher energy intakes than girls. In older children, average daily energy intakes vary between 6.8 MJ/day (1,625 kcal/day) in boys aged 5-8 years and 13.2 MJ/day (3,145 kcal/day) in boys aged 13-15 years, and between 6.1 MJ/day (1,460 kcal/day) in girls aged 10-14 years and 10.0 MJ/day (2,385 kcal/day) in girls aged 13-15 years. In adolescents, observed average energy intakes are between 9.9 MJ/day (2,364 kcal/day) in boys aged 15-17 years and 14.7 MJ/day (3,504 kcal/day) in boys aged 16-18 years, and between 6.8 MJ/day (1,625 kcal/day) in girls aged 15-18 years and 9.9 MJ/day (2,364 kcal/day) in girls aged 15-17 years (see Appendices 3B to 3F).

In adults, average energy intakes vary between 7.1 MJ/day (1,688 kcal/day) and 15.3 MJ/day (3,657 kcal/day) in men and between 5.7 MJ/day (1,373 kcal/day) and 11.4 MJ/day (2,725 kcal/day) in women. Ranges vary from 3.1 to 8.1 MJ/day (747 to 1,940 kcal/day) at the lower (2.5-10<sup>th</sup> percentile) end to 8.9 to 21.0 MJ/day (2,111 to 5,023 kcal/day) at the upper (90-97.5<sup>th</sup> percentile) end of the intake distributions. The lowest energy intakes are usually observed in older age groups (see Appendices 4B to 4E).

#### **4. Overview of dietary reference values and recommendations**

A number of national and international organisations have estimated energy requirements for all age groups and for pregnant and lactating women. They have generally been estimated as TEE, and TEE has been calculated as the product of REE x PAL, or from regression equations in which age, sex, body mass and, where appropriate, height are considered. REE (or BEE or basal metabolic rate, according to the terminology used in the reports) is estimated from different equations, and PAL values used vary between countries.

##### **4.1. Adults**

Most authorities (AFSSA, 2001; D-A-CH, 2012; FAO/WHO/UNU, 2004; Health Council of the Netherlands, 2001; NNR, 2004; SACN, 2011; SCF, 1993) determined average energy requirements using the factorial approach (Appendix 5). Usually REE was estimated using Schofield's predictive equations based on body mass (FAO/WHO/UNU, 1985). However, AFSSA (2001) used the predictive equations by Black (1996), while SACN estimated REE with the equations by Henry (2005), because they predict slightly lower values compared to Schofield's equations and estimate REE with a higher accuracy, as determined by Weijts et al. (2008) in overweight/obese subjects. The body masses entered into these equations were either derived from observed heights in the respective country and calculated for a desirable BMI within the healthy BMI range, were based on mean population body masses (NNR, 2004), or used incremental body masses within a defined body mass range (FAO/WHO/UNU, 2004). The calculated REE values were then multiplied with PAL values ranging between 1.4 and 2.4. Some authorities assumed lower PAL values for older people (AFSSA, 2001; D-A-CH, 2012; Health

Council of the Netherlands, 2001; SACN, 2011; SCF, 1993), used desirable PAL values (D-A-CH, 2012; Health Council of the Netherlands, 2001; SCF, 1993), and/or defined PAL values or ranges of PAL values representing certain lifestyle activity levels. SACN (2011) derived PAL values from two studies in middle-aged US adults in which TEE was measured with the DLW method and REE was either measured (Moshfegh et al., 2008) or calculated (Tooze et al., 2007). The resulting distribution of PAL values was assumed to also represent PAL values of the UK population, and the median PAL value was used to derive average energy requirements according to age and sex.

IoM (2005) collected DLW data on adults separately for those with normal body mass and for overweight/obese subjects. The normal body mass database comprised 169 men and 238 women with a BMI between 18.5 and 25 kg/m<sup>2</sup>. Based on this database, the IoM derived prediction equations of TEE by nonlinear regression analysis taking into account age, sex, height, body mass, and a physical activity constant. Four physical activity constants were defined as equivalents to a range of PAL values appropriate for sedentary, low active, active, and very active lifestyles. Individual PAL values were determined by dividing the measured individual TEE values by the measured or predicted individual REE values, and PAL values less than 1.0 or greater than 2.5 were omitted.

#### 4.2. Infants and children

It is generally accepted that TEE is different for breast-fed and formula-fed infants. Some authorities derived values for formula-fed infants only (AFSSA, 2001; D-A-CH, 2012; SCF, 1993), whereas others estimated energy requirements according to feeding mode (FAO/WHO/UNU, 2004; SACN, 2011) (Appendices 6, 7).

Older estimates of energy requirements of infants were based on measurements of energy intake (FAO/WHO/UNU, 1985; SCF, 1993). More recent estimated energy requirements have been based on measurements of energy expenditure using DLW data from healthy, well-nourished, full-term infants available from 1987 onwards. The DLW database has subsequently been extended to comprise also older infants and young children up to 24 months of age. The energy expended for growth was estimated from changes in body mass and body composition, i.e. gains in protein and fat mass during growth, and was added to the estimated TEE.

The DLW database used by FAO/WHO/UNU (2004) comprised 13 studies with DLW measurements performed on a total of 417 healthy, well-nourished, non-stunted infants aged 0-12 months and growing along the trajectory of the WHO reference standard (1983). Most (11/13) studies were done in the UK, the US, and the Netherlands. Linear regression analysis using body mass as the predictor for TEE was applied. An allowance for energy deposition in tissues during growth was added, which was calculated by gains in protein and fat, and corresponding energy deposition, assuming that the energy equivalents of protein and fat deposition are 23.6 and 38.7 kJ/g (5.64 and 9.25 kcal/g), respectively (Butte et al., 2000b). Since formula-fed infants had higher TEE during the first year of life, separate predictive equations for breast-fed and formula-fed infants were proposed.

For infants and children up to two years of age, the DLW database of IoM (2005) comprised children within the 3<sup>rd</sup> and 97<sup>th</sup> percentile for US body mass-for-length values. A single equation involving only body mass was found suitable to predict TEE in all individuals irrespective of sex. Because of the small sample size of the data used and the limited range of estimated physical activity, PAL was not included in the TEE equation. The IoM calculated the estimated energy requirements (EERs) for infants and very young children as TEE plus energy deposition for growth. The energy requirement for growth was computed from rates of protein and fat deposition in a longitudinal study of infants (0.5-24 months of age) (Butte et al., 2000b), and applied to the 50<sup>th</sup> percentile of gain in body mass for boys and girls of similar ages (Guo et al., 1991). In children aged 0-36 months, a single equation involving only body mass was found suitable to predict TEE in all individuals irrespective of sex (IoM, 2005). EERs were provided for each sex, for each month between one and 35, taking into account reference body masses for the United States (Kuczmarski et al., 2000) and calculated energy deposition allowance (Butte et al., 2000b; Guo et al., 1991).

For older children and adolescents (Appendix 7), FAO/WHO/UNU (2004) derived quadratic predictive equations with body mass as the single predictor from studies of TEE with a total of 801 boys and 808 girls using either the DLW method or HR monitoring (Torun, 2001). Using these predictive equations, TEE was calculated based on the WHO reference values of body mass-for-age (Torun, 2001; WHO, 1983).

Energy deposited in growing tissues was estimated by multiplying the mean daily gain in body mass at each year of age by the average energy deposited in growing tissues (8.6 kJ or 2 kcal per gram of gain in body mass) using the WHO reference values of body mass-for-age (WHO, 1983). A set of values for mean daily energy requirement (MJ or kcal/day) was calculated for each sex, requirements being the sum of energy deposition and TEE. This was then divided by the median body mass at each year to express requirements as energy units per kilogram of body mass.

To account for less or more physically active lifestyles in children aged six years and older, FAO/WHO/UNU recommended to subtract or add 15 % of energy requirements as estimated with the use of the predictive equations valid for children and adolescents with “average” physical activity (FAO/WHO/UNU, 2004; Torun, 2001). Examples of activities performed with less physically active than average lifestyles as well as for those performed with vigorous lifestyles were given.

For US children aged 3-18 years, EERs were calculated for boys and girls separately, because of variations in growth rate and physical activity (IoM, 2005) (Appendix 7). DLW data from the normative database with US children within the 5<sup>th</sup> and 85<sup>th</sup> percentile of BMI were used to develop equations to predict TEE based on a child’s sex, age, height, body mass and PAL category. Energy deposition was computed based on published rates of gain in body mass (Baumgartner et al., 1986) and estimated rates of protein and fat deposition in children (Fomon et al., 1982) and adolescents (Haschke, 1989). An average of 84 kJ/day (20 kcal/day) for energy deposition for children aged 3-8 years, and of 105 kJ/day (25 kcal/day) for children and adolescents aged 9-18 years (IoM, 2005), was therefore added to the calculated TEE. Taking into consideration US reference body masses and heights (Kuczmarski et al., 2000), a set of energy requirement values was proposed for each sex, each age and the four PAL categories.

For children up to nine years of age, other authorities used the factorial approach to derive estimated energy requirements, except for the SCF (1993) who employed measurements of energy intake following the approach of FAO/WHO/UNU (1985) but without adding an allowance of 5 % to allow for a desirable level of physical activity (Appendix 8). Energy expenditure for growth was either accounted for by adding a fixed percentage to the amount of REE x PAL (NNR, 2004), by slightly increasing the PAL value (SACN, 2011), or by adding average amounts of deposited protein and fat, as well as gain in body mass, for the various age groups and considering expenditure for synthesis (AFSSA, 2001; Health Council of the Netherlands, 2001; SCF, 1993).

For older children and adolescents aged 10 to 17 years, other authorities mostly used the factorial approach to derive energy requirements. REE was usually predicted with the equations developed by Schofield (FAO/WHO/UNU, 1985; Schofield et al., 1985), except for SACN (2011) who used the predictive equations of Henry (2005). PAL values to be multiplied with estimated REE were either based on data from Torun et al. (1996) using the DLW technique (D-A-CH, 2012; Health Council of the Netherlands, 2001; NNR, 2004), were based on calculated average energy expenditure with various daily activities (AFSSA, 2001; SCF, 1993), or were derived from a dataset of all published DLW studies of children aged over one year published until 2006 (SACN, 2011).

### 4.3. Pregnancy

Table 2 lists DRVs for pregnant women set by various authorities (referring to energy intakes above the values for non-pregnant women).

**Table 2:** Overview of Dietary Reference Values (DRVs) for additional energy during pregnancy (in addition to those for non-pregnant, non-lactating women)

	SCF (1993)	Health Council of the Netherlands (2001)	NNR (2004)	FAO/WHO/UNU (2004)	IoM (2005)	SACN (2011)	D-A-CH (2012)
1 <sup>st</sup> trimester			negligible	+0.35 MJ/day (+85 kcal/day)	0	/	+1.1 MJ/day (+255 kcal/day)
2 <sup>nd</sup> trimester	+0.75 MJ/day (+180 kcal/day)		+1.56 MJ/day (+350 kcal/day)	+1.2 MJ/day (+285 kcal/day)	+1.4 MJ/day (+340 kcal/day)	/	(whole pregnancy).
3 <sup>rd</sup> trimester	from the 10 <sup>th</sup> week of pregnancy for women of normal body mass	+1.2 MJ/day (+290 kcal/day) (whole pregnancy)	+2.1 MJ/day (+500 kcal/day)	+2.0 MJ/day (+475 kcal/day)	+1.9 MJ/day (+452 kcal/day)	+0.8 MJ/day (+191 kcal/day)	To be adjusted in case of a change in PAL during pregnancy compared to the non-pregnant state

In the FAO/WHO/UNU report (2004) the extra amount of energy required during pregnancy was calculated, assuming a mean gestational gain in body mass of 12 kg (WHO, 1995a), by two methods, using either the cumulative increment in REE during pregnancy or the cumulative increment in TEE, plus the energy deposited as protein and fat. In the calculation using the increment in REE, it was assumed that the efficiency in energy utilisation to synthesise protein and fat was 90 %. Adjustments for efficiency of energy utilisation were not necessary in the calculations that used the increment in TEE, as TEE measured with DLW includes the energy expenditure for synthesis. The estimates of the additional energy required during pregnancy were very similar using either REE or TEE for the calculation: 323 MJ (77,100 kcal) and 320 MJ (76,500 kcal), respectively. These values, which were based on experimental data, differ by only 4 % from the theoretical estimate of 335 MJ (80,000 kcal) made by the 1981 FAO/WHO/UNU expert consultation (FAO/WHO/UNU, 1985). Averaging the two factorial calculations, the extra energy expenditure of pregnancy is 321 MJ (77,000 kcal) divided into 0.35 MJ/day, 1.2 MJ/day and 2.0 MJ/day (85 kcal/day, 285 kcal/day and 475 kcal/day) during the first, second and third trimesters, respectively.

IoM (2005) determined the EERs during pregnancy by adding the TEE of non-pregnant women, a median change in TEE of 33.5 kJ/week (8 kcal/week) and an energy deposition during pregnancy of 753 kJ/day (180 kcal/day) (factorial method). The median change in TEE per gestational week was calculated based on a dataset of pregnant women with normal pre-pregnancy BMIs (18.5-25 kg/m<sup>2</sup>) and longitudinal DLW measurements of TEE throughout pregnancy. It was found that the energy expenditure of pregnancy was not equally distributed over pregnancy. No increase in energy intake was recommended for the first trimester, as TEE was considered to change little and gain in body mass was considered to be minor. For pregnant women aged 19-50 years the EERs were calculated as follows:

$$EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + \text{additional energy expenditure during pregnancy} + \text{energy deposition}$$

$$1^{\text{st}} \text{ trimester: } EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + 0 + 0$$

$$2^{\text{nd}} \text{ trimester: } EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + 0.67 \text{ MJ} + 0.75 \text{ MJ}$$

$$(EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + 160 \text{ kcal} (=8 \text{ kcal/week} \times 20 \text{ weeks}) + 180 \text{ kcal})$$

$$3^{\text{rd}} \text{ trimester: } EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + 1.1 \text{ MJ} + 0.75 \text{ MJ}$$

$$(EER_{\text{pregnant}} = EER_{\text{non-pregnant}} + 272 \text{ kcal} (=8 \text{ kcal/week} \times 34 \text{ weeks}) + 180 \text{ kcal})$$

For pregnant adolescent women aged 14 to 18 years the same equations were applied, taking into account the adolescent  $EER_{\text{non-pregnant}}$  instead of the adult  $EER_{\text{non-pregnant}}$ .

In the SACN report (2011) it was considered that the energy reference values for pregnancy estimated by the factorial method (as in the reports of FAO/WHO/UNU (2004) and IoM (2005)) exceed energy intakes observed in populations of well-nourished pregnant women giving birth to infants with a body mass in the healthy range. Consequently, it was considered that there was no reason to amend the increment of 0.8 MJ/day (191 kcal/day) in the last trimester previously recommended (DoH, 1991). It was also indicated that women entering pregnancy as overweight may not require the increment, but data were insufficient to derive a recommendation for this subgroup.

The Nordic Nutrition Recommendations (2004) referred to estimations of energy requirement during pregnancy based on TEE and total energy deposition. In women of normal body mass, the additional energy requirement was considered to be negligible in the first trimester and increased by 1.5 MJ/day (350 kcal/day) and 2.1 MJ/day (500 kcal/day) in the second and third trimesters, respectively (NNR, 2004).

The German-Swiss-Austrian reference values (D-A-CH, 2012) stated that for the whole duration of pregnancy an additional 300 MJ (71,100 kcal) was needed, and recommended that this be distributed evenly throughout pregnancy. This corresponds to an additional energy intake of 1.1 MJ/day (255 kcal/day). In case of a change in PAL during pregnancy compared to the non-pregnant state, the additional energy intake was to be adjusted accordingly.

AFSSA (2001) did not set any reference values for energy during pregnancy and commented on the spontaneous adaptation of the energy intakes of women during pregnancy and the importance of a weight gain within the recommended range.

The Health Council of the Netherlands (2001) concluded that the average extra energy expenditure of pregnancy was 1.5 MJ/day (359 kcal/day, based on the factorial method and assuming an unchanged pattern of activity). However, as women generally tend to be less physically active during pregnancy, the extra energy requirement during pregnancy was estimated to be 1.2 MJ/day (287 kcal/day), derived from data based on the DLW technique applied in small sets of Swedish, British and US pregnant women.

SCF (1993) provided estimates of the additional daily energy requirements (from the tenth week of pregnancy) according to pre-pregnancy BMI (18.5-19.9, 20.0-25.9, >25.9 kg/m<sup>2</sup>), considering the corresponding gain in body mass (12.5-18 kg, 11.4-16 kg, 7-11.5 kg). Because of possible adjustment in either physical activity or metabolism by the second trimester of pregnancy, the SCF considered it reasonable to halve the supposed extra energy requirement, which therefore would be 0.75 MJ/day (179 kcal/day) from the tenth week of pregnancy for women with a normal pre-pregnancy BMI.

#### **4.4. Lactation**

Table 3 lists DRVs for energy for lactating women set by various organisations (referring to energy intakes above the values for non-pregnant women).



**Table 3:** Overview of Dietary Reference Values for additional energy during lactation (in addition to those for non-pregnant women)

	0-6 months <i>post partum</i>	From 6 months <i>post partum</i> onwards
<b>SCF (1993)</b>	0-1 months: +1.5 MJ/day (+359 kcal/day) 1-2 months: +1.8 MJ/day (+430 kcal/day) 2-3 months: +1.92 MJ/day (+459 kcal/day) 3-6 months: +1.71 MJ/day (+409 kcal/day)	Minor weaning practice from six months: +1.92 MJ/day (+459 kcal/day) Substantial weaning practice from six months: +0.88 MJ/day (+210 kcal/day)
<b>Health Council of the Netherlands (2001)</b>	+2.1 MJ/day (+502 kcal/day)	
<b>NNR (2004)</b>	+2.0 MJ/day (+478 kcal/day)	
<b>FAO/WHO/UNU (2004)</b>	First six months: In well-nourished women with adequate gain in body mass: +2.1 MJ/day (+ 505 kcal/day).	Second six months: variable.
<b>IoM (2005)</b>	First six months: +1.4 MJ/day (+330 kcal/day)	Second six months: +1.7 MJ/day (+400 kcal/day)
<b>SACN (2011)</b>	First six months: +1.4 MJ/day (+330 kcal/day)	Second six months: depends on breast milk intake of infant and maternal body composition
<b>D-A-CH (2012)</b>	First four months: +2.7 MJ/day (+635 kcal/day). After four months: +2.2 MJ/day (+525 kcal/day) in women exclusively breastfeeding); +1.2 MJ/day (+285 kcal/day) in women gradually introducing complementary feeding). To be adjusted in case of change in PAL compared to the non-pregnant state.	

According to FAO/WHO/UNU (2004), total energy requirements during lactation are equal to those of the pre-pregnancy period, plus the additional demands imposed by the need for adequate milk production. For women exclusively breastfeeding during the first six months *post partum*, the mean energy expenditure over the six-month period is 2.8 MJ/day (807 g milk/day x 2.8 kJ/g, and assuming 80 % efficiency) (675 kcal/day). From the age of six months onwards, when infants are partially breast-fed and milk production is on average 550 g/day, the energy expenditure imposed by lactation is 1.9 MJ/day (460 kcal/day). Fat stores accumulated during pregnancy may cover part of the additional energy needs in the first months of lactation. Assuming an energy factor of 27.2 MJ/kg (6,500 kcal/kg) body mass (Butte and Hopkinson, 1998; Butte and King, 2002), the rate of loss in body mass in well-nourished women would correspond to the mobilisation of 27.2 x 0.8 kg/month = 21.8 MJ/month (5,210 kcal/month), or 0.72 MJ/day (170 kcal/day), from body energy stores. This amount of energy can be subtracted from the 2.8 MJ/day (675 kcal/day) required during the first six months of lactation for milk production in well-nourished (but not in undernourished) women. The result, 2.1 MJ/day (500 kcal/day), is similar to the additional energy required when infants are partially breast-fed after six months of lactation. Energy requirements for milk production in the second six months are dependent on rates of milk production, which are highly variable among women and populations.

In the report of IoM (2005), TEE values were derived from DLW data on lactating women with normal pre-pregnancy BMIs (18.5-25 kg/m<sup>2</sup>) and fully breastfeeding their infants at one, two, three, four and six months *post partum*. These TEE values include the energy needed for milk synthesis. A comparison of the measured TEE of lactating women and the TEE calculated from age, height, body mass and PAL (using the IoM prediction equation for adult women) showed that the differences were minimal.

Therefore, using a factorial approach, the IoM estimated the energy requirements during lactation from the requirement for adult women with a normal body mass, taking into account milk energy outputs, and energy mobilisation from tissue stores (loss of body mass). In the first six months *post*



*partum* it was considered that well-nourished lactating women experienced an average loss of body mass of 0.8 kg/month equivalent to 0.72 MJ/day (170 kcal/day). Stability of body mass was assumed after six months *post partum*. The milk energy output was considered to be around 2.1 MJ/day (500 kcal/day) in the first six months and 1.7 MJ/day (400 kcal/day) in the second six months (calculated from the milk production rate and its energy content). For lactating women (19 years or older), the average requirements (ARs) were set as follows:

$$AR_{\text{lactation}} = AR_{\text{pre-pregnancy}} + \text{milk energy output} - \text{energy from body mass loss}$$

$$\begin{aligned} \text{0-6 months: } AR_{\text{lactation}} &= \text{adult } AR_{\text{pre-pregnancy}} + 2.1 \text{ MJ} - 0.72 \text{ MJ} \\ (AR_{\text{lactation}} &= \text{adult } AR_{\text{pre-pregnancy}} + 500 \text{ kcal} - 170 \text{ kcal}) \end{aligned}$$

$$\begin{aligned} \text{7-12 months: } AR_{\text{lactation}} &= \text{adult } AR_{\text{pre-pregnancy}} + 1.7 \text{ MJ} - 0 \\ (AR_{\text{lactation}} &= \text{adult } AR_{\text{pre-pregnancy}} + 400 \text{ kcal} - 0) \end{aligned}$$

For lactating women aged 14 to 18 years the same equations apply, taking into account the adolescent  $AR_{\text{pre-pregnancy}}$  instead of the adult  $AR_{\text{pre-pregnancy}}$ .

SACN (2011) used the same factorial method as IoM (2005) for the first six months of lactation.

The Nordic Nutrition Recommendations (2004) estimated the extra need for energy during lactation based on the energy content of human milk, approximately 2.8 kJ/g (0.67 kcal/g), multiplied by the production of human milk during the whole weaning period. The average mobilisation of fat from stores to satisfy energy needs was taken into account. An additional energy intake of 2.0 MJ/day (478 kcal/day) during lactation was suggested.

The German-Swiss-Austrian reference values (D-A-CH, 2012) calculated an additional energy requirement for lactating women of 2.7 MJ/day (635 kcal/day) for the first four months *post partum*. After four months, a distinction was made between women exclusively breastfeeding and women partially breastfeeding. For the first group, an additional energy requirement of 2.2 MJ/day (525 kcal/day) was estimated, and for the latter 1.2 MJ/day (285 kcal/day). In case the PAL was changed during lactation compared to the pre-pregnant state, the additional energy intake was to be adjusted accordingly.

AFSSA (2001) did not set any reference values for energy during lactation and commented on the adaptation of the energy expenditure during lactation and the use of body energy stores.

The Health Council of the Netherlands (2001) calculated the average extra energy requirement during lactation based on the energy value of human milk plus the energy required to produce it. Considering the total energy content of human milk to be approximately 2.7 kJ/mL (0.65 kcal/mL) and the average amount of milk secreted to be 800 mL/day, the amount of energy secreted via human milk was calculated to be approximately 2.2 MJ/day (525 kcal/day). Assuming an efficiency of conversion of energy from food to human milk of 80 %, the energy expenditure of lactation was considered to be 2.7 MJ/day (635 kcal/day). Taking into account the average decrease in body fat of 0.5 kg per month of lactation, the Health Council of the Netherlands estimated the average extra energy requirement during lactation to be 2.1 MJ/day (500 kcal/day).

SCF (1993) proposed values for additional energy requirements for lactation derived from the UK COMA Committee (DoH, 1991), but applied an efficiency value of 95 % for milk production. In case of full breastfeeding, additional energy requirements were set for intervals from zero to one, one to two, two to three and three to six months *post partum*, taking into account milk volume, energy expenditure and an average allowance for loss of body mass (0.5 kg/month following delivery). From six months onwards, minor or substantial complementary feeding practices were considered separately, taking into account the same previous three parameters (milk volume, energy expenditure and an average allowance for loss of body mass).

## 5. Criteria and approaches for deriving the Average Requirement (AR) for energy

### 5.1. Criteria

#### 5.1.1. Energy balance

Energy balance is achieved when metabolisable energy intake is equal to TEE, which includes the energy deposited in new tissue in growth and in pregnancy and the energy secreted in milk in lactation. A positive energy balance occurs when energy intake is in excess of these requirements, whereas a negative energy balance occurs when energy needs are not met by intake. When energy balance is maintained over a prolonged period, an individual is considered to be in a steady state. This can include short periods during which the day-to-day balance between intake and expenditure is not obtained. Short-term, day-to-day energy imbalances are associated with the deposition and mobilisation of glycogen and fat. In terms of regulation of body mass it is important to consider the overall energy balance over a prolonged period of time.

Within certain limits, humans can adapt to transient or long-term changes in energy intake through various physiological and behavioural responses related to energy expenditure and/or changes in growth. Energy balance is then achieved at a new steady state. However, adjustments to low or high energy intakes entail biological and behavioural penalties, such as reduced growth velocity, loss of lean body mass, excessive accumulation of body fat, increased risk of disease, forced rest periods, and physical or social limitations in performing certain activities and tasks. Therefore, estimated energy requirements should be based on the amounts of energy necessary and sufficient to maintain energy balance in healthy adult men and women who are maintaining a desirable body mass and level of activity (FAO/WHO/UNU, 2004). Correspondingly, the increments in energy requirements for growth, pregnancy and lactation should be ascertained in healthy children and women with, respectively, desirable growth rates and development or desirable courses of pregnancy and lactation. Ageing is accompanied by changes in energy balance. The heterogeneity in the alteration of body mass, body composition, and physical activity during the course of biological ageing should be taken into account in the derivation of the AR for older adults.

#### 5.1.2. Body mass, body mass index (BMI) and body composition

Because mortality and risk of disease increase with both high and low BMI values, a stable body mass within target BMI values is desirable. An obesity task force has defined the healthy BMI of adults to be between 18.5 and 24.9 kg/m<sup>2</sup> (WHO, 2000). BMI values outside this target range have been found to be associated with increased morbidity and mortality. In this Opinion, a BMI of 22 kg/m<sup>2</sup>, as the midpoint of this range of healthy BMI, has been used for the calculation of average energy requirements of adults.

Stable body mass is a simple indicator of the adequacy of energy intake that matches energy expenditure in the long term. The main disadvantages of relying on body mass and BMI are that they do not reliably reflect body fat, which is an independent predictor of disease risk (IoM, 2005; Willett et al., 1999). Although sophisticated techniques are available to measure precisely fat free mass (FFM) and FM of individuals, these techniques have not generally been applied in clinical and epidemiological studies investigating the associations with morbidity and mortality. Therefore, BMI, although only an indirect indicator of body composition, is used to classify underweight and overweight individuals, and as the target parameter for the AR for energy.

BMI has a different relation to fat and muscle mass among the elderly than among younger individuals due to age-related changes in body mass and its composition. There is also a reduction in stature with age of 1-2 cm/decade, which has been reported to begin at about 30 years of age and to become more rapid at older ages (Sorkin et al., 1999). Because of these age-related changes in elderly populations, BMI may not have the same associations with morbidity and mortality as in young to middle-aged adults. As BMI by itself seems to have only limited explanatory power with regard to morbidity and mortality in older persons, the Panel concludes that additional indices such as body composition

(i.e. FM, FFM, muscle mass, fat distribution, age-related changes in body height) should also be considered when deriving AR for energy for older age groups.

There are specific target BMI values for children because desirable BMI changes with age. On average, a rapid increase of the BMI occurs during the first year of life. The BMI subsequently declines, reaches a minimum around four to six years, and then gradually increases up to the end of growth (“adiposity rebound”) (IoM, 2005; Kuczmarski et al., 2000; Rolland-Cachera et al., 2006). Cut-off points to define underweight and overweight can be established by using growth charts of healthy children living in an environment that supports optimal growth and development such as the most recent WHO Child Growth Standards (Butte et al., 2007; WHO, 2007; WHO Multicentre Growth Reference Study Group, 2006). According to the WHO classifications for overweight and obesity in younger children (from birth to five years), children above +1 SD of the age-specific mean BMI are described as being “at risk of overweight”, above +2 SD as overweight, and above +3 SD as obese. For school-aged children and adolescents, growth curves that accord with the WHO Child Growth Standards for preschool children and the BMI cut-offs for adults were constructed with merged data from the 1977 National Center for Health Statistics (NCHS/WHO) growth reference (1-24 years) and data from the under-fives growth standards’ cross-sectional sample (18-71 months). Overlapping of the age ranges allowed to smooth the transition between the two samples (de Onis et al., 2007). For older children, the WHO adolescence BMI-for-age curves at 19 years closely coincide with adult overweight (BMI 25) at +1 SD, and with adult obesity (BMI 30) at +2 SD. As a result, these SD classifications to define overweight and obesity were applied to children aged 5-19 years (de Onis and Lobstein, 2010).

### 5.1.3. Body mass gain in pregnancy

There is substantial variance in reported gestational increases in body mass (Fraser et al., 2011; Herring et al., 2008) which is the major determinant of the incremental energy needs during pregnancy. The WHO Collaborative Study on Maternal Anthropometry and Pregnancy Outcomes identified gestational increases in body mass associated with an optimal ratio of maternal and fetal health outcomes<sup>9</sup> to be between 10–14 kg (mean, 12 kg) (WHO, 1995a).

Both low and excessive gestational increases in body mass are related to adverse outcomes of pregnancy (IoM/NRC, 2009). Higher maternal gestational increases in body mass are associated with a decreased risk of small-for-gestational-age infants (especially among underweight women) but are associated with an increased risk of large-for-gestational-age infants, low 5-minute Apgar scores, gestational diabetes, preeclampsia, failed labour induction, cesarean delivery, *post partum* infection and *post partum* body mass retention; on the other hand, an inadequate gestational increase in body mass increases the risk of fetal death, preterm labour and delivery, and infants with low body mass at birth (DeVader et al., 2007).

Evidence from the scientific literature is consistent in showing that pre-pregnancy BMI is an independent predictor of many adverse outcomes of pregnancy (IoM/NRC, 2009; Kiel et al., 2007; Stotland et al., 2006). Ranges for an increase in body mass have been recommended according to pre-pregnancy BMI (<18.5 kg/m<sup>2</sup>: 12.5-18 kg, 18.5-24.9 kg/m<sup>2</sup>: 11.5-16 kg, 25.0-29.9 kg/m<sup>2</sup>: 7-11.5 kg, ≥30.0 kg/m<sup>2</sup>: 5-9 kg) (IoM/NRC, 2009). However, lower gestational increases in body mass of 2-10 kg in women with a pre-pregnancy BMI between 20 and 24.9 kg/m<sup>2</sup> (Cedergren, 2007) and even moderate losses of body mass in overweight (-0.03 kg/week) and obese (-0.019 kg/week) women (Oken et al., 2009) have been associated with optimal maternal and fetal outcomes.

The Panel concludes that an intake corresponding to ARs for energy for pregnancy based on a target gestational increase in body mass of around 12 kg is most likely to be associated with optimal

<sup>9</sup> For the mother in terms of maternal mortality, complications of pregnancy, labour and delivery, *post partum* weight retention and lactational performance, and for the infant in terms of fetal growth, gestational duration, mortality and morbidity.

maternal and fetal health outcomes in women with pre-pregnant BMIs in the range between 18.5 and 24.9 kg/m<sup>2</sup> (WHO, 1995a).

#### 5.1.4. Physical activity

A certain amount of habitual physical activity is desirable for biological and social well-being. The health benefits of regular physical activity and improved physical fitness are well documented (Blair et al., 2001) and many of the known health benefits of physical exercise result, either directly or indirectly, from the beneficial effects on the maintenance of a healthy body mass and body composition. Regular exercise may help to preserve (Forbes, 2000) or to increase (Teixeira et al., 2003) FFM, in particular muscle mass. Because FFM has a relatively high metabolic activity (see Section 2.3.1.), it is an important determinant of energy expenditure at rest (Halliday et al., 1979).

There is consensus among experts that a habitual PAL of 1.70 or higher is associated with a lower risk of overweight and obesity, cardiovascular disease, diabetes and several types of cancer, osteoporosis, and sarcopenia (FAO/WHO/UNU, 2004).

Habitual physical activity, and hence TEE, decreases after a given age (Black et al., 1996; Roberts, 1996), and in advanced age PAL values can be very low. In free- and independently living healthy Swedish men and women aged 91-96 years, PAL values were on average only 1.38 (Rothenberg et al., 2000). In a cohort of community-dwelling US older adults (aged 70-82 years) who are described as high-functioning, able to independently perform activities of daily living, and with no evidence of life-threatening illnesses, a wide variation of PAL values was observed, with an overall mean PAL value of 1.70 (Moshfegh et al., 2008; Tooze et al., 2007). Some elderly individuals who have remained physically active are even able to maintain high levels of energy expenditure, with PAL values as high as 2.48 (Reilly et al., 1993; Withers et al., 1998). This indicates that the age at which TEE and energy requirements start decreasing depends on individual, social and cultural factors that promote or limit habitual physical activity among older adults. Information on the relationship between PAL and mortality has been published in a prospective study of healthy older adults (aged 70-82 years) (Manini et al., 2006). Over an average of 6.15 years of follow-up, participants in the upper tertile of EEPA (PAL greater than 1.78) had a significantly reduced risk of all-cause mortality than those in the lowest tertile (PAL less than 1.57).

In children, regular physical activity in conjunction with good nutrition is associated with health, adequate growth and well-being, improved academic performance, and probably with lower risk of disease in adult life (Boreham and Riddoch, 2001; Torun and Viteri, 1994; Viteri and Torun, 1981). Children who are physically active explore their environment and interact socially more than their less active counterparts. There may also be a behavioural carry-over into adulthood, whereby active children are more likely to be active adults, with the ensuing health benefits of exercise (Boreham and Riddoch, 2001).

The level of physical activity within a population is very variable and may deviate from that which is desirable. Thus, ARs for energy based on desirable PALs may promote an energy intake exceeding the actual energy expenditure, and thereby favour an undesirable increase in body mass. The Panel concludes that ARs for energy should be given for specified activity levels in consideration of the actual rather than desirable PALs of population groups.

## 5.2. Approaches

In principle there are two approaches for determining the AR for energy:

The **first** one is the factorial method to estimate TEE. It was originally proposed by FAO/WHO/UNU (1985) and adopted by the most recent FAO/WHO/UNU report for calculating energy requirements of adults (FAO/WHO/UNU, 2004). This approach involves the calculation of TEE as PAL x REE, where REE is predicted from anthropometric measures, and PAL can be estimated either from time-allocated lists of daily activities expressed as PAR values or, alternatively, by dividing TEE (measured by the

DLW method) by REE which was measured by indirect calorimetry or calculated with predictive equations. Advantages of this approach are that it accounts for the diversity in body size, body composition and habitual physical activity among adult populations with different geographic, cultural and economic backgrounds, and therefore can be universally applied (FAO/WHO/UNU, 2004).

The **second** approach is to use TEE, as measured by the DLW method, directly to derive regression equations which describe how TEE varies as a function of anthropometric variables (such as body mass and height) for defined population groups. This approach has been applied by FAO/WHO/UNU (FAO/WHO/UNU, 2004) for children and by IoM (2005) for the US Dietary Reference Intake (DRI) values for energy for all population groups except lactating women. For children and non-pregnant adults (IoM, 2005), the level of physical activity was accommodated within the regression by designating an activity constant for each individual calculated from TEE and REE values in the datasets. One of four physical activity constants representing a predefined PAL range was used (sedentary, low active, active, very active). In this way, sex-specific regression equations for the prediction of TEE were identified based on age, body mass, height and physical activity categories.

Although the DLW dataset assembled for the US DRI report (IoM, 2005) includes most of the UK studies published up until the writing of that report, SACN (2011) considered this dataset as not being suitable for their approach because study subjects were not recruited explicitly as a representative sample of the UK or any other adult population; furthermore, several of these DLW studies involved investigations of physical activity measurement devices (e.g. accelerometers), and specifically recruited subjects with relatively high activity lifestyles. Instead, for adults, SACN (2011) considered two studies in which energy expenditure was measured using the DLW method, i.e. the OPEN study (n=451, 40-69 years) (Subar et al., 2003; Tooze et al., 2007) and the Beltsville study (n=476; 30-70 years) (Moshfegh et al., 2008). Both studies were comprised of an urban population with subjects recruited from the Washington DC metropolitan area who were considered as comparable to the current UK population as regards distribution of BMI values and ethnic mixture. However, no objective measures of physical activity were made in either study. Therefore, regression modelling as an approach to derive AR for energy was abandoned by SACN because of the inability of TEE prediction models to account for variation in EEPA.

The Panel notes that in addition to these objections, the normative database from which the regression equations were derived by IoM (2005) includes only a small number of individuals who were not randomly selected (adults aged 19 to 96 years: 238 women, 169 men; children aged 3-18 years: 358 girls, 167 boys). Furthermore, although SACN considered subjects of the OPEN study (Subar et al., 2003, Tooze et al., 2007) and the Beltsville study (Moshfegh et al., 2008) to be comparable to the UK population, in a validation study with DLW measurements in a small adult population in the UK (n=66) PAL values (1.81 and 1.74 for men and women, respectively) were on average higher than those of the OPEN and Beltsville studies (mean PAL value 1.63) (Ruston et al., 2004). This could indicate either recruitment bias or, in fact, differences between the populations. It is also questionable whether this limited number of subjects from an urban population of the Washington DC area is representative of the European population. Moreover, data are lacking for some age groups (18-29 years and >70 years), or these age groups are under-represented and require the performance of interpolation.

Therefore, consistent with SACN (2011), the Panel decided not to use regression modelling to determine AR for energy for children and adults, and to follow the factorial approach which is supported by larger datasets. For similar reasons related to the available DLW data, the Panel decided not to derive PAL values by dividing TEE (from DLW studies) by REE (measured or estimated).



### 5.3. Derivation of energy requirements of various population groups

#### 5.3.1. Adults

For adults, the application of the factorial method for estimating TEE is considered to be the most suitable as it accounts for the diversity in body size, body composition and habitual physical activity among adult populations with different geographic, cultural and economic backgrounds, and therefore allows a universal application (FAO/WHO/UNU, 2004).

##### 5.3.1.1. Calculation of resting energy expenditure (REE)

The Panel calculated REE for men and women aged 18-79 years based on individual body heights measured in nationally representative surveys in 13 EU countries, and corresponding individual body masses calculated to yield a BMI of 22 kg/m<sup>2</sup> (see Table 4).

**Table 4:** Median of measured body heights and body masses of 16,500 men and 19,969 women in 13 EU Member States<sup>10</sup> compared to body masses calculated for a BMI of 22 kg/m<sup>2</sup>

Age (years)	n	Measured body height (cm) Median	Measured body mass <sup>(a)</sup> (kg) Median	Body mass <sup>(b)</sup> (kg) assuming a BMI of 22 kg/m <sup>2</sup> Median
<b>Men</b>				
18 - 29	2,771	178	75.0	69.7
30 - 39	2,971	178	82.0	69.7
40 - 49	3,780	177	82.0	68.5
50 - 59	3,575	175	82.0	67.4
60 - 69	2,611	174	80.0	66.4
70 - 79	792	172	80.0	65.1
<b>Women</b>				
18 - 29	3,589	164	60.0	59.4
30 - 39	3,866	164	63.8	59.2
40 - 49	4,727	163	66.0	58.5
50 - 59	4,066	162	68.0	57.7
60 - 69	2,806	160	67.0	56.3
70 - 79	915	159	63.5	55.6


(a): n values for this variable slightly differ.

(b): Body masses calculated for individual measured body heights assuming a BMI of 22 kg/m<sup>2</sup>.

For the prediction of REE with the equations of Harris-Benedict (1919), Schofield et al. (1985), Mifflin et al. (1990), Müller et al. (2004) and Henry (2005) (see Section 2.4.1.), individual data from 36,469 subjects were used (for details of the database and calculation see Appendices 9 and 11). Figure 1 illustrates the median REE values according to age group and sex obtained with the respective equations. Predicted REE decreases with age for both sexes. For women, the Mifflin predictive equation predicted lowest values in all age groups  $\geq 30$  years. For men, there is no equation that always predicted the lowest values. For both sexes, the Harris-Benedict equation predicted highest values for ages 18-29 and 30-39 years, whereas Schofield predicted highest values for ages 40-49 and 50-59 years.

<sup>10</sup>Bulgaria, Czech Republic, Finland, France, Germany, Ireland, Luxembourg, Poland, Portugal, Slovakia, Spain, The Netherlands, United Kingdom

As is illustrated in Figure 1, the discrepancy in the results for REE calculated with the various prediction equations becomes larger with increasing age (from 5 % at age 18-29 years to 11-13 % at age 70-79 years).



**Figure 1:** REE (median) for adult men (n=16,500) and adult women (n=19,969) calculated with the equations of Harris-Benedict (1919), Schofield et al. (1985), Mifflin et al. (1990), Müller et al. (2004) and Henry (2005) using body heights measured in nationally representative surveys in 13 EU Member States and corresponding body masses to yield a BMI of 22 kg/m<sup>2</sup>

The predictive validity of these equations with regard to older adults has been tested in a sample of free-living older persons. A dataset of measurements of REE by indirect calorimetry in 551 elderly subjects (385 women and 165 men, age range 60-96 years) participating in the GISELA study (Lührmann et al., 2010) was used. Agreement between REE predicted with the equations listed above and measured REE was assessed by the method of Bland-Altman (Bland and Altman, 1987). The results confirm the differences in the accuracy of the various equations (Appendix 10). In the female subjects the equation of Schofield et al. (1985) performed best, followed by the equations of Müller et al. (2004), Henry (2005), Harris-Benedict (1919) and Mifflin et al. (1990), while in the male subjects the equation of Müller et al. (2004) performed best, followed by the equations of Henry (2005), Schofield et al. (1985), Harris-Benedict (1919) and Mifflin et al. (1990). With the exceptions of the equation of Schofield et al. (1985) for females and the equation of Müller et al. (2004) for males, all equations underestimated REE as compared to measured values. This evaluation also confirms that the equation of Mifflin et al. (1990) underestimates REE considerably, at least in GISELA subjects (who, however, are not a representative sample for this age range and can be considered as more active and health conscious than average). The accuracy of REE values predicted by these equations, compared with REE measured by indirect calorimetry varied from 74 % (Schofield equation) to 33 % (Mifflin equation) and from 72 % (Müller equation) to 57 % (Mifflin equation) for female and male subjects, respectively.

The Panel considers that there is presently no equation to accurately predict REE, even at a group level, because of differences in average body mass, height and body composition between populations, the influence of sex and ethnicity on body mass and composition, and the decrease in REE that occurs with ageing.

For this Opinion, the Panel applied all five predictive equations. The range of predicted REE obtained with the equations of Harris and Benedict (1919), Schofield et al. (1985), Mifflin et al. (1990), Müller et al. (2004) and Henry (2005) is obvious from the respective lowest and highest median values calculated with these equations (Table 6). Depending on the equation used, respective results for lowest and highest median REE differ between 0.31 MJ/day (74 kcal/day) and 0.68 MJ/day (162 kcal/day) within a sex and age group. Because FFM is the main determinant of REE, at a given BMI equations yielding values for REE at the lower end may be more appropriate for populations with a higher percentage of body fat, compared to those with a higher percentage of FFM.

#### 5.3.1.2. Selection of physical activity level (PAL) values

FAO/WHO/UNU (1985), as well as SACN (2011), identified the lower limit of PAL to be 1.27, which is consistent with studies performed in a calorimeter on non-ambulatory chair-bound and non-exercising subjects (where PAL values of 1.17-1.27 were observed) (Black, 1996). The lower limit of energy expenditure in subjects performing only the minimal activities associated with daily living is between 1.35 and 1.4 (Alfonzo-Gonzalez et al., 2004; Goran et al., 1994b; SACN, 2011; Warwick, 2006). Minimal activities of daily living are usually confined to activities like eating, personal hygiene, dressing, and walking short distances such as from a bed to a chair, and to different locations in a room and corridor outside the room. The upper limit to human physical activity is that exhibited for limited periods of time by elite endurance athletes and soldiers on field exercises, for whom PAL values between 3 and 4.7 have been reported (Black, 1996; Hoyt and Friedl, 2006). The maximum PAL value associated with a sustainable lifestyle within the general population appears to be about 2.5 (Black et al., 1996; SACN, 2011; Westerterp and Plasqui, 2004) and can be expected in people engaging frequently in heavy physical work or in strenuous leisure activities for several hours (Black, 1996; FAO/WHO/UNU, 2004; Withers et al., 1998). Low active people can increase their PAL significantly by regular exercise. Examples of the extent of changes in PAL associated with various activities can be extracted from studies that imposed a programme of training on free-living people normally undertaking very little strenuous exercise (Bingham et al., 1989; Blaak et al., 1992; Westerterp et al., 1992). For example, in these studies, 60 minutes of brisk walking at between >6 and <7.5 km/h daily resulted in an increase in PAL of 0.2 while 60 minutes of jogging at 9 km/h daily increased PAL value by 0.4.

SACN derived PAL values from the combined dataset of the OPEN (Subar et al., 2003, Tooze et al., 2007) and the Beltsville study (Moshfegh et al., 2008) by dividing TEE by REE measurements (SACN, 2011). There was no evidence of any significant variation of PAL with either body mass or sex. Regression analysis did show that PAL values decrease slightly with age. However, age explained < 1 % of the variance (i.e. PAL = 1.69 at 30 years and 1.63 at 70 years). It was concluded that energy reference values can be defined independently of age at least up to the age of 70 years.

A meta-analysis of studies that involved a total of 319 men and women aged 18-64 years showed a modal value for PAL of 1.60 (range 1.55 to 1.65) for both men and women (Black et al., 1996). For the most part, subjects were from affluent societies in developed countries. Typical sub-populations included students, housewives, white-collar or professional workers, and unemployed or retired individuals; only three people were specifically identified as manual workers. Hence, the authors of the meta-analysis defined the study participants as people living a “predominantly sedentary Western lifestyle”.

SACN (2011) used the distribution of PALs observed in the combined OPEN (Subar et al., 2003; Tooze et al., 2007) and Beltsville (Moshfegh et al., 2008) datasets and defined the median PAL (1.63) as the assumed population activity level, and the 25<sup>th</sup> (1.49) and 75<sup>th</sup> percentile (1.78) boundary PAL values as reference values for population groups of men and women thought to be less or more active than average. Although this approach of deriving PAL values took advantage of the measurement of TEE in free-living conditions by the DLW method, the Panel decided not to adopt it for defining reference PAL values for the population of the EU because of the limitations outlined under Section 5.2..

Furthermore, within the dataset of DLW studies in healthy adults assembled for the SACN report, all of the studies which report PAL values were examined for descriptions of the activities/lifestyles of the subjects. PAL values were assigned to three categories of light, moderate or heavy activity. The values show that the range of PAL values is considerable within subjects classified as exhibiting similar lifestyles, and demonstrate only a weak relationship between lifestyle or self-reported physical activity and PAL.

The Panel therefore decided to apply PAL values of equal steps within the observed range of physical activity levels associated with a sustainable lifestyle for calculating AR for energy. In this way, PAL values can be allocated to lifestyles where values of 1.4, 1.6, 1.8, 2.0 and >2.0 could indicate low active (sedentary), moderately active, active, very active and highly active lifestyles, respectively.

Examples of lifestyles which may roughly correlate with PAL values, keeping in mind that the association may be weak as discussed above, are presented in Table 5 and Appendix 12A, while Appendix 12B lists contributions of various activities to PAL values.

**Table 5:** Examples of lifestyles associated with certain physical activity levels (PALs) estimated over 24 hours

Estimation of PAL values from various types of occupations and leisure-time activities	Time spent (hours)			
	No occupation	Office work	Standing work	Physical work
Sleeping (PAR <sup>(a)</sup> 0.95)	8	8	8	8
Lying quietly and awake, listening to music, watching TV (PAR 1.3)	11	3	3	3
Household tasks (light effort), eating, family interactions (PAR 2.3)	3.5	3.5	3.5	3.5
Outside activities e.g. shopping (PAR 3)	0.5	0.5	0.5	0.5
Commuting by bus/car/train (PAR 1.3)	1	1	1	1
Walking ~5 km/h to work (PAR 3.5)	0	0	0	0
Office work, assuming six hours sitting (PAR 1.5) and two hours standing (PAR 2.5) = Mean PAR 1.75	0	8	0	0
Predominantly standing work, assuming six hours standing-working (PAR 3) and two hours seated (PAR 1.5) = Mean PAR 2.6	0	0	8	0
Physically active work, assuming two hours low active (PAR 1.5), three hours standing-working (PAR 3) and three hours hard physical work (PAR 6) = Mean PAR 3.75	0	0	0	8
<b>Resulting PAL</b>	1.4	1.5	1.8	2.2
Daily exercise time (hours) at PAR 7 (jogging/cycling/dancing/rowing stationary (100 watts)) required to reach a PAL of 1.7	1.2	0.6	0	0
<b>Resulting PAL with active lifestyle</b>				
1.5 hours resting (PAR 1.3) switched to more active behaviour such as 0.5 hour shopping (PAR 3) and one hour of light activity at home (domestic tasks, cooking, light gardening) (PAR 2.3), and walking (one hour) rather than going by car (PAR 3.5 vs. 1.5)	1.5	1.7	2.0	2.3
Daily exercise time (hours) at PAR 7 (jogging/cycling/dancing/rowing stationary (100 watts)) required to reach a PAL of 1.7	0.6	0	0	0

(a): PAR, physical activity ratio.

Based on data from Ainsworth et al. (2011).

### 5.3.1.3. Ranges of Average Requirement (AR) for energy for adults

Estimated ARs of adults in the EU, based on the factorial method using lowest and highest median REE, respectively, calculated as described above (Section 5.3.1.1.), and PAL values of 1.4 through 2.4 in steps of 0.2 increments, are presented in Table 6. The figures illustrate the variability in AR among

the population of the EU as a function of REE and PAL. The impact of the variability in REE, resulting from the use of the different predictive equations, on AR at a given BMI and PAL is in the range of 0.43 MJ/day (104 kcal/day) up to 1.6 MJ/day (382 kcal/day).

**Table 6:** Ranges of Average Requirement (AR) for energy for adults with a BMI of 22 kg/m<sup>2</sup> and at six different physical activity levels (PALs)

Age (years)	Lowest median REE (MJ/day)	Highest median REE (MJ/day)	Range of AR at PAL=1.4 (MJ/day) <sup>(a)</sup>	Range of AR at PAL=1.6 (MJ/day) <sup>(a)</sup>	Range of AR at PAL=1.8 (MJ/day) <sup>(a)</sup>	Range of AR at PAL=2.0 (MJ/day) <sup>(a)</sup>	Range of AR at PAL=2.2 (MJ/day) <sup>(a)</sup>	Range of AR at PAL=2.4 (MJ/day) <sup>(a)</sup>
<b>Men</b>								
18 - 29	7.0	7.4	9.8 - 10.3	11.2 - 11.8	12.6 - 13.3	14.0 - 14.7	15.4 - 16.2	16.8 - 17.7
30 - 39	6.8	7.0	9.5 - 9.8	10.8 - 11.2	12.2 - 12.6	13.5 - 14.0	14.9 - 15.4	16.2 - 16.8
40 - 49	6.6	6.9	9.2 - 9.7	10.5 - 11.1	11.9 - 12.5	13.2 - 13.9	14.5 - 15.3	15.8 - 16.7
50 - 59	6.3	6.9	8.8 - 9.6	10.0 - 11.0	11.3 - 12.4	12.5 - 13.8	13.8 - 15.1	15.0 - 16.5
60 - 69	5.9	6.4	8.3 - 9.0	9.5 - 10.3	10.7 - 11.5	11.8 - 12.8	13.0 - 14.1	14.2 - 15.4
70 - 79	5.5	6.2	7.7 - 8.7	8.8 - 9.9	9.9 - 11.2	11.0 - 12.4	12.2 - 13.6	13.3 - 14.9
<b>Women</b>								
18 - 29	5.6	5.9	7.9 - 8.3	9.0 - 9.5	10.1 - 10.7	11.2 - 11.9	12.4 - 13.0	13.5 - 14.2
30 - 39	5.3	5.7	7.5 - 7.9	8.6 - 9.1	9.6 - 10.2	10.7 - 11.4	11.8 - 12.5	12.8 - 13.6
40 - 49	5.1	5.5	7.2 - 7.7	8.2 - 8.8	9.2 - 9.9	10.2 - 11.1	11.3 - 12.2	12.3 - 13.3
50 - 59	4.8	5.5	6.8 - 7.7	7.7 - 8.8	8.7 - 9.9	9.7 - 11.0	10.6 - 12.1	11.6 - 13.2
60 - 69	4.6	5.0	6.5 - 7.0	7.4 - 8.0	8.3 - 9.1	9.2 - 10.1	10.1 - 11.1	11.1 - 12.1
70 - 79	4.3	5.0	6.0 - 6.9	6.9 - 7.9	7.7 - 8.9	8.6 - 9.9	9.5 - 10.9	10.3 - 11.9

(a): Based on lowest and highest median REE (see Appendix 9).

The ranges in kcal/day of AR for energy for adults are tabled in Appendix 14A.

### 5.3.2. Infants

#### 5.3.2.1. Total energy expenditure (TEE)

Published mean data on the TEE of infants living in developed and developing countries showed that TEE increases linearly with age, and, standardised by body mass, ranges from 255 to 393 kJ/kg (61-94 kcal/kg) per day (Butte, 2005). TEE of breast-fed infants was shown to be lower than that of formula-fed infants (Butte et al., 1990; Butte et al., 2000a; Davies et al., 1990; Jiang et al., 1998), however differences in TEE between the groups diminished after the first year of life (Butte et al., 2000a).

Because of the differences between infants initially breast- or formula-fed for four months after birth, separate regression equations for TEE as a function of body mass were obtained for these two groups (Butte, 2005).

According to Butte (2005), TEE for breast-fed infants can be predicted as follows:

$$\text{TEE (MJ/day)} = -0.635 + 0.388 \text{ kg}; n=195, r=0.87, \text{SEE}=0.453 \text{ MJ/day}$$

$$\text{TEE (kcal/day)} = -152.0 + 92.8 \text{ kg}; \text{SEE}=108 \text{ kcal/day}$$

(n=number of observations; SEE=standard error of estimate)

Butte reported that TEE was 12, 7, 6 and 3 % higher in formula-fed compared to breast-fed infants at 3, 6, 9 and 12 months, respectively, suggesting that energy requirements of formula-fed infants may be slightly higher than those of breast-fed infants. However, the Panel considers that the data on which the equation for initially formula-fed infants is based may no longer be appropriate because of recent significant changes in the composition of infant formula (e.g. a protein to energy ratio closer to human milk), and therefore proposes that the equation for initially breast-fed infants is applied when calculating TEE of formula-fed infants.



### 5.3.2.2. Energy deposition in new tissue

TEE measured using the DLW method includes the energy expended in tissue synthesis, but not the energy deposited in growing tissues. Therefore, the latter should be added when calculating the AR for energy for infants. Energy deposited in new tissue was estimated from a multi-component body composition model (total body water, total body potassium and bone mineral content) (Butte et al., 2000b) based on a modified version of Fomon's term infant reference (Fomon et al., 1982) describing changes in body composition during growth. Estimates of protein and fat gain over three-month periods were used to predict energy accrued per gram of gain in body mass (Table 7).

**Table 7:** Energy content of tissue deposition during the second half of infancy (Butte et al., 2000b; Butte, 2005; FAO/WHO/UNU, 2004)

Age interval (months)	Protein gain (g/day)	Fat mass gain (g/day)	Gain in body mass (g/day)	Energy deposited in growing tissues <sup>(a)</sup> (kJ/g)
<b>Boys</b>				
6-9	2.3	0.5	11.8	6.2
9-12	1.6	1.7	9.1	11.4
<b>Girls</b>				
6-9	2.0	0.8	10.6	7.4
9-12	1.8	1.1	8.7	9.8

(a): Taking into account that 1 g protein = 23.6 kJ; 1 g fat = 38.7 kJ.

The estimates of energy deposited in new tissue are applied to the gain in body mass observed in the WHO Growth Standards for infants (2006) to estimate rates of energy deposition at monthly intervals. These predictions of energy deposited during growth derive from a relatively small study by Butte et al. (2000b) which was validated against other datasets (Butte, 2005). The Panel notes that the evolution of body mass and composition studied by Butte (2005), especially regarding gains in FM and FFM during the first year of life, differs from other studies (de Bruin et al., 1998; Fields et al., 2011; Fomon et al., 1982). Since the impact on energy requirements was only marginal, the Panel decided to use the values proposed by Butte in line with FAO/WHO/UNU (2004) and SACN (2011). It is assumed that these values for the energy deposited in new tissue are appropriate for infants growing according to the WHO body mass velocity values, even though in the original study (Butte et al., 2000b) the pattern of breastfeeding followed was not fully described and the growth of infants did not fully reflect the WHO growth trajectory (WHO Multicentre Growth Reference Study Group, 2006).

### 5.3.3. Children

As with adults, the application of the factorial method for estimating TEE seems the most suitable for children as the advantages mentioned previously make it an approach well-fitted for the European context. Moreover, this approach allows estimating AR for energy for children and adolescents based on body masses and heights from the WHO Growth Standards for children up to two years of age (WHO Multicentre Growth Reference Study Group, 2006) and from recent harmonised growth curves for EU children (van Buuren et al., 2012) (Table 8).

**Table 8:** Median body heights and body masses from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006) and from harmonised growth curves for children in the EU (van Buuren et al., 2012)

Age (years)	Median body height (m) <sup>(a)</sup>		Median body mass (kg) <sup>(a)</sup>	
	Boys	Girls	Boys	Girls
1	0.76	0.74	9.6	8.9
2	0.88	0.86	12.2	11.5
3	0.97	0.96	14.7	14.2
4	1.04	1.03	17.0	16.4
5	1.11	1.10	19.2	18.7
6	1.17	1.16	21.5	21.1
7	1.23	1.22	24.3	23.8
8	1.30	1.28	27.4	26.8
9	1.35	1.34	30.6	30.0
10	1.40	1.40	33.8	33.7
11	1.45	1.46	37.3	37.9
12	1.51	1.52	41.5	42.6
13	1.58	1.58	46.7	47.5
14	1.65	1.61	52.7	51.6
15	1.71	1.63	59.0	54.6
16	1.75	1.64	64.1	56.4
17	1.77	1.64	67.5	57.4

(a): For children aged 1-2 years, data were taken from WHO Multicentre Growth Reference Study Group (2006); for older children, data are from van Buuren et al. (2012).

#### 5.3.3.1. Calculation of resting energy expenditure (REE)

From the available prediction equations for REE in children, the equations from Schofield et al. (1985) and Henry (2005) (Appendix 13) were derived from a large number of subjects covering the age range from 0 to 18 years, and therefore are both used to calculate REE. For the ages 1-2 years, median body masses and heights from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006) were used in the equations to calculate the REE, whereas for children aged 3-17 years, the 50<sup>th</sup> percentiles of recently calculated reference body masses and heights for EU children (van Buuren et al., 2012) were used (for details of the database and computation of reference body heights and body masses see Appendix 11). Because the equations of Schofield et al. (1985) and Henry (2005) have overlapping age bands (0-3, 3-10, 10-18 years) the choice of equation is ambiguous at the age boundaries. Following the approach of SACN (2011) and the observation that the transition of the predicted values for the three age bands is smoother, the REE equation for 3-10 year-olds is used for the 3 year-olds, and the equation for 10-18 year-olds is used for those aged 10 years. The results reveal that REE calculated with these two equations are very similar and differ at most by 0.26 MJ/day (62 kcal/day) in some age and sex groups (Table 9).

#### 5.3.3.2. Selection of physical activity levels (PALs)

PAL values for children and adolescents were derived from measurements of TEE and REE. These values vary considerably according to lifestyle, geographic habitat and socioeconomic conditions, and inter-individual coefficients of variability as high as  $\pm 34\%$  (Torun, 2001) have been reported. As indicated in the FAO/WHO/UNU report (2004), most studies were carried out on random or convenient samples, and therefore may not have captured the full range of potential physical activity.

SACN (2011) derived PAL values from a dataset of all published DLW studies in children aged over one year, including those studies assembled by Torun (2005) and other studies published until 2006. Among these studies, seven were from Sweden, six from the UK and two from the Netherlands. The analysis revealed no influence of sex but an increase in PAL values with age. From an early age, however, there was a wide range of mean PAL values so that variation in PAL at any age was much greater than variation with age itself. Nevertheless, three age groups were identified within which the distribution of PAL values could be observed: 1-3 years, >3-<10 years and 10-18 years. These age

ranges also correspond to the age ranges for which REE prediction equations have been generated by both Schofield et al. (1985) and Henry (2005).

As for adults, SACN (2011) calculated the AR for children using the median (PALs 1.39, 1.57 and 1.73 for ages 1-3, >3-<10 and 10-18 years, respectively), 25<sup>th</sup> percentile (PALs 1.35, 1.42 and 1.66 for ages 1-3, >3-<10 and 10-18 years, respectively) and 75<sup>th</sup> percentile (PALs 1.43, 1.69 and 1.85 for ages 1-3, >3-<10 and 10-18 years, respectively) PAL values.

The Panel decided to rely on these results for defining the ranges of PAL values in children, for the reasons mentioned already in Section 5.3.1.2., and not to use the observed median and centile PAL values but, analogously to adults, to apply PAL values of equal steps within the observed ranges of PALs in the respective age groups for computing AR for energy. Thus, PAL values applied for estimating AR are as follows: 1.4 and 1.6 for the age group 1-3 years; 1.4, 1.6, 1.8, 2.0 and 2.2 for the age group >3- <10 years; and 1.4, 1.6, 1.8, 2.0, 2.2 and 2.4 for the age group 10-18 years (Table 9).

Examples of populations with PAL values at the lower range, or less active than average, are children and adolescents who spend several hours every day at school or in sedentary occupations, do not practise physical sports regularly, generally use motor vehicles for transportation, and spend most leisure time in activities that require little physical effort, such as watching television, reading, using computers or playing without much body displacement. Examples of populations with PAL values at the upper level, vigorous lifestyles, or that are more active than average, are children and adolescents who walk long distances every day or use bicycles for transportation, engage in high energy-demanding occupations, perform high energy-demanding chores for several hours each day, and/or practise sports or exercise that demand a high level of physical effort for several hours, several days of the week (FAO/WHO/UNU, 2004).

#### 5.3.3.3. Energy expenditure of children and adolescents for growth

Energy needs for growth have two components: 1) the energy used to synthesise growing tissues, and 2) the energy deposited in those tissues.

Energy spent in tissue synthesis is part of TEE. Due to the marked fall in deposited energy during the first year of life, the deposited energy accounts for only a relatively small proportion (<2 %) of the total energy needs of children at all ages after the first year of life (see Section 2.3.3.).

The composition of newly accrued tissue mass during growth was based on measurements at one and two years of age (Butte et al., 2000a; Butte et al., 2000b; Butte, 2001). Assuming that the composition of normally growing tissues does not change much between the end of infancy and the onset of puberty, the average amount of energy deposited in growing tissues is about 8.6 kJ (2 kcal) per gram of gain in body mass (Butte et al., 2000b; Butte, 2001; Torun, 2005). Even if this amount of energy was over- or underestimated by 50 %, it would only produce an error of about  $\pm 1$  % in the calculations of energy requirements in childhood and adolescence. In the report by FAO/WHO/UNU (2004), the energy deposited in growing tissues was estimated by multiplying the mean daily body mass gain at each year of age between 1 and 17 years by the average energy deposited in growing tissues. It was estimated that the amount of energy deposited is covered by an average increase in PAL of 1 % (FAO/WHO/UNU, 2004; James and Schofield, 1990).

#### 5.3.3.4. Ranges of Average Requirement (AR) for energy for children and adolescents

Estimated AR of children and adolescents in the EU, based on the factorial method using median REE calculated as described above (Section 5.3.3.1.) and PAL values of 1.4 to 2.4 in incremental steps of 0.2, are presented in Table 9. The figures illustrate the variability in AR among children and adolescents in the EU depending on age, sex and PAL values. The figures also reveal that estimated AR based on REE values calculated with the two equations are very similar and differ only in some age and sex groups by at most 0.62 MJ/day (149 kcal/day).

**Table 9:** Range of Average Requirement (AR) for energy for children and adolescents based on the factorial method for predicting REE, at different physical activity levels (PALs)

Age (years)	REE <sup>(a)</sup> (Henry) (MJ/day)	REE <sup>(a)</sup> (Schofield et al.) (MJ/day)	Range of AR <sup>(b)</sup> at PAL=1.4 (MJ/day)	Range of AR <sup>(b)</sup> at PAL=1.6 (MJ/day)	Range of AR <sup>(b)</sup> at PAL=1.8 (MJ/day)	Range of AR <sup>(b)</sup> at PAL=2.0 (MJ/day)	Range of AR <sup>(b)</sup> at PAL=2.2 (MJ/day)	Range of AR <sup>(b)</sup> at PAL=2.4 (MJ/day)
<b>Boys</b>								
1	2.3	2.3	3.2 - 3.3	3.6 - 3.7				
2	3.0	3.0	4.2 - 4.3	4.8 - 4.9				
3	3.5	3.5	4.9 - 4.9	5.6 - 5.6				
4	3.7	3.7	5.2 - 5.3	6.0 - 6.0	6.7 - 6.8	7.5 - 7.5	8.2 - 8.3	
5	3.9	3.9	5.5 - 5.6	6.3 - 6.4	7.1 - 7.2	7.9 - 8.0	8.7 - 8.8	
6	4.2	4.1	5.8 - 5.9	6.7 - 6.7	7.5 - 7.6	8.4 - 8.4	9.2 - 9.3	
7	4.4	4.4	6.2 - 6.3	7.1 - 7.2	8.0 - 8.1	8.9 - 9.0	9.8 - 9.8	
8	4.7	4.7	6.6 - 6.7	7.6 - 7.6	8.5 - 8.6	9.5 - 9.5	10.4 - 10.5	
9	5.0	5.0	7.0 - 7.0	8.1 - 8.1	9.1 - 9.1	10.1 - 10.1	11.1 - 11.1	
10	5.0	5.3	7.1 - 7.4	8.1 - 8.5	9.1 - 9.6	10.1 - 10.6	11.1 - 11.7	12.1 - 12.7
11	5.3	5.5	7.5 - 7.8	8.5 - 8.9	9.6 - 10.0	10.7 - 11.2	11.8 - 12.3	12.8 - 13.4
12	5.6	5.8	8.0 - 8.3	9.1 - 9.4	10.2 - 10.6	11.4 - 11.8	12.5 - 13.0	13.6 - 14.2
13	6.0	6.2	8.5 - 8.8	9.8 - 10.1	11.0 - 11.3	12.2 - 12.6	13.4 - 13.9	14.6 - 15.1
14	6.5	6.7	9.2 - 9.5	10.5 - 10.8	11.8 - 12.2	13.1 - 13.5	14.5 - 14.9	15.8 - 16.2
15	7.0	7.1	9.9 - 10.1	11.3 - 11.6	12.7 - 13.0	14.1 - 14.4	15.5 - 15.9	16.9 - 17.3
16	7.4	7.5	10.4 - 10.6	11.9 - 12.2	13.4 - 13.7	14.9 - 15.2	16.4 - 16.7	17.9 - 18.2
17	7.6	7.8	10.8 - 11.0	12.3 - 12.5	13.8 - 14.1	15.4 - 15.7	16.9 - 17.3	18.4 - 18.8
<b>Girls</b>								
1	2.1	2.1	2.9 - 3.0	3.3 - 3.4				
2	2.8	2.8	3.9 - 4.0	4.4 - 4.5				
3	3.2	3.2	4.5 - 4.6	5.2 - 5.2				
4	3.5	3.4	4.8 - 4.9	5.5 - 5.6	6.2 - 6.3	6.9 - 7.0	7.6 - 7.7	
5	3.7	3.6	5.1 - 5.2	5.9 - 5.9	6.6 - 6.7	7.3 - 7.4	8.0 - 8.1	
6	3.9	3.8	5.4 - 5.5	6.2 - 6.3	7.0 - 7.1	7.7 - 7.8	8.5 - 8.6	
7	4.1	4.1	5.8 - 5.8	6.6 - 6.7	7.4 - 7.5	8.2 - 8.3	9.0 - 9.2	
8	4.4	4.3	6.1 - 6.2	7.0 - 7.1	7.9 - 7.9	8.7 - 8.8	9.6 - 9.7	
9	4.6	4.6	6.5 - 6.6	7.4 - 7.5	8.3 - 8.4	9.3 - 9.4	10.2 - 10.3	
10	4.7	4.7	6.7 - 6.7	7.6 - 7.7	8.6 - 8.6	9.5 - 9.6	10.5 - 10.5	11.4 - 11.5
11	4.9	5.0	7.0 - 7.1	8.0 - 8.1	9.0 - 9.1	10.0 - 10.1	11.0 - 11.1	12.0 - 12.1
12	5.2	5.3	7.3 - 7.5	8.4 - 8.6	9.4 - 9.6	10.5 - 10.7	11.5 - 11.8	12.6 - 12.8
13	5.4	5.6	7.7 - 7.9	8.8 - 9.0	9.9 - 10.1	11.0 - 11.2	12.1 - 12.4	13.2 - 13.5
14	5.6	5.8	8.0 - 8.2	9.1 - 9.3	10.2 - 10.5	11.4 - 11.7	12.5 - 12.8	13.7 - 14.0
15	5.8	5.9	8.2 - 8.4	9.3 - 9.6	10.5 - 10.8	11.7 - 12.0	12.8 - 13.2	14.0 - 14.4
16	5.9	6.0	8.3 - 8.5	9.5 - 9.7	10.6 - 10.9	11.8 - 12.1	13.0 - 13.3	14.2 - 14.5
17	5.9	6.0	8.3 - 8.6	9.5 - 9.8	10.7 - 11.0	11.9 - 12.2	13.1 - 13.4	14.3 - 14.7

(a): REE, resting energy expenditure computed from Henry and Schofield et al. equations (see Appendix 13) and based on anthropometric data shown in Table 8.

(b): Based on REE predicted with both equations, and taking into account energy expenditure for growth by increasing PAL by 1 %.

The range of Average Requirements (ARs) for energy for children and adolescents based on the factorial method using the equations of Schofield et al. (1985) and Henry (2005) for predicting REE, at different PAL values and expressed in kcal/day, is tabled in Appendix 14B.

### 5.3.4. Pregnancy

The additional energy requirement for pregnancy arises from increases in maternal and feto-placental tissue mass, the rise in energy expenditure attributable to increased REE (see Section 2.3.4.), and changes in physical activity. TEF has been shown to be unchanged (Bronstein et al., 1995; Nagy and King, 1984; Spaaij et al., 1994b) or lower (Schutz et al., 1988) than for non-pregnant women, and therefore is not considered in the determination of the additional AR for energy for pregnancy.

### 5.3.4.1. Energy requirement for the increase in tissue mass during pregnancy

Based on the findings that gestational increases in body mass between 10 and 14 kg were associated with optimal maternal and fetal health outcomes (WHO, 1995a) (see Section 5.1.3.), in this Opinion, assuming a pre-pregnancy BMI within the recommended range, a mean gestational increase in body mass of 12 kg is taken as a basis for the calculation of the energy requirement for the increase in tissue mass.

The corresponding protein and fat gains associated with a mean body mass gain of 12 kg (range 10 to 14 kg) observed in the WHO collaborative study would be 597 g (range 497 to 696 g) and 3.7 kg (range 3.1 to 4.4 kg), respectively (FAO/WHO/UNU, 2004). Based on an energy value of 23.6 kJ/g (5.65 kcal/g) for protein deposited, and 38.7 kJ/g (9.25 kcal/g) for fat deposited, this would result in an energy storage of 14.1 MJ (3,370 kcal) for protein and 144.8 MJ (34,600 kcal) for fat (Table 10).

The accretion of tissue mass is not distributed equally throughout the gestational period. The deposition of protein occurs primarily in the second (20 %) and third trimesters (80 %). Assuming that the rate of fat deposition follows the same pattern as the rate of gestational body mass gain, 11 %, 47 % and 42 % of fat is deposited in the first, second and third trimesters, respectively (IoM, 1990). Accordingly, the daily requirement of energy for protein and fat deposition is estimated as 0 and 202 kJ (0 and 48 kcal), 30 and 732 kJ (7 and 175 kcal), and 121 and 654 kJ (29 and 156 kcal) throughout the first, second and third trimesters, respectively (FAO/WHO/UNU, 2004).

### 5.3.4.2. Calculation of additional AR for energy for tissue deposition in pregnancy

As discussed in Section 2.3.4., on average EEPA is not significantly increased during pregnancy. Thus, apart from the energy stored in newly synthesised tissues, the increase in TEE during pregnancy is mainly due to the increase in REE. The cumulative increment of TEE as estimated with the DLW technique was 161.4 MJ (38,560 kcal). When subtracting from this value the energy estimated for the efficiency of energy utilisation of 15.9 MJ (3,800 kcal), which is included in the measurement of TEE by DLW, the remaining cumulative TEE of 145.5 MJ (34,760 kcal) is nearly equal to the estimated cumulative increase of REE (147.8 MJ (35,330 kcal), see Section 2.3.4.). Table 10 reports on the additional energy expenditure during pregnancy.

**Table 10:** Additional energy expenditure of pregnancy in women with an average gestational increase in body mass of 12 kg <sup>(a)</sup> (adapted from FAO/WHO/UNU (2004))

<b>A. Rates of tissue deposition</b>				
	<b>1<sup>st</sup> trimester</b>	<b>2<sup>nd</sup> trimester</b>	<b>3<sup>rd</sup> trimester</b>	<b>Total deposition</b>
	<b>g/day</b>	<b>g/day</b>	<b>g/day</b>	<b>g/280 days</b>
Body mass gain	17	60	54	12,000
Protein deposition <sup>(b)</sup>	0	1.3	5.1	597
Fat deposition <sup>(b)</sup>	5.2	18.9	16.9	3,741
<b>B. Additional energy expenditure of pregnancy estimated from the increment in TEE and energy deposition</b>				
	<b>1<sup>st</sup> trimester</b>	<b>2<sup>nd</sup> trimester</b>	<b>3<sup>rd</sup> trimester</b>	<b>Energy expenditure</b>
	<b>kJ/day</b>	<b>kJ/day</b>	<b>kJ/day</b>	<b>during whole pregnancy</b>
				<b>MJ</b>
Protein deposition <sup>(b)</sup>	0	30	121	14.1
Fat deposition <sup>(b)</sup>	202	732	654	144.8
Total energy expenditure	85	350	1,300	161.4
Total energy expenditure plus energy content of protein and fat deposited	287	1,112	2,075	320.2

(a): Calculated as suggested by Butte and King (2002). Increase in body mass and tissue deposition in first trimester computed from last menstrual period (i.e. an interval of 79 days). Second and third trimesters computed as 280/3 = 93 days each.

(b): Protein and fat deposition estimated from longitudinal studies of body composition during pregnancy, and an energy value of 23.6 kJ (5.65 kcal)/g protein deposited, and 38.7 kJ (9.25 kcal)/g fat deposited.



### 5.3.5. Lactation

The AR for energy during lactation is estimated from TEE, milk energy output, and energy mobilisation from tissue stores that have been accumulated during pregnancy. Compared with non-pregnant, non-lactating women, there are no significant changes in REE, efficiency in work performance, or TEE (Butte and King, 2002), and in most societies women resume their usual level of physical activity in the first month *post partum* or shortly thereafter (Goldberg et al., 1991; Panter-Brick, 1993; Roberts et al., 1982; Tuazon et al., 1987; van Raaij et al., 1990).

TEE of lactating women can be calculated either by the factorial method as described above for non-pregnant and non-lactating women, or from DLW measurements. TEEs of lactating women have been measured by the DLW method in five studies (Butte et al., 2001; Forsum et al., 1992; Goldberg et al., 1991; Kopp-Hoolihan et al., 1999; Lovelady et al., 1993). Measurements were performed at various stages of lactation (one to six months); however, there are several potential sources of error using the DLW method in lactation studies, which may be attributed to isotope exchange and sequestration that occurs during the *de novo* synthesis of milk fat and lactose, and to increased water flux into milk (Butte et al., 2001). Underestimation of carbon dioxide by 1.0 to 1.3 % may theoretically occur due to the export of exchangeable hydrogen bound to solids in milk (IDECG, 1990). This underestimation may increase to 1.5 to 3.4 % due to <sup>2</sup>H sequestration. Furthermore, the number of subjects in these studies was rather small (9 to 24). Therefore, in this Opinion, the Panel based the estimation of the additional AR for energy during lactation on the factorial method.

Mean milk intakes of infants through six months *post partum* measured by the test-weighing technique were 769 g/day for women exclusively breastfeeding (Butte and King, 2002). Correction of the mean milk intakes for the infant's insensible water loss (assumed to be equal to 5 %) gives a mean milk secretion over the first six months *post partum* of 807 g/day (FAO/WHO/UNU, 2004) for exclusively breastfeeding women.

In well-nourished women it has been estimated that on average the equivalent of 0.72 MJ/day (170 kcal/day) of tissue stores may be utilised to support lactation during the first six months *post partum* (Butte and King, 2002), based on a rate of body mass loss of 0.8 kg per month (Butte and Hopkinson, 1998). This will vary depending on the amount of fat deposited during pregnancy, and on the lactation pattern and duration.

During the second half of infancy and the second year of life, volumes of breast milk intake are highly variable and depend on energy intake from complementary foods (FAO/WHO/UNU, 2004). In one study in which up to 12 infants from the US were still breast-fed during the second half of infancy, breast milk intakes had a range of 486 to 963 mL/day at seven months, 288 to 1,006 mL/day at eight months, 242 to 889 mL/day at nine months, 143 to 896 mL/day at 10 months, 132 to 861 mL/day at 11 months and 73 to 772 mL/day at 12 months (Neville et al., 1988). In another study with 40 children from an industrialised country, mean breast milk intake in the second year of life (12-23 months) was 448±251 g/day (WHO, 1998).

## 6. Key data on which to base dietary reference values (DRVs)

The Panel decided to define only one DRV for energy, namely the AR, and to use the factorial method based on REE x PAL to obtain the average energy requirements for adults, children and adolescents. For infants, TEE was derived by regression equations based on DLW measurements. The additional energy requirements associated with growth during infancy, childhood and adolescence, and with pregnancy, were accounted for by estimates of the energy content of the newly-acrued tissue mass, as well as of the energy for its synthesis. For the additional energy requirement during lactation, milk energy output and energy mobilisation from tissue stores accumulated during pregnancy were taken into account. As explained in Section 5, different equations and/or databases can be used, and this would lead to a range of ARs for various situations (see Tables 6 and 9). However, for ease of use, the Panel decided to propose only one AR for a defined age and sex group with a healthy body mass and

for PAL values selected to approximate qualitatively defined situations, i.e. low active (sedentary), moderately active, active and very active lifestyles.

## **6.1. Adults**

In this Opinion, the AR for energy for adults is based on predicted REE and PAL (see Section 5.3.1).

### **6.1.1. Calculation of resting energy expenditure (REE)**

Although several predictive equations may be appropriate for estimating REE of various populations (as outlined in Sections 2.4. and 5.3.1.), for practical reasons the Panel decided to calculate REE as a function of age, sex, body mass and height by means of only one set of equations, namely those of Henry (2005). These equations were chosen because the underlying database is, at present, the most comprehensive as regards number of subjects, their nationalities and age groups. As described in Section 5.3.1.1., measured heights (obtained in nationally representative surveys of adults in 13 EU countries) and corresponding body masses to yield a BMI of 22 kg/m<sup>2</sup> were used to calculate REE (see Table 12). Because of a lack of anthropometric data from EU countries for age groups beyond 79 years of age, the Panel decided not to calculate AR for adults from 80 years onwards.

### **6.1.2. Selection of physical activity level (PAL) values**

From the range of observed PAL values, the Panel decided to use PAL values of 1.4, 1.6, 1.8 and 2.0 to reflect low active (sedentary), moderately active, active and very active lifestyles, respectively, and proposes to apply these PAL values in the factorial method to determine ARs for energy (Table 12). However, the Panel notes that for population groups which are highly active, PAL values above 2.0 may be more appropriate (see Section 5.3.1.2.).

Available data indicate that it is difficult to generalise on the energy requirements of older adults (see Section 5.1.4.). However, in advanced age with reduced mobility, it can be assumed that PAL values are likely to be lower than in younger adults.

## **6.2. Infants**

Exclusive breastfeeding to the age of about six months with continued breastfeeding as part of a progressively varied diet after six months is nutritionally adequate for most healthy infants born at term (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2009). For infants during the first half year of life (until six months of age), energy requirements are considered to be equal to the energy supply from human milk.

The Panel decided to use the equation for estimation of TEE derived from data of initially breast-fed infants (see Section 5.3.2.1.). Energy requirements during infancy were estimated from TEE measured by the DLW method in healthy, full-term infants initially breast-fed for four months after birth and with adequate body mass, plus the energy needs for growth (Table 11, the equivalent table using energy units in kcal is given in Appendix 15). Median body masses from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006) were used to derive the AR for energy for infants growing along the trajectory of this standard. Estimates of energy deposition were based on measured protein and fat gains (see Section 5.3.2.2.).

**Table 11:** Derivation of the Average Requirement (AR) for energy for infants aged 7-11 months

Age (months)	Body mass <sup>(a)</sup> (kg)	Gain in body mass <sup>(b)</sup> (g/day)	Energy deposition <sup>(c)</sup> (kJ/g)	Energy deposition <sup>(d)</sup> (kJ/day)	TEE <sup>(e)</sup> (kJ/day)	AR <sup>(f)</sup> (kJ/day)	AR (kJ/kg per day)
<b>Boys</b>							
7	8.3	11.9	6.2	73.8	2,585	2,659	320
8	8.6	10.5	6.2	65.3	2,702	2,767	322
9	8.9	9.5	6.2	58.9	2,818	2,877	323
10	9.2	8.6	11.4	98.4	2,935	3,033	330
11	9.4	8.1	11.4	92.3	3,012	3,105	330
<b>Girls</b>							
7	7.6	11.5	7.4	84.9	2,314	2,399	316
8	7.9	10.4	7.4	76.7	2,430	2,507	317
9	8.2	9.1	7.4	67.3	2,547	2,614	319
10	8.5	8.2	9.8	80.0	2,663	2,743	323
11	8.7	7.8	9.8	76.1	2,741	2,817	324

(a): 50<sup>th</sup> percentile of WHO Growth Standards

(b): Calculation from one-month body mass increments from 50<sup>th</sup> percentile of WHO Growth Standards, assuming that one month = 30 days

(c): see Table 7

(d): Body mass gain × energy accrued in normal growth

(e): Total energy expenditure (TEE) (MJ/day) =  $-0.635 + 0.388 \times \text{body mass (kg)}$

(f): AR = TEE + energy deposition.

### 6.3. Children and adolescents

In this Opinion, ARs for energy for children and adolescents are based on predicted REE and PAL adjusted for growth.

#### 6.3.1. Calculation of resting energy expenditure (REE)

Although, in principle, both the equations of Schofield et al. (1985) and Henry (2005) are considered as appropriate for the estimation of REE for children and adolescents, for practical reasons and because the results obtained with these equations are very similar, only the equations of Henry (2005) are applied for the estimation of REE values to calculate ARs for energy for children and adolescents as described in Section 5.3.3.1. The Henry equations were chosen because their database comprises more subjects than the one underlying the Schofield equations (see Table 1). For the calculations, median body masses and heights from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006) were used for children aged 1-2 years, and from harmonised growth curves of children in EU countries (van Buuren et al., 2012) for children from three years onwards.

#### 6.3.2. Selection of physical activity level (PAL) values

From the range of observed PAL values in children and adolescents (see Section 5.3.3.2.), the Panel chose to use the following PAL values: 1.4 for the age group 1-3 years; 1.4, 1.6 and 1.8 for the age group >3-<10 years; and 1.6, 1.8 and 2.0 for the age group 10-18 years (Table 14).

In this Opinion, energy expenditure for growth is accounted for by a 1 % increase in PAL values for each age group.

### 6.4. Pregnancy

The extra amount of energy required during pregnancy is calculated using the cumulative increment in TEE plus the energy deposited as protein and fat (see Section 5.3.4.). Based on these data, the average extra energy requirement for pregnancy is 320 MJ (76,530 kcal) divided into approximately 0.29 MJ/day, 1.1 MJ/day and 2.1 MJ/day (70 kcal/day, 260 kcal/day and 500 kcal/day) during the first, second and third trimesters, respectively (Tables 10 and 15).

A high variability is seen in the rates of gestational increase in body mass and energy expenditure of pregnant women depending on differences in pre-pregnant body mass and composition, lifestyle and underlying nutritional status. The coefficient of variability of the cumulative increase in REE was 16 % between studies, but the variability between women in each study was higher, namely 45 to 70 % in many cases (WHO/FAO/UNU, 2004). This variability should be taken into account when using the AR for additional energy intake during pregnancy as a DRV on an individual basis.

## **6.5. Lactation**

For exclusive breastfeeding during the first six months of life, the mean energy expenditure of lactation over the six month period is 2.8 MJ/day (670 kcal/day) based on a mean milk production of 807 g/day, an energy density of milk of 2.8 kJ/g (0.67 kcal/g), and an energetic efficiency of 80 %. Energy mobilisation from tissues in the order of 0.72 MJ/day (170 kcal/day) (Butte and King, 2002) may contribute to this energy expenditure and reduce the additional energy requirement during lactation to 2.1 MJ/day (500 kcal/day) over pre-pregnancy requirements (Table 15).

During the second half of infancy and the second year of life, volumes of breast milk secreted are highly variable and depend on an infant's energy intake from complementary foods. Thus, the Panel decided not to propose a DRV for additional energy intake for women lactating beyond the first six months after birth.

## CONCLUSIONS

The Panel concludes that one DRV for energy, namely an AR, can be derived for adults, infants and children, and pregnant and lactating women. For infants, this is based on estimates of TEE determined with DLW studies, whereas for children and adults TEE is determined factorially from estimates of REE using the predictive equations of Henry (2005) and the energy needed for various levels of physical activity. For pregnant and lactating women, the additional energy needed for the deposition of newly formed tissue, and for milk output, is derived from data acquired with the DLW method, or from factorial estimates, respectively. Summary tables with the proposed AR expressed as kcal/day can be found in Appendix 16.

The application of these values requires consideration of the objective, such as dietary assessment (for groups or individuals), dietary planning (for groups or individuals specifying also the goal for body mass: maintenance, increase, decrease), labelling dietary reference values, and development of food-based dietary guidelines, and a need to define and characterise the target population (homogeneous, heterogeneous, in relation to age, sex, physical activity, body mass). The detailed information provided in Section 5 should help with adapting the values to specific objectives and populations/individuals.

**Table 12:** Summary of Average Requirement (AR) for energy for adults

Age (years)	REE <sup>(a)</sup> (MJ/day)	AR at PAL=1.4 (MJ/day)	AR at PAL=1.6 (MJ/day)	AR at PAL=1.8 (MJ/day)	AR at PAL=2.0 (MJ/day)
<b>Men</b>					
18 - 29	7.0	9.8	11.2	12.6	14.0
30 - 39	6.8	9.5	10.8	12.2	13.5
40 - 49	6.7	9.3	10.7	12.0	13.4
50 - 59	6.6	9.2	10.5	11.9	13.2
60 - 69	6.0	8.4	9.6	10.9	12.1
70 - 79	5.9	8.3	9.5	10.7	11.9
<b>Women</b>					
18 - 29	5.6	7.9	9.0	10.1	11.2
30 - 39	5.4	7.6	8.7	9.8	10.8
40 - 49	5.4	7.5	8.6	9.7	10.7
50 - 59	5.3	7.5	8.5	9.6	10.7
60 - 69	4.9	6.8	7.8	8.8	9.7
70 - 79	4.8	6.8	7.7	8.7	9.6

(a): REE, resting energy expenditure predicted with the equations of Henry (2005) using body mass and height. Because these have overlapping age bands (18-30 years, 30-60 years,  $\geq 60$  years) (see Appendix 2), the choice of equation is ambiguous at the age boundaries. The REE equations for 18-30 year-olds are used for adults aged 18-29 years, the equations for 30-60 year-olds are used for adults aged 30-39, 40-49 and 50-59 years, and the equations for  $\geq 60$  year-olds are used for adults aged 60-69 and 70-79 years.



**Table 13:** Summary of Average Requirement (AR) for energy for infants

Age	AR (MJ/day)		AR (MJ/kg body mass per day)	
	Boys	Girls	Boys	Girls
7 months	2.7	2.4	0.32	0.32
8 months	2.8	2.5	0.32	0.32
9 months	2.9	2.6	0.32	0.32
10 months	3.0	2.7	0.33	0.32
11 months	3.1	2.8	0.33	0.32

**Table 14:** Summary of Average Requirement (AR) for energy for children and adolescents

Age (years)	REE <sup>(a)</sup> (MJ/day)	AR <sup>(b)</sup> at PAL <sup>(c)</sup> =1.4 (MJ/day)	AR <sup>(b)</sup> at PAL=1.6 (MJ/day)	AR <sup>(b)</sup> at PAL=1.8 (MJ/day)	AR <sup>(b)</sup> at PAL=2.0 (MJ/day)
<b>Boys</b>					
1	2.3	3.3			
2	3.0	4.3			
3	3.5	4.9			
4	3.7	5.3	6.0	6.8	
5	3.9	5.6	6.4	7.2	
6	4.2	5.9	6.7	7.6	
7	4.4	6.3	7.2	8.1	
8	4.7	6.7	7.6	8.6	
9	5.0	7.0	8.1	9.1	
10	5.0		8.1	9.1	10.1
11	5.3		8.5	9.6	10.7
12	5.6		9.1	10.2	11.4
13	6.0		9.8	11.0	12.2
14	6.5		10.5	11.8	13.1
15	7.0		11.3	12.7	14.1
16	7.4		11.9	13.4	14.9
17	7.6		12.3	13.8	15.4
<b>Girls</b>					
1	2.1	3.0			
2	2.8	4.0			
3	3.2	4.6			
4	3.5	4.9	5.6	6.3	
5	3.7	5.2	5.9	6.7	
6	3.9	5.5	6.3	7.1	
7	4.1	5.8	6.7	7.5	
8	4.4	6.2	7.1	7.9	
9	4.6	6.6	7.5	8.4	
10	4.7		7.6	8.6	9.5
11	4.9		8.0	9.0	10.0
12	5.2		8.4	9.4	10.5
13	5.4		8.8	9.9	11.0
14	5.6		9.1	10.2	11.4
15	5.8		9.3	10.5	11.7
16	5.9		9.5	10.6	11.8
17	5.9		9.5	10.7	11.9

(a): REE, resting energy expenditure computed with the predictive equations of Henry (2005) using median heights and body masses from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006) (for children aged 1-2 years) or from harmonised growth curves of children in the EU (van Buuren et al., 2012) (for children aged 3-17 years). Because these have overlapping age bands (0-3, 3-10, 10-18 years), the choice of equation is ambiguous at the age boundaries. The REE equation for 3-10 year-olds is used for the 3 year-olds and the equation for 10-18 year-olds is used for those aged 10 years.

(b): Taking into account a coefficient of 1.01 for growth.

(c): PAL, physical activity level.

**Table 15:** Summary of Average Requirement (AR) for energy for pregnant and lactating women (in addition to the AR for non-pregnant women)

	AR (MJ/day)
<b>Pregnant women</b>	
1 <sup>st</sup> trimester	+ 0.29
2 <sup>nd</sup> trimester	+ 1.1
3 <sup>rd</sup> trimester	+ 2.1
<b>Lactating women</b>	
0-6 months <i>post partum</i>	+ 2.1

## RECOMMENDATIONS FOR RESEARCH

The Panel proposes that the impact of differences in body composition (i.e. FM and FFM) on REE in relation to age, ethnicity and possible other factors should be explored further so that predictive equations for REE could be adjusted to take this into account. In the future, imaging techniques (such as whole body magnetic resonance imaging and echocardiography methods) and specific metabolic rates of major tissues and organs may allow the development of organ/tissue-based predictive equations for REE with a higher accuracy compared to predictive equations for REE based on body mass (index), or on FM and FFM.

The Panel suggests further research particularly on body composition in infants, especially regarding gains in FFM and FM during the first year of life in relation to energy intakes.

Further research is desirable on energy requirements and body composition in relation to pregnancy outcomes for mother and offspring, especially in relation to maternal fat gain and its retention after pregnancy.

Since a precise estimate of the EEPA is essential to determine energy requirements, the Panel stresses the need for the standardisation of the conditions for recording activity expenditure in order to generate reliable and reproducible values for EEPA taking into account sex, age and physiological status (such as pregnancy). Based on such values, future research on the relationship between the amount and intensity of physical activity, maintenance of body mass and long term health is needed in order to advise on PALs for different population groups.

There is a paucity of data regarding REE and TEE of older adults ( $\geq 80$  years). As this age group is an increasing subpopulation in EU countries the Panel suggests that future research addresses this gap. In particular, PALs of older age groups likely to promote maintenance of mobility and reduce the risk for morbidity and mortality need to be identified.

For a more precise estimate of energy requirements at the EU level, the Panel suggests generating and collecting more DLW data, in conjunction with REE measurements, in healthy adults and children in the EU who are representative of various geographical regions, and including individuals of all ages with a broad range of PALs corresponding to well-defined lifestyles.

## REFERENCES

- Abrasheva M, Angelova K, Baikova D, Duleva V, Ivanova L, Jordanov B, Ovcharova D, Petrova S, Vatrlova K and Vukov M, 1998. Dietary and nutritional status survey of the population in Bulgaria. National Center of Hygiene, Medical Ecology and Nutrition, 230 pp.
- AFSSA (Agence Française de Sécurité Sanitaire des Aliments), 2001. Apports nutritionnels conseillés pour la population française [Nutritional recommendations for the French population]. Editions Tec&Doc, Paris, France, 605 pp.
- AFSSA (Agence Française de Sécurité Sanitaire des Aliments), 2009. Étude Individuelle Nationale des Consommations Alimentaires 2 (INCA 2) (2006-2007). Rapport. [Individual and National Study on Food Consumption 2 (INCA 2) 2006-2007. Report]. 228 pp.
- Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, O'Brien WL, Bassett DR, Jr., Schmitz KH, Emplaincourt PO, Jacobs DR, Jr. and Leon AS, 2000. Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise*, 32, S498-504.
- Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DRJ, Tudor-Locke C, Greer JL, Vezina J, Whitt-Glover MC and Leon AS, 2011. The Compendium of Physical Activities Tracking Guide. Healthy Lifestyles Research Center, College of Nursing & Health Innovation, Arizona State University. Available from: <https://sites.google.com/site/compendiumofphysicalactivities/> Accessed November 2012.
- al-Adsani H, Hoffer LJ and Silva JE, 1997. Resting energy expenditure is sensitive to small dose changes in patients on chronic thyroid hormone replacement. *Journal of Clinical Endocrinology and Metabolism*, 82, 1118-1125.
- Alfonzo-Gonzalez G, Doucet E, Almeras N, Bouchard C and Tremblay A, 2004. Estimation of daily energy needs with the FAO/WHO/UNU 1985 procedures in adults: comparison to whole-body indirect calorimetry measurements. *European Journal of Clinical Nutrition*, 58, 1125-1131.
- Alkerwi A, Sauvageot N, Donneau AF, Lair ML, Couffignal S, Beissel J, Delagardelle C, Wagener Y, Albert A and Guillaume M, 2010. First nationwide survey on cardiovascular risk factors in Grand-Duchy of Luxembourg (ORISCAV-LUX). *BMC Public Health*, 10, 468.
- Amcoff E, Edberg A, Enghardt Barbieri H, Lindroos AK, Nälsén C, Pearson M and Warensjö Lemming E, 2012. Riksmaten – vuxna 2010–11. Livsmedels- och näringsintag bland vuxna i Sverige [Riksmaten - adults 2010-11. The food and nutrient intake among adults in Sweden]. Livsmedelsverket, 180 pp.
- Amirkalali B, Hosseini S, Heshmat R and Larijani B, 2008. Comparison of Harris Benedict and Mifflin-ST Jeor equations with indirect calorimetry in evaluating resting energy expenditure. *Indian Journal of Medical Sciences*, 62, 283-290.
- Andersen E, Hutchings B, Jansen J and Nyholm M, 1982. Højde og vægt hos danske børn. [Heights and weights of Danish children] *Ugeskrift for Læger* [The Journal of the Danish Medical Association], 144, 1760-1765.
- Andersen N, Fagt S, Groth M, Hartkopp H, Møller A, Ovesen L and Warming D, 1996. Danskernes kostvaner 1995. Hovedresultater [The Danish diet 1995. Main results]. Levnedsmiddelstyrelsen, Søborg, Denmark, 298 pp.
- Avdičová M, Hrubá F and Námešná J, 2005. Surveillance of Chronic diseases in the Republic. CINDI programme in the SR. Regional Authority of Public Health in Banská Bystrica and WHO - Regional Office for Europe. Banská Bystrica: Dali BB, s.r.o: 153.
- Bates E, Lennox A, Bates C and Swan G, 2011. National Diet and Nutrition Survey. Headline results from Years 1 and 2 (combined) of the Rolling Programme (2008/2009 - 2009/2010). A Survey carried out on the behalf of the Food Standards Agency and the Department of Health, 68 pp.

- Baumgartner RN, Roche AF and Himes JH, 1986. Incremental growth tables: supplementary to previously published charts. *American Journal of Clinical Nutrition*, 43, 711-722.
- Belgardt BF and Bruning JC, 2010. CNS leptin and insulin action in the control of energy homeostasis. *Annals of the New York Academy of Sciences*, 1212, 97-113.
- Benedict F, 1932. The racial element in human metabolism. *American Journal of Physical Anthropology*, 16, 463-473.
- Bingham SA, Goldberg GR, Coward WA, Prentice AM and Cummings JH, 1989. The effect of exercise and improved physical fitness on basal metabolic rate. *British Journal of Nutrition*, 61, 155-173.
- Biro L, Regoly-Merei A, Nagy K, Peter S, Arato G, Szabo C, Martos E and Antal M, 2007. Dietary habits of school children: representative survey in metropolitan elementary schools. Part two. *Annals of Nutrition and Metabolism*, 51, 454-460.
- Bisdee JT, James WP and Shaw MA, 1989. Changes in energy expenditure during the menstrual cycle. *British Journal of Nutrition*, 61, 187-199.
- Bitar A, Vermorel M, Fellmann N, Bedu M, Chamoux A and Coudert J, 1996. Heart rate recording method validated by whole body indirect calorimetry in 10-yr-old children. *Journal of Applied Physiology*, 81, 1169-1173.
- Blaak EE, Westerterp KR, Bar-Or O, Wouters LJ and Saris WH, 1992. Total energy expenditure and spontaneous activity in relation to training in obese boys. *American Journal of Clinical Nutrition*, 55, 777-782.
- Black AE, Coward WA, Cole TJ and Prentice AM, 1996. Human energy expenditure in affluent societies: an analysis of 574 doubly-labelled water measurements. *European Journal of Clinical Nutrition*, 50, 72-92.
- Black AE, 1996. Physical activity levels from a meta-analysis of doubly labeled water studies for validating energy intake as measured by dietary assessment. *Nutrition Reviews*, 54, 170-174.
- Blair SN, Cheng Y and Holder JS, 2001. Is physical activity or physical fitness more important in defining health benefits? *Medicine and Science in Sports and Exercise*, 33, S379-399.
- Bland JM and Altman DG, 1987. Statistical methods for assessing agreement between measurements. *Biochimica clinica*, 11, 399-404.
- Blokstra A, Vissink P, Venmans LMAJ, Holleman P, van der Schouw YT, Smit HA and Verschuren WMM, 2011. Nederland de Maat genomen, 2009-2010. Monitoring van risicofactoren in de algemene bevolking. RIVM report 260152001/2011. National Institute for Public Health and the Environment, 27 pp.
- Boreham C and Riddoch C, 2001. The physical activity, fitness and health of children. *Journal of Sports Sciences*, 19, 915-929.
- Boullata J, Williams J, Cottrell F, Hudson L and Compher C, 2007. Accurate determination of energy needs in hospitalized patients. *Journal of the American Dietetic Association*, 107, 393-401.
- Bray GA, Whipp BJ, Koyal SN and Wasserman K, 1977. Some respiratory and metabolic effects of exercise in moderately obese men. *Metabolism: Clinical and Experimental*, 26, 403-412.
- Bronstein MN, Mak RP and King JC, 1995. The thermic effect of food in normal-weight and overweight pregnant women. *British Journal of Nutrition*, 74, 261-275.
- Brouwer E, 1957. On simple formulae for calculating the heat expenditure and the quantities of carbohydrate and fat oxidized in metabolism of men and animals, from gaseous exchange (Oxygen intake and carbonic acid output) and urine-N. *Acta Physiologica et Pharmacologica Neerlandica*, 6, 795-802.

- Buchholz AC, Rafii M and Pencharz PB, 2001. Is resting metabolic rate different between men and women? *British Journal of Nutrition*, 86, 641-646.
- Butte NF, Wong WW, Ferlic L, Smith EO, Klein PD and Garza C, 1990. Energy expenditure and deposition of breast-fed and formula-fed infants during early infancy. *Pediatric Research*, 28, 631-640.
- Butte NF, 1996. Energy requirements of infants. *European Journal of Clinical Nutrition*, 50 Suppl 1, S24-36.
- Butte NF, Henry CJ and Torun B, 1996. Report on the working group on energy requirements of infants, children and adolescents. *European Journal of Clinical Nutrition*, 50 Suppl 1, S188-189.
- Butte NF and Hopkinson JM, 1998. Body composition changes during lactation are highly variable among women. *Journal of Nutrition*, 128, 381S-385S.
- Butte NF, Hopkinson JM, Mehta N, Moon JK and Smith EO, 1999. Adjustments in energy expenditure and substrate utilization during late pregnancy and lactation. *American Journal of Clinical Nutrition*, 69, 299-307.
- Butte NF, Wong WW, Hopkinson JM, Heinz CJ, Mehta NR and Smith EO, 2000a. Energy requirements derived from total energy expenditure and energy deposition during the first 2 y of life. *American Journal of Clinical Nutrition*, 72, 1558-1569.
- Butte NF, Hopkinson JM, Wong WW, Smith EO and Ellis KJ, 2000b. Body composition during the first 2 years of life: an updated reference. *Pediatric Research*, 47, 578-585.
- Butte NF, 2001. Energy requirements of infants. Background paper prepared for the joint FAO/WHO/UNU Expert consultation on Energy in Human nutrition.
- Butte NF, Wong WW and Hopkinson JM, 2001. Energy requirements of lactating women derived from doubly labeled water and milk energy output. *Journal of Nutrition*, 131, 53-58.
- Butte NF and King JC, 2002. Energy requirements during pregnancy and lactation. Energy background paper prepared for the joint FAO/WHO/UNU Consultation on Energy in Human Nutrition.
- Butte NF, Lopez-Alarcon MG and Garza C, 2002. Nutrient adequacy of exclusive breastfeeding for the term infant during the first six months of life. *World Health Organization*, 57 pp.
- Butte NF, 2005. Energy requirements of infants. *Public Health Nutrition*, 8, 953-967.
- Butte NF and King JC, 2005. Energy requirements during pregnancy and lactation. *Public Health Nutrition*, 8, 1010-1027.
- Butte NF, Garza C and de Onis M, 2007. Evaluation of the feasibility of international growth standards for school-aged children and adolescents. *Journal of Nutrition*, 137, 153-157.
- Cedergren MI, 2007. Optimal gestational weight gain for body mass index categories. *Obstetrics and Gynecology*, 110, 759-764.
- Ceesay SM, Prentice AM, Day KC, Murgatroyd PR, Goldberg GR, Scott W and Spurr GB, 1989. The use of heart rate monitoring in the estimation of energy expenditure: a validation study using indirect whole-body calorimetry. *British Journal of Nutrition*, 61, 175-186.
- Cifková R and Škodová Z, 2004. Dlouhodobé trendy hlavních rizikových faktorů kardiovaskulárních onemocnění v české populaci [Longitudinal trends in major cardiovascular risk factors in the Czech population]. *Casopis Lekarů Ceskyh [Czech Medical Journal]*, 143, 219-226.
- Cikrikci E, Gokbel H and Bediz CS, 1999. Basal metabolic rates of Turkish women during pregnancy. *Annals of Nutrition and Metabolism*, 43, 80-85.
- Cole T, 2002. The Oxford Brookes BMR database - a reanalysis. Report commissioned by FAO for the joint FAO/WHO/UNU Expert Consultation on Energy in Human Nutrition.



- Compher C, Frankenfield D, Keim N, Roth-Yousey L and Evidence Analysis Working G, 2006. Best practice methods to apply to measurement of resting metabolic rate in adults: a systematic review. *Journal of the American Dietetic Association*, 106, 881-903.
- Coward W and Cole T, 1991. The doubly labeled water method for the measurement of energy expenditure in humans: risks and benefits. In: *New Techniques in Nutritional Research*. Eds Whitehead R, Prentice A. Academic, San Diego/London, USA/UK, 139–176.
- Cynober L, Alix E, Arnaud-Battandier F, Bonnefoy M, Brocker P, Cals MJ, Cherbut C, Coplo C, Ferry M, Ghisolfi-Marque A, Kravtchenko T, Lesourd B, Mignot C, P. PM and 2000. Apports nutritionnels conseillés chez la personne âgée [Nutritional intake in the elderly]. *Nutrition clinique et métabolisme*, 14, 3s-60s.
- D-A-CH (Deutsche Gesellschaft für Ernährung - Österreichische Gesellschaft für Ernährung - Schweizerische Gesellschaft für Ernährungsforschung - Schweizerische Vereinigung für Ernährung), 2012. Referenzwerte für die Nährstoffzufuhr [Reference values for nutrient intakes]. Umschau Braus Verlag, Frankfurt am Main, Germany, 251 pp.
- Daly JM, Heymsfield SB, Head CA, Harvey LP, Nixon DW, Katzeff H and Grossman GD, 1985. Human energy requirements: overestimation by widely used prediction equation. *American Journal of Clinical Nutrition*, 42, 1170-1174.
- Danforth E, Jr. and Burger A, 1984. The role of thyroid hormones in the control of energy expenditure. *Clinics in Endocrinology and Metabolism*, 13, 581-595.
- Davies PSW, Ewing G, Coward WA and Lucas A, 1990. Energy metabolism in breast-fed and formula-fed infants. In: *Breast-feeding, nutrition, infection and infant growth in developed and emerging countries*. Eds Atkinson SA, Hanson LÅ, Chandra RK. Arts Biomedical, St. John's, Newfoundland, Canada, 521.
- de Boer E, Hulshof K and ter Doest D, 2006. Voedselconsumptie van jonge peuters [Food consumption of young children]. TNO report V6269, 37 pp.
- de Bruin NC, Degenhart HJ, Gal S, Westerterp KR, Stijnen T and Visser HK, 1998. Energy utilization and growth in breast-fed and formula-fed infants measured prospectively during the first year of life. *American Journal of Clinical Nutrition*, 67, 885-896.
- de Groot LC, Boekholt HA, Spaaij CK, van Raaij JM, Drijvers JJ, van der Heijden LJ, van der Heide D and Hautvast JG, 1994. Energy balances of healthy Dutch women before and during pregnancy: limited scope for metabolic adaptations in pregnancy. *American Journal of Clinical Nutrition*, 59, 827-832.
- De Lorenzo A, Tagliabue A, Andreoli A, Testolin G, Comelli M and Deurenberg P, 2001. Measured and predicted resting metabolic rate in Italian males and females, aged 18-59 y. *European Journal of Clinical Nutrition*, 55, 208-214.
- de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C and Siekmann J, 2007. Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization*, 85, 660-667.
- de Onis M and Lobstein T, 2010. Defining obesity risk status in the general childhood population: which cut-offs should we use? *International Journal of Pediatric Obesity*, 5, 458-460.
- De Vriese S, Huybrechts I, Moreau M and van Oyen H, 2006. De Belgische Voedselconsumptiepeiling 1 – 2004 [The Belgian food consumption survey 1 - 2004]. Afdeling Epidemiologie, Wetenschappelijk Instituut Volksgezondheid, Depotnummer : D/2006/2505/17, IPH/EPI REPORTS N° 2006 - 016. Available from: <https://www.wiv-isp.be/epidemiepinl/foodnl/table04.htm>, accessed November 2012.
- Deharveng G, Charrondiere UR, Slimani N, Southgate DA and Riboli E, 1999. Comparison of nutrients in the food composition tables available in the nine European countries participating in

- EPIC. European Prospective Investigation into Cancer and Nutrition. *European Journal of Clinical Nutrition*, 53, 60-79.
- Department of Health and Social Security, 1977. The composition of mature human milk. Report on Health and Social Subjects no. 12. HMSO, London, UK, 58 pp.
- DeVader SR, Neeley HL, Myles TD and Leet TL, 2007. Evaluation of gestational weight gain guidelines for women with normal prepregnancy body mass index. *Obstetrics and Gynecology*, 110, 745-751.
- do Carmo I, Dos Santos O, Camolas J, Vieira J, Carreira M, Medina L, Reis L, Myatt J and Galvao-Teles A, 2008. Overweight and obesity in Portugal: national prevalence in 2003-2005. *Obesity Reviews*, 9, 11-19.
- DoH (Department of Health), 1991. Dietary reference values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. HMSO, London, UK, 212 pp.
- Durnin JV, McKillop FM, Grant S and Fitzgerald G, 1987. Energy requirements of pregnancy in Scotland. *Lancet*, 2, 897-900.
- Duro D, Rising R, Cole C, Valois S, Cedillo M and Lifshitz F, 2002. New equations for calculating the components of energy expenditure in infants. *Journal of Pediatrics*, 140, 534-539.
- EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2009. Scientific Opinion on the appropriate age for introduction of complementary feeding of infants. *EFSA Journal*, 7(12):1423, 19 pp.
- EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010. Scientific Opinion on principles for deriving and applying Dietary Reference Values. *EFSA Journal*, 8(3):1458, 30 pp.
- EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2012. Scientific Opinion on Dietary Reference Values for protein. *EFSA Journal*, 10(2):2557, 66 pp.
- Elia M, 1991. Energy equivalents of CO<sub>2</sub> and their importance in assessing energy expenditure when using tracer techniques. *American Journal of Physiology*, 260, E75-88.
- Elia M, 1992. Organ and tissue contributions to metabolic rate. In: *Energy Metabolism: Tissue Determinants and Cellular Corollaries*. Eds Kinney JM, Tucker HN. Raven, New York, USA, 61-80.
- Elia M and Cummings JH, 2007. Physiological aspects of energy metabolism and gastrointestinal effects of carbohydrates. *European Journal of Clinical Nutrition*, 61 Suppl 1, S40-74.
- Elmadfa I, Meyer A, Nowak V, Hasenegger V, Putz P, Verstraeten R, Remaut-DeWinter AM, Kolsteren P, Dostalova J, Dlouhy P, Trolle E, Fagt S, Biloft-Jensen A, Mathiessen J, Velsing Groth M, Kambek L, Gluskova N, Voutilainen N, Erkkila A, Vernay M, Krems C, Strassburg A, Vasquez-Caicedo AL, Urban C, Naska A, Efstathopoulou E, Oikonomou E, Tsiotas K, Bountziouka V, Benetou V, Trichopoulou A, Zajkas G, Kovacs V, Martos E, Heavey P, Kelleher C, Kennedy J, Turrini A, Selga G, Sauka M, Petkeviciene J, Klumbiene J, Holm Totland T, Andersen LF, Halicka E, Rejman K, Kowrygo B, Rodrigues S, Pinhao S, Ferreira LS, Lopes C, Ramos E, Vaz Almeida MD, Vlad M, Simcic M, Podgrajsek K, Serra Majem L, Roman Vinas B, Ngo J, Ribas Barba L, Becker V, Fransen H, Van Rossum C, Ocke M and Margetts B, 2009a. *European Nutrition and Health Report 2009*. Forum of Nutrition, 62, 1-412.
- Elmadfa I, Freisling H, Nowak V, Hofstädter D, Hasenegger V, Ferge M, Fröhler M, Fritz K, Meyer AL, Putz P, Rust P, Grossgut R, Mischek D, Kiefer I, Schätzer M, Spanblöchel J, Sturtzel B, Wagner K-H, Zilberszac A, Vojir F and Plsek K, 2009b. *Österreichischer Ernährungsbericht 2008 [Austrian Nutrition Report 2008]*. Institut für Ernährungswissenschaften der Universität Wien, Bundesministerium für Gesundheit, 454 pp.

- Enghardt-Barbieri H, Pearson M and Becker W, 2006. Riksmaten – Barn 2003. Livsmedels – och näringsintag bland barn i Sverige [Riksmaten - Children 2003. The food and nutritional intake among children in Sweden]. Livsmedelsverket, 216 pp.
- FAO (Food and Agriculture Organization of the United Nations), 1971. The adoption of Joules as unit of energy. Working paper for the Joint FAO/WHO/UNU Expert Consultation on Energy and Protein Requirements. Available at: <http://www.fao.org/docrep/meeting/009/ae906e/ae906e17.htm>. Accessed November 2012.
- FAO (Food and Agriculture Organization of the United Nations), 2003. Food energy – methods of analysis and conversion factors. Report of a technical workshop, Rome, 3–6 December 2002. FAO Food and Nutrition Paper 77, 93 pp.
- FAO/WHO/UNU (Food and Agriculture Organization of the United Nations/World Health Organization/United Nations University), 1985. Energy and protein requirements. Report of a Joint WHO/FAO/UNU Expert Consultation, Rome, 5-17 October 1981. World Health Organization Technical Report Series 724, 206 pp.
- FAO/WHO/UNU (Food and Agriculture Organization of the United Nations/World Health Organization/United Nations University), 2004. Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation: Rome, 17–24 October 2001. FAO food and nutrition technical report series, 103 pp.
- Ferro-Luzzi A, 1987. The application of energy costs to activities and overall energy expenditure over 24 hours (including BMR levels) of the elderly. Report to FAO.
- Fields DA, Gilchrist JM, Catalano PM, Gianni ML, Roggero PM and Mosca F, 2011. Longitudinal body composition data in exclusively breast-fed infants: a multicenter study. *Obesity* (Silver Spring), 19, 1887-1891.
- Firouzbakhsh S, Mathis RK, Dorchester WL, Oseas RS, Groncy PK, Grant KE and Finklestein JZ, 1993. Measured resting energy expenditure in children. *Journal of Pediatric Gastroenterology and Nutrition*, 16, 136-142.
- Fomon SJ, Haschke F, Ziegler EE and Nelson SE, 1982. Body composition of reference children from birth to age 10 years. *American Journal of Clinical Nutrition*, 35, 1169-1175.
- Forbes GB, 2000. Body fat content influences the body composition response to nutrition and exercise. *Annals of the New York Academy of Sciences*, 904, 359-365.
- Forsum E, Sadurskis A and Wager J, 1988. Resting metabolic rate and body composition of healthy Swedish women during pregnancy. *American Journal of Clinical Nutrition*, 47, 942-947.
- Forsum E, Kabir N, Sadurskis A and Westerterp K, 1992. Total energy expenditure of healthy Swedish women during pregnancy and lactation. *American Journal of Clinical Nutrition*, 56, 334-342.
- Forsum E and Löf M, 2007. Energy metabolism during human pregnancy. *Annual Review of Nutrition*, 27, 277-292.
- Frankenfield D, Roth-Yousey L and Compher C, 2005. Comparison of predictive equations for resting metabolic rate in healthy nonobese and obese adults: a systematic review. *Journal of the American Dietetic Association*, 105, 775-789.
- Fraser A, Tilling K, Macdonald-Wallis C, Hughes R, Sattar N, Nelson SM and Lawlor DA, 2011. Associations of gestational weight gain with maternal body mass index, waist circumference, and blood pressure measured 16 y after pregnancy: the Avon Longitudinal Study of Parents and Children (ALSPAC). *American Journal of Clinical Nutrition*, 93, 1285-1292.
- Fredriks AM, van Buuren S, Burgmeijer RJF, Verloove-Vanhorick SP and Wit JM, 1998. Nederlandse groeidiagrammen 1997 in historisch perspectief [Dutch growth diagrams 1997 in historical perspective]. In: *De vierde Landelijke Groeistudie (1997). Presentatie nieuwe groeidiagrammen*.

- [The 4th nationwide growth study (1997). Presentation of new growth charts]. Bureau Boerhaave Commissie. Ed Wit JM. Rijksuniversiteit Leiden, Leiden, The Netherlands, 1-14.
- Fredriks AM, van Buuren S, Burgmeijer RJ, Meulmeester JF, Beuker RJ, Brugman E, Roede MJ, Verloove-Vanhorick SP and Wit JM, 2000a. Continuing positive secular growth change in The Netherlands 1955-1997. *Pediatric Research*, 47, 316-323.
- Fredriks AM, van Buuren S, Wit JM and Verloove-Vanhorick SP, 2000b. Body index measurements in 1996-7 compared with 1980. *Archives of Disease in Childhood*, 82, 107-112.
- Freeman JV, Cole TJ, Chinn S, Jones PR, White EM and Preece MA, 1995. Cross sectional stature and weight reference curves for the UK, 1990. *Archives of Disease in Childhood*, 73, 17-24.
- Frigerio C, Schutz Y, Whitehead R and Jequier E, 1991. A new procedure to assess the energy requirements of lactation in Gambian women. *American Journal of Clinical Nutrition*, 54, 526-533.
- Gabrijelčič Blenkuš M, Gregorič M, Tivadar B, Koch V, Kostanjevec S, Turk Fajdiga V, Žalar A, Lavtar D, Kuhar D and Rozman U, 2009. Prehrambene navade odraslih prebivalcev Slovenije z vidika varovanja zdravja [Food habits of the adult population of Slovenia in terms of health]. Inštitut za varovanje zdravja Republike Slovenije, Pedagoška faculteta, Univerza v Ljubljani, 186 pp.
- Gallagher D, Visser M, Wang Z, Harris T, Pierson RN, Jr. and Heymsfield SB, 1996. Metabolically active component of fat-free body mass: influences of age, adiposity, and gender. *Metabolism: Clinical and Experimental*, 45, 992-997.
- Gallagher D, Ruts E, Visser M, Heshka S, Baumgartner RN, Wang J, Pierson RN, Pi-Sunyer FX and Heymsfield SB, 2000. Weight stability masks sarcopenia in elderly men and women. *American Journal of Physiology - Endocrinology and Metabolism*, 279, E366-375.
- Gallagher D, Albu J, He Q, Heshka S, Boxt L, Krasnow N and Elia M, 2006. Small organs with a high metabolic rate explain lower resting energy expenditure in African American than in white adults. *American Journal of Clinical Nutrition*, 83, 1062-1067.
- Gannon B, DiPietro L and Poehlman ET, 2000. Do African Americans have lower energy expenditure than Caucasians? *International Journal of Obesity and Related Metabolic Disorders*, 24, 4-13.
- Garza C and Butte NF, 1986. Energy concentration of human milk estimated from 24-h pools and various abbreviated sampling schemes. *Journal of Pediatric Gastroenterology and Nutrition*, 5, 943-948.
- Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Sawyer MB, Ashford J and Black AE, 1991. Longitudinal assessment of the components of energy balance in well-nourished lactating women. *American Journal of Clinical Nutrition*, 54, 788-798.
- Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Wensing C, Black AE, Harding M and Sawyer M, 1993. Longitudinal assessment of energy expenditure in pregnancy by the doubly labeled water method. *American Journal of Clinical Nutrition*, 57, 494-505.
- Goran MI, Poehlman ET and Danforth E, Jr., 1994a. Experimental reliability of the doubly labeled water technique. *American Journal of Physiology*, 266, E510-515.
- Goran MI, Calles-Escandon J, Poehlman ET, O'Connell M and Danforth E, Jr., 1994b. Effects of increased energy intake and/or physical activity on energy expenditure in young healthy men. *Journal of Applied Physiology*, 77, 366-372.
- Guo SM, Roche AF, Fomon SJ, Nelson SE, Chumlea WC, Rogers RR, Baumgartner RN, Ziegler EE and Siervogel RM, 1991. Reference data on gains in weight and length during the first two years of life. *Journal of Pediatrics*, 119, 355-362.
- Halliday D, Hesp R, Stalley SF, Warwick P, Altman DG and Garrow JS, 1979. Resting metabolic rate, weight, surface area and body composition in obese women. *International Journal of Obesity*, 3, 1-6.

- Hamill PV, Drizd TA, Johnson CL, Reed RB and Roche AF, 1977. NCHS growth curves for children birth-18 years. United States. Vital and Health Statistics. Series 11: Data from the National Health Survey, i-iv, 1-74.
- Harper ME, Green K and Brand MD, 2008. The efficiency of cellular energy transduction and its implications for obesity. *Annual Review of Nutrition*, 28, 13-33.
- Harrington KE, Robson PJ, Kiely M, Livingstone MB, Lambe J and Gibney MJ, 2001. The North/South Ireland Food Consumption Survey: survey design and methodology. *Public Health Nutrition*, 4, 1037-1042.
- Harris JA and Benedict FG, 1919. A biometric study of basal metabolism in man. Carnegie Institution of Washington; Publication No. 279, Washington, D. C., USA, 266 pp.
- Haschke F, 1989. Body composition during adolescence. Body composition measurements in Infants and Children: Report of the 98th Ross conference on pediatric research. Ross Laboratories, 76-83.
- Hasson RE, Howe CA, Jones BL and Freedson PS, 2011. Accuracy of four resting metabolic rate prediction equations: effects of sex, body mass index, age, and race/ethnicity. *Journal of Science and Medicine in Sport*, 14, 344-351.
- Health Council of the Netherlands, 2001. Dietary Reference Intakes: energy, proteins, fats and digestible carbohydrates. Publication no. 2001/19, 168 pp.
- Henderson L, Gregory J and Swan G, 2002. National Diet and Nutrition Survey: adults aged 19 to 64 years, Volume 1: Types and quantities of foods consumed. The Stationery Office (TSO), 88 pp.
- Henry C, 2001. Basal metabolic rate studies in humans: measurement and application. Background document prepared for the joint FAO/WHO/UNU Expert Consultation on Energy in Human Nutrition, 2001.
- Henry CJ and Rees DG, 1991. New predictive equations for the estimation of basal metabolic rate in tropical peoples. *European Journal of Clinical Nutrition*, 45, 177-185.
- Henry CJ, 2005. Basal metabolic rate studies in humans: measurement and development of new equations. *Public Health Nutrition*, 8, 1133-1152.
- Heppner KM, Habegger KM, Day J, Pfluger PT, Perez-Tilve D, Ward B, Gelfanov V, Woods SC, DiMarchi R and Tschop M, 2010. Glucagon regulation of energy metabolism. *Physiology and Behavior*, 100, 545-548.
- Herring SJ, Oken E, Haines J, Rich-Edwards JW, Rifas-Shiman SL, Kleinman Sc DK and Gillman MW, 2008. Misperceived pre-pregnancy body weight status predicts excessive gestational weight gain: findings from a US cohort study. *BMC Pregnancy Childbirth*, 8, 54.
- Hervey GR and Tobin G, 1983. Luxusconsumption, diet-induced thermogenesis and brown fat: a critical review. *Clinical Science*, 64, 7-18.
- Hester SN, Hustead DS, Mackey AD, Singhal A and Marriage BJ, 2012. Is the macronutrient intake of formula-fed infants greater than breast-fed infants in early infancy? *Journal of Nutrition and Metabolism*, 2012, 891201.
- Hilbig A and Kersting M, 2006. Effects of age and time on energy and macronutrient intake in German infants and young children: results of the DONALD study. *Journal of Pediatric Gastroenterology and Nutrition*, 43, 518-524.
- Hofman A, Boerlage PA and Bots ML, 1995. Prevalentie van chronische ziekten bij ouderen; het ERGO-onderzoek. [Prevalence of chronic diseases in the elderly, the Rotterdam study]. *Nederlands Tijdschrift voor Geneeskunde [Dutch Journal of Medicine]*, 139, 1975-1978.
- Hofsteenge GH, Chinapaw MJ, Delemarre-van de Waal HA and Weijs PJ, 2010. Validation of predictive equations for resting energy expenditure in obese adolescents. *American Journal of Clinical Nutrition*, 91, 1244-1254.



- Holm Totland T, Kjerpeseth Melnæs B, Lundberg-Hallén N, Helland-Kigen KM, Lund-Blix NA, Borch Myhre J, Wetting Johansen AM, Bjørge Løken E and Frost Andersen L, 2012. Norkost 3. En landsomfattende kostholdsundersøkelse blant menn og kvinner i Norge i alderen 18-70 år, 2010-11 [Norkost 3 A nationwide nutrition survey among men and women in Norway aged 18-70 years, 2010-11]. Universitetet i Oslo, Mattilsynet og Helsedirektoratet, 70 pp.
- Hoyt RW and Friedl KE, 2006. Field studies of exercise and food deprivation. *Current Opinion in Clinical Nutrition and Metabolic Care*, 9, 685-690.
- Hunter GR, Weinsier RL, Darnell BE, Zuckerman PA and Goran MI, 2000. Racial differences in energy expenditure and aerobic fitness in premenopausal women. *American Journal of Clinical Nutrition*, 71, 500-506.
- Huybrechts I and De Henauw S, 2007. Energy and nutrient intakes by pre-school children in Flanders-Belgium. *British Journal of Nutrition*, 98, 600-610.
- Hytten FE and Chamberlain G, 1980. *Clinical Physiology in Obstetrics*. Blackwell Scientific Publications, Oxford, UK, 518 pp.
- IDECG (International Dietary Energy Consultancy Group), 1990. The doubly-labelled water method for measuring energy expenditure: technical recommendations for use in humans. A consensus report. NAHRES-4, International Atomic Energy Agency, 301 pp.
- Illingworth PJ, Jung RT, Howie PW, Leslie P and Isles TE, 1986. Diminution in energy expenditure during lactation. *British Medical Journal (Clinical Research Edition)*, 292, 437-441.
- IoM (Institute of Medicine), 1990. *Nutrition During Pregnancy: Part I: Weight Gain, Part II: Nutrient Supplements*. National Academy Press, Washington, D. C., USA, 468 pp.
- IoM (Institute of Medicine), 1991. *Nutrition during lactation*. National Academy Press, Washington, D. C., USA, 310 pp.
- IoM (Institute of Medicine), 1992. *Nutrition Issues in Developing Countries*. National Academy Press, Washington, D. C., USA, 189 pp.
- IoM (Institute of Medicine), 2005. *Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids (Macronutrients)*. National Academies Press, Washington D. C., USA, 1357 pp.
- IoM/NRC (Institute of Medicine/National Research Council), 2009. *Weight Gain During Pregnancy: Reexamining the Guidelines*. The National Academies Press, Washington D. C., USA, 854 pp.
- IUNA (Irish Universities Nutrition Alliance), 2011. *National Adult Nutrition Survey*.
- IUNA (Irish Universities Nutrition Alliance), a. *National Pre-school Nutrition Survey*. Available from: <http://www.iuna.net/>. Accessed on November 2012.
- IUNA (Irish Universities Nutrition Alliance), b. *National Children's Food Survey 2003-2004*. Available from: <http://www.iuna.net/>. Accessed on November 2012.
- IUNA (Irish Universities Nutrition Alliance), c. *The National Teens' Food Survey 2005-2006*. Available from: <http://www.iuna.net/>. Accessed on November 2012.
- James WTP, Ralph A and Ferro-Luzzi A, 1989. Energy need of the elderly. A new approach. Vol. 6: nutrition, ageing and the elderly. In: *Human nutrition: a comprehensive treatise*. Eds Munro HN, Danford DE. Plenum press, New York, USA, 129-151.
- James WTP and Schofield C, 1990. *Human energy requirements. A manual for planners and nutritionists*. Oxford Medical Publications under arrangement with FAO, Oxford, UK, 172 pp.
- Jiang Z, Yan Q, Su Y, Acheson KJ, Thelin A, Piguet-Welsch C, Ritz P and Ho ZC, 1998. Energy expenditure of Chinese infants in Guangdong Province, south China, determined with use of the doubly labeled water method. *American Journal of Clinical Nutrition*, 67, 1256-1264.

- Joffe R, Ozolinš G, Šantare D, Bartkevics V, L. M and Briška I, 2009. Latvijas iedzīvotāju visaptverošais partikas paterina petijums, 2007-2009 [The National Food Consumption Survey of Latvia, 2007 – 2009]. Nacionalais diagnostikas centrs, Partikas un veterināra dienesta Partikas centrs, 115 pp.
- Johansson L and Sovoll K, 1999. Norkost, 1997. Landsomfattende kostholdsundersøkelse blant menn og kvinner i alderen 16-79 år [Norkost, 1997. Nationwide dietary survey among men and women aged 16-79 years]. Rapport nr.2/1999. Statens råd för ernæring og fysisk aktivitet, 84 pp.
- Khalaj-Hedayati K, Bosity-Westphal A, Muller MJ and Dittmar M, 2009. Validation of the BIOPAC indirect calorimeter for determining resting energy expenditure in healthy free-living older people. *Nutrition research*, 29, 531-541.
- Kiel DW, Dodson EA, Artal R, Boehmer TK and Leet TL, 2007. Gestational weight gain and pregnancy outcomes in obese women: how much is enough? *Obstetrics and Gynecology*, 110, 752-758.
- Kiely M, Flynn A, Harrington KE, Robson PJ and Cran G, 2001. Sampling description and procedures used to conduct the North/South Ireland Food Consumption Survey. *Public Health Nutrition*, 4, 1029-1035.
- Klausen B, Toubro S and Astrup A, 1997. Age and sex effects on energy expenditure. *American Journal of Clinical Nutrition*, 65, 895-907.
- Knudtzon J, Waaler PE, Skjærven R, Solberg LK and Steen J, 1988. Nye norske percentil kurver for høyde, vekt og hodeomkrets for alderen 0 17 år [New Norwegian percentile curves for height, weight and head circumference of aged 0 17 years]. *Tidsskrift for den Norske lægeforening [Journal of the Norwegian Medical Association]*, 26, 2125-2135.
- Kobe H, Krzysnik C and Mis NF, 2011. Under- and Over-Reporting of Energy Intake in Slovenian Adolescents. *Journal of Nutrition Education and Behavior*, 1-10.
- Kopp-Hoolihan LE, van Loan MD, Wong WW and King JC, 1999. Longitudinal assessment of energy balance in well-nourished, pregnant women. *American Journal of Clinical Nutrition*, 69, 697-704.
- Kozak LP, 2010. Brown fat and the myth of diet-induced thermogenesis. *Cell Metabolism*, 11, 263-267.
- Kratěnová J, Lustigová M, Malý M and Žejglicová K, 2007. Health status and selected parameters of demographic and health statistics. Summary Report – 2006. In: *Environmental Health Monitoring System in the Czech Republic*. Ed Puklová V. National Institute of Public Health, Prague, Prague, Czech Republic, 61-75.
- Krems C, Luhrmann PM, Strassburg A, Hartmann B and Neuhauser-Berthold M, 2005. Lower resting metabolic rate in the elderly may not be entirely due to changes in body composition. *European Journal of Clinical Nutrition*, 59, 255-262.
- Kristiansen AL and Andersen LF, 2009. Småbarnskost – 2 år. Landsomfattende kostholdsundersøkelse blant 2 år gamle barn - Småbarnskost 2007 [Nationwide diet survey among 2-year-old children]. IS-1731, Helsedirektoratet, Mattilsynet og Universitetet i Oslo, 94 pp.
- Kuczumarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Roche AF and Johnson CL, 2000. CDC growth charts: United States. *Advance Data*, 1-27.
- Kyttälä P, Ovaskainen M, Kronberg-Kippilä C, Erkkola M, Tapanainen H, Tuokkola J, Veijola R, Simell O, Knip M and Virtanen SM, 2008. The Diet of Finnish Preschoolers. B32/2008, National Public Health Institute, 158 pp.
- Kyttälä P, Erkkola M, Kronberg-Kippilä C, Tapanainen H, Veijola R, Simell O, Knip M and Virtanen SM, 2010. Food consumption and nutrient intake in Finnish 1-6-year-old children. *Public Health Nutrition*, 13, 947-956.

- Levine JA, Schleusner SJ and Jensen MD, 2000. Energy expenditure of nonexercise activity. *American Journal of Clinical Nutrition*, 72, 1451-1454.
- Levine JA, 2004a. Non-exercise activity thermogenesis (NEAT). *Nutrition Reviews*, 62, S82-97.
- Levine JA, 2004b. Nonexercise activity thermogenesis (NEAT): environment and biology. *American journal of physiology. Endocrinology and metabolism*, 286, E675-685.
- Levine JA and Kotz CM, 2005. NEAT--non-exercise activity thermogenesis--egocentric & geocentric environmental factors vs. biological regulation. *Acta Physiologica Scandinavica*, 184, 309-318.
- Livesey G, Smith T, Eggum BO, Tetens IH, Nyman M, Roberfroid M, Delzenne N, Schweizer TF and Decombaz J, 1995. Determination of digestible energy values and fermentabilities of dietary fibre supplements: a European interlaboratory study in vivo. *British Journal of Nutrition*, 74, 289-302.
- Lopes C, Oliveira A, Santos AC, Ramos E, Gaio AR, Severo M and Barros H, 2006. Consumo alimentar no Porto. Faculdade de Medecina da Universidade do Porto, 172 pp.
- Lovelady CA, Meredith CN, McCrory MA, Nommsen LA, Joseph LJ and Dewey KG, 1993. Energy expenditure in lactating women: a comparison of doubly labeled water and heart-rate-monitoring methods. *American Journal of Clinical Nutrition*, 57, 512-518.
- Lucas A, Ewing G, Roberts SB and Coward WA, 1987. How much energy does the breast fed infant consume and expend? *British Medical Journal (Clinical Research Edition)*, 295, 75-77.
- Lührmann PM, Herbert BM and Neuhäuser-Berthold M, 2001. Effects of fat mass and body fat distribution on resting metabolic rate in the elderly. *Metabolism: Clinical and Experimental*, 50, 972-975.
- Lührmann PM, Bender R, Edelmann-Schafer B and Neuhäuser-Berthold M, 2009. Longitudinal changes in energy expenditure in an elderly German population: a 12-year follow-up. *European Journal of Clinical Nutrition*, 63, 986-992.
- Lührmann PM, Edelmann-Schafer B and Neuhäuser-Berthold M, 2010. Changes in resting metabolic rate in an elderly German population: cross-sectional and longitudinal data. *Journal of Nutrition, Health and Aging*, 14, 232-236.
- Lusk G, 1928. *The Element of the Science of Nutrition*. W. B. Saunders Company, London, UK, 844 pp.
- Maffeis C, Schutz Y, Micciolo R, Zocante L and Pinelli L, 1993. Resting metabolic rate in six- to ten-year-old obese and nonobese children. *Journal of Pediatrics*, 122, 556-562.
- Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, Tylavsky F, Bauer DC, Goodpaster BH and Harris TB, 2006. Daily activity energy expenditure and mortality among older adults. *JAMA*, 296, 171-179.
- Manios Y, 2006. Design and descriptive results of the "Growth, Exercise and Nutrition Epidemiological Study In preSchoolers": the GENESIS study. *BMC Public Health*, 6, 32.
- Manios Y, Grammatikaki E, Papoutsou S, Liarigkovinos T, Kondaki K and Moschonis G, 2008. Nutrient intakes of toddlers and preschoolers in Greece: the GENESIS study. *Journal of the American Dietetic Association*, 108, 357-361.
- Matthys C, De Henauw S, Devos C and De Backer G, 2003. Estimated energy intake, macronutrient intake and meal pattern of Flemish adolescents. *European Journal of Clinical Nutrition*, 57, 366-375.
- Melzer K, Laurie Karsegard V, Genton L, Kossovsky MP, Kayser B and Pichard C, 2007. Comparison of equations for estimating resting metabolic rate in healthy subjects over 70 years of age. *Clinical Nutrition*, 26, 498-505.

- Mensink GBM, Heseker H, Richter A, Stahl A and Vohmann C, 2007. Forschungsbericht: Ernährungsstudie als KiGGS-Modul (EsKiMo) [Research report: nutritional study as KiGGS module]. 143 pp.
- Merrill AL and Watt BK, 1973. Energy value of foods – basis and derivation. Handbook no. 74. United States Department of Agriculture, Washington D. C., USA, 105 pp.
- Merten C, Ferrari P, Bakker M, Boss A, Hearty A, Leclercq C, Lindtner O, Tlustos C, Verger P, Volatier JL and Arcella D, 2011. Methodological characteristics of the national dietary surveys carried out in the European Union as included in the European Food Safety Authority (EFSA) Comprehensive European Food Consumption Database. Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment, 28, 975-995.
- Michaelsen KF, Larsen PS, Thomsen BL and Samuelson G, 1994. The Copenhagen Cohort Study on Infant Nutrition and Growth: breast-milk intake, human milk macronutrient content, and influencing factors. American Journal of Clinical Nutrition, 59, 600-611.
- Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA and Koh YO, 1990. A new predictive equation for resting energy expenditure in healthy individuals. American Journal of Clinical Nutrition, 51, 241-247.
- Molnar D, Jeges S, Erhardt E and Schutz Y, 1995. Measured and predicted resting metabolic rate in obese and nonobese adolescents. Journal of Pediatrics, 127, 571-577.
- Moreira P, Santos S, Padrao P, Cordeiro T, Bessa M, Valente H, Barros R, Teixeira V, Mitchell V, Lopes C and Moreira A, 2010. Food patterns according to sociodemographics, physical activity, sleeping and obesity in Portuguese children. International Journal of Environmental Research and Public Health, 7, 1121-1138.
- Moshfegh AJ, Rhodes DG, Baer DJ, Murayi T, Clemens JC, Rumpler WV, Paul DR, Sebastian RS, Kuczynski KJ, Ingwersen LA, Staples RC and Cleveland LE, 2008. The US Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. American Journal of Clinical Nutrition, 88, 324-332.
- Motil KJ, Montandon CM and Garza C, 1990. Basal and postprandial metabolic rates in lactating and nonlactating women. American Journal of Clinical Nutrition, 52, 610-615.
- MRI (Max Rubner Institut), 2008a. Ergebnisbericht, Teil 1. Nationale Verzehrsstudie II [Report, Part 1. National Nutrition Survey II]. Max Rubner Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel, 193 pp.
- MRI (Max Rubner Institut), 2008b. Ergebnisbericht, Teil 2. Nationale Verzehrsstudie II [Report, Part 2. National Nutrition Survey II]. Max Rubner Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel, 307 pp.
- Müller MJ, Bosy-Westphal A, Kutzner D and Heller M, 2002. Metabolically active components of fat-free mass and resting energy expenditure in humans: recent lessons from imaging technologies. Obesity Reviews, 3, 113-122.
- Müller MJ, Bosy-Westphal A, Klaus S, Kreymann G, Luhrmann PM, Neuhauser-Berthold M, Noack R, Pirke KM, Platte P, Selberg O and Steiniger J, 2004. World Health Organization equations have shortcomings for predicting resting energy expenditure in persons from a modern, affluent population: generation of a new reference standard from a retrospective analysis of a German database of resting energy expenditure. American Journal of Clinical Nutrition, 80, 1379-1390.
- Nagy LE and King JC, 1984. Postprandial energy expenditure and respiratory quotient during early and late pregnancy. American Journal of Clinical Nutrition, 40, 1258-1263.
- Nedergaard J, Bengtsson T and Cannon B, 2007. Unexpected evidence for active brown adipose tissue in adult humans. American journal of physiology. Endocrinology and metabolism, 293, E444-452.

- Neville MC, Keller R, Seacat J, Lutes V, Neifert M, Casey C, Allen J and Archer P, 1988. Studies in human lactation: milk volumes in lactating women during the onset of lactation and full lactation. *American Journal of Clinical Nutrition*, 48, 1375-1386.
- Nhung BT, Khan NC, Hop LT, Lien DT, Le DS, Hien VT, Kunii D, Sakai T, Nakamori M and Yamamoto S, 2005. FAO/WHO/UNU equations overestimate resting metabolic rate in Vietnamese adults. *European Journal of Clinical Nutrition*, 59, 1099-1104.
- NNR (Nordic Nutrition Recommendations), 2004. Integrating nutrition and physical activity. Nordic Council of Ministers, Copenhagen, Denmark, 436 pp.
- Ocké MC, van Rossum CTM, Fransen HP, Buurma EJM, de Boer EJ, Brants HAM, Niekerk EM, van der Laan JD, Drijvers JJMM and Ghameshlou Z, 2008. Dutch National Food Consumption Survey - Young Children 2005/2006. RIVM Report 350070001/2008, National Institute for Public Health and the Environment, 105 pp.
- Oken E, Kleinman KP, Belfort MB, Hammitt JK and Gillman MW, 2009. Associations of gestational weight gain with short- and longer-term maternal and child health outcomes. *American Journal of Epidemiology*, 170, 173-180.
- Ortega RM, Lopez-Sobaler AM, Ballesteros JM, Perez-Farinos N, Rodriguez-Rodriguez E, Aparicio A, Perea JM and Andres P, 2011. Estimation of salt intake by 24 h urinary sodium excretion in a representative sample of Spanish adults. *British Journal of Nutrition*, 105, 787-794.
- Owen OE, Kavle E, Owen RS, Polansky M, Caprio S, Mozzoli MA, Kendrick ZV, Bushman MC and Boden G, 1986. A reappraisal of caloric requirements in healthy women. *American Journal of Clinical Nutrition*, 44, 1-19.
- Owen OE, Holup JL, D'Alessio DA, Craig ES, Polansky M, Smalley KJ, Kavle EC, Bushman MC, Owen LR, Mozzoli MA and et al., 1987. A reappraisal of the caloric requirements of men. *American Journal of Clinical Nutrition*, 46, 875-885.
- Panter-Brick C, 1993. Seasonality of energy expenditure during pregnancy and lactation for rural Nepali women. *American Journal of Clinical Nutrition*, 57, 620-628.
- Paturi M, Tapanainen H, Reinivuo H and Pietinen P, 2008. The National FINDiet 2007 Survey. Report B23/2008, KTL-National Public Health Institute, 230 pp.
- Pedersen AN, Fagt S, Groth MV, Christensen T, Biloft-Jensen A, Matthiessen J, Lyhne Andersen N, Kørup K, Hartkopp H, Hess Ygil K, Hinsch HJ, Saxholt E and Trolle E, 2010. Danskernes kostvaner 2003-2008. Hovedresultater [Dietary habits in Denmark 2003-2008. Main results]. DTU Fødevareinstituttet, 200 pp.
- Petrova S and Angelova K, 2006. Food-based dietary guidelines for Bulgarian adults – scientific background for development and formulation. *Advances in Bulgarian Science*, 4, 19-33.
- Piers LS, Diggavi SN, Thangam S, van Raaij JM, Shetty PS and Hautvast JG, 1995. Changes in energy expenditure, anthropometry, and energy intake during the course of pregnancy and lactation in well-nourished Indian women. *American Journal of Clinical Nutrition*, 61, 501-513.
- Pietinen P, Paturi M, Reinivuo H, Tapanainen H and Valsta LM, 2010. FINDIET 2007 Survey: energy and nutrient intakes. *Public Health Nutrition*, 13, 920-924.
- Pomerleau J, McKee M, Robertson A, Kadziauskiene K, Abaravicius A, Vaask S, Pudule I and Grinberga D, 2001. Macronutrient and food intake in the Baltic republics. *European Journal of Clinical Nutrition*, 55, 200-207.
- Prentice AM and Prentice A, 1988. Energy costs of lactation. *Annual Review of Nutrition*, 8, 63-79.
- Prentice AM, Goldberg GR, Davies HL, Murgatroyd PR and Scott W, 1989. Energy-sparing adaptations in human pregnancy assessed by whole-body calorimetry. *British Journal of Nutrition*, 62, 5-22.



- Prentice AM, 1995. Alcohol and obesity. *International Journal of Obesity and Related Metabolic Disorders*, 19 Suppl 5, S44-50.
- Prentice AM, Spaaij CJ, Goldberg GR, Poppitt SD, van Raaij JM, Totton M, Swann D and Black AE, 1996a. Energy requirements of pregnant and lactating women. *European Journal of Clinical Nutrition*, 50 Suppl 1, S82-110.
- Prentice AM, Goldberg GR, Murgatroyd PR and Cole TJ, 1996b. Physical activity and obesity: problems in correcting expenditure for body size. *International Journal of Obesity and Related Metabolic Disorders*, 20, 688-691.
- Raben A, Agerholm-Larsen L, Flint A, Holst JJ and Astrup A, 2003. Meals with similar energy densities but rich in protein, fat, carbohydrate, or alcohol have different effects on energy expenditure and substrate metabolism but not on appetite and energy intake. *American Journal of Clinical Nutrition*, 77, 91-100.
- Ramirez-Zea M, 2005. Validation of three predictive equations for basal metabolic rate in adults. *Public Health Nutrition*, 8, 1213-1228.
- RCPCH, 2011. UK-WHO Growth Charts: early years. Royal College of Paediatrics and Child Health. Available at: <http://www.rcpch.ac.uk/growthcharts>. Accessed on November 2012.
- Reilly JJ, Lord A, Bunker VW, Prentice AM, Coward WA, Thomas AJ and Briggs RS, 1993. Energy balance in healthy elderly women. *British Journal of Nutrition*, 69, 21-27.
- Reilly JJ, Ashworth S and Wells JC, 2005. Metabolisable energy consumption in the exclusively breast-fed infant aged 3--6 months from the developed world: a systematic review. *British Journal of Nutrition*, 94, 56-63.
- Roberts SB, Paul AA, Cole TJ and Whitehead RG, 1982. Seasonal changes in activity, birth weight and lactational performance in rural Gambian women. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 76, 668-678.
- Roberts SB, 1996. Energy requirements of older individuals. *European Journal of Clinical Nutrition*, 50 Suppl 1, S112-117.
- Roberts SB and Dallal GE, 2005. Energy requirements and aging. *Public Health Nutrition*, 8, 1028-1036.
- Roberts SB and Rosenberg I, 2006. Nutrition and aging: changes in the regulation of energy metabolism with aging. *Physiological Reviews*, 86, 651-667.
- Rodler I, Bíró L, Greiner E, Zajkás G, Szórád I, Varga A, Domonkos A, Ágoston H, Balázs A, Mozsáry E, Vitrai J, Hermann D, Boros J, Németh R and Kéki Z, 2005. Táplálkozási vizsgálat Magyarországon, 2003–2004. Energia- és makrotápanyagbevitel [Dietary survey in Hungary, 2003–2004. Energy and macro-nutrient intake]. *Orvosi Hetilap [Hungarian Medical Journal]*, 146, 1781–1789.
- Rodriguez G, Moreno LA, Sarria A, Fleta J and Bueno M, 2002. Resting energy expenditure in children and adolescents: agreement between calorimetry and prediction equations. *Clinical Nutrition*, 21, 255-260.
- Rolland-Cachera MF, Deheeger M, Maillot M and Bellisle F, 2006. Early adiposity rebound: causes and consequences for obesity in children and adults. *International Journal of Obesity*, 30 Suppl 4, S11-17.
- Rothenberg EM, Bosaeus IG, Westertorp KR and Steen BC, 2000. Resting energy expenditure, activity energy expenditure and total energy expenditure at age 91-96 years. *British Journal of Nutrition*, 84, 319-324.
- Roza AM and Shizgal HM, 1984. The Harris Benedict equation reevaluated: resting energy requirements and the body cell mass. *American Journal of Clinical Nutrition*, 40, 168-182.

- Ruston D, Hoare J, Henderson L, Gregory J, Bates C, Prentice A, Birch M, Swan G and Farron M, 2004. The National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 4: Nutritional Status (anthropometry and blood analytes), blood pressure and physical activity. TSO, London, 166 pp.
- Saarela T, Kokkonen J and Koivisto M, 2005. Macronutrient and energy contents of human milk fractions during the first six months of lactation. *Acta Paediatrica*, 94, 1176-1181.
- SACN (Scientific Advisory Committee on Nutrition), 2011. Dietary reference values for energy. TSO, 220 pp.
- Sadurskis A, Kabir N, Wager J and Forsum E, 1988. Energy metabolism, body composition, and milk production in healthy Swedish women during lactation. *American Journal of Clinical Nutrition*, 48, 44-49.
- SCF (Scientific Committee on Food), 1993. Report on nutrient and energy intakes for the European Community, 31st Series. Food - Science and Technique, European Commission, 255 pp.
- Schoeller DA, 1988. Measurement of energy expenditure in free-living humans by using doubly labeled water. *Journal of Nutrition*, 118, 1278-1289.
- Schoeller DA, 2007. Making indirect calorimetry a gold standard for predicting energy requirements for institutionalized patients. *Journal of the American Dietetic Association*, 107, 390-392.
- Schofield WN, Schofield C and James WTP, 1985. Basal metabolic rate: Review and prediction, together with an annotated bibliography of source material. *Human Nutrition. Clinical Nutrition*, 39C Suppl 1, 1-96.
- Schulz LO and Schoeller DA, 1994. A compilation of total daily energy expenditures and body weights in healthy adults. *American Journal of Clinical Nutrition*, 60, 676-681.
- Schutz Y, Bessard T and Jequier E, 1984. Diet-induced thermogenesis measured over a whole day in obese and nonobese women. *American Journal of Clinical Nutrition*, 40, 542-552.
- Schutz Y, Golay A and Jequier E, 1988. 24 h energy expenditure (24-EE) in pregnant women with a standardised activity level. *Experientia*, 44, A31.
- Seale JL, Rumpler WV and Moe PW, 1991. Description of a direct-indirect room-sized calorimeter. *American Journal of Physiology*, 260, E306-320.
- Serra-Majem L, Ribas-Barba L, Salvador G, Jover L, Raido B, Ngo J and Plasencia A, 2007. Trends in energy and nutrient intake and risk of inadequate intakes in Catalonia, Spain (1992-2003). *Public Health Nutrition*, 10, 1354-1367.
- Sette S, Le Donne C, Piccinelli R, Arcella D, Turrini A and Leclercq C, 2010. The third Italian National Food Consumption Survey, INRAN-SCAI 2005-06 - Part 1: Nutrient intakes in Italy. *Nutrition, Metabolism and Cardiovascular Diseases*, 1-11.
- Sharp TA, Bell ML, Grunwald GK, Schmitz KH, Sidney S, Lewis CE, Tolan K and Hill JO, 2002. Differences in resting metabolic rate between white and African-American young adults. *Obesity Research*, 10, 726-732.
- Shetty PS, Henry CJ, Black AE and Prentice AM, 1996. Energy requirements of adults: an update on basal metabolic rates (BMRs) and physical activity levels (PALs). *European Journal of Clinical Nutrition*, 50 Suppl 1, S11-23.
- Silva JE, 2006. Thermogenic mechanisms and their hormonal regulation. *Physiological Reviews*, 86, 435-464.
- Smit HA, Verschuren WMM and Bueno de Mesquita HB, 1994. Monitoring van risicofactoren en gezondheid in Nederland (MORGEN-project): Doelstellingen en werkwijze [Monitoring of risk factors and health in the Netherlands (MORGEN-project): Objectives and methods]. RIVM-rapport nummer 263 200 001, 104 pp.

- Sorkin JD, Muller DC and Andres R, 1999. Longitudinal change in height of men and women: implications for interpretation of the body mass index: the Baltimore Longitudinal Study of Aging. *American Journal of Epidemiology*, 150, 969-977.
- Spaaij CJ, van Raaij JM, de Groot LC, van der Heijden LJ, Boekholt HA and Hautvast JG, 1994a. Effect of lactation on resting metabolic rate and on diet- and work-induced thermogenesis. *American Journal of Clinical Nutrition*, 59, 42-47.
- Spaaij CJ, van Raaij JM, Van der Heijden LJ, Schouten FJ, Drijvers JJ, De Groot LC, Boekholt HA and Hautvast JG, 1994b. No substantial reduction of the thermic effect of a meal during pregnancy in well-nourished Dutch women. *British Journal of Nutrition*, 71, 335-344.
- Stock MJ and Rothwell NJ, 1981. Sympathetic control of brown adipose tissue in the regulation of body weight. *Biochemical Society Transactions*, 9, 525-527.
- Stotland NE, Cheng YW, Hopkins LM and Caughey AB, 2006. Gestational weight gain and adverse neonatal outcome among term infants. *Obstetrics and Gynecology*, 108, 635-643.
- Subar AF, Kipnis V, Troiano RP, Midthune D, Schoeller DA, Bingham S, Sharbaugh CO, Trabulsi J, Runswick S, Ballard-Barbash R, Sunshine J and Schatzkin A, 2003. Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study. *American Journal of Epidemiology*, 158, 1-13.
- Szponar L, Sekula W, Nelson M and Weisell RC, 2001. The Household Food Consumption and Anthropometric Survey in Poland. *Public Health Nutrition*, 4, 1183-1186.
- Szponar L, Sekula W, Rychlik E, Ołtarzewski M and Figurska K, 2003. Badania indywidualnego spożycia żywności i stanu odżywienia w gospodarstwach domowych [Household food consumption and anthropometric survey]. Instytut Żywności i Żywienia, Warsaw, Poland, 900 pp.
- Teixeira PJ, Going SB, Houtkooper LB, Metcalfe LL, Blew RM, Flint-Wagner HG, Cussler EC, Sardinha LB and Lohman TG, 2003. Resistance training in postmenopausal women with and without hormone therapy. *Medicine and Science in Sports and Exercise*, 35, 555-562.
- Tennefors C, Coward WA, Hernell O, Wright A and Forsum E, 2003. Total energy expenditure and physical activity level in healthy young Swedish children 9 or 14 months of age. *European Journal of Clinical Nutrition*, 57, 647-653.
- Thomson MA, Bucolo S, Quirk P and Shepherd RW, 1995. Measured versus predicted resting energy expenditure in infants: a need for reappraisal. *Journal of Pediatrics*, 126, 21-27.
- TNO/LUMC (Nederlandse Organisatie voor toegepast-natuurwetenschappelijk onderzoek/Leids Universitair Medisch Centrum), 1998. Groeidiagrammen van de vierde landelijke groeistudie [Growth charts for the fourth nationwide growth study]. Bohn Stafleu Van Loghum.
- Tooze JA, Schoeller DA, Subar AF, Kipnis V, Schatzkin A and Troiano RP, 2007. Total daily energy expenditure among middle-aged men and women: the OPEN Study. *American Journal of Clinical Nutrition*, 86, 382-387.
- Torun B and Viteri FE, 1994. Influence of exercise on linear growth. *European Journal of Clinical Nutrition*, 48 Suppl 1, S186-189.
- Torun B, Davies PS, Livingstone MB, Paolisso M, Sackett R and Spurr GB, 1996. Energy requirements and dietary energy recommendations for children and adolescents 1 to 18 years old. *European Journal of Clinical Nutrition*, 50 Suppl 1, S37-80.
- Torun B, 2001. Energy requirements of children and adolescents. Background paper prepared for the joint FAO/WHO/UNU Expert consultation on Energy in Human Nutrition, 2001.
- Torun B, 2005. Energy requirements of children and adolescents. *Public Health Nutrition*, 8, 968-993.
- Troiano RP, Frongillo EA, Jr., Sobal J and Levitsky DA, 1996. The relationship between body weight and mortality: a quantitative analysis of combined information from existing studies. *International Journal of Obesity and Related Metabolic Disorders*, 20, 63-75.

- Tuazon MA, van Raaij JM, Hautvast JG and Barba CV, 1987. Energy requirements of pregnancy in the Philippines. *Lancet*, 2, 1129-1131.
- Valencia ME, McNeill G, Brockway JM and Smith JS, 1992. The effect of environmental temperature and humidity on 24 h energy expenditure in men. *British Journal of Nutrition*, 68, 319-327.
- van Beusekom CM, Zeegers TA, Martini IA, Velvis HJ, Visser GH, van Doormaal JJ and Muskiet FA, 1993. Milk of patients with tightly controlled insulin-dependent diabetes mellitus has normal macronutrient and fatty acid composition. *American Journal of Clinical Nutrition*, 57, 938-943.
- van Buuren S, Schönbeck Y and van Dommelen P, 2012. CT/EFSA/NDA/2010/01: Collection, collation and analysis of data in relation to reference heights and reference weights for female and male children and adolescents (0-18 years) in the EU, as well as in relation to the age of onset of puberty and the age at which different stages of puberty are reached in adolescents in the EU. Technical Report submitted to EFSA, 59 pp.
- van Raaij JM, Vermaat-Miedema SH, Schonk CM, Peek ME and Hautvast JG, 1987. Energy requirements of pregnancy in The Netherlands. *Lancet*, 2, 953-955.
- van Raaij JM, Schonk CM, Vermaat-Miedema SH, Peek ME and Hautvast JG, 1990. Energy cost of walking at a fixed pace and self-paced before, during, and after pregnancy. *American Journal of Clinical Nutrition*, 51, 158-161.
- van Raaij JM, Schonk CM, Vermaat-Miedema SH, Peek ME and Hautvast JG, 1991. Energy cost of lactation, and energy balances of well-nourished Dutch lactating women: reappraisal of the extra energy requirements of lactation. *American Journal of Clinical Nutrition*, 53, 612-619.
- van Rossum CTM, Fransen HP, Verkaik-Kloosterman J, Buurma-Rethans EJM and Ocké MC, 2011. Dutch National Food Consumption Survey 2007-2010: Diet of children and adults aged 7 to 69 years. RIVM Report number: 350050006/2011, National Institute for Public Health and the Environment, 148 pp.
- Vander Weg MW, Watson JM, Klesges RC, Eck Clemens LH, Slawson DL and McClanahan BS, 2004. Development and cross-validation of a prediction equation for estimating resting energy expenditure in healthy African-American and European-American women. *European Journal of Clinical Nutrition*, 58, 474-480.
- Vaughan L, Zurlo F and Ravussin E, 1991. Aging and energy expenditure. *American Journal of Clinical Nutrition*, 53, 821-825.
- Vaz M, Karaolis N, Draper A and Shetty P, 2005. A compilation of energy costs of physical activities. *Public Health Nutrition*, 8, 1153-1183.
- Viteri F and Torun B, 1981. Nutrition, physical activity and growth. In: *The biology of normal human growth*. Eds Ritzén M, Aperia A, Hall K, Larsson A, Zetterberg A, Zetterström R. Raven Press, New York, USA, 265-273.
- Wahrlich V, Anjos LA, Going SB and Lohman TG, 2006. Validation of the VO2000 calorimeter for measuring resting metabolic rate. *Clinical Nutrition*, 25, 687-692.
- Walsberg GE and Hoffman TC, 2005. Direct calorimetry reveals large errors in respirometric estimates of energy expenditure. *Journal of Experimental Biology*, 208, 1035-1043.
- Wang Z, Ying Z, Bosy-Westphal A, Zhang J, Schautz B, Later W, Heymsfield SB and Muller MJ, 2010. Specific metabolic rates of major organs and tissues across adulthood: evaluation by mechanistic model of resting energy expenditure. *American Journal of Clinical Nutrition*, 92, 1369-1377.
- Warwick PM, 2006. Factorial estimation of daily energy expenditure using a simplified method was improved by adjustment for excess post-exercise oxygen consumption and thermic effect of food. *European Journal of Clinical Nutrition*, 60, 1337-1340.

- Webb P, 1986. 24-hour energy expenditure and the menstrual cycle. *American Journal of Clinical Nutrition*, 44, 614-619.
- Weijs PJ, Kruizenga HM, van Dijk AE, van der Meij BS, Langius JA, Knol DL, Strack van Schijndel RJ and van Bokhorst-de van der Schueren MA, 2008. Validation of predictive equations for resting energy expenditure in adult outpatients and inpatients. *Clinical Nutrition*, 27, 150-157.
- Weijs PJ, 2008. Validity of predictive equations for resting energy expenditure in US and Dutch overweight and obese class I and II adults aged 18-65 y. *American Journal of Clinical Nutrition*, 88, 959-970.
- Weijs PJ and Vansant GA, 2010. Validity of predictive equations for resting energy expenditure in Belgian normal weight to morbid obese women. *Clinical Nutrition*, 29, 347-351.
- Weir JB, 1949. New methods for calculating metabolic rate with special reference to protein metabolism. *The Journal of physiology*, 109, 1-9.
- Wells JCK, 1994. Energy metabolism in breast-fed and formula-fed infants. Thesis (PhD), University of Cambridge, Cambridge, UK.
- Westerterp KR, Meijer GA, Janssen EM, Saris WH and Ten Hoor F, 1992. Long-term effect of physical activity on energy balance and body composition. *British Journal of Nutrition*, 68, 21-30.
- Westerterp KR and Goran MI, 1997. Relationship between physical activity related energy expenditure and body composition: a gender difference. *International Journal of Obesity and Related Metabolic Disorders*, 21, 184-188.
- Westerterp KR and Plasqui G, 2004. Physical activity and human energy expenditure. *Current Opinion in Clinical Nutrition and Metabolic Care*, 7, 607-613.
- Weyand PG, Smith BR, Puyau MR and Butte NF, 2010. The mass-specific energy cost of human walking is set by stature. *Journal of Experimental Biology*, 213, 3972-3979.
- Weyer C, Snitker S, Bogardus C and Ravussin E, 1999. Energy metabolism in African Americans: potential risk factors for obesity. *American Journal of Clinical Nutrition*, 70, 13-20.
- WHO (World Health Organization), 1983. Measuring change in nutritional status. Guidelines for assessing the nutritional impact of supplementary feeding programmes for vulnerable groups. 100 pp.
- WHO (World Health Organization), 1985. Energy and protein requirements: Report of a joint FAO/WHO/UNU expert consultation. WHO Technical Report Series No. 724, 206 pp.
- WHO (World Health Organization), 1995a. Maternal anthropometry and pregnancy outcomes. A WHO Collaborative Study. *Bulletin of the World Health Organization*, 73 Suppl, 1-98.
- WHO (World Health Organization), 1995b. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. WHO Technical Report Series. No. 854, 452 pp.
- WHO (World Health Organization), 1998. Complementary feeding of young children in developing countries. A review of current scientific knowledge. WHO/NUT/98.1, 248 pp.
- WHO (World Health Organization), 2000. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. WHO Technical Report Series No. 894, 253 pp.
- WHO (World Health Organization), 2007. Growth reference data for 5-19 years. Available from: <http://www.who.int/growthref/en/>. Accessed on November 2012.
- WHO Multicentre Growth Reference Study Group (World Health Organization), 2006. WHO (World Health Organization) Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. 336 pp.
- WHO Working Group on Infant Growth (World Health Organization), 1994. An evaluation of infant growth. WHO/NUT/94.8, 85 pp.



- Wijers SL, Saris WH and van Marken Lichtenbelt WD, 2009. Recent advances in adaptive thermogenesis: potential implications for the treatment of obesity. *Obesity Reviews*, 10, 218-226.
- Willett WC, Dietz WH and Colditz GA, 1999. Guidelines for healthy weight. *New England Journal of Medicine*, 341, 427-434.
- Wilms B, Schmid SM, Ernst B, Thurnheer M, Mueller MJ and Schultes B, 2010. Poor prediction of resting energy expenditure in obese women by established equations. *Metabolism: Clinical and Experimental*, 59, 1181-1189.
- Withers RT, Smith DA, Tucker RC, Brinkman M and Clark DG, 1998. Energy metabolism in sedentary and active 49- to 70-yr-old women. *Journal of Applied Physiology*, 84, 1333-1340.
- Wouters-Adriaens MP and Westerterp KR, 2006. Basal metabolic rate as a proxy for overnight energy expenditure: the effect of age. *British Journal of Nutrition*, 95, 1166-1170.
- Yang X, Li M, Mao D, Zeng G, Zhuo Q, Hu W, Piao J, Yang X and Huang C, 2010. Basal energy expenditure in southern Chinese healthy adults: measurement and development of a new equation. *British Journal of Nutrition*, 104, 1817-1823.
- Zuntz N, 1897. Über den Stoffverbrauch des Hundes bei Muskelarbeit. *Pflügers Archiv für die gesammte Physiologie des Menschen und der Thiere*, 68, 191-211.

APPENDICES

APPENDIX 1: ENERGY CONTENT OF HUMAN MILK FROM HEALTHY MOTHERS OF TERM INFANTS IN THE EU

Reference	n (samples)	Country	Stage of lactation	Energy content (mean ± SD)		Gross energy (GE) or metabolisable energy (ME)	Method of analysis
				kJ/g	kcal/g		
Department of Health and Social Security (1977) <sup>(a)</sup>	14	UK	NA	2.8 ± 0.5	0.67 ± 0.12	Estimated ME	NA
Lucas et al. (1987)	12	UK	5 weeks 11 weeks	2.48 ± 0.21 <sup>(b, c)</sup> 2.58 ± 0.21 <sup>(b, c)</sup>	0.59 ± 0.05 <sup>(b, c)</sup> 0.62 ± 0.05 <sup>(b, c)</sup>	ME	Metabolisable energy intake per day measured by DLW/milk volume intake per day
Wells (1994) <sup>(a)</sup>	21	UK		2.5 ± 0.5	0.60 ± 0.12	ME	NA
van Beusekom et al. (1993)	5	NL	0-5 days 6-10 days >10 days	2.51 ± 0.57 <sup>(b, c, d)</sup> 2.60 ± 0.22 <sup>(b, c, d)</sup> 2.59 ± 0.29 <sup>(b, c, d)</sup>	0.60 ± 0.14 <sup>(b, c, d)</sup> 0.62 ± 0.05 <sup>(b, c, d)</sup> 0.62 ± 0.07 <sup>(b, c, d)</sup>	ME	Calculated from carbohydrate, protein, and fat content
de Bruin et al. (1998)	23	NL	1 month 2 months 4 months	2.81 <sup>(b)</sup> 2.72 <sup>(b)</sup> 2.59 <sup>(b)</sup>	0.67 <sup>(b)</sup> 0.65 <sup>(b)</sup> 0.62 <sup>(b)</sup>	GE	Calculated from carbohydrate, protein, and fat content
Michaelsen et al. (1994)	(96)	DK	2-9 months	3.17 ± 0.45 <sup>(b)</sup>	0.76 ± 0.11 <sup>(b)</sup>	GE	Calculated from carbohydrate, protein, and fat content
Saarela et al. (2005)	53 (253)	FIN	1 week 1 month 2 months 3 months 4 months 5 months 6 months	2.92 ± 0.41 <sup>(b)</sup> 3.00 ± 0.40 <sup>(b)</sup> 2.98 ± 0.43 <sup>(b)</sup> 2.84 ± 0.51 <sup>(b)</sup> 2.79 ± 0.53 <sup>(b)</sup> 2.72 ± 0.44 <sup>(b)</sup> 2.79 ± 0.47 <sup>(b)</sup>	0.70 ± 0.10 <sup>(b)</sup> 0.72 ± 0.10 <sup>(b)</sup> 0.71 ± 0.10 <sup>(b)</sup> 0.68 ± 0.12 <sup>(b)</sup> 0.67 ± 0.13 <sup>(b)</sup> 0.65 ± 0.11 <sup>(b)</sup> 0.67 ± 0.11 <sup>(b)</sup>	GE	Calculated from lactose, protein, and fat content
Sadurskis et al. (1988)	23	SWE	2 months	2.68 ± 0.33	0.64 ± 0.08	GE	Bomb calorimeter

(a): cited from Reilly et al. (2005).

(b): calculated using a correction for the density of milk of 1.032 g/mL according to Neville et al. (1988).

(c): mean ± standard error.

(d): calculated by Hester et al. (2012).

NA, not available; UK, United Kingdom; NL, the Netherlands; DK, Denmark; FIN, Finland; SWE, Sweden.

**APPENDIX 2: PREDICTIVE EQUATIONS FOR REE IN ADULTS**

Author	Year	Number of subjects	Age range or mean	BMI range or mean	Remarks on large database	Sex	Statistics and cross validation	REE predictive equations
Harris and Benedict <sup>(a)</sup> (kcal/day) (USA)	(1919)	239	27 (M) 31 (F)	-	-	136 M 103 F	M: r=0.86, CL=±211 kcal <sup>a</sup> F: r=0.77, CL=±212 kcal	M: BM x 13.7516 + HTCM x 5.0033 – AGE x 6.755+66.473 F: BM x 9.5634 + HTCM x 1.8496 – AGE x 4.6756+655.0955
Schofield et al. <sup>(b)</sup> (body mass) (MJ/day)	(1985)	7,173, including children (about 11,000 values, including group mean values)	n=4,814 >18 y	mean BMIs of the 6 groups: between 21 and 24	n=3,388 Italians (47 %), n=615 tropical residents, n=322 Indians; 114 published studies, most European and North American subjects (Italian, closed circuit calorimetry)	4,809 M 2,364 F	M: r=0.65, SE=0.64, n=2879 M: r=0.60, SE=0.70, n=646 M: r=0.71, SE=0.69, n=50 F: r=0.73, SE=0.49, n=829 F: r=0.68, SE=0.47, n=372 F: r=0.68, SE=0.45, n=38	M: AGE 18-30 y: 0.063 x BM + 2.896 M: AGE 30-60 y: 0.048 x BM + 3.653 M: AGE ≥60 y: 0.049 x BM + 2.459 F: AGE 18-30 y: 0.062 x BM + 2.036 F: AGE 30-60 y: 0.034 x BM + 3.538 F: AGE ≥60 y: 0.038 x BM + 2.755
Schofield et al. <sup>(b)</sup> (body mass and height) (MJ/day)	(1985)	7,173, including children (about 11,000 values, including group mean values)	n=4,814 >18 y	mean BMIs of the 6 groups: between 21 and 24	n=3,388 Italians (47 %), n=615 tropical residents, n=322 Indians; 114 published studies, most European and North American subjects (Italian, closed circuit calorimetry)	4,809 M 2,364 F	M: r=0.65, SE=0.64, n=2879 M: r=0.60, SE=0.70, n=646 M: r=0.74, SE=0.66, n=50 F: r=0.73, SE=0.49, n=829 F: r=0.68, SE=0.47, n=372 F: r=0.73, SE=0.43, n=38	M: AGE 18-30 y: 0.063 x BM – 0.042 x HTM + 2.953 M: AGE 30-60 y: 0.048 x BM – 0.011 x HTM + 3.67 M: AGE ≥60 y: 0.038 x BM + 4.068 x HTM – 3.491 F: AGE 18-30 y: 0.057 x BM + 1.184 x HTM + 0.411 F: AGE 30-60 y: 0.034 x BM + 0.006 x HTM + 3.53 F: AGE ≥60 y: 0.033 x BM + 1.917 x HTM + 0.074
FAO <sup>(b)</sup> (body mass) (MJ/day)	(1985)	This report mentions that the equations are based on Schofield et al (1985); however, the figures in Schofield et al. (1985) differ slightly from those of the FAO because additional data were included by the authors of that analysis after the FAO report was compiled.					M: r=0.65, SD=0.632 M: r=0.60, SD=0.686 M: r=0.79, SD=0.619 F: r=0.72, SD=0.506 F: r=0.70, SD=0.452 F: r=0.74, SD=0.452	M: AGE 18–30 y: 0.0640 x BM + 2.84 M: AGE 30–60 y: 0.0485 x BM + 3.67 M: AGE ≥60 y: 0.0565 x BM + 2.04 F: AGE 18–30 y: 0.0615 x BM + 2.08 F: AGE 30–60 y: 0.0364 x BM + 3.47 F: AGE ≥60 y: 0.0439 x BM + 2.49

Author	Year	Number of subjects	Age range or mean	BMI range or mean	Remarks on large database	Sex	Statistics and cross validation	REE predictive equations
FAO <sup>(b)</sup> (body mass and height) (MJ/day)	(1985)	This report mentions that the equations are based on Schofield et al (1985); however, the figures in Schofield et al. (1985) differ slightly from those of the FAO because additional data were included by the authors of that analysis after the FAO report was compiled.					M: r=0.65, RSD=0.632 M: r=0.60, RSD=0.686 M: r=0.84, RSD=0.552 F: r=0.73, RSD=0.502 F: r=0.70, RSD=0.452 F: r=0.82, RSD=0.393	M: AGE 18–30 y: 0.0644 x BM – 0.1130 x HTM + 3.0 M: AGE 30–60 y: 0.0472 x BM + 0.0669 x HTM + 3.769 M: AGE ≥60 y: 0.0368 x BM + 4.7195 x HTM – 4.481 F: AGE 18–30 y: 0.0556 x BM + 1.3974 x HTM + 0.146 F: AGE 30–60 y: 0.0364 x BM – 0.1046 x HTM + 3.619 F: AGE ≥60 y: 0.0385 x BM + 2.6652 x HTM – 1.264
Owen et al. (kcal/day) (USA)	(1987) (M)	104	18-82 y (M)	20.4-58.7 (M)	-	60 M (including 16 obese, BMI>30 kg/m <sup>2</sup> )	M: r=0.75	M: BM x 10.2 + 879
	(1986) (F)		18-65 (F)	18.2-49.6 (F)		44 F (including 16 obese, BMI>30 kg/m <sup>2</sup> )	F: r=0.74	F: BM x 7.18 + 795
Mifflin et al. (kcal/day) (USA)	(1990)	498 (264 normal body mass, 234 obese)	19-78 y	17-42	-	251 M (129 normal body mass, 122 obese), 247 F (135 normal body mass, 112 obese)	R <sup>2</sup> =0.71	9.99 x BM + 6.25 x HTCM – 4.92 x AGE + 166 x SEX - 161
De Lorenzo et al. (kJ/day) (Italy)	(2001)	320	18–59 y	Mean: about 27 (range: 18.6–40)	-	127 M, 193 F	M: R <sup>2</sup> =0.597, SEE=650 kJ/day F: R <sup>2</sup> =0.597, SEE=581 kJ/day	M: 53.284 x BM + 20.957 x HTCM – 23.859 x AGE + 487 F: 46.322 x BM + 15.744 x HTCM – 16.66 x AGE + 944
Müller et al. <sup>(b)</sup> (MJ/day) (Germany)	(2004)	2,528 (development of equation in sub- population 1: n=1,046)	5–91 y	Mean BMI: 27	-	1,027 M, 1,501 F (development: 388 M, 658 F)	Development: R <sup>2</sup> =0.73, SEE=0.83. Cross-validation in sub- population 2: n=1,059 (410 M, 649 F) <sup>e</sup> , r=0.83	0.047 x BM – 0.01452 x AGE + 1.009 x SEX + 3.21
Müller et al. <sup>(b)</sup> (BMI <sup>(d)</sup> , MJ/day) (Germany)	(2004)	2,528 (development of equation in sub- population 1: n=1,046)	5–91 y	Mean BMI: 27	-	1,027 M, 1,501 F (development: 388 M, 658 F)	Development: R <sup>2</sup> =0.52, SEE=0.79 (n=444, for BMI >18.5 to 25). Cross-validation in sub- population 2: r=0.72. Development: R <sup>2</sup> =0.62, SEE=0.77 (n=266, for BMI >25 to <30). Cross-validation in sub- population 2: r=0.79. Development: R <sup>2</sup> =0.75, SEE=0.91 (n=278, for BMI ≥30). Cross- validation in sub-population 2: r=0.84	BMI >18.5 to 25: 0.02219 x BM +0.02118 x HTCM – 0.01191 x AGE + 0.884 x SEX + 1.233 BMI >25 to <30: 0.04507 x BM - 0.01553 x AGE + 1.006 x SEX + 3.407 BMI ≥30: 0.05 x BM - 0.01586 x AGE + 1.103 x SEX + 2.924

Author	Year	Number of subjects	Age range or mean	BMI range or mean	Remarks on large database	Sex	Statistics and cross validation	REE predictive equations
Müller et al. <sup>(b)</sup> (FFM <sup>(d)</sup> , MJ/day) (Germany)	(2004)	2,528 (development of equation in sub-population 1: n=1,046)	5–91 y	Mean BMI: 27	-	1,027 M, 1,501 F (development: 388 M, 658 F),	Development: R <sup>2</sup> =0.71, SEE=0.77. Cross-validation in sub-population 2: n=1,059 (410 M, 649 F) <sup>c</sup> : r=0.83	0.05192 x FFM + 0.04036 x FM + 0.869 x SEX - 0.01181 x AGE + 2.992
Müller et al. <sup>(b)</sup> (BMI <sup>(f)</sup> and FFM <sup>(d)</sup> , MJ/day) (Germany)	(2004)	2,528 (development of equation in sub-population 1: n=1,046)	5–91 y	Mean BMI: 27	-	1,027 M, 1,501 F (development: 388 M, 658 F),	Development: R <sup>2</sup> =0.54, SEE=0.78, (n=444, for BMI > 18.5 to 25). Cross-validation in sub-population 2: r=0.75. Development: R <sup>2</sup> =0.65, SEE=0.62 (n=266, for BMI >25 to <30). Cross-validation in sub-population 2: r=0.79 Development: R <sup>2</sup> =0.70, SEE=0.87 (n=278, for BMI ≥30). Cross-validation in sub-population 2: r=0.84.	BMI > 18.5 to 25: 0.0455 x FFM + 0.0278 x FM + 0.879 x SEX - 0.01291 x AGE + 3.634  BMI >25 to <30: 0.03776 x FFM + 0.03013 x FM + 0.93 x SEX - 0.01196 x AGE + 3.928  BMI ≥30: 0.05685 x FFM + 0.04022 x FM + 0.808 x SEX - 0.01402 x AGE + 2.818
Vander Weg et al. (Memphis equation) (kJ/day) (USA)	(2004)	471 women (development of equation in sub-population 1: 239 women)	18-39 y	Mean BMI: 25.2		471 women (205 African American, 266 European American)	Development: 239 women (97 African American, 142 European American); adjusted R <sup>2</sup> : 0.51. Cross-validation in sub-population 2: 232 women (108 African American, 124 European American); adjusted R <sup>2</sup> : 0.55 for African American, 0.31 for European American.	616.93 - 14.9 x AGE + 35.12 x BM + 19.83 x HTCM - 271.88 x ETHNICITY
Henry <sup>(b)</sup> (body mass) (MJ/day)  (Oxford Database)	(2005)	10,552 (10,502) including children	-	-	166 separate investigations, only individual data points; all Italian, closed circuit data excluded; 4,018 subjects from the tropics included	5,794 M          4,702 F (4,708)	M: r= 0.760, SE=0.652; n=2,821  M: r=0.742, SE=0.693; n=1,010  M: r=0.776, SE=0.685; n=534  M: r=0.766, SE=0.697; n=270  M: r=0.779, SE=0.667; n=264  F: r=0.700, SE=0.564; n=1,664  F: r=0.690, SE=0.581; n=1,023  F: r=0.786, SE=0.485; n=334 (340)  F: r=0.796, SE=0.476; n=185  F: r=0.746, SE=0.518; n=155	M: AGE 18–30 y: 0.0669 x BM + 2.28  M: AGE 30–60 y: 0.0592 x BM + 2.48  M: AGE ≥60 y: 0.0563 x BM + 2.15  M: AGE 60-70 y: 0.0543 x BM + 2.37  M: AGE ≥70 y: 0.0573 x BM + 2.01  F: AGE 18–30 y: 0.0546 x BM + 2.33  F: AGE 30–60 y: 0.0407 x BM + 2.90  F: AGE ≥60 y: 0.0424 x BM + 2.38  F: AGE 60-70 y: 0.0429 x BM + 2.39  F: AGE ≥70 y: 0.0417 x BM + 2.41



Author	Year	Number of subjects	Age range or mean	BMI range or mean	Remarks on large database	Sex	Statistics and cross validation	REE predictive equations
Henry <sup>(b)</sup>	(2005)	10,552 <i>(10502)</i>	-	-	166 separate investigations, only individual data points; all Italian closed circuit data excluded; 4,018 subjects from the tropics included.	5,794 M	M: r=0.764, SE=0.648; n=2,816	M: AGE 18–30 y: 0.0600 x BM + 1.31 x HTM + 0.473
(body mass and height)		including children					M: r=0.756, SE= 0.678; n=1,006	M: AGE 30–60 y: 0.0476 x BM + 2.26 x HTM - 0.574
(MJ/day)						4,702 F	M: r=0.789, SE=0.668; n= 533	M: AGE ≥60 y: M: 0.0478 x BM + 2.26 x HTM - 1.07
(Oxford database)							F: r=0.724, SE=0.542; n=1,655	F: AGE 18–30 y: 0.0433 x BM + 2.57 x HTM - 1.18
							F: r=0.713, SE=0.564; n=1,023	F: AGE 30–60 y: 0.0342 x BM + 2.10 x HTM - 0.0486
							F: r=0.805, SE=0.472; n=324	F: AGE ≥60 y: 0.0356 x BM + 1.76 x HTM + 0.0448

CL, confidence limits; ETHNICITY (African American = 1, European American = 0); F, female; FFM, fat free mass (kg); FM, fat mass (kg); HTCMT, height in cm; HTM, height in meter; M, male; r, correlation coefficient; SD, standard deviation; SEX (M=1, F=0); SE, standard error; SEE, standard error of estimate; BM, body mass in kg.

<sup>(a)</sup> From: (Roza and Shizgal, 1984) (not the original publication).

<sup>(b)</sup> Equations are also available for children (FAO/WHO/UNU, 1985; Henry, 2005; Müller et al., 2004; Schofield et al., 1985).

<sup>(c)</sup> Equations are also available for BMI ≤18.5, either with body mass, age and sex, or with FFM and FM.

<sup>(d)</sup> Body composition method: bioimpedance analysis (different equations, multicentre study).

<sup>(e)</sup> Including n=482 (BMI >18.5-25), n=267 (BMI >25 to <30), n=261 (BMI ≥30).

<sup>(f)</sup> Equation also available for BMI ≤18.5.

<sup>(g)</sup> Figures given in italics differ from those in the publication but are assumed to be as such after recalculation of the figures, as also stated in Ramirez-Zea (2005) for total number per sex.

**APPENDIX 3A: POPULATION, METHODS AND PERIOD OF DIETARY ASSESSMENT IN CHILDREN AND ADOLESCENTS IN EUROPEAN COUNTRIES**

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Austria</b>	Boys and girls aged 7-9 years	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Boys and girls aged 10-14 years	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Boys and girls aged 14-19 years	24-hour recall	2003-2004	(Elmadfa et al., 2009a; Elmadfa et al., 2009b). <i>Mainly from a large Viennese sample.</i>
<b>Belgium</b>	Boys and girls aged 2.5-3 years	3-day record	2002-2003	(Huybrechts and De Henauw, 2007). <i>Data collected in Flanders.</i>
	Boys and girls aged 4-6.5 years	3-day record	2002-2003	(Huybrechts and De Henauw, 2007). <i>Data collected in Flanders.</i>
	Boys and girls aged 13-15 years	7-day record	1997	(Matthys et al., 2003). <i>Data collected in the region of Ghent in Flanders.</i>
	Boys and girls aged 15-18 years	Two non consecutive 24-hour recalls	2004	(De Vriese et al., 2006)
<b>Bulgaria</b>	Boys and girls aged 1-3 years	24-hour recall	1998	(Abrashveva et al., 1998)
	Boys and girls aged 3-6 years	24-hour recall	1998	(Abrashveva et al., 1998)
	Boys and girls aged 6-10 years	24-hour recall	1998	(Abrashveva et al., 1998)
	Boys and girls aged 10-14 years	24-hour recall	1998	(Abrashveva et al., 1998)
	Boys and girls aged 14-18 years	24-hour recall	1998	(Abrashveva et al., 1998)
<b>Czech Republic</b>	Boys and girls aged 4-6 years	48-hour recall	2007	(Elmadfa et al., 2009a)
	Boys and girls aged 7-9 years	48-hour recall	2007	(Elmadfa et al., 2009a)
<b>Denmark</b>	Boys and girls aged 1-3 years	7-day record	1995	(Andersen et al., 1996)
	Boys and girls aged 4-5 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 6-9 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 10-13 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 14-17 years	7-day record	2003-2008	(Pedersen et al., 2010)
<b>Finland</b>	Children aged 1 year	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 2 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 3 years	3 day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 4 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 6 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
<b>France</b>	Boys and girls aged 4-6 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 7-9 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 10-14 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 15-18 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Germany</b>	Infants aged 12 months	3-day records	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 18 months	3-day records	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 2 years	3-day records	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 3 years	3-day records	1989-2003	(Hilbig and Kersting, 2006).
	Boys and girls aged 6 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 7-9 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 10-11 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 12 years	Dietary history (over the last 4 weeks)	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 13-14 years	Dietary history (over the last 4 weeks)	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 15-17 years	Dietary history (over the last 4 weeks)	2006	(Mensink et al., 2007)
<b>Greece</b>	Boys and girls aged 1-5 years	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006; Manios et al., 2008)
	Boys and girls aged 12-24 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006; Manios et al., 2008)
	Boys and girls aged 25-36 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006; Manios et al., 2008)
	Boys and girls aged 37-48 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006; Manios et al., 2008)
	Boys and girls aged 49-60 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006; Manios et al., 2008)
<b>Hungary</b>	Boys and girls aged 11-14 years	3-day record	2005-2006	(Biro et al., 2007). <i>Data collected in Budapest.</i>
<b>Ireland</b>	Boys and girls aged 1 year	4-day record	2010-2011	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 2 years	4-day record	2010-2011	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 3 years	4-day record	2010-2011	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 4 years	4-day record	2010-2011	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 5-8 years	7-day record	2003-2004	(IUNA (Irish Universities Nutrition Alliance), b)
	Boys and girls aged 9-12 years	7-day record	2003-2004	(IUNA (Irish Universities Nutrition Alliance), b)
	Boys and girls aged 13-14 years	7-day record	2005-2006	(IUNA (Irish Universities Nutrition Alliance), c)
	Boys and girls aged 15-17 years	7-day record	2005-2006	(IUNA (Irish Universities Nutrition Alliance), c)
<b>Italy</b>	Boys and girls 0-<3 years	3-day record	2005-2006	(Sette et al., 2010)
	Boys and girls 3-<10 years	3-day record	2005-2006	(Sette et al., 2010)
	Boys and girls 10-<18 years	3-day record	2005-2006	(Sette et al., 2010)
<b>Latvia</b>	Boys and girls aged 7-16 years	Two non-consecutive 24-hour dietary recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
<b>The Netherlands</b>	Infants aged 9 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Infants aged 12 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Children aged 18 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Boys and girls aged 2-3 years	2-day record (independent days)	2005-2006	(Ocké et al., 2008)
	Boys and girls aged 4-6 years	2-day record (independent days)	2005-2006	(Ocké et al., 2008)
	Boys and girls aged 7-8 years	Two non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Boys and girls aged 9-13 years	Two non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Boys and girls aged 14-18 years	Two non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Norway</b>	Children aged 2 years	Food frequency questionnaire	2007	(Kristiansen and Andersen, 2009)
	Boys and girls aged 4 years	4-day record	2000	(Elmadfa et al., 2009a)
	Boys and girls aged 9 years	4-day record	2000	(Elmadfa et al., 2009a)
	Boys and girls aged 13 years	4-day record	2000	(Elmadfa et al., 2009a)
	Boys and girls aged 16-19 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
<b>Poland</b>	Boys and girls aged 1-3 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 4-6 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 7-9 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 10-12 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 13-15 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 16-18 years	24-hour recall	2000	(Szponar et al., 2003)
<b>Portugal</b>	Boys and girls aged 5-10 years	Food frequency questionnaire	2006-2007	(Moreira et al., 2010). <i>Data collected in Porto.</i>
<b>Slovenia</b>	Boys and girls aged 14-16 years	Food frequency questionnaire	2003-2005	(Kobe et al., 2011)
<b>Spain</b>	Boys and girls aged 10-14 years	Two non-consecutive 24-hour recalls	2002-2003	(Elmadfa et al., 2009a). <i>Data collected in Catalonia.</i>
	Boys and girls aged 15-18 years	Two non-consecutive 24-hour recalls	2002-2003	(Elmadfa et al., 2009a). <i>Data collected in Catalonia.</i>
<b>Sweden</b>	Boys and girls aged 4 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
	Boys and girls aged 8-9 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
	Boys and girls aged 11-12 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
<b>United Kingdom</b>	Boys and girls aged 1.5-3 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Boys and girls aged 4-10 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Boys and girls aged 11-18 years	4-day food diary	2008-2010	(Bates et al., 2011)

mo, months

**APPENDIX 3B: ENERGY INTAKE OF CHILDREN AGED ~1-3 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Infants and/or young children (both sexes)</b>										
<b>Bulgaria</b>	1-3	154	5.9 <sup>1</sup>	3.0 <sup>1</sup>	5.4 <sup>1</sup>		1,401	705	1,299	
<b>Germany</b>	12 mo	432 <sup>2,3</sup>	0.35 <sup>4</sup>	0.06 <sup>4</sup>						
	18 mo	478 <sup>2</sup>	0.33 <sup>4</sup>	0.06 <sup>4</sup>						
	2	458 <sup>2</sup>	0.32 <sup>4</sup>	0.05 <sup>4</sup>						
	3	427 <sup>2</sup>	0.31 <sup>4</sup>	0.05 <sup>4</sup>						
<b>Ireland</b>	1	126	4.2	0.9	4.2		1,005	222	1,000	
	2	124	4.7	1.2	4.6		1,122	281	1,105	
	3	126	4.8	0.9	4.8		1,148	213	1,144	
<b>Italy</b>	0-<3	52	4.7	1.8	4.4	1.9-8.0	1,113	419	1,057	457-1,905
<b>The Netherlands</b>	9 mo	333	4.1	0.7	4.0	3.2-5.0 <sup>5</sup>				
	12 mo	306	4.5	0.7	4.4	3.7-5.4 <sup>5</sup>				
	18 mo	302	4.9	0.8	4.8	4.0-5.9 <sup>5</sup>				
<b>United Kingdom</b>	1.5-3	219	4.8	1.2	4.7	2.7-7.0 <sup>6</sup>	1,127	280	1,113	649-1,678 <sup>6</sup>
<b>Young children</b>										
<b>Boys</b>										
<b>Belgium</b>	2.5-3	102	6.5	1.1	6.5					
<b>Denmark</b>	1-3	129	6.9							
<b>Finland</b>	1 <sup>3</sup>	257	3.9	0.7			938	158		
	2	112	4.6	1.0			1,107	234		
	3	236	5.4	1.0			1,279	236		
<b>Greece</b>	12-24 mo	100	5.4	0.9			1,277	211		
	25-36 mo	274	5.8	1.0			1,395	228		
	37-48 mo	488	6.0	1.0			1,442	237		
<b>The Netherlands</b>	2-3	327	5.8		5.7	4.2-7.5	1,375		1,363	1,000-1,792
<b>Norway</b>	2	829	5.9	1.5						
<b>Poland</b>	1-3	70	5.9	2.2	5.5		1,407	524	1,318	
<b>Girls</b>										
<b>Belgium</b>	2.5-3	95	5.8	0.9	5.7					
<b>Denmark</b>	1-3	149	6.4							
<b>Finland</b>	1 <sup>3</sup>	198	3.6	0.6			863	132		
	2	118	4.5	0.9			1,077	213		
	3	235	5.0	1.0			1,211	234		
<b>Greece</b>	12-24 mo	107	5.2	0.8			1,247	179		
	25-36 mo	226	5.6	0.9			1,338	219		
	37-48 mo	434	5.8	1.0			1,379	237		
<b>The Netherlands</b>	2-3	313	5.5		5.4	4.1-7.2	1,308		1,288	971-1708
<b>Norway</b>	2	826	5.5	1.5						
<b>Poland</b>	1-3	48	5.4	1.6	5.3		1,283	378	1,277	

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 3A, unless stated otherwise. mo, months.

<sup>1</sup>Calculated from values in kcal.

<sup>2</sup>Number of 3-day records.

<sup>3</sup>Breast-fed infants not included.

<sup>4</sup>MJ/kg body mass (mean body mass of boys and girls, at 12 mo (whether breast-fed or not): 10.1 and 9.3 kg; 18 mo: 11.8 and 11.0 kg; 2 years: 13.2 and 12.3 kg; 3 years: 15.6 and 14.7 kg, respectively). Underreporters excluded.

<sup>4</sup>P10-P90.

<sup>5</sup>P2.5-P97.5.



**APPENDIX 3C: ENERGY INTAKE OF CHILDREN AGED ~4-6 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Boys</b>										
Belgium	4-6.5	236	6.4	0.9	6.4					
Czech Republic	4-6	641	6.5	1.3						
Denmark	4-5	81	7.7	1.5	7.6	5.7-10.5				
Finland	4	307	5.8	1.1			1,388	258		
	6	364	6.7	1.2			1,599	278		
France	4-6	164	6.3	0.1						
Germany	6	106	7.2	1.4	7.3	4.8-9.8	1,712	332	1,738	1,145-2,341
Greece	49-60 mo	356	6.2	0.1			1,475	296		
The Netherlands	4-6	327	6.7		6.6	5.3-8.2	1,587		1,579	1,252-1,951
Norway	4	206	6.3	1.5						
Poland	4-6	82	7.9	2.4	7.5		1,890	562	1,800	
Sweden	4	302	6.5	1.2	6.5	4.5-8.8	1,556	298	1,546	1,086-2,097
United Kingdom	4-10	210	6.7	1.3	6.6	4.3-9.7 <sup>1</sup>	1,591	314	1,573	1,021-2,301 <sup>1</sup>
<b>Girls</b>										
Belgium	4-6.5	228	5.9	0.9	5.9					
Czech Republic	4-6	446	6.5	1.3						
Denmark	4-5	78	7.0	1.6	6.8	5.2-9.7				
Finland	4	247	5.5	1.0			1,302	233		
	6	349	6.0	1.1			1,431	256		
France	4-6	162	6.3	0.1						
Germany	6	102	6.3	1.3	6.2	3.8-8.7	1,511	320	1,471	912-2,071
Greece	49-60 mo	389	5.9	0.1			1,414	260		
The Netherlands	4-6	312	6.2		6.2	4.7-7.8	1,479		1,470	1,123-1,866
Norway	4	185	6.1	1.2						
Poland	4-6	84	7.1	2.4	7.0		1,698	582	1,663	
Sweden	4	288	6.1	1.2	6.1	4.2-7.9	1,454	289	1,450	1,000-1,895
United Kingdom	4-10	213	6.4	1.3	6.5	3.8-8.9 <sup>1</sup>	1,519	314	1,531	900-2,114 <sup>1</sup>
<b>Both sexes</b>										
Bulgaria	3-6	199	7.4 <sup>2</sup>	3.1 <sup>2</sup>	6.8 <sup>2</sup>		1,759	735	1,628	
Ireland	4	124	5.3	1.0	5.2		1,264	240	1,241	
Italy	3-<10	193	8.0	2.0	8.0	4.8-11.5	1,914	488	1,906	1,138-2,750

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 3A, unless stated otherwise.  
mo, months.

<sup>1</sup>P2.5-P97.5.

<sup>2</sup>Calculated from values in kcal.

**APPENDIX 3D: ENERGY INTAKE OF CHILDREN AGED ~7-9 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Boys</b>										
Austria	7-9	146	6.9	1.9						
Czech Republic	7-9	940	7.6	2.8						
Denmark	6-9	172	8.8	2.2	8.4	6.2-12.7				
France	7-9	160	7.6	0.2						
Germany	7-9	321	7.8	1.6	7.8	5.5-10.6	1,867	371	1,850	1,312-2,514
Ireland	5-8	145	6.8	1.5	6.8	4.6-9.6	1,625	359	1,608	1,106-2,287
Latvia	7-16	295	8.2 <sup>1</sup>				1,948			
The Netherlands	7-8	153			8.1	5.3-11.6			1,929	1,267-2,753
Norway	9	402	8.6	2.0						
Poland	7-9	101	9.1	2.9	9.1		2,184	695	2,167	
Portugal	5-10	985	9.7 <sup>1</sup>	2.7 <sup>1</sup>			2,327	647		
Sweden	8-9	444	8.1	1.8	8.0	5.5-11.2	1,927	423	1,901	1,311-2,682
<b>Girls</b>										
Austria	7-9	134	6.3	1.6						
Czech Republic	7-9	765	7.6	2.8						
Denmark	6-9	151	7.8	1.6	7.7	5.5-10.8				
France	7-9	144	6.9	0.2						
Germany	7-9	308	7.0	1.4	7.0	4.5-9.5	1,663	333	1,669	1,075-2,271
Ireland	5-8	151	6.4	1.2	6.2	4.6-8.4	1,517	278	1,467	1,105-1,985
Latvia	7-16	277	6.9 <sup>1</sup>				1,660			
The Netherlands	7-8	151			8.4	5.9-11.3			2,011	1,409-2,706
Norway	9	408	7.7	2.0						
Poland	7-9	103	8.0	2.5	7.8		1,921	592	1,843	
Portugal	5-10	991	9.1 <sup>1</sup>	2.5 <sup>1</sup>			2,177	593		
Sweden	8-9	445	7.2	1.5	7.1	4.8-9.6	1,719	360	1,699	1,139-2,301
<b>Both sexes</b>										
Bulgaria	6-10	235	9.5 <sup>1</sup>	3.8 <sup>1</sup>	9.1 <sup>1</sup>		2,277	900	2,179	

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 3A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

**APPENDIX 3E: ENERGY INTAKE OF CHILDREN AGED ~10-14 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Boys</b>										
Austria	10-14	248	7.0	2.0						
Belgium	13-15	74	10.6	2.1						
Bulgaria	10-14	167	11.1 <sup>1</sup>	4.5 <sup>1</sup>	10.2 <sup>1</sup>		2,659	1,071	2,450	
Denmark	10-13	164	9.3	2.5	9.3	5.9-12.7				
France	10-14	160	8.7	0.2						
Germany	10-11	199	8.0	1.8	7.6	5.4-11.3	1,908	436	1,813	1,297-2,682
	12	114	10.6	3.2	10.4	6.1-18.1	2,522	769	2,470	1,455-4,316
	13-14	214	11.7	3.8	11.4	6.3-18.4	2,803	917	2,726	1,503-4,383
Hungary	11-14	124	10.4	1.9			2,489	453		
Ireland	9-12	148	8.0	1.6	7.9	5.8-10.5	1,890	369	1,871	1,383-2,495
	13-14	95	9.0	2.1	8.9	5.8-12.9	2,137	502	2,103	1,398-3,073
Italy	10-<18	108	10.8	3.1	10.6	6.8-15.5	2,576	744	2,540	1,630-3,709
The Netherlands	9-13	351			9.8	6.6-13.7			2,330	1,576-3,253
Norway	13	590	9.5	3.5						
Poland	10-12	128	10.3	3.4	10.1		2,468	821	2,414	
	13-15	218	13.2	4.6	12.7		3,145	1,092	3,027	
Spain	10-14	66	9.8	1.7						
Sweden	11-12	517	7.8	2.2	7.6	4.5-11.8	1,864	518	1,814	1,075-2,811
United Kingdom	11-18	238	8.5	2.1	8.1	4.5-12.7 <sup>2</sup>	2,007	508	1,916	1,074-3,019 <sup>2</sup>
<b>Girls</b>										
Austria	10-14	239	6.1	1.7						
Belgium	13-15	89	8.0	2.0						
Bulgaria	10-14	180	9.3 <sup>1</sup>	3.7 <sup>1</sup>	9.0 <sup>1</sup>		2,225	881	2,143	
Denmark	10-13	196	7.9	2.3	7.8	4.5-11.3				
France	10-14	144	7.5	0.1						
Germany	10-11	198	7.6	1.6	7.7	5.2-10.3	1,808	394	1,842	1,234-2,444
	12	103	9.3	3.2	8.3	4.2-14.7	2,222	763	1,986	1,007-3,508
	13-14	230	9.5	2.7	9.3	5.6-14.1	2,277	651	2,224	1,332-3,352
Hungary	11-14	111	9.2	1.5			2,195	358		
Ireland	9-12	150	7.0	1.4	6.9	4.6-9.4	1,654	333	1,649	1,089-2,227
	13-14	93	7.0	1.6	7.0	4.3-9.9	1,674	377	1,667	1,009-2,356
Italy	10-<18	139	8.7	2.2	8.7	5.0-12.5	2,091	532	2,081	1,187-2,999
The Netherlands	9-13	352			8.4	5.9-11.3			2,010	1,408-2,705
Norway	13	515	8.1	2.6						
Poland	10-12	121	8.9	2.7	8.8		2,124	646	2,098	
	13-15	134	10.0	3.7	9.7		2,385	882	2,308	
Spain	10-14	53	8.4	0.9						
Sweden	11-12	499	6.9	1.9	6.7	4.0-10.1	1,650	453	1,613	958-2,410
United Kingdom	11-18	215	6.9	1.7	6.9	3.6-10.3 <sup>2</sup>	1,637	413	1,637	850-2,437 <sup>2</sup>

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 3A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

<sup>2</sup>P2.5-97.5.

**APPENDIX 3F: ENERGY INTAKE OF ADOLESCENTS AGED ~15-18 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Boys</b>										
Austria	14-19	1,527	11.5	3.0						
Belgium	15-18	405	11.0	2.6	10.8		2,639	631	2,592	
Bulgaria	14-18	178	11.9 <sup>1</sup>	4.1	11.1 <sup>1</sup>		2,842	974	2,657	
Denmark	14-17	101	10.1	3.2	10.5	5.0-14.9				
France	15-18	181	10.2	0.4						
Germany	15-17	294	14.3	5.4	13.4	8.0-23.0	3,414	1,286	3,202	1,905-5,498
Ireland	15-17	129	9.9	2.5	9.7	6.1-14.6	2,344	595	2,314	1,459-3,473
The Netherlands	14-18	352			11.0	7.7-15.0			2,622	1,830-3,580
Norway	16-19	92	13.9							
Poland	16-18	130	14.7	4.8	14.1		3,504	1,130	3,380	
Slovenia	14-16	1,085	12.8				3,053			
Spain	15-18	61	10.7	2.0						
<b>Girls</b>										
Austria	14-19	1,422	8.5	2.2						
Belgium	15-18	401	7.7	1.6	7.6		1,844	373	1,817	
Bulgaria	14-18	190	9.0 <sup>1</sup>	3.4 <sup>1</sup>	8.3 <sup>1</sup>		2,149	824	1,994	
Denmark	14-17	134	7.4	2.3	7.1	4.3-11.2				
France	15-18	222	6.8	0.2						
Germany	15-17	317	9.9	3.8	9.3	5.4-16.2	2,364	916	2,228	1,284-3,853
Ireland	15-17	124	7.2	2.1	7.0	4.0-10.9	1,712	491	1,663	952-2,599
The Netherlands	14-18	354			8.4	5.9-11.3			2,008	1,406-2,703
Norway	16-19	86	9.1							
Poland	16-18	122	9.4	3.7	8.8		2,237	887	2,108	
Slovenia	14-16	1,346	9.8				2,332			
Spain	15-18	57	7.9	1.1						

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 3A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

**APPENDIX 4A: POPULATION, METHODS AND PERIOD OF DIETARY ASSESSMENT IN ADULTS IN EUROPEAN COUNTRIES**

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Austria</b>	Men and women aged 19-64 years	24-hour recall	2005-2006	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Men and women aged 65 years and over	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
<b>Belgium</b>	Men and women aged 19-59 years	Two non consecutive 24-hour recalls	2004-2005	(De Vriese et al., 2006)
	Men and women aged 60-74 years	Two non consecutive 24-hour recalls	2004-2005	(De Vriese et al., 2006)
	Men and women aged 75 years and over	Two non consecutive 24-hour recalls	2004-2005	(De Vriese et al., 2006)
<b>Bulgaria</b>	Men and women aged 18-30 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Men and women aged 30-60 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Men and women aged 60-75 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Men and women aged >75 years	24-hour recall	1998	(Abrasheva et al., 1998)
<b>Czech Republic</b>	Men and women aged 19-64 years	24-hour recall	2000-2001	(Cifková and Škodová, 2004; Elmadfa et al., 2009a)
<b>Denmark</b>	Men and women aged 18-75 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 18-24 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 25-34 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 35-44 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 45-54 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 55-64 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Men and women aged 65-75 years	7-day record	2003-2008	(Pedersen et al., 2010)
<b>Estonia</b>	Men and women aged 19-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Men and women aged 19-34 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Men and women aged 35-49 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Men and women aged 50-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
<b>Finland</b>	Men and women aged 25-64 years	48-hour recall	2007	(Paturi et al., 2008; Pietinen et al., 2010)
	Men and women aged 25-34 years	48-hour recall	2007	(Paturi et al., 2008)
	Men and women aged 35-44 years	48-hour recall	2007	(Paturi et al., 2008)
	Men and women aged 45-54 years	48-hour recall	2007	(Paturi et al., 2008)
	Men and women aged 55-64 years	48-hour recall	2007	(Paturi et al., 2008)
	Men and women aged 65-75 years	48-hour recall	2007	(Paturi et al., 2008)
<b>France</b>	Men and women aged 19-64 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)
	Men and women aged 65-74 years	Three 24-hour recalls	2006-2007	(Elmadfa et al., 2009a)
<b>Germany</b>	Men and women aged 19-80 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)
	Men and women aged 19-24 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)
	Men and women aged 25-34 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)
	Men and women aged 35-50 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)
	Men and women aged 51-64 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)
	Men and women aged 65-80 years	24-hour recall + Dietary history	2005-2006	(MRI, 2008b)



Country	Population	Dietary assessment method	Year of survey	Reference
<b>Greece</b>	Men and women aged 19-64 years	Food frequency questionnaire + 24-hour recall in subgroup	1994-1999	(Elmadfa et al., 2009a)
	Men and women aged 65 years and over	Food frequency questionnaire	1994-1999	(Elmadfa et al., 2009a)
<b>Hungary</b>	Men and women aged 18-59 years	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Men and women aged 18-34 years	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Men and women aged 35-59 years	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Men and women aged 60 years and over	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
<b>Ireland</b>	Men and women aged 18-64 years	4-day record	2008-2010	(IUNA, 2011)
	Men and women aged 18-35 years	4-day record	2008-2010	(IUNA, 2011)
	Men and women aged 36-50 years	4-day record	2008-2010	(IUNA, 2011)
	Men and women aged 51-64 years	4-day record	2008-2010	(IUNA, 2011)
	Men and women aged 65-90 years	4-day record	2008-2010	(IUNA, 2011)
<b>Italy</b>	Men and women aged 18-<65years	3-day record	2005-2006	(Sette et al., 2010)
	Men and women aged 65 and over	3-day record	2005-2006	(Sette et al., 2010)
<b>Latvia</b>	Men and women aged 17-26 years	Two non-consecutive 24-hour recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
	Men and women aged 27-36 years	Two non-consecutive 24-hour recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
	Men and women aged 37-46 years	Two non-consecutive 24-hour recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
	Men and women aged 47-56 years	Two non-consecutive 24-hour recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
	Men and women aged 57-64 years	Two non-consecutive 24-hour recalls + food frequency questionnaire	2008	(Joffe et al., 2009)
<b>Lithuania</b>	Men and women aged 19-64 years	24-hour recall	2007	(Elmadfa et al., 2009a)
<b>The Netherlands</b>	Men and women aged 19-30 years	Two non-consecutive 24-hour recalls	2007-2010	(van Rossum et al., 2011)
	Men and women aged 31-50 years	Two non-consecutive 24-hour recalls	2007-2010	(van Rossum et al., 2011)
	Men and women aged 51-69 years	Two non-consecutive 24-hour recalls	2007-2010	(van Rossum et al., 2011)

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Norway</b>	Men and women aged 18-70 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
	Men and women aged 18-29 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
	Men and women aged 30-39 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
	Men and women aged 40-49 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
	Men and women aged 50-59 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
	Men and women aged 60-70 years	Two randomly distributed 24-hour dietary recalls + food propensity questionnaire	2010-2011	(Holm Totland et al., 2012)
<b>Poland</b>	Men and women aged 19-25 years	24-hour recall	2000	(Szponar et al., 2003)
	Men and women aged 26-60 years	24-hour recall	2000	(Szponar et al., 2003)
	Men and women aged 61 years and over	24-hour recall	2000	(Szponar et al., 2003)
<b>Portugal</b>	Men and women aged 18-≥65 years	Food frequency questionnaire	1999-2003	(Elmadfa et al., 2009a; Lopes et al., 2006). <i>Data collected in Porto.</i>
	Men and women aged 18-39 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006). <i>Data collected in Porto.</i>
	Men and women aged 40-49 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006). <i>Data collected in Porto.</i>
	Men and women aged 50-64 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006). <i>Data collected in Porto.</i>
	Men and women aged 65 years and over	Food frequency questionnaire	1999-2003	(Elmadfa et al., 2009a; Lopes et al., 2006). <i>Data collected in Porto.</i>
<b>Romania</b>	Men and women aged 19-64 years	Personal interview	2006	(Elmadfa et al., 2009a)
	Men and women aged 65 years and over	Personal interview	2006	(Elmadfa et al., 2009a)
<b>Slovenia</b>	Men and women aged 18-65 years	Food frequency questionnaire and 24-hour recall	2007-2008	(Gabrijelčič Blenkuš et al., 2009)
<b>Spain</b>	Men and women aged 18-24 years	Two non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Men and women aged 25-44 years	Two non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Men and women aged 45-64 years	Two non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Men and women aged 65-75 years	Two non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
<b>Sweden</b>	Men and women aged 18-80 years	4-day record	2010-2011	(Amcoff et al., 2012)
	Men and women aged 18-30 years	4-day record	2010-2011	(Amcoff et al., 2012)
	Men and women aged 31-44 years	4-day record	2010-2011	(Amcoff et al., 2012)
	Men and women aged 45-64 years	4-day record	2010-2011	(Amcoff et al., 2012)
	Men and women aged 65-80 years	4-day record	2010-2011	(Amcoff et al., 2012)
<b>United Kingdom</b>	Men and women aged 19-64 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Men and women aged 65 years and over	4-day food diary	2008-2010	(Bates et al., 2011)

**APPENDIX 4B: ENERGY INTAKE OF ADULTS AGED ~19-65 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Men</b>										
Austria	19-64	778	9.0	3.1						
Belgium	19-59	413	10.8	3.0	10.4		2,578	720	2,495	
Czech Republic	19-64	1,046	12.4	3.7						
Denmark	18-75	1,569	10.4	2.9	10.3	7.0-14.3 <sup>1</sup>				
Estonia	19-64	900	9.6	4.8			2,278	1,144		
Finland	25-64	730	9.2	3.0			2,206	705		
France	19-64	852	10.0	0.1						
Germany	19-64	4,912	11.0	4.3						
Greece	19-64	8,365	10.4	3.0						
Hungary	18->60	473	11.7	2.4			2,792	570		
Ireland	18-64	634	10.1	2.7	10.0		2,397	650	2,377	
Italy	18-<65	1,068	10.0	2.7	9.8	6.2-14.6	2,390	650	2,332	1,471-3,499
Lithuania	19-65	849	10.3	4.3						
Norway	18-70	862	10.9	3.4	10.5	NA-17.3				
Portugal	18-≥65	917	9.9	2.3			2,367	560	2,300	1,551-3,369
Romania	19-64	177	13.9	5.2						
Slovenia	18-65	NA	13.1 <sup>2</sup> 9.0 <sup>3</sup>							
Sweden	18-80	792	9.4	2.8	9.3	5.1-13.7				
United Kingdom	19-64	346	9.2	3.0	8.9	4.7-17.1 <sup>4</sup>	2,200	706	2,112	1,115-4,058 <sup>4</sup>
<b>Women</b>										
Austria	19-64	1,345	7.5	2.5						
Belgium	19-59	460	7.0	1.9	6.8		1,680	447	1,637	
Czech Republic	19-64	1,094	9.7	3.0						
Denmark	18-75	1,785	7.9	2.1	7.9	5.4-10.5 <sup>1</sup>				
Estonia	19-64	1,115	6.9	3.2			1,640	766		
Finland	25-64	846	6.8	2.0			1,620	483		
France	19-64	1,499	7.2	0.1						
Germany	19-64	6,016	8.1	2.5						
Greece	19-64	12,034	8.3	2.4						
Hungary	18->60	706	9.2	1.8			2,205	429		
Ireland	18-64	640	7.2	2.0	7.2		1,725	482	1,706	
Italy	18-<65	1,245	8.1	2.2	8.0	4.9-11.8	1,939	526	1,909	1,162-2,827
Lithuania	18-65	1,087	7.4	3.0						
Norway	18-70	925	8.0	2.4	7.8	NA-12.0				
Portugal	18-≥65	1,472	8.7	2.1			2,079	494	2,040	1,352-2,953
Romania	19-64	341	11.4	4.9						
Slovenia	18-65	NA	11.3 <sup>2</sup> 7.5 <sup>3</sup>							
Sweden	18-80	1,005	7.4	2.1	7.3	4.3-11.0				
United Kingdom	19-64	461	6.9	2.0	6.7	3.1-11.4 <sup>4</sup>	1,638	477	1,604	747-2,700 <sup>4</sup>

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 4A, unless stated otherwise.

<sup>1</sup>P10-P90.

<sup>2</sup>Food frequency questionnaire.

<sup>3</sup>24-hour recall.

<sup>4</sup>P2.5-P97.5.

NA, not available.

**APPENDIX 4C: ENERGY INTAKE OF ADULTS AGED ~19-34 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Men</b>										
<b>Bulgaria</b>	18-30	208	11.8 <sup>1</sup>	5.1 <sup>1</sup>	10.8 <sup>1</sup>		2,820	1,231	2,590	
<b>Denmark</b>	18-24	105	11.1	3.1	10.8	6.4-15.8				
	25-34	234	11.3	2.9	11.0	7.0-16.1				
<b>Estonia</b>	19-34	396	10.3 <sup>1</sup>	5.3 <sup>1</sup>			2,464	1,255		
<b>Finland</b>	25-34	137	9.9	3.2			2,362	764		
<b>Germany</b>	19-24	510	12.0 <sup>1</sup>	0.20 <sup>1,2</sup>	11.2 <sup>1</sup>	6.1-21.0 <sup>1</sup>	2,872	48.12 <sup>2</sup>	2,680	1,452-5,023
	25-34	690	11.6 <sup>1</sup>	0.17 <sup>1,2</sup>	10.8 <sup>1</sup>	6.3-19.6 <sup>1</sup>	2,783	41.36 <sup>2</sup>	2,581	1,505-4,692
<b>Hungary</b>	18-34	136	12.4	2.3			2,965	551		
<b>Ireland</b>	18-35	276	10.7	2.9	10.6		2,553	664	2,540	
<b>Latvia</b>	17-26	191	10.0 <sup>1</sup>				2,394			
	27-36	116	10.0 <sup>1</sup>				2,393			
<b>The Netherlands</b>	19-30	356			11.5	8.1-15.6			2,573	1,940-3,731
<b>Norway</b>	18-29	138	12.8	4.0						
	30-39	136	11.5	3.5						
<b>Poland</b>	19-25	191	15.3	4.6	15.1		3,657	1,090	3,613	
<b>Portugal</b>	18-39	179					2,496	584	2,427	1,622-3,577
<b>Spain</b>	18-24	127	10.0 <sup>1</sup>				2,384			
	25-44	326	9.4 <sup>1</sup>				2,242			
<b>Sweden</b>	18-30	132	9.4	3.5	9.5	4.7-13.7	2,246	830	2,259	1,122-3,283
<b>Women</b>										
<b>Bulgaria</b>	18-30	204	8.2 <sup>1</sup>	3.2 <sup>1</sup>	7.5 <sup>1</sup>		1,954	758	1,789	
<b>Denmark</b>	18-24	150	8.2	2.3	8.1	4.9-12.2				
	25-34	340	8.3	2.2	8.3	4.9-11.8				
<b>Estonia</b>	19-34	459	7.4 <sup>1</sup>	3.4 <sup>1</sup>			1,760	801		
<b>Finland</b>	25-34	180	7.2	2.2			1,711	525		
<b>Germany</b>	19-24	510	8.4 <sup>1</sup>	0.13 <sup>1,2</sup>	8.0 <sup>1</sup>	4.8-13.3 <sup>1</sup>	1,996	30.69 <sup>2</sup>	1,914	1,141-3,171
	25-34	972	8.5 <sup>1</sup>	0.09 <sup>1,2</sup>	8.0 <sup>1</sup>	4.9-13.2 <sup>1</sup>	2,031	21.11 <sup>2</sup>	1,929	1,165-3,151
<b>Hungary</b>	18-34	176	9.5	1.7			2,280	407		
<b>Ireland</b>	18-35	255	7.5	2.3	7.4		1,783	542	1,762	
<b>Latvia</b>	17-26	187	7.1 <sup>1</sup>				1,690			
	27-36	90	6.4 <sup>1</sup>				1,523			
<b>The Netherlands</b>	19-30	347			8.4	5.9-11.3			1,999	1,399-2,693
<b>Norway</b>	18-29	143	8.1	2.5						
	30-39	169	8.4	2.4						
<b>Poland</b>	19-25	211	8.2	3.2	7.8		1,957	763	1,872	
<b>Portugal</b>	18-39	299					2,141	515	2,096	1,409-3,109
<b>Spain</b>	18-24	182	7.8 <sup>1</sup>				1,869			
	25-44	376	7.2 <sup>1</sup>				1,714			
<b>Sweden</b>	18-30	202	7.6	2.3	7.7	3.9-11.1	1,819	538	1,836	926-2,660

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 4A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

<sup>2</sup>SE.

**APPENDIX 4D: ENERGY INTAKE OF ADULTS AGED ~35-64 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Men</b>										
<b>Bulgaria</b>	30-60	224	11.7 <sup>1</sup>	3.8 <sup>1</sup>	11.5 <sup>1</sup>		2,788	904	2,747	
<b>Denmark</b>	35-44	318	11.1	3.1	10.9	6.2-16.6				
	45-54	336	10.3	2.7	10.3	6.4-14.8				
	55-64	336	9.9	2.7	9.6	5.7-14.8				
<b>Estonia</b>	35-49	319	9.2 <sup>1</sup>	4.6 <sup>1</sup>			2,190	1,103		
	50-64	185	8.5 <sup>1</sup>	3.7 <sup>1</sup>			2,033	873		
<b>Finland</b>	35-44	177	9.5	3.2			2,277	806		
	45-54	190	9.2	3.4			2,202	603		
	55-64	226	8.6	2.5			2,061	636		
<b>Germany</b>	35-50	2,079	11.0 <sup>1</sup>	0.08 <sup>1,2</sup>	10.5 <sup>1</sup>	6.0 <sup>1</sup> -17.9 <sup>1</sup>	2,640	19.29 <sup>2</sup>	2,509	1,435-4,271
	51-64	1,633	10.0 <sup>1</sup>	0.08 <sup>1,2</sup>	9.6 <sup>1</sup>	5.4 <sup>1</sup> -16.1 <sup>1</sup>	2,400	19.60 <sup>2</sup>	2,297	1,301-3,843
<b>Hungary</b>	35-59	199	12.0	2.2			2,862	533		
<b>Ireland</b>	36-50	205	9.7	2.5	9.6		2,322	591	2,310	
	51-64	153	9.3	2.4	9.1		2,217	581	2,157	
<b>Latvia</b>	37-46	136	9.7 <sup>1</sup>				2,319			
	47-56	155	9.3 <sup>1</sup>				2,230			
	57-64	108	8.9 <sup>1</sup>				2,121			
<b>The Netherlands</b>	31-50	348			11.1	7.7-15.1			2,647	1,848-3,611
	51-69	351			10.0	6.9-13.9			2,390	1,637-3,309
<b>Norway</b>	40-49	179	10.6	3.1						
	50-59	192	10.4	3.1						
<b>Poland</b>	26-60	865	13.0	4.4	12.6		3,114	1,056	3,019	
<b>Portugal</b>	40-49	197	10.3 <sup>1</sup>				2,453	530	2,406	1,679-3,372
	50-64	295	9.9 <sup>1</sup>				2,354	561	2,300	1,591-3,271
<b>Spain</b>	45-64	265	8.4 <sup>1</sup>				2,018			
<b>Sweden</b>	31-44	183	9.8	2.4	9.9	6.2-13.6	2,343	573	2,362	1,480-3,246
	45-64	308	9.4	2.8	9.4	5.1-14.2	2,254	674	2,252	1,209-3,403
<b>Women</b>										
<b>Bulgaria</b>	30-60	224	8.2 <sup>1</sup>	3.0 <sup>1</sup>	7.9 <sup>1</sup>		1,956	724	1,891	
<b>Denmark</b>	35-44	412	8.3	2.2	8.2	4.9-12.1				
	45-54	359	7.6	1.9	7.8	4.6-10.5				
	55-64	326	7.5	1.9	7.3	4.9-10.6				
<b>Estonia</b>	35-49	376	6.7 <sup>1</sup>	3.2 <sup>1</sup>			1,605	765		
	50-64	280	6.2 <sup>1</sup>	2.8 <sup>1</sup>			1,491	676		
<b>Finland</b>	35-44	211	7.1	2.1			1,687	497		
	45-54	232	6.7	1.9			1,601	461		
	55-64	223	6.3	1.8			1,502	433		
<b>Germany</b>	35-50	2,694	8.2 <sup>1</sup>	0.05 <sup>1,2</sup>	7.8 <sup>1</sup>	4.6 <sup>1</sup> -12.8 <sup>1</sup>	1,948	11.74 <sup>2</sup>	1,870	1,098-3,066
	51-64	1,840	7.8 <sup>1</sup>	0.05 <sup>1,2</sup>	7.5 <sup>1</sup>	4.6 <sup>1</sup> -11.9 <sup>1</sup>	1,856	13.10 <sup>2</sup>	1,793	1,092-2,837
<b>Hungary</b>	35-59	295	9.4	1.9			2,237	443		
<b>Ireland</b>	36-50	232	7.1	1.9	7.1		1,696	444	1,684	
	51-64	153	7.0	1.7	7.1		1,674	416	1,682	
<b>Latvia</b>	37-46	136	6.5 <sup>1</sup>				1,562			
	47-56	149	6.7 <sup>1</sup>				1,608			
	57-64	109	6.4 <sup>1</sup>				1,530			
<b>The Netherlands</b>	31-50	351			8.2	5.7-11.1			1,956	1,361-2,644
	51-69	353			7.8	5.3-10.6			1,849	1,268-2,525
<b>Norway</b>	40-49	256	8.1	2.4						
	50-59	193	7.9	2.3						
<b>Poland</b>	26-60	1,997	8.4	3.0	8.1		1,997	721	1,927	
<b>Portugal</b>	40-49	340	9.0 <sup>1</sup>				2,160	478	2,127	1,488-2,959
	50-64	494	8.8 <sup>1</sup>				2,102	498	2,065	1,382-3,012



Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)				
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95	
Spain	45-64	337	6.6 <sup>1</sup>				1,573				
Sweden	31-44	247	7.6	2.2	7.6	4.3-11.3	1,820	517	1,813	1,027-2,709	
	45-64	358	7.3	2.1	7.2	4.3-10.7	1,755	510	1,711	1,039-2,553	

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 4A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

<sup>2</sup>SE.

**APPENDIX 4E: ENERGY INTAKE OF ADULTS AGED ~65 YEARS AND OVER IN EUROPEAN COUNTRIES**

Country	Age (years)	n	Energy (MJ/day)				Energy (kcal/day)			
			mean	SD	P50	P5 – P95	mean	SD	P50	P5 – P95
<b>Men</b>										
Austria	65+	147	7.7	2.4						
Belgium	60-74	416	9.1	2.2	8.9		2,172	525	2,129	
	75+	389	8.3	2.2	8.0		1,993	527	1,923	
Bulgaria	60-75	186	10.2 <sup>1</sup>	3.8 <sup>1</sup>	9.7 <sup>1</sup>		2,431	919	2,319	
	76+	101	9.0 <sup>1</sup>	3.2 <sup>1</sup>	8.6 <sup>1</sup>		2,153	757	2,064	
Denmark	65-75	240	9.5	2.5	9.2	5.8-13.7				
Finland	65-74	229	7.7	2.3			1,848	554		
France	65-74	130	9.0	0.2						
Germany	65-80	1,469	9.2 <sup>1</sup>	0.07 <sup>1,2</sup>	8.9 <sup>1</sup>	5.3-13.8 <sup>1</sup>	2,191	16.32 <sup>2</sup>	2,129	1,263-3,303
Greece	65+	2,508	8.5	2.5			2,018	600.6		
Hungary	60+	138	10.5	2.3			2,519	546		
Ireland	65+	106	8.3	2.6	8.0		1,983	630	1,905	
Italy	65+	202	9.6	2.3	9.5	6.2-13.6	2,296	556	2,267	1,471-3,241
Norway	60-70	217	9.9	2.9						
Poland	61+	226	10.6	3.6	10.4		2,524	860	2,493	
Portugal	65+	246	9.3	2.2			2,219	530	2,161	1,455-3,206
Romania	65+	177	13.0	4.1						
Spain	65-75	122	7.1 <sup>1</sup>				1,688			
Sweden	65-80	169	8.7	2.3	8.5	5.2-13.0	2,083	550	2,036	1,246-3,100
United Kingdom	65+	96	8.3	2.1	8.3	3.7-11.8 <sup>3</sup>	1,976	511	1,973	882-2,801 <sup>3</sup>
<b>Women</b>										
Austria	65+	202	7.1	1.7						
Belgium	60-74	406	6.7	1.6	6.5		1,597	387	1,564	
	75+	355	6.2	1.5	6.1		1,482	351	1,462	
Bulgaria	60-75	194	8.1 <sup>1</sup>	2.6 <sup>1</sup>	7.7 <sup>1</sup>		1,926	613	1,848	
	76+	113	7.6 <sup>1</sup>	2.7 <sup>1</sup>	7.6 <sup>1</sup>		1,807	636	1,814	
Denmark	65-75	198	7.4	1.9	7.3	4.5-10.7				
Finland	65-74	234	5.9	1.7			1,412	414		
France	65-74	219	6.7	0.1						
Germany	65-80	1,562	7.3 <sup>1</sup>	0.05 <sup>1,2</sup>	7.1 <sup>1</sup>	4.4-10.9 <sup>1</sup>	1,753	12.47 <sup>2</sup>	1,708	1,044-2,610
Greece	65+	3,600	6.8	2.1			1,620	491.7		
Hungary	60+	235	8.8	1.7			2,110	412		
Ireland	65+	120	6.5	1.6	6.3		1,555	382	1,508	
Italy	65+	316	7.7	2.0	7.6	4.6-11.4	1,834	486	1,828	1,094-2,732
Norway	60-70	164	7.4	2.2						
Poland	61+	365	8.3	2.8	8.0		1,974	658	1,917	
Portugal	65+	339	8.0	1.9			1,910	444	1,878	1,226-2,736
Romania	65+	341	10.9	3.4						
Spain	65-75	122	5.7 <sup>1</sup>				1,373			
Sweden	65-80	198	7.1	1.8	7.0	4.6-10.5	1,703	432	1,670	1,095-2,500
United Kingdom	65+	128	6.4	1.3	6.2	4.1-8.9 <sup>3</sup>	1,522	319	1,470	980-2,111 <sup>3</sup>

NB: Values in MJ and kcal are indicated as published in the reports listed in Appendix 4A, unless stated otherwise.

<sup>1</sup>Calculated from values in kcal.

<sup>2</sup>SE.

<sup>3</sup>P2.5-P97.5.

**APPENDIX 5: OVERVIEW OF THE APPROACHES USED BY SELECTED AUTHORITIES FOR THE ESTIMATION OF AVERAGE REQUIREMENTS (ARs) FOR ENERGY FOR ADULTS**

	REE equations	Age ranges for calculations	Body mass	PAL values	Source of PAL values	Method to estimate daily average energy requirements	Comments
<b>SCF (1993)</b>	Schofield's equations based on body mass, according to age and sex (FAO/WHO/UNU, 1985). Specific equations for adults $\geq 60$ y based on some of the original data collected by Schofield et al., data on Scottish men and data on Italian men and women (Ferro-Luzzi, 1987; James et al., 1989): Men: 60-74 y: REE (MJ/day)=0.0499 x body mass +2.93; $\geq 75$ y: REE (MJ/day)=0.035 x body mass + 3.43. Women: 60-74 y: REE (MJ/day)=0.0386 x body mass +2.88; $\geq 75$ y: REE (MJ/day)=0.0410 x body mass + 2.61	18-29 y, 30-59 y, 60-74 y, $\geq 75$ y	Observed European values taken from 11 studies (representative national samples and specific surveys), and weighted for the total number of adults in each age group in each country. Calculated body masses (from observed heights and a desirable BMI of 22).	Average PAL values varying between 1.33 and 2.10, including or without desirable physical activities, determined for each sex, according to age (18-59 y, 60-74 y, $\geq 75$ y) and for observed body masses.	(FAO/WHO/UNU, 1985; Ferro-Luzzi, 1987; James and Schofield, 1990)	PAL x REE	ARs: i) for each sex, age group, 5 kg increase in body mass (between 60 and 80 kg for men and 45 to 70 kg for women) and for each 0.1 increase in PAL varying between 1.4 and 2.2. ii) for each sex, on average as well as for each age group, for either the desirable or the actual median body mass, and for average PALs with or without desirable physical activity.
<b>AFSSA (2001)</b>	Black's equations (Black et al., 1996): Men (MJ/day): $0.963 \times \text{body mass}^{0.48} \times \text{height}^{0.50} \times \text{age}^{-0.13}$ Women (MJ/day): $1.083 \times \text{body mass}^{0.48} \times \text{height}^{0.50} \times \text{age}^{-0.13}$	20-30 y, 31-40 y, 41-50 y, 51-60 y, 61-70 y	Body mass (5 kg increase, from 55 to 100 kg for men, 45 to 90 kg for women) and height values for a BMI of 22 kg/m <sup>2</sup>	1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3	Several publications	PAL x REE	ARs for each sex, each age range, body mass and height for a BMI of 22 kg/m <sup>2</sup> , and each 0.1 increase in PAL value. To be corrected for BMI value (to decrease AR by 1 % for each 1 kg/m <sup>2</sup> exceeding the BMI of 22 kg/m <sup>2</sup> , and to increase by 1 % for each 1 kg/m <sup>2</sup> lower than the BMI of 22 kg/m <sup>2</sup> ). 1.5 and 1.8 times REE for active and healthy elderly subjects (Black, 1996; Cynober et al., 2000; Roberts, 1996). No conclusions for elderly subjects aged $\geq 80$ y

	REE equations	Age ranges for calculations	Body mass	PAL values	Source of PAL values	Method to estimate daily average energy requirements	Comments
<b>Health Council of the Netherlands (2001)</b>	Schofield's equations based on body mass, according to age and sex (FAO/WHO/UNU, 1985), modified for the older age groups (SCF, 1993).	19-30 y, 31-50 y, 51-70 y, ≥70 y	Calculated from observed Dutch height values (Hofman et al., 1995; Smit et al., 1994) and a BMI of 22.5 (18-50 y), 24 (51-70 y) and 25 (≥71 y) (Troiano et al., 1996; WHO, 1995b)	At the low average PAL in NL: 1.7 (19-50 y), 1.6 (51-70 y), and 1.5 (≥71 y). At the adequate PAL: 1.9 (19-50 y), 1.8 (51-70 y), and 1.7 (≥71 y).	DLW data (Black et al., 1996)	PAL x REE	ARs for each sex, age group, and for PAL values accounting either for the low average level of physical activity in the Netherlands or for an adequate level of physical activity.
<b>FAO/WHO/UNU (2004)</b>	Schofield's equations (1985)	18-30 y, 30-60 y, >60 y	Every 5 kg increase in body mass (between 50 and 90 kg for men, between 45 and 85 kg for women)	3 PAL ranges associated with a population's lifestyle: 1.40-1.69 (sedentary or light activity), 1.70-1.99 (active or moderately active), and 2.00-2.40 (vigorous or vigorously active) For calculations: 1.45, 1.60, 1.75, 1.90, 2.05 and 2.20	-	PAL x REE	Several sets of values, i.e. for men and women, for three age ranges, six PAL values, and for every 5 kg increase in body mass, and expressed as MJ/day or kcal/day, as well as per kg of body mass
<b>NNR (2004)</b>	Schofield's equations based on body mass, according to age and sex (FAO/WHO/UNU, 1985), modified for the older age groups (SCF, 1993).	18-30 y, 31-60 y, 61-74 y, ≥75 y	Mean corrected reference body masses for each sex and age range between 18 and 74 y, calculated based on mean population body masses in Denmark (1995), Sweden (1997-98) and Finland (Finrisk, 2000) (adjusted for individuals with a BMI ≠ 18.5-25.0). For the age range ≥75 y, AR was calculated using reference body masses for each sex by subtracting 1 kg from the body masses used for the age group 61-74 y	AR estimates proposed for three PAL values: 1.4 (sedentary), 1.6 (normal), 1.8 (active)	Studies using DLW measurements (Ainsworth et al., 2000; Black et al., 1996)	PAL x REE	ARs assuming normal body mass and energy balance, for each sex, each age range, and three PAL values

REE equations	Age ranges for calculations	Body mass	PAL values	Source of PAL values	Method to estimate daily average energy requirements	Comments
<b>IoM (2005)</b> /	≥19 y	Tables of heights and body masses of men and women, corresponding to BMIs of 18.5, 22.5 and 25 kg/m <sup>2</sup> . Reference body masses for a BMI of 22.5 kg/m <sup>2</sup> for men and a BMI of 21.5 kg/m <sup>2</sup> for women corresponding to the 50 <sup>th</sup> percentile among 19-y-old subjects (Kuczmarski et al., 2000) Calculations done for body masses for a BMI of 18.5 and for a BMI of 24.99, for each 0.05 m increase in height (varying between 1.45 and 1.95 m).	1.0-1.39 (sedentary), 1.4-1.59 (low active), 1.6-1.89 (active), 1.9-2.49 (very active).  PA coefficient for the equations = 1.00 (for sedentary men and women), 1.11 (for low active men), 1.12 (for low active women), 1.25 (for active men), 1.27 (for active women), 1.48 (for very active men), 1.45 (for very active women).	A dataset of adults with normal body mass using DLW measurements (IoM, 2005)	Men: AR [kcal/day] = 662 – (9.53 x age [y]) + PA x (15.91 x body mass [kg] + 539.6 x height [m]) (n=169, SE fit=284.5 kcal, R <sup>2</sup> =0.75)  Women: AR [kcal/day]=354 – (6.91 x age [y]) + PA x (9.36 x body mass [kg] + 726 x height [m]) (n=238, SE fit=231.6, R <sup>2</sup> =0.74)	ARs for 30-year-old men and women of various heights (between 1.45 and 1.95 m), with BMIs of 18.5 and 24.99 kg/m <sup>2</sup> and the corresponding body masses. For each year below 30: to add 7 kcal/day for women and 10 kcal/day for men. For each year above 30: to subtract 7 kcal/day for women and 10 kcal/day for men.
<b>SACN (2011)</b> Henry's equations (Henry, 2005) based on body mass and height, according to age and sex.	19-24 y, 25-34 y, 35-44 y, 45-54 y, 65-74 y, ≥75 y, all adults	Calculated from British height values (Health Survey for England, data for 2009) and a BMI of 22.5	Median PAL of 1.63 for both sexes and all ages. For older adults with reduced mobility or not in good health, the PAL value of 1.49 (=25 <sup>th</sup> centile) may be used. PAL values of 1.49 for “less active” and of 1.78 for “more active” correspond to 25 <sup>th</sup> and 75 <sup>th</sup> centile of PAL distribution. Examples are also given of the changes in PAL associated with increased activity.	Median, 25 <sup>th</sup> and 75 <sup>th</sup> percentiles from DLW studies (Moshfegh et al., 2008; Subar et al., 2003; Toozee et al., 2007) in US populations with similar levels of overweight and obesity and similar ethnic composition as the UK population. Exclusion of subjects with PAL <1.27 (n=38) and >2.5 (n=1).	PAL x REE	Energy requirements for men and women for seven age categories and all men / all women: population AR (median value), and energy requirements for less active and more active people. Values for each sex and each age group, at observed mean height-for-age values and body masses corresponding to a BMI of 22.5 kg/m <sup>2</sup> . Energy reference values for older adults with maintained general health and mobility: unlikely to differ from younger adults. For the extreme elderly, likely PAL of 1.49 (25 <sup>th</sup> centile) or lower (e.g. 1.38 observed in some otherwise healthy elderly subjects (Rothenberg et al., 2000)).



	REE equations	Age ranges for calculations	Body mass	PAL values	Source of PAL values	Method to estimate daily average energy requirements	Comments
<b>D-A-CH (2012)</b>	Based on Schofield's equations (FAO/WHO/UNU, 1985), according to each sex	19-<25 y 25-<51 y 51-<65 y ≥65 y	Calculated from German height values and a BMI of 22 (women), 24 (men)	<p>PAL values provided for different work or free time activities:</p> <p>1.2 (exclusively sedentary or bed-bound), 1.4-1.5 (exclusively seated work with little or no physical activity during leisure time), 1.6-1.7 (seated work, but occasionally also including work standing and moving around), 1.8-1.9 (Work including both standing and moving around), 2.0-2.4 (very strenuous work).</p> <p>PAL values used for calculations:</p> <p>- desired physical activity: 1.75 (15-24 y), 1.70 (25-50 y), 1.60 (51 y and older); low physical activity (1.45), high physical activity (2.2) - 1.4, 1.6, 1.8, 2.0</p>	(Black et al., 1996; SCF, 1993; Shetty et al., 1996)	PAL x REE	ARs for both sexes and for four age groups

**APPENDIX 6: OVERVIEW OF THE APPROACHES TO ESTIMATE AVERAGE REQUIREMENTS (ARs) FOR ENERGY FOR INFANTS AND YOUNG CHILDREN OF SELECTED COUNTRIES AND AUTHORITIES OTHER THAN FAO/WHO/UNU AND IOM**

	Age range	AR calculation	Body mass used for AR calculations	Comments
<b>SCF (1993)</b>	0-36 months	Adapted from British ARs for infants and young children (DoH, 1991)	Rounded British average body masses (except for age 1 month: US data (Hamill et al., 1977))	ARs only intended for formula-fed infants, at 1, 3, 6, 9, 12, 18, 24, 30 and 36 months and for each sex
<b>Health Council of the Netherlands (2001)</b>	Infants	24 h energy consumption (based on DLW data (Butte et al., 2000a; de Bruin et al., 1998)) + deposited energy for growth (calculated considering Dutch body masses, body's protein and fat percentages at the age-group boundaries)	Dutch reference body masses (Fredriks et al., 1998; Fredriks et al., 2000a; Fredriks et al., 2000b; TNO/LUMC, 1998)	ARs, in MJ/day per kg of body mass and in MJ per day, without distinction on sex, for 0-2, 3-5, 6-11 months
<b>AFSSA (2001)</b>	Infants	Energy expenditure per kg body mass x body mass values (according to sex) + deposited energy for growth (values differing for boys and girls and based on mean daily rates of protein and lipid deposition) (Butte, 1996)	Origin of body mass values not specified	ARs only intended for formula-fed infants, for each sex and for each one month increase in age. Values for the first two months corrected for the digestibility of feeding formulae
<b>NNR (2004)</b>	0-23 months	Energy expenditure (DLW data on healthy children, (Butte et al., 1996; Butte et al., 2000a; Tennefors et al., 2003)) + deposited energy for growth	Values based on the mean reference values from Denmark (Andersen et al., 1982), Norway (Knudtzon et al., 1988) and the Swedish (2000) and Finnish (1993) growth charts (values used only for the summary table for 0-17 y)	ARs valid for both breast-fed and formula-fed infants, per kg of body mass, at 1, 3, 6, 12, 18 months
<b>SACN (2011)</b>	Infants	FAO/WHO/UNU (2004) and Butte (2005) : TEE (equations as a function of body mass distinguishing breast-fed infants, formula-fed infants, and infants with mixed or unknown feeding (Butte, 2005)) + deposited energy for growth based on measured protein and fat gains (Butte et al., 2000b; Fomon et al., 1982) applied to UK body mass increments (UK-WHO Growth Standards)	UK-WHO Growth Standards (RCPCH, 2011)	ARs distinguishing breast-fed and formula-fed infants as well as infants with mixed or unknown feeding, for each sex and each one month increase in age
<b>D-A-CH (2012)</b>	Infants	Used the approach of Butte (1996)	Reference body masses based on median values for US infants	ARs for 0-<4 and 4-<12 months, for formula-fed infants

**APPENDIX 7: OVERVIEW OF THE APPROACHES OF FAO/WHO/UNU (2004) AND IOM (2005) TO ESTIMATE AVERAGE REQUIREMENTS (ARS) FOR ENERGY FOR INFANTS, CHILDREN AND ADOLESCENTS**

	FAO/WHO/UNU (2004)	IoM (2005)
<b>Age</b>	<b>Infants</b>	<b>0-36 months</b>
Method of calculation of TEE	Simple linear regression on body mass (kg). All infants (kcal/day): $-99.4 + 88.6 \times \text{body mass}$ (n=320, r=0.85, SEE=109 kcal/day). Breast-fed (kcal/day): $-152.0 + 92.8 \times \text{body mass}$ (n=195, r=0.87, SEE=108 kcal/day). Formula-fed (kcal/day): $-29.0 + 82.6 \times \text{body mass}$ (n=125, r=0.85, SEE=110 kcal/day).	Simple linear regression on body mass (kg). (kcal/day): $-100 + 89 \times \text{body mass}$
Source of data for the calculation	DLW data (Butte, 2001)	DLW data (IoM, 2005)
Body mass used for AR calculations	Median body mass-for-age (WHO (1994) pooled breast-fed dataset)	US reference body masses (Kuczmarski et al., 2000)
Calculation of energy deposition for growth	From gains in protein and fat and the corresponding energy deposition (Butte et al., 2000b), considering the median body mass gain according to age	From gains in protein and fat and the corresponding energy deposition (Butte et al., 2000b), considering the median body mass gain according to age (Guo et al., 1991)
AR calculation	ARs = TEE + energy deposition during growth	ARs = TEE + energy deposition during growth
Comments	ARs with or without distinction of sex, with or without distinction of breast-fed and formula-fed infants, and for each month of age	ARs for each sex and for each month of age
<b>Age</b>	<b>1-18 y</b>	<b>3-18 y</b>
Method of calculation of TEE	Quadratic equations with body mass as the single predictor  Boys (kcal/day): $310.2 + 63.3 \times \text{body mass} - 0.263 \times \text{body mass}^2$ (n=801, r=0.982, r <sup>2</sup> =0.964, SEE=124 kcal/day)  Girls (kcal/day): $263.4 + 65.3 \times \text{body mass} - 0.454 \times \text{body mass}^2$ (n=808, r=0.955, r <sup>2</sup> =0.913, SEE=155 kcal/day)  For children between 1 and 2 years, TEE estimates were reduced by 7% as the predicted values would have been otherwise 7% higher than the actual measurements of TEE (Butte, 2001)	Nonlinear regression analysis, with age, height and body mass, considering sex and 4 categories of physical activity coefficients (for sedentary, low active, active, very active subjects)  Boys (kcal/day) = $88.5 - (61.9 \times \text{age} + \text{PA} \times (26.7 \times \text{body mass} + 903 \times \text{height}))$ (SE fit=82.6, R <sup>2</sup> =0.98)  Girls (kcal/day) = $135.3 - (30.8 \times \text{age} + \text{PA} \times (10.0 \times \text{body mass} + 934 \times \text{height}))$ (SE fit=96.7, R <sup>2</sup> =0.95).
Source of data for the calculation	Derived from data on DLW and HR monitoring (Torun, 2001)	Derived from data on DLW (IoM, 2005)
Body mass used for AR calculations	Median body masses at the mid-point of each year (WHO reference values of body mass-for-age (1983))	US reference body masses and heights (Kuczmarski et al., 2000)
Calculation of energy deposition for growth	Mean daily body mass gain at each year of age (between 1-2 y and 17-18 y) (WHO, 1983) x average energy deposited in growing tissues (8.6 kJ/g of body mass gain, calculated considering estimated rates of protein and fat deposition) (Butte et al., 2000a; Butte, 2001)	Median daily rates of gain in body mass at each year of age (between 3.5 and 17.5 y) (Baumgartner et al., 1986) x energy deposited in growing tissues (calculated considering estimated rates of protein and fat deposition (Fomon et al., 1982; Haschke, 1989))
AR calculation	AR = TEE + energy deposition for growth	AR = TEE + energy deposition for growth
Comments	3 calculated sets of values: i) in absolute values, for each sex and for each one year increase, ii) per kg of body mass (AR divided by the median body mass at each year), for each sex and for each one year increase, iii) for 1-5 y, considering only moderate physical activity, and for children 6-18 y considering moderate physical activity (after calculation of "average" PAL values, by dividing TEE by calculated REE (Schofield et al., 1985)), light and heavy physical activity.	Calculated for each sex, for each one year increase in age, and the 4 PAL categories (defined as for adults: sedentary, low active, active, very active)

**APPENDIX 8: OVERVIEW OF THE APPROACHES TO ESTIMATE DAILY AVERAGE REQUIREMENTS (ARS) FOR ENERGY FOR CHILDREN AND ADOLESCENTS OF SELECTED COUNTRIES AND AUTHORITIES OTHER THAN FAO/WHO/UNU AND IOM**

	Method of calculation of REE	Body mass used for AR calculations	PAL values	Calculation of energy deposition for growth	AR calculation	Comments
<b>SCF (1993): 3-9 y, 10-18 y</b>	<b>3-9 y:</b> NA (use of intake data) <b>10-18 y:</b> REE equations (Schofield et al., 1985)	Average body masses from 9 European countries, weighted on the basis of each country's population at a given age	<b>3-9 y:</b> NA <b>10-18 y:</b> values for moderate physical activity, for 10-13 y (1.65 for boys, 1.55 for girls) and 14-18 y (1.58 for boys, 1.50 for girls) based on (FAO/WHO/UNU, 1985)	<b>3-9 y:</b> NA <b>10-18 y:</b> calculated based on a total energy cost of growth of 21 kJ per g of daily body mass gain, values between 0.03 and 0.35 MJ/day, differing for boys and girls	<b>3-9 y:</b> average body mass for boys and girls x energy intake per kg of BM, without the 5% increment proposed by FAO/WHO/UNU (1985) <b>10-18 y:</b> approach of FAO/WHO/UNU (1985): REE x PAL + deposited energy for growth	<b>3.5-9.5 y:</b> ARs, for each sex and one year increase in age between 3.5 and 9.5 y. <b>10.5-17.5 y:</b> ARs for each sex and one year increase between 10.5 and 17.5 y
<b>Health Council of the Netherlands (2001): 1-18 y</b>	REE equations ((Schofield et al., 1985) based on body mass)	Dutch reference body masses (Fredriks et al., 1998; Fredriks et al., 2000a; Fredriks et al., 2000b; TNO/LUMC, 1998)	DLW data (Torun et al., 1996) 1-3 y: 1.5; 4-8 y: 1.6; 9-13 y: 1.8; girls, 14-18 y: 1.7; boys, 14-18 y: 1.8	Accretion expenditure of growth calculated from Dutch BM, body's protein and fat percentages at the age group limits. Values between 0.05 and 0.13 MJ/day, differing for each sex.	AR = REE x average PAL + deposited energy for growth	ARs, for each sex and age range (1-3 y, 4-8 y, 9-13 y, 14-18 y) and PAL value.
<b>AFSSA (2001): 1-9 y, 10-18 y</b>	<b>1-9 y:</b> NA (use of energy expenditure from DLW data (Torun et al., 1996)) <b>10-18 y:</b> REE equations based on height and body mass (FAO/WHO/UNU, 1985)	Average body mass for age (origin of body mass values not specified)	<b>1-9 y:</b> 3 PALs varying with age: average, low, high (French DLW data, 1999). Average value varying with age: 1.5 for 2-3 y, 1.55 for 4 y, 1.6 for 5 y, 1.75 for 6-9 y. <b>10-18 y:</b> 9 PAL values for each 0.1 increase in PAL between 1.4 and 2.2	Average energy stored in tissues, considering deposited protein and fat and body mass gain, generally differing for boys and girls	<b>1-9 y, for average PAL:</b> BM x energy expenditure per kg body mass (based on DLW data) + deposited energy for growth (corrected values for low and high PALs) <b>10-18 y:</b> REE x PAL + deposited energy for growth	<b>1-9 y:</b> ARs, for each sex, each one year increase in age and each PAL. <b>10-18 y:</b> ARs, for each sex, BM (between 30 and 80 kg for boys, 30-70 kg for girls), each PAL and the average BMI of each age. ARs corrected only for girls 10-18 y according to BMI above or below the average value.

	Method of calculation of REE	Body mass used for AR calculations	PAL values	Calculation of energy deposition for growth	AR calculation	Comments
<b>NNR (2004): 2-5 y, 6-9 y, 10-17 y</b>	<b>2-5 y:</b> NA (use of energy expenditure from DLW data (Torun et al., 1996)) <b>6-9 y:</b> use of published values (Torun et al., 1996) calculated from REE (FAO/WHO/UNU, 1985) and a moderate PAL value. <b>10-17 y:</b> REE equations ((Schofield et al., 1985) based on body mass)	Values based on the mean reference values from Denmark (Andersen et al., 1982), Norway (Knudtzon et al., 1988) and the Swedish (2000) and Finnish (1993) growth charts	<b>2-5 y:</b> NA. <b>6-9 y:</b> moderate physical activity considering the evaluation of PAL values based on DLW, HR monitoring and activity-time allocation studies (Torun et al., 1996) <b>10-17 y:</b> 3 PAL categories (Torun et al., 1996). Light activity: for girls, 1.50 (10-13 y), 1.45 (14-17 y), for boys, 1.55 (10-13 y), 1.60 (14-17 y). Moderate activity: for girls, 1.70 (10-13 y), 1.65 (14-17 y), for boys, 1.75 (10-13 y), 1.80 (14-17 y). Heavy activity: for girls, 1.90 (10-13 y), 1.85 (14-17 y), for boys, 1.95 (10-13 y), 2.05 (14-17 y).	-	<b>2-5 y:</b> energy expenditure (DLW data on healthy children (Torun et al., 1996) + deposited energy for growth (2%)) <b>6-9 y:</b> use of published values (Torun et al., 1996) <b>10-17 y:</b> REE x PAL	<b>2-9 y:</b> ARs per kg of BM, for each sex and each one year increase. <b>10-17 y:</b> ARs per kg of BM, for each sex, each one year increase, and each PAL category (light, moderate, heavy). <b>0-17 y:</b> ARs in MJ/day, for each sex, for 0-1, 3, 6, 12 months, then one year increase, considering average Nordic BM for age, moderate physical activity and the AR per kg of body mass previously calculated
<b>SACN (2011): 1-18 y</b>	Henry's equations (Henry, 2005) based on body mass and height	<b>1-4 y:</b> median body masses and heights indicated by the growth standards (RCPCH, 2011) <b>5-18 y:</b> median British body masses and heights (UK 1990 references) (Freeman et al., 1995)	Median, 25 <sup>th</sup> and 75 <sup>th</sup> percentiles of PAL values adjusted for growth (in terms of a 1 % increase, compilation of published DLW data) without distinction of sex: for age ranges 1-3 y (1.36, 1.40, 1.45), 4-9 y (1.43, 1.58, 1.70), 10-18 y (1.68, 1.75, 1.86).	Adjustments of PAL values for growth in terms of 1 % increase.	REE x PAL	Energy requirements for each sex and each one year increase in age: population AR (calculated with median PAL value) and energy requirements for less active (25 <sup>th</sup> percentile of PALs), and more active (75 <sup>th</sup> percentile of PALs) subjects
<b>D-A-CH (2012): 1-18 y</b>	<b>1-&lt;15 y:</b> NA (use of energy expenditure from DLW data (Torun et al., 1996)) <b>15-&lt;19 y:</b> REE equations (FAO/WHO/UNU, 1985) based on body mass	<b>1-&lt;15 y:</b> Reference body masses based on median values for US children <b>15-&lt;19 y:</b> German data, calculating body masses from body heights and a BMI of 22 kg/m <sup>2</sup> for men and 21 kg/m <sup>2</sup> for women	<b>1-&lt;15 y:</b> moderate physical activity considering the evaluation of PAL values based on DLW, HR monitoring and activity-time allocation studies (Torun et al., 1996) <b>15-&lt;19 y:</b> moderate physical activity: 1.75; PALs of 1.4, 1.6, 1.8, 2.0 also used	<b>1-&lt;15 y:</b> NA <b>15-&lt;19 y:</b> NA	<b>1-&lt;15 y:</b> Used the approach of Torun et al. (1996) <b>15-&lt;19 y:</b> REE x PAL	ARs, for each sex and age range: 1-<4 y, 4-<7 y, 7-<10 y, 10-<13 y, 13-<15 y, 15-<19 y

y, years; REE, resting energy expenditure; PAL, physical activity level; NA, not applicable.

**APPENDIX 9: REE CALCULATED WITH FIVE MOST USED PREDICTIVE EQUATIONS USING MEASURED HEIGHTS FROM SURVEYS IN 13 EU MEMBER STATES AND BODY MASSES TO YIELD A BMI OF 22**

Age (years)	n	REE (MJ/day) estimated with Mifflin et al. 1990 Median (P5-P95)	REE (MJ/day) estimated with Henry 2005 Median (P5-P95)	REE (MJ/day) estimated with Müller et al. 2004 Median (P5-P95)	REE (MJ/day) estimated with Harris-Benedict (1919) Median (P5-P95)	REE (MJ/day) estimated with Schofield et al. 1985 Median (P5-P95)	Lowest and highest median REE (MJ/day)	REE (kcal/day) estimated with Mifflin et al. 1990 Median (P5-P95)	REE (kcal/day) estimated with Henry 2005 Median (P5-P95)	REE (kcal/day) estimated with Müller et al. 2004 Median (P5-P95)	REE (kcal/day) estimated with Harris-Benedict (1919) Median (P5-P95)	REE (kcal/day) estimated with Schofield et al. 1985 Median (P5-P95)	Lowest and highest median REE (kcal/day)
<b>Men</b>													
18 - 29	2,771	7.1 (6.4-7.8)	7.0 (6.3-7.7)	7.2 (6.7-7.6)	7.4 (6.6-8.2)	7.3 (6.7-7.8)	7.0; 7.4	1,702 (1,534-1,868)	1,674 (1,506-1,836)	1,712 (1,609-1,818)	1,761 (1,573-1,943)	1,737 (1,602-1,870)	1,674; 1,761
30 - 39	2,971	6.8 (6.2-7.6)	6.8 (6.1-7.5)	7.0 (6.6-7.5)	7.0 (6.3-7.9)	7.0 (6.6-7.5)	6.8; 7.0	1,637 (1,487-1,823)	1,621 (1,466-1,796)	1,666 (1,575-1,784)	1,675 (1,506-1,879)	1,672 (1,577-1,783)	1,621; 1,675
40 - 49	3,780	6.6 (6.0-7.3)	6.7 (6.0-7.4)	6.8 (6.4-7.3)	6.7 (5.9-7.5)	6.9 (6.5-7.4)	6.6; 6.9	1,574 (1,423-1,755)	1,599 (1,445-1,781)	1,625 (1,531-1,737)	1,592 (1,421-1,792)	1,659 (1,564-1,774)	1,574; 1,659
50 - 59	3,575	6.3 (5.6-7.0)	6.6 (5.9-7.3)	6.6 (6.2-7.0)	6.3 (5.5-7.1)	6.9 (6.5-7.3)	6.3; 6.9	1,499 (1,337-1,669)	1,578 (1,417-1,741)	1,537 (1,474-1,681)	1,496 (1,314-1,688)	1,645 (1,547-1,748)	1,496; 1,645
60 - 69	2,611	6.0 (5.4-6.6)	6.0 (5.3-6.6)	6.4 (6.0-6.8)	5.9 (5.2-6.6)	6.1 (5.2-6.8)	5.9; 6.4	1,438 (1,279-1,589)	1,440 (1,258-1,589)	1,531 (1,437-1,625)	1,416 (1,243-1,583)	1,457 (1,241-1,631)	1,416; 1,531
70 - 79	792	5.7 (5.1-6.4)	5.9 (5.2-6.6)	6.2 (5.8-6.7)	5.5 (4.8-6.3)	6.0 (5.2-6.8)	5.5; 6.2	1,365 (1,208-1,537)	1,416 (1,252-1,574)	1,482 (1,388-1,590)	1,320 (1,148-1,517)	1,429 (1,233-1,614)	1,320; 1,482
80 - 89	55	5.4 (4.8-5.5)	5.8 (5.1-5.9)	6.0 (5.6-6.1)	5.2 (4.4-5.3)	5.8 (5.0-5.9)	5.2; 6.0	1,295 (1,137-1,323)	1,375 (1,229-1,402)	1,437 (1,337-1,453)	1,236 (1,050-1,266)	1,379 (1,205-1,412)	1,236; 1,437
90+	12	5.2 (4.2-5.7)	5.8 (4.8-6.3)	5.8 (5.2-6.1)	4.9 (3.7-5.4)	5.8 (4.6-6.5)	4.9; 5.8	1,243 (995-1,354)	1,389 (1,146-1,515)	1,398 (1,252-1,466)	1,160 (890-1,281)	1,396 (1,105-1,544)	1,160; 1,398



Age (years)	n	REE (MJ/day) estimated with Mifflin et al. 1990 Median (P5-P95)	REE (MJ/day) estimated with Henry 2005 Median (P5-P95)	REE (MJ/day) estimated with Müller et al. 2004 Median (P5-P95)	REE (MJ/day) estimated with Harris-Benedict (1919) Median (P5-P95)	REE (MJ/day) estimated with Schofield et al. 1985 Median (P5-P95)	Lowest and highest median REE (MJ/day)	REE (kcal/day) estimated with Mifflin et al. 1990 Median (P5-P95)	REE (kcal/day) estimated with Henry 2005 Median (P5-P95)	REE (kcal/day) estimated with Müller et al. 2004 Median (P5-P95)	REE (kcal/day) estimated with Harris-Benedict (1919) Median (P5-P95)	REE (kcal/day) estimated with Schofield et al. 1985 Median (P5-P95)	Lowest and highest median REE (kcal/day)
<b>Women</b>													
18 - 29	3,589	5.6 (5.0-6.3)	5.6 (5.0-6.3)	5.7 (5.3-6.1)	5.9 (5.5-6.4)	5.7 (5.2-6.4)	5.6; 5.9	1,342 (1,203-1,502)	1,346 (1,208-1,509)	1,352 (1,267-1,449)	1,416 (1,324-1,521)	1,372 (1,245-1,525)	1,342; 1,416
30 - 39	3,866	5.3 (4.8-6.0)	5.4 (5.0-5.9)	5.5 (5.2-5.9)	5.7 (5.3-6.1)	5.6 (5.3-5.8)	5.3; 5.7	1,278 (1,150-1,438)	1,296 (1,186-1,418)	1,308 (1,232-1,404)	1,357 (1,273-1,462)	1,327 (1,269-1,394)	1,278; 1,357
40 - 49	4,727	5.1 (4.6-5.8)	5.4 (4.9-5.9)	5.3 (5.0-5.7)	5.5 (5.1-5.9)	5.5 (5.3-5.8)	5.1; 5.5	1,224 (1,095-1,381)	1,285 (1,178-1,418)	1,270 (1,191-1,367)	1,309 (1,221-1,413)	1,321 (1,264-1,394)	1,224; 1,321
50 - 59	4,066	4.8 (4.3-5.4)	5.3 (4.8-5.8)	5.1 (4.8-5.5)	5.2 (4.9-5.6)	5.5 (5.2-5.8)	4.8; 5.5	1,154 (1,018-1,299)	1,274 (1,157-1,384)	1,223 (1,141-1,311)	1,249 (1,159-1,344)	1,315 (1,253-1,375)	1,154; 1,315
60 - 69	2,806	4.6 (4.0-5.2)	4.9 (4.5-5.3)	5.0 (4.6-5.3)	5.0 (4.6-5.4)	5.0 (4.6-5.5)	4.6; 5.0	1,102 (966-1,232)	1,164 (1,068-1,279)	1,187 (1,106-1266)	1,202 (1,109-1,288)	1,195 (1,099-1,309)	1,102; 1,202
70 - 79	915	4.3 (3.7-4.9)	4.8 (4.4-5.3)	4.8 (4.4-5.1)	4.8 (4.4-5.2)	5.0 (4.5-5.4)	4.3; 5.0	1,027 (887-1,180)	1,154 (1,054-1,268)	1,138 (1,055-1,230)	1,138 (1,046-1,241)	1,185 (1,086-1,298)	1,027; 1,185
80 - 89	88	4.0 (3.5-4.4)	4.7 (4.3-5.1)	4.6 (4.3-4.8)	4.5 (4.2-4.8)	4.8 (4.5-5.2)	4.0; 4.8	955 (839-1,059)	1,124 (1,035-1,216)	1,091 (1,021-1,150)	1,078 (1,004-1,140)	1,155 (1,066-1,246)	955; 1,155
90+	4	3.4 (3.4-3.9)	4.4 (4.4-4.8)	4.2 (4.2-4.5)	4.1 (4.1-4.4)	4.6 (4.6-4.9)	3.4; 4.6	813 (813-932)	1,064 (1,064-1,144)	1,000 (1,000-1,072)	971 (971-1,052)	1,095 (1,095-1,175)	813; 1,095

The respective predictive equations based on body mass and height were used, where available (see Appendix 2). For Müller et al., the equation based on body mass for subpopulation 1 was used (Müller et al., 2004).

**APPENDIX 10: COMPARISON OF MEASURED REE OF GISELA SUBJECTS (LAST AVAILABLE MEASUREMENTS) WITH REE CALCULATED WITH VARIOUS PREDICTIVE EQUATIONS**

<b>Women (n=386, 61-96 years, BMI 15.9-43.6)</b>		<b>Median 5<sup>th</sup>-95<sup>th</sup> Percentile</b>				
<b>REE (kJ/day) measured</b>		5,590 4,516-7,092				
<b>REE (kJ/day) predicted using the equation(s) of</b>	<b>Bias</b>	<b>Upper limit of agreement</b>	<b>Lower limit of agreement</b>	<b>Accuracy (± 10 %) (n/%)</b>	<b>R<sup>2</sup></b>	
Schofield (1985)	26	1,041	- 989	285/74 %	0.57	
Müller et al. (2004)	- 263	749	- 1,275	280/73%	0.58	
Henry (2005)	- 311	710	- 1,332	281/73 %	0.58	
Harris-Benedict (1919)	- 364	648	- 1,376	259/67 %	0.57	
Mifflin et al. (1990)	- 795	235	- 1,825	127/33 %	0.57	
<b>Men (n=165, 60-92 years, BMI 18.8-47.4)</b>		<b>Median 5-95<sup>th</sup> Percentile</b>				
<b>REE (kJ/day) measured</b>		6,674 5,595-8,880				
<b>REE (kJ/day) predicted using the equation(s) of</b>	<b>Bias</b>	<b>Upper limit of agreement</b>	<b>Lower limit of agreement</b>	<b>Accuracy (± 10 %) (n/%)</b>	<b>R<sup>2</sup></b>	
Müller et al. (2004)	43	1,284	- 1,199	118/72 %	0.57	
Henry (2005)	- 203	1,081	- 1,487	117/71 %	0.53	
Schofield (1985)	- 276	1,117	- 1,668	115/70 %	0.45	
Harris-Benedict (1919)	- 494	779	- 1,767	96/58 %	0.56	
Mifflin et al. (1990)	- 540	733	- 1,813	94/57 %	0.56	

Bias = mean of differences (in kJ) of calculated REE versus measured REE; Upper limit of agreement = Bias + (1.96 x SD); Lower limit of agreement = Bias - (1.96 x SD); Accuracy: estimated as the number and percentage of subjects that have a REE predicted by the equation within 10 % of the measured REE.

## APPENDIX 11: REFERENCE BODY HEIGHTS AND BODY MASSES FOR INFANTS, CHILDREN AND ADULTS

### Infants and children

For the calculation of the average energy requirement, reference body masses and reference body heights are required. It has previously been recommended to develop a database with reference body masses and heights that are representative for the total population in the EU (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010). Concurrently, harmonised growth references for height, body mass and body mass index (BMI) at the EU level were calculated (van Buuren et al., 2012) using existing data available from the individual EU Member States and covering the period of 1990-2011. The coverage of the population in the EU was 90.1 % for height-for-age, 87.5 % for body mass-for-age, and 85.2 % for BMI-for-age. The proposed harmonised EU growth references are used in this Opinion for the ages 3-17 years. For infants and children up to two years of age, data were taken from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006). Median body heights and body masses for children aged 1-17 years are shown in Table 8.

### Adults

For the report on nutrient and energy intakes for the European Community by the SCF (1993), weighted median body masses of European men and women derived from the pooling of national data from a limited number of EU Member States were used. These data are relatively old and not necessarily representative of the newer EU Member States. For this Opinion, an attempt was made to gather more recent anthropometric data, to account for possible secular changes and the increase in size of the EU. For Bulgaria (Petrova and Angelova, 2006), Finland (Paturi et al., 2008), France (AFSSA, 2009), Germany (MRI, 2008a, 2008b), Ireland (Harrington et al., 2001; Kiely et al., 2001), Poland (Szponar et al., 2001; Szponar et al., 2003), Spain (AESAN) (Ortega et al., 2011) and the United Kingdom (Henderson et al., 2002) individual data on measured body heights and body masses from representative surveys were already available to EFSA via the Comprehensive Food Consumption Database (Merten et al., 2011). Various other countries for which such data may be available were identified with the help of the European Commission Directorate General – SANCO and WHO Regional Office in Europe. Following a request for data submission, such data were received from the Netherlands (Health examination survey in the Netherlands 2009-2010) (Blokstra et al., 2011), Portugal (do Carmo et al., 2008), Slovakia (CINDI 2008) (Avdičová et al., 2005), Luxembourg (Alkerwi et al., 2010) and the Czech Republic (HELEN Study: Health, Life Style and Environment 2004-2005) (Kratěnová et al., 2007). The overall population coverage, i.e. the number of inhabitants in these 13 EU Member States relative to all EU citizens, is equal to 66-71 % for age groups between 18 and 69 years, 43 % for the age group 70-79 years, and even lower for age groups 80-89 years and  $\geq 90$  years.

Weighting factors were used in order to take into account the population size of the respective country for which data were available. Weighting factors were obtained for both sexes by dividing, for each country, the population size of the age categories by the number of subjects included in the survey. Information on the population by country, age category and sex were extracted from the EUROSTAT website (<http://epp.eurostat.ec.europa.eu>), and are referred to 2010. Body masses were calculated for a BMI of 22 kg/m<sup>2</sup> using measured body heights.

Median measured body heights and body masses, as well as body masses for a BMI of 22 kg/m<sup>2</sup>, based on data obtained in the 13 EU Member States are listed in Table 4.

**APPENDIX 12A: EXAMPLES OF RELATIONSHIPS REPORTED BETWEEN LIFESTYLE, ACTIVITY AND PHYSICAL ACTIVITY LEVEL (PAL)**

Intensity of activities	PAL	Examples
Chair-bound or bedridden	1.2 – 1.3	Fragile, invalid people
Seated work with no option of moving around and little or no strenuous leisure activity	1.4 – 1.5	Office workers
Seated work with discretion, requirement to move around, little or no strenuous leisure activity	1.6 – 1.7	Students, white-collar or professional workers
Standing work	1.8 – 1.9	Homemakers, shop assistants, waiters, craftsmen
Significant amounts of sports or strenuous leisure activity (30-60 minutes 4-5 times/week)	+ 0.3	
Strenuous work or highly active leisure	2.0 – 2.4	Building labourers, miners, lumberjacks, competitive athletes

Adapted from Black et al. (1996).

**APPENDIX 12B: CONTRIBUTION OF VARIOUS ACTIVITIES TO PHYSICAL ACTIVITY LEVELS (PALS)**

Effect of various activities on PAL	Physical activity ratio (PAR)	3 h/week	5 h/week	1 h/day
		Addition to PAL		
<b>Home and leisure activities</b>				
Light activity (knitting, sewing) while sitting	1.3	0.02	0.04	0.05
Standing fidgeting	1.8	0.03	0.05	0.08
Playing piano	2.3	0.04	0.07	0.10
Household tasks, moderate effort	3.5	0.06	0.10	0.15
Walking (3.2 km/h)	2.8	0.05	0.08	0.12
Cycling (leisurely)	4	0.07	0.12	0.17
Gardening	3.8	0.07	0.11	0.16
Hunting general	5	0.09	0.15	0.21
Fishing general	3	0.05	0.09	0.13
<b>Sports</b>				
Stretching, Hatha Yoga	2.5	0.04	0.07	0.10
Billiards	2.5	0.04	0.07	0.10
Weight lifting/Body building vigorously	6	0.11	0.18	0.25
Ballet, modern, or jazz, general, rehearsal or class	5	0.09	0.15	0.21
Dancing aerobic low impact	5	0.09	0.15	0.21
Dancing aerobic high impact	7.3	0.13	0.22	0.30
Bicycling/rowing stationary 100 watts	7	0.13	0.21	0.29
Bicycling/rowing stationary 150 watts	8.5	0.15	0.25	0.35
Swimming leisurely	6	0.11	0.18	0.25
Swimming crawl 50 m/min	8.3	0.15	0.25	0.35
Swimming crawl 75 m/min	10	0.18	0.30	0.42
Walking 6.4 km/h	5	0.09	0.15	0.21
Running 8 km/h	8.3	0.15	0.25	0.35
Running 16 km/h	14.5	0.26	0.43	0.60
Running cross country	9	0.16	0.27	0.38
Rock or mountain climbing	8	0.14	0.24	0.33
Roller skating	7	0.13	0.21	0.29

Effect of various activities on PAL	Physical activity ratio (PAR)	Addition to PAL		
		3 h/week	5 h/week	1 h/day
Tennis double	4.5	0.08	0.13	0.19
Tennis single	8	0.14	0.24	0.33
Cycling 16-19 km/h	6.8	0.12	0.20	0.28
Cycling 19-22 km/h	8	0.14	0.24	0.33
Cycling 22-25 km/h	10	0.18	0.30	0.42
Squash	12	0.21	0.36	0.50
Golf	4.8	0.09	0.14	0.20

Based on published values of metabolic equivalents from Ainsworth et al. (2011), used as a proxy of PAR.

**APPENDIX 13: SELECTED PREDICTIVE EQUATIONS FOR REE IN CHILDREN AND ADOLESCENTS**

**Predictive equations for REE for children and adolescents from Schofield (1985) using body mass (BM, in kg) and height (H, in m)**

Age (years)	Boys				Girls			
	MJ/day (kcal/day)	n	SE	r	MJ/day (kcal/day)	n	SE	r
0-3	0.0007 BM + 6.349 H - 2.584 (0.167 BM + 1517.4 H - 617.6)	162	0.243	0.97	0.068 BM + 4.281 H - 1.730 (16.25 BM + 1023.2 H - 413.5)	137	0.216	0.97
3-10	0.082 BM + 0.545 H + 1.736 (19.6 BM + 130.3 H + 414.9)	338	0.280	0.83	0.071 BM + 0.677 H + 1.553 (16.97 BM + 161.8 H + 371.2)	413	0.290	0.81
10-18	0.068 BM + 0.574 H + 2.157 (16.25 BM + 137.2 H + 515.5)	734	0.439	0.93	0.035 BM + 1.948 H + 0.837 (8.365 BM + 465 H + 200)	575	0.453	0.82

n, number of individuals; SE, standard error; r, correlation coefficient of the linear regression.

**Predictive equations for REE for children and adolescents from Henry (2005) using body mass (BM, in kg) and height (H, in m)**

Age (years)	Boys				Girls			
	MJ/day (kcal/day)	n	SE	r	MJ/day (kcal/day)	n	SE	r
0-3	0.118 BM + 3.59 H - 1.55 (28.2 BM + 859 H - 371)	246	0.246	0.96	0.127 BM + 2.94 H - 1.20 (30.4 BM + 703 H - 287)	201	0.232	0.96
3-10	0.0632 BM + 1.31 H + 1.28 (15.1 BM + 74.2 H + 306) <sup>(1)</sup>	289	0.322	0.84	0.0666 BM + 0.878 H + 1.46 (15.9 BM + 210 H + 349)	403	0.357	0.83
10-18	0.0651 BM + 1.11 H + 1.25 (15.6 BM + 266 H + 299)	863	0.562	0.86	0.0393 BM + 1.04 H + 1.93 (9.40 BM + 249 H + 462)	1,063	0.521	0.76

n, number of individuals; SE, standard error; r, correlation coefficient of the linear regression.

<sup>(1)</sup> likely error in the cited formula, so in this Opinion the respective formula for MJ/day was used and the results for kcal/day obtained after conversion.



**APPENDIX 14A: RANGES OF AVERAGE REQUIREMENT (AR) FOR ENERGY FOR ADULTS BASED ON THE FACTORIAL METHOD AND PREDICTING REE WITH FIVE MOST USED EQUATIONS**

Age (years)	Lowest median REE (kcal/day)	Highest median REE (kcal/day)	Range of AR at PAL=1.4 <sup>(1)</sup> (kcal/day)	Range of AR at PAL=1.6 <sup>(1)</sup> (kcal/day)	Range of AR at PAL=1.8 <sup>(1)</sup> (kcal/day)	Range of AR at PAL=2.0 <sup>(1)</sup> (kcal/day)	Range of AR at PAL=2.2 <sup>(1)</sup> (kcal/day)	Range of AR at PAL=2.4 <sup>(1)</sup> (kcal/day)
<b>Men</b>								
18 - 29	1,674	1,761	2,338 - 2,466	2,672 - 2,818	3,006 - 3,170	3,340 - 3,522	3,674 - 3,875	4,008 - 4,227
30 - 39	1,621	1,675	2,264 - 2,344	2,588 - 2,679	2,911 - 3,014	3,235 - 3,349	3,558 - 3,684	3,881 - 4,019
40 - 49	1,574	1,659	2,204 - 2,322	2,519 - 2,654	2,834 - 2,986	3,148 - 3,317	3,463 - 3,649	3,778 - 3,981
50 - 59	1,496	1,645	2,094 - 2,304	2,393 - 2,633	2,692 - 2,962	2,991 - 3,291	3,290 - 3,620	3,590 - 3,949
60 - 69	1,416	1,531	1,982 - 2,144	2,265 - 2,450	2,549 - 2,756	2,832 - 3,062	3,115 - 3,369	3,398 - 3,675
70 - 79	1,320	1,482	1,848 - 2,075	2,112 - 2,371	2,376 - 2,667	2,640 - 2,964	2,904 - 3,260	3,169 - 3,556
<b>Women</b>								
18 - 29	1,342	1,416	1,878 - 1,983	2,147 - 2,266	2,415 - 2,549	2,683 - 2,832	2,952 - 3,116	3,220 - 3,399
30 - 39	1,278	1,357	1,789 - 1,899	2,045 - 2,171	2,300 - 2,442	2,556 - 2,713	2,812 - 2,985	3,067 - 3,256
40 - 49	1,224	1,321	1,713 - 1,849	1,958 - 2,114	2,203 - 2,378	2,448 - 2,642	2,692 - 2,906	2,937 - 3,170
50 - 59	1,154	1,315	1,616 - 1,841	1,847 - 2,104	2,077 - 2,367	2,308 - 2,630	2,539 - 2,893	2,770 - 3,156
60 - 69	1,102	1,202	1,542 - 1,682	1,762 - 1,923	1,983 - 2,163	2,203 - 2,403	2,423 - 2,644	2,644 - 2,884
70 - 79	1,027	1,185	1,438 - 1,659	1,643 - 1,896	1,849 - 2,133	2,054 - 2,370	2,260 - 2,607	2,465 - 2,844

<sup>(1)</sup> Based on lowest and highest median REE (see Appendix 9).

**APPENDIX 14B: RANGES OF AVERAGE REQUIREMENT (AR) FOR ENERGY FOR CHILDREN AND ADOLESCENTS BASED ON THE FACTORIAL METHOD AND PREDICTING REE WITH TWO PREDICTIVE EQUATIONS**

Age (years)	REE (kcal/day) (Henry)	REE (kcal/day) (Schofield)	Range of AR at PAL=1.4 (kcal/day)	Range of AR at PAL=1.6 (kcal/day)	Range of AR at PAL=1.8 (kcal/day)	Range of AR at PAL=2.0 (kcal/day)	Range of AR at PAL=2.2 (kcal/day)	Range of AR at PAL=2.4 (kcal/day)
<b>Boys</b>								
1	550	533	753 - 777	861 - 888				
2	727	717	1,013 - 1,028	1,158 - 1,175				
3	830	829	1,172 - 1,174	1,339 - 1,341				
4	888	884	1,249 - 1,256	1,428 - 1,436	1,606 - 1,615	1,785 - 1,794	1,963 - 1,974	
5	942	935	1,322 - 1,332	1,511 - 1,522	1,700 - 1,712	1,889 - 1,903	2,078 - 2,093	
6	996	988	1,398 - 1,409	1,597 - 1,610	1,797 - 1,811	1,997 - 2,013	2,196 - 2,214	
7	1,059	1,052	1,487 - 1,497	1,700 - 1,711	1,912 - 1,925	2,125 - 2,139	2,337 - 2,353	
8	1,126	1,121	1,585 - 1,592	1,811 - 1,819	2,037 - 2,046	2,264 - 2,274	2,490 - 2,501	
9	1,191	1,191	1,683 - 1,684	1,924 - 1,925	2,164 - 2,165	2,405 - 2,406	2,645 - 2,647	
10	1,196	1,257	1,691 - 1,777	1,933 - 2,031	2,174 - 2,285	2,416 - 2,539	2,658 - 2,793	2,899 - 3,047
11	1,264	1,321	1,788 - 1,868	2,043 - 2,135	2,298 - 2,401	2,554 - 2,668	2,809 - 2,935	3,065 - 3,202
12	1,345	1,397	1,902 - 1,976	2,174 - 2,258	2,445 - 2,540	2,717 - 2,822	2,989 - 3,104	3,260 - 3,387
13	1,444	1,491	2,041 - 2,108	2,333 - 2,409	2,625 - 2,710	2,916 - 3,011	3,208 - 3,313	3,500 - 3,614
14	1,555	1,598	2,199 - 2,259	2,513 - 2,582	2,828 - 2,905	3,142 - 3,228	3,456 - 3,550	3,770 - 3,873
15	1,670	1,709	2,362 - 2,416	2,699 - 2,761	3,036 - 3,107	3,374 - 3,452	3,711 - 3,797	4,048 - 4,142
16	1,761	1,797	2,489 - 2,542	2,845 - 2,905	3,201 - 3,268	3,556 - 3,631	3,912 - 3,994	4,268 - 4,357
17	1,819	1,856	2,572 - 2,624	2,940 - 2,999	3,307 - 3,374	3,675 - 3,748	4,042 - 4,123	4,409 - 4,498
<b>Girls</b>								
1	503	488	690 - 712	789 - 813				
2	669	657	930 - 946	1,062 - 1,082				
3	775	767	1,084 - 1,096	1,239 - 1,253				
4	826	816	1,154 - 1,168	1,319 - 1,335	1,483 - 1,502	1,648 - 1,668	1,813 - 1,835	
5	877	866	1,224 - 1,239	1,399 - 1,417	1,574 - 1,594	1,749 - 1,771	1,924 - 1,948	
6	928	917	1,297 - 1,312	1,482 - 1,500	1,667 - 1,687	1,852 - 1,875	2,037 - 2,062	
7	984	973	1,376 - 1,392	1,572 - 1,591	1,769 - 1,790	1,956 - 1,989	2,162 - 2,187	

Age (years)	REE (kcal/day) (Henry)	REE (kcal/day) (Schofield)	Range of AR at PAL=1.4 (kcal/day)	Range of AR at PAL=1.6 (kcal/day)	Range of AR at PAL=1.8 (kcal/day)	Range of AR at PAL=2.0 (kcal/day)	Range of AR at PAL=2.2 (kcal/day)	Range of AR at PAL=2.4 (kcal/day)
8	1,045	1,034	1,461 - 1,477	1,670 - 1,688	1,879 - 1,899	2,088 - 2,110	2,297 - 2,321	
9	1,107	1,097	1,551 - 1,566	1,773 - 1,790	1,994 - 2,013	2,216 - 2,237	2,437 - 2,461	
10	1,125	1,133	1591 - 1,602	1,818 - 1,831	2,046 - 2,059	2,273 - 2,288	2,500 - 2,517	2,728 - 2,746
11	1,181	1,198	1669 - 1,694	1,908 - 1,936	2,146 - 2,177	2,385 - 2,419	2,623 - 2,661	2,862 - 2,903
12	1,240	1,266	1754 - 1,790	2,004 - 2,046	2,255 - 2,301	2,505 - 2,557	2,756 - 2,813	3,006 - 3,069
13	1,299	1,331	1837 - 1,882	2,099 - 2,150	2,361 - 2,419	2,624 - 2,688	2,886 - 2,957	3,149 - 3,226
14	1,346	1,381	1903 - 1,952	2,175 - 2,231	2,447 - 2,510	2,719 - 2,789	2,991 - 3,068	3,262 - 3,347
15	1,379	1,415	1950 - 2,001	2,228 - 2,287	2,507 - 2,573	2,786 - 2,859	3,064 - 3,145	3,343 - 3,430
16	1,398	1,434	1,977 - 2,028	2,259 - 2,318	2,542 - 2,608	2,824 - 2,898	3,107 - 3,187	3,389 - 3,477
17	1,409	1,446	1,992 - 2,044	2,277 - 2,336	2,562 - 2,628	2,846 - 2,920	3,131 - 3,212	3,416 - 3,504

**APPENDIX 15: DERIVATION OF THE AVERAGE REQUIREMENT (AR) FOR ENERGY FOR INFANTS AGED 7-11 MONTHS**

Age (months)	Body mass <sup>(1)</sup> (kg)	Gain in body mass <sup>(2)</sup> (g/day)	Energy deposition <sup>(3)</sup> (kcal/g)	Energy deposition <sup>(4)</sup> (kcal/day)	TEE <sup>(5)</sup> (kcal/day)	AR <sup>(6)</sup> (kcal/day)	AR (kcal/kg per day)
<b>Boys</b>							
7	8.3	11.9	1.5	17.6	618	636	76
8	8.6	10.5	1.5	15.6	646	661	77
9	8.9	9.5	1.5	14.1	674	688	77
10	9.2	8.6	2.7	23.5	701	725	79
11	9.4	8.1	2.7	22.1	720	742	79
<b>Girls</b>							
7	7.6	11.5	1.8	20.3	553	573	76
8	7.9	10.4	1.8	18.3	581	599	76
9	8.2	9.1	1.8	16.1	609	625	76
10	8.5	8.2	2.3	19.1	636	656	77
11	8.7	7.8	2.3	18.2	655	673	77

<sup>(1)</sup> 50<sup>th</sup> percentile of WHO Growth Standards.

<sup>(2)</sup> Calculation from 1-month body mass increments from 50<sup>th</sup> percentile of WHO Growth Standards, assuming that 1 month = 30 days.

<sup>(3)</sup> see Table 7.

<sup>(4)</sup> Body mass gain × energy accrued in normal growth.

<sup>(5)</sup> Total energy expenditure (TEE) (kcal/day) = -152.0 + 92.8 × body mass (kg).

<sup>(6)</sup> AR = TEE + energy deposition.

**APPENDIX 16: SUMMARY OF AVERAGE REQUIREMENT (AR) FOR ENERGY EXPRESSED IN KCAL/DAY**

**Summary of Average Requirement (AR) for energy for adults**

Age (years)	REE <sup>(1)</sup> (kcal/day)	AR at PAL=1.4 (kcal/day)	AR at PAL=1.6 (kcal/day)	AR at PAL=1.8 (kcal/day)	AR at PAL=2.0 (kcal/day)
<b>Men</b>					
18 - 29	1,674	2,338	2,672	3,006	3,340
30 - 39	1,621	2,264	2,588	2,911	3,235
40 - 49	1,599	2,234	2,553	2,873	3,192
50 - 59	1,578	2,204	2,519	2,834	3,149
60 - 69	1,440	2,017	2,305	2,593	2,882
70 - 79	1,416	1,984	2,267	2,550	2,834
<b>Women</b>					
18 - 29	1,346	1,878	2,147	2,415	2,683
30 - 39	1,296	1,813	2,072	2,331	2,590
40 - 49	1,285	1,798	2,055	2,312	2,569
50 - 59	1,274	1,783	2,037	2,292	2,547
60 - 69	1,164	1,628	1,861	2,093	2,326
70 - 79	1,154	1,614	1,844	2,075	2,305

<sup>(1)</sup> REE, resting energy expenditure predicted with the equations of Henry (2005) using body mass and height. Because these have overlapping age bands (18-30 years, 30-60 years, ≥60 years) (see Appendix 2), the choice of equation is ambiguous at the age boundaries. The REE equations for 18-30 year-olds are used for adults aged 18-29 years, the equations for 30-60 year-olds are used for adults aged 30-39, 40-49 and 50-59 years, and the equations for ≥60 year-olds are used for adults aged 60-69 and 70-79 years.

**Summary of Average Requirement (AR) for energy for infants**

Age	AR (kcal/day)		AR (kcal/kg body mass per day)	
	Boys	Girls	Boys	Girls
7 months	636	573	76	76
8 months	661	599	77	76
9 months	688	625	77	76
10 months	725	656	79	77
11 months	742	673	79	77

### Summary of Average Requirement (AR) for energy for children and adolescents

Age (years)	REE <sup>(1)</sup> (kcal/day)	AR <sup>(2)</sup> at PAL <sup>(3)</sup> =1.4 (kcal/day)	AR <sup>(2)</sup> at PAL=1.6 (kcal/day)	AR <sup>(2)</sup> at PAL=1.8 (kcal/day)	AR <sup>(2)</sup> at PAL=2.0 (kcal/day)
<b>Boys</b>					
1	550	777			
2	727	1,028			
3	830	1,174			
4	888	1,256	1,436	1,615	
5	942	1,332	1,522	1,712	
6	996	1,409	1,610	1,811	
7	1,059	1,497	1,711	1,925	
8	1,126	1,592	1,819	2,046	
9	1,191	1,684	1,925	2,165	
10	1,196		1,933	2,174	2,416
11	1,264		2,043	2,298	2,554
12	1,345		2,174	2,445	2,717
13	1,444		2,333	2,625	2,916
14	1,555		2,513	2,828	3,142
15	1,670		2,699	3,036	3,374
16	1,761		2,845	3,201	3,556
17	1,819		2,940	3,307	3,675
<b>Girls</b>					
1	503	712			
2	669	946			
3	775	1,096			
4	826	1,168	1,335	1,502	
5	877	1,239	1,417	1,594	
6	928	1,312	1,500	1,687	
7	984	1,392	1,591	1,790	
8	1,045	1,477	1,688	1,899	
9	1,107	1,566	1,790	2,013	
10	1,125		1,818	2,046	2,273
11	1,181		1,908	2,146	2,385
12	1,240		2,004	2,255	2,505
13	1,299		2,099	2,361	2,624
14	1,346		2,175	2,447	2,719
15	1,379		2,228	2,507	2,786
16	1,398		2,259	2,542	2,824
17	1,409		2,277	2,562	2,846

<sup>(1)</sup> REE, resting energy expenditure computed with the predictive equations of Henry. Because the equations of Henry have overlapping age bands (0-3, 3-10, 10-18 years), the choice of equation is ambiguous at the age boundaries. The REE equation for 3-10 year-olds is used for the 3 year-olds and the equation for 10-18 year-olds is used for those aged 10 years.

<sup>(2)</sup> Taking into account a coefficient of 1.01 for growth.

<sup>(3)</sup> PAL, physical activity level

### Summary of Average Requirement (AR) for energy for pregnant and lactating women (in addition to the AR for non-pregnant women)

	AR (kcal/day)
Pregnant women	
1 <sup>st</sup> trimester	+70
2 <sup>nd</sup> trimester	+260
3 <sup>rd</sup> trimester	+500
Lactating women	+500
0-6 months <i>post partum</i>	



## GLOSSARY AND ABBREVIATIONS

AESAN	Agencia Española de Seguridad Alimentaria y Nutrición
AFSSA	Agence Française de Sécurité Sanitaire des Aliments
AR	Average requirement
ATP	Adenosin-triphosphate
BEE	Basal energy expenditure
BM	Body mass
BMI	Body mass index
cal	Calorie
CINDI	Countrywide Integrated Noncommunicable Diseases Intervention
COMA	Committee on Medical Aspects of Food Policy
CV	Coefficient of variation
d	Day
D-A-CH	Deutschland-Austria-Confoederatio Helvetica
DLW	Doubly-labelled water
DoH	Department of Health
DRI	Dietary reference intake
DRV	Dietary reference value
EC	European Commission
EEPA	Energy expenditure of physical activity
EER	Estimated energy requirement
EFSA	European Food Safety Authority
EU	European Union
FAO	Food and Agriculture Organization
FFM	Fat-free mass
FM	Fat mass
GE	Gross energy

GISELA	Giessener Senioren Langzeitstudie
HELEN	Health, Life Style and Environment
HR	Heart rate
IDECG	International Dietary Energy Consultancy Group
IE	Ingested energy
IoM	U.S. Institute of Medicine of the National Academy of Sciences
ME	Metabolisable energy
mo	Month
n	Number of individuals
NA	Not available
NCHS	National Center for Health Statistics
NNR	Nordic Nutrition Recommendations
NRC	National Research Council
OPEN	Observing Protein and Energy Nutrition
PA	Physical activity
PAL	Physical activity level
PAR	Physical activity ratio
PRI	Population reference intake
REE	Resting energy expenditure
SACN	Scientific Advisory Committee on Nutrition
SCF	Scientific Committee on Food
SD	Standard deviation
SE	Standard error
SEE	Standard error of estimate
SI	International System of Units
TEE	Total energy expenditure
TEF	Termic effect of food
UK	United Kingdom

UNU	United Nations University
US	United States
VCO <sub>2</sub>	Carbon dioxide production
VO <sub>2</sub>	Oxygen consumption
WHO	World Health Organization
y	year