

## Regular Article

## PLATELETS AND THROMBOPOIESIS

Identification of a new dysfunctional platelet P2Y<sub>12</sub> receptor variant associated with bleeding diathesis

Anna Lecchi,<sup>1</sup> Cristina Razzari,<sup>2</sup> Silvia Paoletta,<sup>3</sup> Arnaud Dupuis,<sup>4</sup> Lea Nakamura,<sup>5</sup> Philippe Ohlmann,<sup>4</sup> Christian Gachet,<sup>4</sup> Kenneth A. Jacobson,<sup>3</sup> Barbara Zieger,<sup>5</sup> and Marco Cattaneo<sup>2</sup>

<sup>1</sup>Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Ca' Granda - Ospedale Maggiore Policlinico, Milan, Italy; <sup>2</sup>Divisione di Medicina 3, Ospedale San Paolo, Dipartimento di Scienze della Salute - Università degli Studi di Milano, Milan, Italy; <sup>3</sup>Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD; <sup>4</sup>Unité Mixte de Recherche\_S949, Inserm, Université de Strasbourg, Etablissement Français du Sang-Alsace, Strasbourg, France; and <sup>5</sup>Department of Pediatrics and Adolescent Medicine, University Medical Center Freiburg, Freiburg, Germany

## Key Points

- Two patients with bleeding diathesis had dysfunctional platelet P2Y<sub>12</sub>R for ADP, attributable to homozygous His187Gln mutation.
- These studies delineate a region of transmembrane 5 of P2Y<sub>12</sub>R that is required for normal function after ADP binding.

Defects of the platelet P2Y<sub>12</sub> receptor (P2Y<sub>12</sub>R) for adenosine diphosphate (ADP) are associated with increased bleeding risk. The study of molecular abnormalities associated with inherited qualitative defects of the P2Y<sub>12</sub>R protein is useful to unravel structure-function relationships of the receptor. We describe the case of 2 brothers, sons of first cousins, with lifelong history of abnormal bleeding, associated with dysfunctional P2Y<sub>12</sub>R and a previously undescribed missense mutation in the encoding gene. ADP (4-20 μM)-induced aggregation of patients' platelets was markedly reduced and rapidly reversible. Other agonists induced borderline-normal aggregation. Inhibition of vasodilator-stimulated phosphoprotein phosphorylation and prostaglandin E<sub>1</sub>-induced increase in cyclic adenosine monophosphate (cAMP) by ADP was impaired, whereas inhibition of cAMP increase by epinephrine was normal. [<sup>3</sup>H]PSB-0413, a selective P2Y<sub>12</sub>R antagonist, bound to a normal number of binding sites; however, its affinity, and that of the agonists ADP and 2-methylthio-adenosine-5'-diphosphate, was reduced. Patients' DNA showed a homozygous c.847T>A substitution that

changed the codon for His-187 to Gln (p.His187Gln). Crystallographic data and molecular modeling studies indicated that His187 in transmembrane 5 is important for agonist and nucleotide antagonist binding and located in a region undergoing conformational changes. These studies delineate a region of P2Y<sub>12</sub>R required for normal function after ADP binding. (*Blood*. 2015; 125(6):1006-1013)

## Introduction

P2Y<sub>12</sub>, which maps to chromosome 3q21-q25, is 1 of the 2 receptors for adenosine diphosphate (ADP) that are expressed by platelets.<sup>1,2</sup> It is a G inhibitory protein (Gi)-coupled receptor that plays an important role both in hemostasis and thrombosis. This role is demonstrated by the observations that patients with inherited defects of the receptor display a bleeding diathesis<sup>2,3</sup> and patients with clinical manifestations of atherothrombosis are protected from major adverse cardiovascular events by P2Y<sub>12</sub> receptor (P2Y<sub>12</sub>R) inhibitors.<sup>4</sup>

Congenital P2Y<sub>12</sub>R defects are autosomal recessive disorders, associated with qualitative or quantitative abnormalities of the receptor.<sup>2</sup> Affected patients display a bleeding diathesis of variable severity, characterized by mucocutaneous bleeding and excessive postsurgical or posttraumatic blood loss. The most typical platelet function abnormality of the disorder, which should raise the diagnostic suspicion of the disease, is the failure by ADP, even at very high concentrations (>10 μM), to induce full and irreversible platelet aggregation.<sup>2,5</sup> Tests that evaluate the degree of inhibition of adenylyl cyclase by ADP, by measuring either the platelet levels of cyclic

adenosine monophosphate (cAMP) or the phosphorylation of vasodilator-stimulated phosphoprotein (VASP) after the exposure of platelets to prostaglandin (PG) E<sub>1</sub> or other stimuli for adenylyl cyclase, should be used to confirm the diagnosis.<sup>2,6</sup>

The study of molecular abnormalities associated with inherited qualitative defects of the P2Y<sub>12</sub>R protein is useful to unravel structure-function relationships of the receptor. Mutations in the region that spans the transition between transmembrane helix (TM) 6 and extracellular loop (EL) 3 are associated with receptor dysfunction despite normal ligand binding, suggesting that this region of the molecule plays a role in signal transduction.<sup>7-9</sup> A heterozygous mutation, predicting a lysine to glutamate (p.Lys174Glu) substitution in the P2Y<sub>12</sub>R, was identified in 1 patient with reduced and reversible aggregation in response to ADP and an ~50% reduction in binding of agonist radioligand [<sup>3</sup>H]2-methylthio-adenosine-5'-diphosphate (2MeSADP).<sup>10</sup> Considering that Lys174 is situated in the EL2 of the P2Y<sub>12</sub>R, adjacent to Cys175, which may be important for the integrity of the ADP binding site on the receptor,<sup>11</sup> and that a hemagglutinin-tagged

Submitted July 26, 2013; accepted November 21, 2014. Prepublished online as *Blood* First Edition paper, November 26, 2014; DOI 10.1182/blood-2013-07-517896.

B.Z. and M.C. contributed equally to this study.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

p.Lys174Glu P2Y<sub>12</sub>R variant showed surface expression in Chinese hamster ovary cells, it was hypothesized that the p.Lys174Glu mutation inhibits ADP binding to the receptor. Recently reported crystallographic data<sup>12,13</sup> do not show a direct involvement of Lys174 in agonist binding but rather a role in stabilizing the receptor conformation through interresidue ionic interactions and/or in affecting the approach of the ligand to its binding site. In one patient, with no personal history of abnormal bleeding, reduced expression of P2Y<sub>12</sub>R was associated with the heterozygous mutation p.Pro341Ala, in a putative postsynaptic density 95/disc large/zonula occludens-1-binding motif, most likely as a consequence of significantly compromised P2Y<sub>12</sub>R recycling.<sup>14</sup>

Here, we report the case of an index patient and his brother, with an inherited dysfunctional P2Y<sub>12</sub>R associated with a normal number of platelet binding sites but reduced affinity for the P2Y<sub>12</sub>R antagonist radioligand [<sup>3</sup>H]2-Propylthioadenosine-5'-adenylic acid (1,1-chloro-1-phosphonomethyl-1-phosphonyl)anhydride ([<sup>3</sup>H]PSB-0413),<sup>15,16</sup> ADP, and 2MeSADP, and with homozygous c.847T>A substitution, which changed the codon for His187 to Gln (p.His187Gln).

## Methods

### Patients and controls

The proband (III-7 in Figure 1) is a 45-year-old man living in Germany, who belongs to a large family of Turkish origin (Figure 1). He presented with recurrent epistaxis, postoperative bleeding after tooth extraction, thyroid explantation because of thyroid hyperplasia, and spleen explantation, consequent to spleen rupture in a car accident. He had normal platelet count ( $307 \times 10^9/L$ ), screening coagulation tests (prothrombin time and activated partial thromboplastin time), and plasma levels of von Willebrand factor antigen and ristocetin cofactor activity. Serum thromboxane B<sub>2</sub> levels and the platelet contents of serotonin, ADP, adenosine triphosphate (ATP), and fibrinogen were normal. Blood samples for genetic studies were available from his parents, who are first cousins, and 9 relatives (2 grandparents and 7 brothers and sisters), with negative, mild, or severe bleeding history. The 57-year-old brother, with severe bleeding history (III-1, Figure 1), also provided blood samples for platelet function studies.

A total of 36 healthy subjects, with no personal history of abnormal bleeding, were also studied. All subjects gave consent to the study.

### Materials

[<sup>3</sup>H]PSB-0413 was prepared by catalytic hydrogenation with tritium gas (GE Healthcare, Buckinghamshire, UK) of the propargyl precursor PSB-0412 as described.<sup>15</sup> ADP, 2MeSADP, collagen, the thromboxane/prostaglandin endoperoxide analog 9,11-dideoxy-11,9-epoxymethano-prostaglandin F<sub>2</sub>, thrombin receptor activating peptide 6, PGE<sub>1</sub>, and PGI<sub>2</sub> were from Sigma (St. Louis, MO). Apyrase was purified from potatoes.<sup>17</sup> Commercial preparations of luciferin/luciferase reagent were used to measure the platelet ATP and ADP contents (ATP Assay Kit; BioOrbit Oy, Turku, Finland), and platelet secretion concurrently with platelet aggregation (Chronolume; Chrono-log Corp., Havertown, PA). Serum thromboxane B<sub>2</sub> was measured by a commercially available enzymatic immunoassay (Thromboxane B<sub>2</sub> EIA kit; Cayman Chemical Co., Ann Arbor, MI).

### Preparation of platelet-rich plasma (PRP) and washed platelet suspensions

For studies with PRP, 9 volumes of blood were drawn into 1 volume of 109 mmol/L trisodium citrate and then centrifuged at 200g for 10 minutes.<sup>18,19</sup> The supernatant PRP was transferred into a clean plastic tube; the platelet counts in PRP samples were adjusted to  $2.5 \times 10^{11}/L$  in experiments performed in Freiburg (Germany) and not adjusted to a predefined value in experiments performed in Milan (Italy) and Strasbourg (France).<sup>18,20</sup> For the

preparation of washed platelet suspensions, 6 volumes of blood were drawn into 1 volume of acid-citrate-dextrose anticoagulant, centrifuged at 200g for 10 minutes to obtain PRP, which was used to prepare twice-washed platelet suspensions, according to the method described by Mustard et al,<sup>21</sup> with the exception that 500 nmol/L PGI<sub>2</sub> was added during the first and second washes. Platelet counts in washed platelet suspensions were adjusted to  $3 \times 10^{11}/L$ .

### Studies of platelet aggregation and secretion

The first screening of platelet aggregation in patient III-7 was performed in Freiburg (Germany), using the light transmission aggregometer APACT 4 (Labor Fibrinometer, Ahrensburg, Germany), and collagen (Nycomed, Austria) and ADP (MP Biomedicals) as agonists. Two separate sets of experiments were subsequently done in Milan (Italy), where platelet aggregation and secretion were studied simultaneously by lumi-aggregometry. Samples of PRP (0.45 mL) were incubated with 50  $\mu$ L luciferin/luciferase reagent at 37°C for 30 seconds and stirred at 1000 rpm in a lumi-aggregometer (Lumi-aggregometer; Chrono-log Corp.). After incubation, 10  $\mu$ L of an aggregating agent was added, and the aggregation and ATP secretion tracings were recorded for 3 minutes. Platelet aggregation of patient III-1 was studied in Strasbourg (France) in an APACT 4004 aggregometer.

### Binding of fluorescein-conjugated fibrinogen to activated platelets

To measure the binding of fibrinogen to activated platelets, aliquots of 100  $\mu$ L PRP ( $5 \times 10^{10}/L$ ) were mixed with fluorescein-conjugated fibrinogen (final concentration 150  $\mu$ g/mL) and were stimulated with different concentrations of ADP (0.25–2.0  $\mu$ mol/L). After 3 minutes incubation, 100  $\mu$ L 1% formaldehyde was added at room temperature. Platelets were then washed with phosphate-buffered saline and then diluted in 500  $\mu$ L phosphate-buffered saline for analysis on FACS Calibur (Becton Dickinson, Germany). Histograms were obtained from 5000 cells, and the data were analyzed using the software CellQuest Pro (Becton Dickinson). Mean fluorescence intensities (MFIs) vs cell number were expressed in a linear mode.


### Binding of [<sup>3</sup>H]PSB-0413 to washed platelets

Binding experiments were performed using [<sup>3</sup>H]PSB-0413, which is a tritiated derivative of a selective nucleotide antagonist of the P2Y<sub>12</sub>R, AR-C67085MX [2-propylthioadenosine-5'-adenylic acid (1,1-chloro-1-phosphonomethyl-1-phosphonyl)anhydride], and  $9 \times 10^7$  washed platelets.<sup>15</sup> Nonspecific binding was defined in the presence of 1 mM ADP. Washed platelets were incubated with the ligand at 37°C for 5 minutes; then bound and free radioactivity was separated by filtration through Whatman GF/B glass-fiber filters. Filters were then washed with  $5 \times 2$  mL of ice-cold washing buffer (Tris HCl, 50 mM, pH 7.5; EDTA, 1 mM; MgCl<sub>2</sub>, 5 mM; NaCl, 100 mM). Filter-bound radioactivity was counted in 2-mL liquid scintillation counter. Assays were performed in triplicate in 3 independent experiments, 2 in III-7 and 1 in III-1.

Competition experiments were performed in triplicate using a fixed concentration of [<sup>3</sup>H]PSB-0413 (200 nM) and increasing concentrations of ADP ( $10^{-10}$ – $10^{-2}$  M) or 2MeSADP ( $10^{-11}$ – $10^{-4}$  M) at 37°C for 5 minutes.

### Measurement of platelet cAMP

Platelet cAMP was measured by a radioisotopic assay, using a commercially available kit (Cyclic AMP [<sup>3</sup>H] assay system; Amersham International, UK). Duplicate samples of 1 mL citrated PRP containing 1 mM theophylline were incubated with Tyrode's buffer and PGE<sub>1</sub> (1  $\mu$ M), PGE<sub>1</sub> and ADP or epinephrine (0.1 and 1.0  $\mu$ M), or Tyrode's buffer alone in a control mixture. After incubation at 37°C (2 minutes), 1 mL of 5% trichloroacetic acid was added, and the samples were snap-frozen in dry ice and methanol, thawed at ambient temperature, and then shaken at 4°C for 45 minutes. After centrifugation at 4°C for 30 minutes, the supernatant was extracted 3 times with 5 mL of water-saturated ether, dried under a stream of nitrogen at 60°C, and stored at –20°C. Before assay, the samples were reconstituted with 0.05 mol/L Tris buffer containing 4 mmol/L EDTA.



### VASP phosphorylation analysis

The VASP phosphorylation analysis was performed immediately after blood collection, using Platelet VASP (Diagnostica Stago, Asnières, France) according to the manufacturer's instructions. Briefly, blood samples were incubated with PGE<sub>1</sub> and ADP, alone and in combination, for 10 minutes and fixed with paraformaldehyde. Platelets were then permeabilized with a nonionic detergent, labeled with a primary monoclonal antibody (16C2), followed by a secondary fluorescein isothiocyanate-conjugated polyclonal goat anti-mouse Ig antibody. Analyses were performed using a Becton Dickinson (Plymouth, UK) FACS Calibur flow cytometer. The platelet population was identified from its forward and side scatter distribution, and 10 000 platelets were gated. Platelet reactivity index (PRI) was calculated from the MFI, reflecting VASP phosphorylation, of samples incubated with PGE<sub>1</sub> or PGE<sub>1</sub> plus ADP, according to the following formula:  $PRI = [(MFI_{(PGE_1 + ADP)} - MFI_{(PGE_1)}) / MFI_{(PGE_1)}] \times 100$ . The analysis was completed in ~30 minutes following blood sampling.

### Genetic studies

Genomic DNA was isolated from peripheral blood lymphocytes using standard procedures. The entire coding sequence of the P2Y<sub>12</sub>R gene was amplified by a single polymerase chain reaction (PCR). PCR was performed using the primers (forward) 5'-CCTTAGGCTGAAAATAACCATCCTC3' and (reverse) 5'-GCGCTTTGCTTAAACGAGTTCTGAAC3', with DreamTaq Green Polymerase, 10× DreamTaq Green (Fermentas ThermoFisher Scientific Inc., Waltham, MA) and 2'-deoxynucleoside 5'-triphosphate mix (Invitrogen-Life Technologies Corp., Carlsbad, CA). The amplified fragments were subjected to DNA sequence analysis by ABI 3730xl 96-capillary DNA Analyzers (Eurofins MWG Operon, Huntsville, AL).

### Molecular modeling of the P2Y<sub>12</sub>R

The recently resolved crystallographic structures of the human P2Y<sub>12</sub>R in complex with the antagonist AZD1283 (Protein Data Bank ID: 4NTJ)<sup>12</sup> and with the full agonist 2MeSADP (Protein Data Bank ID: 4PXZ)<sup>13</sup> were used in this study. Models of the P2Y<sub>12</sub>R Gln-187 variant, based on the agonist-bound P2Y<sub>12</sub>R structure, were built by mutating His187 to Gln directly in the structure and then by modifying its conformation by selecting 5 different, suitable rotamers.


All P2Y<sub>12</sub>R structures were prepared using the Protein Preparation Wizard tool<sup>22</sup> implemented in the Schrödinger suite,<sup>23</sup> adding all the hydrogen atoms and the missing side chains of residues whose backbone coordinates were observed in the structure. The orientation of polar hydrogens was optimized, the protein protonation states were adjusted, and the overall structures were minimized with harmonic restraints on the heavy atoms, to remove strain. Then, all the heterogroups and water molecules were deleted.

The SiteMap tool of the Schrödinger suite was used to identify potential binding sites in the structures. Molecular docking of PSB-0413 at the agonist-bound P2Y<sub>12</sub>R crystal and at the P2Y<sub>12</sub> Gln-187 variant models was performed by means of the Glide package<sup>24</sup> from the Schrödinger suite.<sup>23</sup> In particular, a Glide Grid was centered on the centroid of residues located within 6 Å from the previously identified cavity. The Glide Grid was built using an inner box (ligand diameter midpoint box) of 14 Å × 14 Å × 14 Å and an outer box (box within which all the ligand atoms must be contained) that extended 20 Å in each direction from the inner one. Docking of the ligand was performed in the rigid binding sites using the XP (extra precision) procedure. The top-scoring docking poses of the ligand were subjected to visual inspection and analysis of protein-ligand interactions to select the final binding conformations.

## Results

### Studies in patient III-7 (propositus)

**Platelet aggregation and secretion.** The platelets from patient III-7 changed shape normally following stimulation with 1 μM ADP (not shown) and underwent aggregation in response to 4-20 μM ADP that was markedly lower than normal and rapidly reversible (Figure 2). The aggregation of these platelets in response to other platelet agonists was normal (not shown). Very similar results were obtained in experiments performed in Freiburg (Figure 2) and in Milan (not shown). The ATP secretion from patient platelets was absent after stimulation by ADP, and normal after stimulation by other agonists (Table 1).



**Figure 2.** ADP-induced platelet aggregation in PRP from a normal control and from patient III-7. Numbers below the aggregation tracings indicate the concentration of ADP added ( $\mu$ M). These experiments were performed using the light transmission aggregometer APACT 4 (Labor Fibrinometer).

**Binding of fluorescein-conjugated fibrinogen to ADP-activated platelets.** Compared with healthy controls, the amount of fluorescein-conjugated fibrinogen bound to platelets upon stimulation with ADP (0.25–2.0  $\mu$ mol/L) was severely impaired (Figure 3).

**Inhibition of PGE<sub>1</sub>-induced increase in platelet cAMP.** The basal level of platelet cAMP was normal (12.2 pmol/10<sup>9</sup> platelets; normal range: 6–17.9) and normally increased  $\sim$ 10-fold after stimulation with 1  $\mu$ M PGE<sub>1</sub>. ADP, in a concentration-dependent manner, inhibited the production of cAMP by normal platelets exposed to PGE<sub>1</sub>, but this response was greatly impaired in the patient's platelets (Table 2). In contrast to ADP, epinephrine normally inhibited the production of cAMP in the patient's platelets exposed to PGE<sub>1</sub> (Table 2), suggesting that the defect could be attributed to an abnormality in the platelet P2Y<sub>12</sub>R responsible for specifically coupling ADP stimulation to adenylyl cyclase inhibition.

**VASP phosphorylation assay.** The PRI value of patient's platelets was 30% in an experiment done in Milan and 1% in a subsequent experiment done in Freiburg, compared with  $78.7 \pm 6.7\%$  in platelets from 31 normal subjects.

**Binding of [<sup>3</sup>H]PSB-0413 to washed platelet suspensions.** The binding of [<sup>3</sup>H]PSB-0413 to patient platelets was measured in 2 separate experiments and compared with that observed in normal volunteers. The specific binding was saturable in both cases, and Scatchard plot analysis yielded a linear fit. The number of binding sites per platelet (B<sub>max</sub>) was comparable between normal and patient platelets (Table 3); however, the dissociation constant (K<sub>D</sub>) values were higher with patient platelets than with normal platelets (Table 3).

**Characterization of the P2Y<sub>12</sub>R gene.** The observation that the patient platelets had a normal number of binding sites for [<sup>3</sup>H]PSB-0413 in spite of a severely impaired function of the P2Y<sub>12</sub>R suggested that a dysfunctional receptor was being synthesized in normal amounts. To identify the underlying structural changes, we

analyzed the coding sequence of P2Y<sub>12</sub>R from DNA fragments generated in a PCR reaction. DNA from the patient showed a homozygous c.847T>A substitution that changed the codon for His-187 to Gln (p.His187Gln), located in the TM5 portion of the receptor.

### Family studies

For 11 of 20 first- and second-degree relatives of the proband, it was possible to retrieve information on the individual bleeding history and blood samples for DNA extraction and genotyping. Seven subjects (the grandparents, the parents, 2 sisters, and 1 brother) were heterozygous for p.His187Gln, 1 brother and 2 sisters had wild-type (WT) P2Y<sub>12</sub>R, and 1 brother had homozygous p.His187Gln. The bleeding history was negative in all subjects, except in 2 sisters (1 with heterozygous p.His187Gln P2Y<sub>12</sub>R, 1 with WT genotype) who had a mild bleeding history, and in the brother (III-1) with homozygous p.His187Gln P2Y<sub>12</sub>R, who had a severe bleeding history.

Patient III-1 underwent extensive platelet function studies in Strasbourg (France), including platelet aggregation both in PRP and washed platelet suspensions, inhibition of PGE<sub>1</sub>-induced increase in platelet cAMP levels by ADP, and VASP phosphorylation assay. The results of these tests were superimposable to those obtained in the proband and compatible with severe defect of P2Y<sub>12</sub>R function. The VASP-P PRI was 1% in 2 separate experiments, similarly to that obtained in patient III-7 in Freiburg; ADP did not inhibit PGE<sub>1</sub>-induced increase in cAMP up to a concentration of 100  $\mu$ M. Similarly to the proband, binding studies of [<sup>3</sup>H]PSB-0413 to his platelets revealed that the B<sub>max</sub> was normal (693 sites per platelet), whereas the K<sub>D</sub> value was higher than normal (19.0 nM) and comparable to the mean K<sub>D</sub> value calculated in the 2 experiments with the proband's platelets. The binding affinity (K<sub>i</sub>) values for both ADP and 2MeSADP for bound radioligand were higher in patient III-1 than in a healthy control (Figure 4).

Finally, the results of preliminary experiments using 1321N1 astrocytoma cells expressing p.His187Gln P2Y<sub>12</sub>R were similar to those obtained in patients' platelets: compared with 1321N1 astrocytoma cells expressing WT P2Y<sub>12</sub>R, we observed decreased affinity for [<sup>3</sup>H]PSB-0413 and a right shift of the displacement curves using ADP and 2MeSADP (data not shown).


**Table 1. Secretion of platelet ATP induced by different agonists in patient III-7**

Agonist	Released ATP (nmol/10 <sup>6</sup> platelets)		
	Patient III-7, experiment 1	Patient III-7, experiment 2	Normal range*
ADP (4 $\mu$ M)	Undetectable	Undetectable	0.02–0.98 (n = 96)
ADP (20 $\mu$ M)	Undetectable	Undetectable	0.04–0.61 (n = 59)
Collagen (2 $\mu$ g/mL)	0.46	0.47	0.17–0.93 (n = 62)
U46619 (0.5 $\mu$ M)	0.02	0.03	0.02–1.03 (n = 72)
TRAP-6 (10 $\mu$ M)	Not done	0.17	0.01–1.08 (n = 41)

PRP containing trisodium citrate as anticoagulant was stimulated by different agonists at the indicated concentrations.

TRAP-6, thrombin receptor activating peptide 6; U46619, thromboxane/prostaglandin endoperoxide analog 9,11-dideoxy-11,9-epoxymethano-prostaglandin F<sub>2</sub>.

\*The normal ranges are defined by the 2.5 and 97.5 percentiles of the distribution of values obtained in healthy subjects.



**Figure 3. Binding of fluorescein-conjugated fibrinogen to ADP-activated platelets from a healthy control and patient III-7.** Fluorescein-conjugated (FITC) fibrinogen binding to PRP stimulated with the indicated concentrations of ADP was measured by flow cytometry (see "Methods" for details). FITC, fluorescein isothiocyanate.

### Molecular modeling studies

To predict the implications of the p.His187Gln substitution at a molecular level, we performed molecular modeling studies of the P2Y<sub>12</sub>R. Two recently published crystallographic structures of the human P2Y<sub>12</sub>R,<sup>12,13</sup> in complex with both agonist and antagonist, resulted to be very useful for this study. Moreover, models of the P2Y<sub>12</sub>R p.His187Gln variant were built using the agonist-bound P2Y<sub>12</sub>R crystal structure, to analyze the possible structural differences because of this sequence variation.

TM5 in both P2Y<sub>12</sub>R crystal structures is unusually straight and tilted, because of the absence of proline (eg, position 5.50 following the Ballesteros-Weinstein numbering system)<sup>25</sup> or glycine residues in this helix in other G-protein-coupled receptors that destabilize its straight  $\alpha$ -helical conformation. Previous modeling attempts of the P2Y<sub>12</sub>R based on the available crystallographic templates are imprecise because of this difference.

In both structures, the point of mutation is located in the upper part of TM5 facing inward toward the binding site, in proximity of the ligand. Comparison of the 2 different crystallographic structures, agonist and antagonist bound, revealed large conformational changes occurring in TM6 and TM7 but also a slightly different orientation of TM5. Consequently, the conformations of the residues in this helix, including His187 (5.36), differ in the 2 structures. In the WT antagonist-bound P2Y<sub>12</sub>R structure, the side chain of His187 is more oriented toward TM4 and is delimiting the binding site making hydrophobic interactions with the ligand AZD1283. In the WT agonist-bound P2Y<sub>12</sub>R structure, the side chain of His187 (5.36) is directly interacting through an H-bond with the 2'-hydroxyl group of 2MeSADP (Figure 5). Moreover, the aromatic ring of this histidine is perpendicular to the adenine core and can stabilize it through hydrophobic interactions. Docking of PSB-0413 at this WT P2Y<sub>12</sub>R structure closely resembles the binding mode of 2MeSADP (Figure 5). In fact, the adenine bases of the 2 ligands occupy a common aromatic binding site, forming a  $\pi$ - $\pi$  interaction with Tyr105 (3.33) and H-bonds with Asn191 (5.40). The 2-propylthio group of PSB-0413 extends deeper in the same hydrophobic pocket occupied by the 2-methylthio group of the crystallized agonist and delimited by Phe106 (3.34), Met152 (4.53), Leu155 (4.56), Ser156 (4.57), Val190 (5.39), and Cys194 (5.43). The ribose 2'- and 3'-hydroxyl groups form H-bonding interactions with Lys179 (EL2) and His187 (5.36) and with Lys179 (EL2) and the backbone carbonyl of Cys97 (3.25), respectively. The noncleavable triphosphate-like chain of PSB-0413 is directed toward TM2 and forms numerous

hydrophilic and ionic interactions with Lys80 (2.60), Arg93 (3.21), Arg256 (6.55), Tyr259 (6.58), and Lys280 (7.35).

When docked at the P2Y<sub>12</sub>R p.His187Gln variant models, PSB-0413 showed the same orientation in the binding site and similar interactions as in the WT receptor (data not shown). However, depending on the rotamer of the side chain of Gln187, the H-bonding network involving the 2'-hydroxyl group can differ. In fact, in some rotamers Gln187 does not form any H-bond with the ligand, while in others it can act as H-bond acceptor or H-bond donor.

### Discussion

Congenital defects and pharmacologic inhibition of the P2Y<sub>12</sub>R for ADP are associated with an increased bleeding risk, demonstrating the important role of this receptor in platelet function and normal hemostasis.<sup>26</sup> Most of the congenital defects of P2Y<sub>12</sub>R characterized to date are associated with a marked decrease of the platelet binding sites for ADP, which has been attributed to mutations that disrupt the synthesis of the receptor,<sup>2,26-30</sup> to alterations in the ligand binding site,<sup>10</sup> or to compromised P2Y<sub>12</sub>R recycling.<sup>14</sup> A recently described patient displayed abnormalities in P2Y<sub>12</sub>R signaling, traffic, and surface expression, which was associated with the substitution of Arg122 with a cysteine.<sup>31</sup> In 1 kindred, a case of a normal expression of the receptor that was associated with severely defective signal transduction was described.<sup>7</sup> Nucleotide sequence analysis of the P2Y<sub>12</sub> gene revealed that the patient was a compound heterozygote for 2 missense mutations that caused the substitution p.Arg256Gln (in TM6) and p.Arg265Trp (in EL3), suggesting that the structural integrity of TM6 and EL3 regions is necessary for normal receptor function. A heterozygous mutation in the same region of the molecule, p.Pro258Thr, was later described in a patient with mild bleeding disorder and severely impaired ADP-induced platelet aggregation, confirming that this region of the molecule is important for receptor function.<sup>8</sup> The recently reported crystallographic data for the human P2Y<sub>12</sub>R support these findings revealing that significant conformational changes occur in the TM6-EL3 region between the agonist- and antagonist-bound structures.<sup>12,13</sup>

In the present report, we describe the case of a patient and his brother with lifelong bleeding disorder associated with defective P2Y<sub>12</sub>R-dependent platelet function, associated with normal expression of the receptor but decreased affinity for its ligand. Based on the results of the VASP phosphorylation assay<sup>6</sup> and on the observation that ADP, up to 100  $\mu$ M, failed to inhibit adenylyl cyclase, we may conclude that the function of the mutant receptor is completely suppressed. The moderately reduced PRI value (30%) initially measured by the VASP-P assay in patient III-7 was not reproduced in a second experiment


**Table 2. Effects of ADP and epinephrine on PGE<sub>1</sub> (1  $\mu$ M)-induced increase of cAMP levels in platelets from patient III-7 and healthy controls**

	cAMP (% of PGE <sub>1</sub> -induced)		
	Patient III-7, experiment 1	Patient III-7, experiment 2	Healthy controls* (n = 21)
ADP (0.1 $\mu$ M)	90.0	92.8	78.62 $\pm$ 11.58
ADP (1.0 $\mu$ M)	107.0	100.0	44.41 $\pm$ 11.88
Epinephrine (1 $\mu$ M)	22.0	26.0	30.36 $\pm$ 13.94

PRP containing trisodium citrate as anticoagulant was exposed to PGE<sub>1</sub> (1  $\mu$ M) in the presence or absence of ADP or epinephrine at the indicated concentrations for 2 minutes at 37°C (see "Methods" for details).

\*Means  $\pm$  SD.

**Figure 4.** Competition of ADP and 2MeSADP for binding of [<sup>3</sup>H]PSB-0413 to washed platelets from patient III-1 and a healthy control. Higher concentrations of either ADP or 2MeSADP were necessary to displace [<sup>3</sup>H]PSB-0413 from the patient's platelets, compared with control.




(PRI = 1%) and in 2 experiments performed in patient III-1, suggesting that the results of the first experiment were likely inaccurate.

The bleeding histories of our proband and his brother were clinically relevant and comparable to that of patients with severe P2Y<sub>12</sub>R deficiency, associated with homozygous base pair deletions in the open reading frame, resulting in frame shifts and premature truncation of the protein.<sup>5</sup> The study of many relatives of the proband, whose parents were first cousins, depicts a typical inheritance pattern of recessive autosomal diseases. The bleeding history of relatives with heterozygous

p.His187Gln P2Y<sub>12</sub>R was essentially absent, comparable to that of subjects with the WT receptor, confirming previous observations.<sup>2</sup>

In order to analyze the structural implications of the p.His187Gln mutation, molecular modeling studies were performed. Recently published crystallographic structures of the P2Y<sub>12</sub>R<sup>12,13</sup> revealed that His187 in TM5 is oriented toward the inside of the receptor within the ligand binding site. In particular, the 2MeSADP-bound structure showed a clear role of His187 (5.36) in agonist binding because of the formation of a H-bond with the ligand's 2'-hydroxyl



**Figure 5.** Binding site of the human P2Y<sub>12</sub>R. Comparison of the crystallographic pose of 2MeSADP (green carbons) (A) and the docking pose of PSB-0413 (cyan carbons) (B) at the WT agonist-bound P2Y<sub>12</sub>R crystal structure. Side chains of some amino acids important for ligand recognition are highlighted (gray carbon sticks). H-bonding and ionic interactions are pictured as red dotted lines. Nonpolar hydrogen atoms are not displayed. In panel A, ribbon regions corresponding to residues whose mutation is associated with a receptor dysfunction previously reported in patients are highlighted in pink and labeled with pink letters as follows: A, Lys174; B, Arg256; Pro258; and C, Arg265.

**Table 3. Binding studies of [<sup>3</sup>H]PSB-0413 on intact platelets from the proband (III-7) and his brother (III-1) and 10 healthy controls**

	Patient III-7, experiment 1	Patient III-7, experiment 2	Patient III-1	Healthy controls* (n = 10)
B <sub>max</sub> (sites per platelet)	285	436	693	425 ± 50
K <sub>D</sub> (nM)	8.4	33.0	19.0	3.3 ± 0.6

\*Means ± SD.

group and hydrophobic interactions with adenine. PSB-0413 docking predicted a binding mode similar to the crystallographic pose of the nucleotide agonist. Therefore, the predicted involvement of His187 (5.36) also in PSB-0413 binding can explain the decreased affinity of [<sup>3</sup>H]PSB-0413 for the patient's platelets, considering that the H187Q mutation can induce suboptimal H-bonding interactions with the ligand and attenuated stabilizing hydrophobic interactions. Moreover, the conformational changes highlighted by the comparison of the agonist- and antagonist-bound P2Y<sub>12</sub>R structures, involving mainly TM6 and TM7 and partially TM5, suggest that this region of the receptor is important for the activation process. Also, mutagenesis studies have shown the importance of residues located in the TM6-EL3 region for the functional integrity of the P2Y<sub>12</sub>R. Therefore, the p.His187Gln substitution could alter the interresidue interactions in this region as compared with WT, potentially interfering with conformational rearrangements occurring during the activation process.

In conclusion, His187 (5.36) is important for agonist and nucleotide antagonist (ie, PSB-0413) binding, as highlighted by crystallographic data and docking results, and located in a region undergoing conformational changes. Therefore, we have determined a structural context for the functional consequences of the p.His187Gln substitution in the P2Y<sub>12</sub>R.

## References

- Hollopeter G, Jantzen HM, Vincent D, et al. Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature*. 2001;409(6817):202-207.
- Cattaneo M. The platelet P2Y<sub>12</sub> receptor for adenosine diphosphate: congenital and drug-induced defects. *Blood*. 2011;117(7):2102-2112.
- Podda G, Femia EA, Pugliano M, Cattaneo M. Congenital defects of platelet function. *Platelets*. 2012;23(7):552-563.
- Cattaneo M. New P2Y<sub>12</sub> inhibitors. *Circulation*. 2010;121(1):171-179.
- Cattaneo M, Lecchi A, Randi AM, McGregor JL, Mannucci PM. Identification of a new congenital defect of platelet function characterized by severe impairment of platelet responses to adenosine diphosphate. *Blood*. 1992;80(11):2787-2796.
- Zighetti ML, Carpani G, Sinigaglia E, Cattaneo M. Usefulness of a flow cytometric analysis of intraplatelet VASP phosphorylation for the detection of patients with genetic defects of the platelet P2Y<sub>12</sub> receptor for ADP. *J Thromb Haemost*. 2010;8(10):2332-2334.
- Cattaneo M, Zighetti ML, Lombardi R, et al. Molecular bases of defective signal transduction in the platelet P2Y<sub>12</sub> receptor of a patient with congenital bleeding. *Proc Natl Acad Sci USA*. 2003;100(4):1978-1983.
- Remijn JA, IJsseldijk MJ, Strunk AL, et al. Novel molecular defect in the platelet ADP receptor P2Y<sub>12</sub> of a patient with haemorrhagic diathesis. *Clin Chem Lab Med*. 2007;45(2):187-189.
- Cattaneo M. Molecular defects of the platelet P2 receptors. *Purinergic Signal*. 2011;7(3):333-339.
- Daly ME, Dawood BB, Lester WA, et al. Identification and characterization of a novel P2Y<sub>12</sub> variant in a patient diagnosed with type 1 von Willebrand disease in the European MCMDM-1VWD study. *Blood*. 2009;113(17):4110-4113.
- Algaier I, Jakubowski JA, Asai F, von Kügelgen I. Interaction of the active metabolite of prasugrel, R-138727, with cysteine 97 and cysteine 175 of the human P2Y<sub>12</sub> receptor. *J Thromb Haemost*. 2008;6(11):1908-1914.
- Zhang K, Zhang J, Gao ZG, et al. Structure of the human P2Y<sub>12</sub> receptor in complex with an antithrombotic drug. *Nature*. 2014;509(7498):115-118.
- Zhang J, Zhang K, Gao ZG, et al. Agonist-bound structure of the human P2Y<sub>12</sub> receptor. *Nature*. 2014;509(7498):119-122.
- Nisar S, Daly ME, Federici AB, et al. An intact PDZ motif is essential for correct P2Y<sub>12</sub> purinoceptor traffic in human platelets. *Blood*. 2011;118(20):5641-5651.
- El-Tayeb A, Griessmeier KJ, Müller CE. Synthesis and preliminary evaluation of [<sup>3</sup>H]PSB-0413, a selective antagonist radioligand for platelet P2Y<sub>12</sub> receptors. *Bioorg Med Chem Lett*. 2005;15(24):5450-5452.
- Ohlmann P, Lecchi A, El-Tayeb A, Müller CE, Cattaneo M, Gachet C. The platelet P2Y<sub>12</sub> receptor under normal and pathological conditions. Assessment with the radiolabeled selective antagonist [<sup>3</sup>H]PSB-0413. *Purinergic Signal*. 2013;9(1):59-66.
- Molnar J, Lorand L. Studies on apyrases. *Arch Biochem Biophys*. 1961;93:353-363.
- Cattaneo M, Cerletti C, Harrison P, et al. Recommendations for the standardization of light transmission aggregometry: a consensus of the working party from the platelet physiology subcommittee of SSC/ISTH. *J Thromb Haemost*. 2013;11(6):1183-1189.
- Femia EA, Pugliano M, Podda G, Cattaneo M. Comparison of different procedures to prepare platelet-rich plasma for studies of platelet aggregation by light transmission aggregometry. *Platelets*. 2012;23(1):7-10.
- Cattaneo M, Lecchi A, Zighetti ML, Lussana F. Platelet aggregation studies: autologous platelet-poor plasma inhibits platelet aggregation when added to platelet-rich plasma to normalize platelet count. *Haematologica*. 2007;92(5):694-697.
- Mustard JF, Perry DW, Ardlie NG, Packham MA. Preparation of suspensions of washed platelets from humans. *Br J Haematol*. 1972;22(2):193-204.
- Sastry GM, Adzhigirey M, Day T, Annabhimoju R, Sherman W. Protein and ligand preparation: parameters, protocols, and influence on virtual screening enrichments. *J Comput Aided Mol Des*. 2013;27(3):221-234.
- Schrödinger Suite. New York, NY: Schrödinger, LLC; 2013.
- Friesner RA, Banks JL, Murphy RB, et al. Glide: a new approach for rapid, accurate docking and

## Acknowledgment

This work was supported by the National Institutes of Health National Institute of Diabetes and Digestive Kidney Diseases Intramural Research Program (K.A.J. and S.P.).

## Authorship

Contribution: A.L. performed platelet function studies, analysis of the data, and final approval of the manuscript; C.R. performed DNA analysis, analysis of the data, and final approval of the manuscript; S.P. performed molecular modeling, analysis of the data, and final approval of the manuscript; A.D. and L.N. performed platelet function studies, analysis of the data, and final approval of the manuscript; P.O. performed ligand binding studies, analysis of the data, and final approval of the manuscript; C.G. performed platelet function studies, ligand binding studies, analysis of the data, and final approval of the manuscript; K.A.J. performed molecular modeling, analysis of the data, drafting of part of the manuscript, and final approval of the manuscript; B.Z. performed clinical management of the patients, platelet function studies, analysis of the data, and final approval of the manuscript; and M.C. performed coordination and design of the study, analysis of the data, drafting of the manuscript and final approval of the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Marco Cattaneo, Unità di Medicina 3, Ospedale San Paolo – Università degli Studi di Milano, Via di Rudini 8 20142 Milano, Italy; e-mail: marco.cattaneo@unimi.it.

- scoring. 1. Method and assessment of docking accuracy. *J Med Chem.* 2004;47(7):1739-1749.
25. Ballesteros JA, Weinstein H. Integrated methods for the construction of three-dimensional models and computational probing of structure–function relations in G protein-coupled receptors. In: Sealfon SC, Conn PM, eds. *Methods in Neurosciences*. San Diego, CA: Academic Press; 1995:366-428.
26. Cattaneo M. Bleeding manifestations of congenital and drug-induced defects of the platelet P2Y<sub>12</sub> receptor for adenosine diphosphate. *Thromb Haemost.* 2011;105(suppl 1):S67-S74.
27. Nurden P, Savi P, Heilmann E, et al. An inherited bleeding disorder linked to a defective interaction between ADP and its receptor on platelets. Its influence on glycoprotein IIb-IIIa complex function. *J Clin Invest.* 1995;95(4):1612-1622.
28. Cattaneo M, Lecchi A, Lombardi R, Gachet C, Zighetti ML. Platelets from a patient heterozygous for the defect of P2CYC receptors for ADP have a secretion defect despite normal thromboxane A<sub>2</sub> production and normal granule stores: further evidence that some cases of platelet 'primary secretion defect' are heterozygous for a defect of P2CYC receptors. *Arterioscler Thromb Vasc Biol.* 2000;20(11):E101-E106.
29. Fontana G, Ware J, Cattaneo M. Haploinsufficiency of the platelet P2Y<sub>12</sub> gene in a family with congenital bleeding diathesis. *Haematologica.* 2009;94(4):581-584.
30. Shiraga M, Miyata S, Kato H, et al. Impaired platelet function in a patient with P2Y<sub>12</sub> deficiency caused by a mutation in the translation initiation codon. *J Thromb Haemost.* 2005;3(10):2315-2323.
31. Patel YM, Lordkipanidzé M, Lowe GC, et al. A novel mutation in the P2Y<sub>12</sub> receptor and a function-reducing polymorphism in protease-activated receptor 1 in a patient with chronic bleeding. *J Thromb Haemost.* 2014;12(5):716-725.



blood

2015 125: 1006-1013

doi:10.1182/blood-2013-07-517896 originally published  
online November 26, 2014

## Identification of a new dysfunctional platelet P2Y<sub>12</sub> receptor variant associated with bleeding diathesis

Anna Lecchi, Cristina Razzari, Silvia Paoletta, Arnaud Dupuis, Lea Nakamura, Philippe Ohlmann, Christian Gachet, Kenneth A. Jacobson, Barbara Zieger and Marco Cattaneo

---

Updated information and services can be found at:

<http://www.bloodjournal.org/content/125/6/1006.full.html>

Articles on similar topics can be found in the following Blood collections

[Platelets and Thrombopoiesis](#) (554 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:

[http://www.bloodjournal.org/site/misc/rights.xhtml#repub\\_requests](http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests)

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>