Marina Samoylova Nicola Piovella Michael Holynski Philippe Wilhelm Courteille Romain Bachelard

World Scientific Book - $9in \times 6in$

Paper-mh

World Scientific Book - $9in \times 6in$

Paper-mh

World Scientific Book - $9in \times 6in$

Paper-mh

Contents

Marina Samoylova Nicola Piovella Michael Holynski Philippe Wilhelm Courteille Romain Bachelard

1.	One-o	mensional photonic band gaps in optical lattices	7
	1.1	Introduction	7
	1.2	Modeling	10
		1.2.1 Scalar microscopic theory	11
		1.2.2 Transfer matrix formalism	22
		1.2.3 Vectorial microscopic theory	27
		1.2.4 Photonic band gaps for three-level atoms	30
	1.3	Measurement of a photonic band gap	35
		1.3.1 Description of the experiment	36
		1.3.2 Reaching the thick grating regime	37
		1.3.3 Interpretation of the observations	38
	1.4	Finite size effects	38
		1.4.1 Applicability of the TM formalism to experiment	39
		1.4.2 Suppression of absorption	42
		1.4.3 Modelling finite-size effects	43
	1.5	Conclusions and Perspectives	45
	.6	Appendix: Derivation of the Transfer Matrix T	47
	.7	Appendix: Vectorial model	50
	.8	Appendix: Vectorial field	56

vi

My Book Title

Bibliography

Chapter 1

One-dimensional photonic band gaps in optical lattices

1.1 Introduction

While the scattering of light by a single point-like particle (i.e., one much smaller than the wavelength of the light) is a well-known process, the scattering by a dense assembly of particles is a rich and still open field. The *cooperation* of particles in scattering of coherent light was first described by Dicke [Dicke (1954)], and subsequently termed Dicke superradiance: provided the density is significant, each particle reemits a wave that is in phase with those emitted by its neighbours, resulting in a constructive interference phenomenon.

The advent of laser cooling has led to the development of cold atomic physics and boosted the interest in coherent, superradiant phenomena. Low temperatures provide almost motionless particles thus avoiding Doppler broadening, an important contribution to decoherence, and facilitating cooperation between close particles. At even lower temperature, Bose-Einstein condensates correspond to a state in which all the particles phase match, with an overall very low momentum spread.

For these cold systems, the cooperativity corresponds to the capacity of the atoms to affect the radiation, i.e., for a high cooperativity each atom receives a significant quantity of radiation from the other atoms, in addition to the external laser source. High optical densities can, for example, give rise to inhomogeneities in the radiation pressure force acting on the atoms, resulting in distortion of the cloud [Walker *et al.* (1990); Sesko *et al.* (1991)]. The phenomenon of collectivity should be considered an asset as it is a coherent effect that can be controlled: for instance, inhomogeneities in the radiation pressure field have been used to define a new compression scheme [Khaykovich and Davidson (1999)].

Paper-mh

My Book Title

Among the collective effects of light scattering, *single photon* scattering by an ensemble of atoms is particularly fascinating. Superradiance of single photons is possible due to the quantum state of the system where each atom has a probability to absorb the unique photon present in the system, and the *synchronized* radiation of the atoms is at the origin of the superradiance. For example, in the case of a cloud of atoms prepared in the "timed Dicke" state, cooperativity makes the atoms emit a superradiant light pulse in a narrow forward cone [Scully and Svidzinsky (2009)]. This single-photon cooperative process also gives rise to a collective Lamb shift [Röhlsberger *et al.* (2010); Scully and Svidzinsky (2010); Keaveney *et al.* (2012)], as well as to other collective frequency shifts [Friedberg *et al.* (1973)].

Cooperative single photon scattering has been studied experimentally with ultra-cold atomic clouds [Bux *et al.* (2010); Bienaimé *et al.* (2010); Bender *et al.* (2010)], where cooperativity manifests in a collapse or an increase of the radiation pressure force acting on the center of mass [Courteille *et al.* (2010)]. This effect is particularly strong when resonance conditions known as Mie resonances are satisfied [Bachelard *et al.* (2012b)]. In general, both the scattered radiation intensity and the radiation pressure force contain signatures of collective scattering and each can serve as a powerful experimental tool for investigating the role of cooperativity in the atom-radiation interaction [Bachelard *et al.* (2012a); Bienaimé *et al.* (2013)].

The greatest potential of using ultra-cold atomic clouds to scatter light may reside in ordered configurations. A typical tool for analyzing periodic structures is Bragg scattering, the diffraction pattern resulting from the interference of the radiation emitted by a periodic array of scatterers excited by an incident light field. This has been used to study various periodic systems, such as crystals, molecules or even artificial photonic bandgap materials. Periodic lattice geometries can also be realized with ultra-cold atomic ensembles by confining the atoms within an optical standing wave. If the array is large enough the interference pattern is constructive only in the precisely defined directions that satisfy the Bragg condition. In this sense Bragg scattering is a particular case of cooperative scattering [Friedberg et al. (1973); Scully et al. (2006); Courteille et al. (2010)] from an ordered structure in the limit of small optical densities, where every photon is scattered once at most. In this limit, the Bragg radiation pattern actually corresponds to the structure factor of the atomic cloud.

One-dimensional photonic band gaps in optical lattices

Bragg scattering experiments using optical lattices have been performed (see Fig. 1.1) in the thin grating regime, where the optical density of the cloud is so low that multiple light scattering events between the stacked atomic layers are rare. In the thick grating regime, characterized by multiple reflections of the incident light, application of specific light frequencies or irradiation angles is expected to give rise to *photonic band gaps* (PBG) also known as *stop bands* or *forbidden bands*. Clearly the collective scattering picture conserves its validity in the optically dense regime of multiple scattering and cooperative effects are responsible for the formation of forbidden photonic bands [Samoylova *et al.* (2013)].




Fig. 1.1 Experimental scheme to study light scattering on one-dimensional optical lattices. A far red detuned retro-reflected *lattice laser* creates a standing wave which acts as a one-dimensional periodic trap for the atoms. A *probe laser* is used to probe the cooperative scattering properties of the lattice in the vicinity of the atomic resonance.

Photonic band gaps for atomic clouds in optical lattices have been predicted in several geometries using a variety of techniques. The transfer matrix (TM) formalism was initially used to calculate the photonic band structure of a 1D array of disk-shaped lattice sites [Deutsch *et al.* (1995)], similar to the situation described in Refs. [Slama *et al.* (2005a,b, 2006)]. This lattice geometry recently led to the first experimental observation of photonic band gaps in an optical lattice [Schilke *et al.* (2011, 2012a)].

In higher dimensions, the Bloch-Floquet model has been used to predict the occurrence of omnidirectional photonic band gaps, for either diamond lattices [Antezza and Castin (2009)], simple cubic or face-centered cubic lattices [Yu (2011)]. However, the generation of omnidirectional photonic bands in 3D optical lattices is technically challenging and has not yet been demonstrated in experiment.

The present paper is devoted to the theory of PBGs in one-dimensional

Paper-mh

My Book Title

optical lattices. It presents results obtained from a microscopic model that describes the collective scattering by a finite collection of point-like scatterers. They are compared to analytical results obtained from the TM formalism which treats scattering from perfect one-dimensional systems. In particular, we describe and discuss finite-size effects and effects due to disorder which the TM formalism cannot capture. Furthermore, we present a microscopic model treating light as a vector field, which is beyond the capability of the TM formalism. Another important part of this work concerns the possibility to manipulate the PBG by tuning the atom-radiation interaction using, for example, an external magnetic field.

Section 1.2 is devoted to the theories of one-dimensional PBG: In Section 1.2.1, we present a microscopic theory where atoms are described as point-like scatterers. We show that it efficiently captures the scattering properties of the optical lattice including the one-dimensional PBG. In Section 1.2.2, the TM formalism is introduced and derived directly from the microscopic theory. The agreements between these approaches, as well as the limitations of each, are discussed. Section 1.2.3 generalizes the microscopic theory by taking into account the vectorial nature of light. Section 1.2.4 completes the theoretical part with an extension of the theory of PBGs to three-level systems and a discussion of the potential role of electromagnetically induced transparency (EIT) schemes in PBGs.

In Section 1.3 we briefly present and discuss a recent and currently unique experiment which demonstrates a PBG in an optical lattice. Naturally, the experiment has peculiarities which are difficult or impossible to take into account in idealizing theoretical models, such as the TM formalism. Among them are disorder and finite-size effects. In Section 1.4 we discuss a number of these effects and show many of them are included in the microscopic theory.

1.2 Modeling

Following the pioneering work of [Deutsch *et al.* (1995)] and until recently 1D PBGs in optical lattices have only been studied using the TM formalism: the atomic layers are assimilated into a dielectric medium with an infinite radius. The density is generally assumed to be perfectly constant throughout the layer, or at least constant transversally and stepwise longitudinally. Although these approximations may seem to lack rigor, they

World Scientific Book - 9in x 6in

Paper-mh

One-dimensional photonic band gaps in optical lattices

have proved to be in good agreement with the experimental measurements of 1D PBG in optical lattices [Schilke *et al.* (2011)].

An alternative approach has been used to characterize the band structure of 3D optical lattices [van Coevorden *et al.* (1996); Antezza and Castin (2009)]. Working in the point-dipole approximation and decomposing the electric field into Bloch waves, Maxwell equations allow calculation of the dispersion relation for the propagation of light within the lattice as well as the local density of states. However, the expansion into Bloch waves is based on the assumption of an infinite lattice which is not appropriate for the optical lattices that are currently being realized experimentally.

Recently, a microscopic theory, inspired by theories of collective scattering by a collection of point-like scatterers [Courteille *et al.* (2010); Bienaimé *et al.* (2010); Bux *et al.* (2010); Bienaime *et al.* (2011)], was introduced by [Samoylova *et al.* (2013)] to describe PBGs. It was shown to successfully capture the photonic properties of optical lattices, such as spontaneous emission, Bragg scattering and PBG.

This section will be devoted to the introduction of this microscopic theory (Sec. 1.2.1), and its connection to TM formalism (Sec. 1.2.2). A formal bridge is demonstrated and numerical results of both theories are also compared with a particular focus on PBG properties.

1.2.1 Scalar microscopic theory

Model

Here a model that describes the interaction of a collection of static twolevel atoms with a scalar wave is presented. It accounts for the fact that each atom is affected by the radiation of all the other atoms and so it describes multiple scattering and collective effects. This model has been used to predict several features of collectivity in disordered clouds, such as the modification of the radiation pressure force [Courteille *et al.* (2010)], Mie scattering and resonances [Bender *et al.* (2010); Bachelard *et al.* (2012b)] or the collective Lamb shift [Bienaime *et al.* (2011)]. While the model is formally the same for ordered atoms, as first discussed in Ref. [Samoylova *et al.* (2013)], the scattering properties of optical lattices are very different.

The atomic cloud is described as a system of two-level (g and e) atoms, with resonant frequency ω_a and position \mathbf{r}_j , which are driven by a uniform laser beam with electric field amplitude E_0 , frequency ω_0 and wave vector

Paper-mh

My Book Title

 $\mathbf{k}_0 = (\omega_0/c) \hat{\mathbf{e}}_z$. The atom-laser interaction is described by the Hamiltonian:

$$\hat{H} = \frac{m\omega_0}{2} \sum_{j=1} \left[\hat{\sigma}_j e^{i(\Delta_0 t - \mathbf{k}_0 \cdot \mathbf{r}_j)} + \text{h.c.} \right]$$

$$+ \hbar \sum_{j=1}^N \sum_{\mathbf{k}} g_k \left(\hat{\sigma}_j e^{-i\omega_a t} + \hat{\sigma}_j^{\dagger} e^{i\omega_a t} \right) \left[\hat{a}_{\mathbf{k}}^{\dagger} e^{i(\omega_k t - \mathbf{k} \cdot \mathbf{r}_j)} + \hat{a}_{\mathbf{k}} e^{-i(\omega_k t - \mathbf{k} \cdot \mathbf{r}_j)} \right],$$

$$(1.1)$$

where $\Omega_0 = dE_0/\hbar$ is the Rabi frequency of the incident laser field and $\Delta_0 = \omega_0 - \omega_a$ is the detuning between the laser and the atomic transition. For simplicity scalar light is considered in (1.1), with a vectorial form of the model being discussed later in Sec. 1.2.3. The case of three-level atoms is discussed in Sec. 1.2.4.

In (1.1), $\hat{\sigma}_j = |g_j\rangle\langle e_j|$ is the lowering operator for the *j*th atom, $\hat{a}_{\mathbf{k}}$ is the photon annihilation operator and $g_k = (d^2\omega_a^2/2\hbar\epsilon_0\omega_k V_\nu)^{1/2}$ is the single-photon Rabi frequency, where *d* is the electric-dipole transition matrix element and V_ν is the photon volume. The special case of a low-intensity laser, where a single photon from mode \mathbf{k} can be assumed to be present in the system, was extensively investigated in Refs. [Friedberg *et al.* (1973); Scully *et al.* (2006); Svidzinsky *et al.* (2008)]. In this case the system (atoms+photons) is described by a state of the form [Svidzinsky *et al.* (2010)]:

$$|\Psi\rangle = \alpha(t)|g_1\dots g_N\rangle|0\rangle_{\mathbf{k}} + e^{-i\Delta_0 t}\sum_{j=1}^N \beta_j(t)|g_1\dots e_j\dots g_N\rangle|0\rangle_{\mathbf{k}}$$
(1.2)

+
$$\sum_{\mathbf{k}} \gamma_{\mathbf{k}}(t) |g_1 \dots g_N\rangle |1\rangle_{\mathbf{k}} + \sum_{\mathbf{k}} \sum_{j \neq m} \epsilon_{j,m,\mathbf{k}}(t) |g_1 \dots g_j \dots g_m \dots g_N\rangle |1\rangle_{\mathbf{k}}.$$

The first term in (1.2) corresponds to the initial ground state without photons, the second term is the sum over the states where a single atom has been excited by the classical field, and the third term corresponds to the atoms that returned to the ground state having emitted a photon in the mode **k**. Hence, the square modulus of α , β_j and $\gamma_{\mathbf{k}}$ represent respectively the probability that the is no photon in the system and no excited atom, the probability that atom j is excited (and no photon), and the probability to have a photon in mode **k** (and all atoms in the ground state). Finally, $|\epsilon_{j,m,\mathbf{k}}|^2$ describes the probability of having two excited atoms and one virtual photon in mode **k** with 'negative' energy. This last term of (1.2) is due to the counter-rotating terms in the Hamiltonian (1.1) and disappears when the rotating wave approximation is made. The scattering process using the latter approximation was studied in several references [Mazets and Kurizki (2007); Courteille *et al.* (2010); Bienaimé *et al.*

World Scientific Book - 9in x 6in

Paper-mh

One-dimensional photonic band gaps in optical lattices

(2010); Skipetrov and Goetchy (2011); Bachelard *et al.* (2011)], but the importance of counter-rotating terms during the relaxation process was pointed out in Ref. [Manassah (2012a)].

The dynamics of each component of the state (1.2) are deduced from the Schrödinger equation:

$$\frac{\partial |\Psi(t)\rangle}{\partial t} = -\frac{i}{\hbar} \hat{H} |\Psi(t)\rangle.$$
(1.3)

Hence, the Hamiltonian (1.1) leads to the following set of differential equations:

$$\dot{\alpha}(t) = -\frac{i}{2}\Omega_0 e^{i\Delta_0 t} \sum_{j=1}^N \beta_j(t) e^{-i\mathbf{k}_0 \cdot \mathbf{r}_j} , \qquad (1.4)$$

$$\dot{\beta}_{j}(t) = i\Delta_{0}\beta_{j} - \frac{i}{2}\Omega_{0}\alpha(t)e^{-i\Delta_{0}t + i\mathbf{k}_{0}\cdot\mathbf{r}_{j}} - i\sum_{\mathbf{k}}g_{k}\gamma_{\mathbf{k}}(t)e^{-i(\omega_{k}-\omega_{0})t + i\mathbf{k}\cdot\mathbf{r}_{j}}$$

$$-i\sum_{\mathbf{k}}g_{k}e^{-i(\omega_{k}+\omega_{a}-\Delta_{0})t}\sum_{m\neq j}\epsilon_{j,m,\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}_{j}} , \qquad (1.5)$$

$$\dot{\gamma}_{\mathbf{k}}(t) = -ig_k e^{i(\omega_k - \omega_0)t} \sum_{j=1}^N \beta_j(t) e^{-i\mathbf{k}\cdot\mathbf{r}_j} .$$
(1.6)

$$\dot{\epsilon}_{j,m,\mathbf{k}}(t) = -ig_k e^{i(\omega_k + \omega_a - \Delta_0)t} \left[e^{-i\mathbf{k}\cdot\mathbf{r}_j}\beta_m + e^{-i\mathbf{k}\cdot\mathbf{r}_m}\beta_j \right].$$
(1.7)

Then integrating Eqs. (1.6) and (1.7) over time with $\gamma_{\mathbf{k}}(0) = 0$ and $\epsilon_{j,m,\mathbf{k}}(0) = 0$ and substituting them in to (1.5), we obtain N coupled equations describing the dynamics of the atomic dipoles:

$$\dot{\beta}_{j}(t) = i\Delta_{0}\beta_{j} - \frac{i}{2}\Omega_{0}\alpha e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}$$

$$-\sum_{\mathbf{k}}g_{k}^{2}\sum_{m=1}^{N}e^{i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}\int_{0}^{t}e^{-i(\omega_{k}-\omega_{0})(t-t')}\beta_{m}(t')dt'$$

$$-\sum_{\mathbf{k}}g_{k}^{2}\int_{0}^{t}dt'e^{i(\Delta_{0}-\omega_{k}-\omega_{a})(t-t')}$$

$$\times \left[\sum_{m\neq j}e^{-i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}\beta_{m}(t') + (N-1)\beta_{j}(t')\right].$$

$$(1.8)$$

The interaction with the vacuum field yields diagonal terms with m = j whose real part corresponds to the single-atom decay term and imaginary part to the self-energy shift, and off-diagonal terms with $m \neq j$ which correspond to the atom-atom interaction mediated by the photon. By separating

14

My Book Title

the two contributions, we can write:

$$\dot{\beta}_{j}(t) = i\Delta_{0}\beta_{j} - \frac{i}{2}\Omega_{0}\alpha e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}$$

$$-\sum_{\mathbf{k}}g_{k}^{2}\int_{0}^{t}d\tau \left[e^{-i(\omega_{k}-\omega_{0})\tau} + e^{i(\Delta_{0}-\omega_{k}-\omega_{a})\tau}(N-1)\right]\beta_{j}(t-\tau)$$

$$-\sum_{\mathbf{k}}g_{k}^{2}\sum_{m\neq j}e^{i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}$$

$$\times \int_{0}^{t}d\tau \left[e^{-i(\omega_{k}-\omega_{0})\tau} + e^{i(\Delta_{0}-\omega_{k}-\omega_{a})\tau}\right]\beta_{m}(t-\tau).$$
(1.10)

We assume a smooth density of modes, so the discrete sum $\sum_{\mathbf{k}}$ can be replaced by the volume integral $[V_{\nu}/(2\pi)^3] \int d\mathbf{k}$. In the linear regime $\alpha \approx 1$ and in the Markov approximation, valid if the decay time is larger than the photon time-of-flight through the atomic cloud, in the integrals of (1.10) we can replace $\beta_j(t-\tau)$ with $\beta_j(t)$ and extend the upper integration limit to infinity, so that (1.10) is approximated by

$$\dot{\beta}_{j}(t) = i\Delta_{0}\beta_{j} - \frac{i}{2}\Omega_{0}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}} - \frac{V_{\nu}}{(2\pi)^{3}}\int d\mathbf{k}g_{k}^{2}\int_{0}^{\infty}d\tau \left[e^{-i(\omega_{k}-\omega_{0})\tau} + e^{-i(\omega_{k}+2\omega_{a}-\omega_{0})\tau}(N-1)\right]\beta_{j}(t) - \frac{V_{\nu}}{(2\pi)^{3}}\int d\mathbf{k}g_{k}^{2}\sum_{m\neq j}e^{i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})} \times \int_{0}^{\infty}d\tau \left[e^{-i(\omega_{k}-\omega_{0})\tau} + e^{-i(\omega_{k}+2\omega_{a}-\omega_{0})\tau}\right]\beta_{m}(t).$$
(1.11)

The imaginary part of the self-field contribution (third term of the right hand side of (1.11)) consists of the self-energy shift of the atom in the excited state plus the self-energy contribution of the atom in the ground state. The effect is an adjustment to the transition frequency ω_a , which is assumed to already include the shift. It can not be computed realistically using our model, which treats the atoms as point particles, and is disregarded in the present approach. After performing the integration over τ the real part of the self-field contribution yields:

$$\frac{V_{\nu}}{2\pi} \int_0^\infty dk k^2 g_k^2 \left[\delta(\omega_k - \omega_0) + (N - 1)\delta(\omega_k + 2\omega_a - \omega_0) \right] = \frac{\Gamma}{2}, \quad (1.12)$$

where $\omega_k = ck$ and $\Gamma = d^2 \omega_0^3 / (2\pi c^3 \hbar \epsilon_0)$ is the single-atom spontaneous decay rate in the radiation scalar theory. The last term in (1.12) arises from the counter-rotating wave terms of the Hamiltonian (1.1) and does

One-dimensional photonic band gaps in optical lattices

not contribute as it corresponds to a negative photon energy $\omega_k \approx -\omega_a$. Using (1.12) in (1.11) we obtain:

$$\dot{\beta}_{j}(t) = \left(i\Delta_{0} - \frac{\Gamma}{2}\right)\beta_{j} - \frac{i}{2}\Omega_{0}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}} - \frac{V_{\nu}}{(2\pi)^{3}}\int d\mathbf{k}g_{k}^{2}\sum_{m\neq j}e^{i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}$$
$$\times \int_{0}^{\infty}d\tau \left[e^{-i(\omega_{k}-\omega_{0})\tau} + e^{-i(\omega_{k}+2\omega_{a}-\omega_{0})\tau}\right]\beta_{m}(t).$$
(1.13)

The last term on the right hand side of (1.13) can be calculated as follows:

$$\frac{V_{\nu}}{(2\pi)^3} \int d\mathbf{k} g_k^2 e^{i\mathbf{k}\cdot(\mathbf{r}_j-\mathbf{r}_m)} \int_0^\infty d\tau \left[e^{-i(\omega_k-\omega_0)\tau} + e^{-i(\omega_k+2\omega_a-\omega_0)\tau} \right] \\
= \frac{c\Gamma}{\pi k_0} \int_0^\infty d\tau \cos(\omega_0\tau) \int_0^\infty dk k \frac{\sin(kr_{jm})}{kr_{jm}} e^{-ick\tau},$$
(1.14)

where $r_{jm} = |\mathbf{r}_j - \mathbf{r}_m|$ and we assumed $\omega_0 \approx \omega_a$. We use the integral expression

$$\int_0^\infty dkk \frac{\sin(kr)}{kr} e^{-ick\tau} = \frac{1}{2cr} \left[\frac{1}{\tau + r/c - i\epsilon} - \frac{1}{\tau - r/c - i\epsilon} \right]$$

where $\epsilon \to 0^+$. Changing the sign of the integration variable τ in the first term, (1.14) becomes

$$-\frac{\Gamma}{2\pi k_0 r_{jm}} \int_{-\infty}^{\infty} d\tau \frac{\cos(\omega_0 \tau)}{\tau - r_{jm}/c + i\epsilon} = \frac{\Gamma}{2} \frac{\exp(ik_0 r_{jm})}{ik_0 r_{jm}}.$$
 (1.15)

Finally, the scattering problem reduces to the set of differential equations [Scully and Svidzinsky (2009); Bachelard *et al.* (2011); Bienaime *et al.* (2011)]:

$$\dot{\beta}_j = \left(i\Delta_0 - \frac{\Gamma}{2}\right)\beta_j - i\frac{\Omega_0}{2}e^{i\mathbf{k}_0\cdot\mathbf{r}_j} - \frac{\Gamma}{2}\sum_{m\neq j}\frac{\exp(ik_0|\mathbf{r}_j - \mathbf{r}_m|)}{ik_0|\mathbf{r}_j - \mathbf{r}_m|}\beta_m(1.16)$$

The kernel in the last term of (1.16) has a real component, $-(\Gamma/2) \sum_{m\neq j} \operatorname{sinc}(k_0 |\mathbf{r}_j - \mathbf{r}_m|)$, describing the *collective* atomic decay, and an imaginary component, $i(\Gamma/2) \sum_{m\neq j} \cos(k_0 |\mathbf{r}_j - \mathbf{r}_m|)/(k_0 |\mathbf{r}_j - \mathbf{r}_m|)$, which contains the collective Lamb shift due to short range interactions between atoms induced by the electromagnetic field [Scully and Svidzinsky (2009, 2010); Röhlsberger *et al.* (2010); Keaveney *et al.* (2012)].

In what follows we will focus on the stationary state of (1.16), which reads

$$(2\Delta_0 + i\Gamma)\beta_j = \Omega_0 e^{i\mathbf{k}_0 \cdot \mathbf{r}_j} - \Gamma \sum_{m \neq j} \frac{\exp(ik_0|\mathbf{r}_j - \mathbf{r}_m|)}{k_0|\mathbf{r}_j - \mathbf{r}_m|}\beta_m.$$
(1.17)

Paper-mh

My Book Title

Practically, the value of the atomic dipoles β_j in the stationary regime is evaluated numerically by inverting the linear problem (1.17), since it is easily cast in the form $M\vec{\beta} = e^{i\mathbf{k}_0\cdot\vec{\mathbf{r}}}$, where $\vec{\beta}$ and $\vec{\mathbf{r}}$ refer to the vectors of β_j and \mathbf{r}_j . From the β_j , the observables described in the next sections are easily calculated.

While we adopted a quantum, single-photon treatment to derive a description of the collective scattering a classical interpretation is also possible. In fact, (1.16) also describes the dynamics of atomic dipoles driven by the *total* electric field, the sum of the incident field and of the field radiated by the other dipoles. Indeed, the last term of ((1.16)) gives the emission of spherical waves by the latter, as described by Huygens principle. Furthermore, model (1.16) has also been derived through another classical approach which treats the two-level atoms as weakly excited classical harmonic oscillators [Svidzinsky *et al.* (2010); Prasad and Glauber (2010)].

Radiation field

The scattered radiation is provided by the positive-frequency component of the electric field

$$E_S(\mathbf{r}, t) = \sum_{\mathbf{k}} \mathcal{E}_k \gamma_{\mathbf{k}}(t) e^{i\mathbf{k}\cdot\mathbf{r} - i\omega_k t}, \qquad (1.18)$$

where $\mathcal{E}_k = (\hbar \omega_k / 2\epsilon_0 V_{\nu})^{1/2}$ is the single-photon electric field. We then integrate (1.6) over time, with $\gamma_{\mathbf{k}}(0) = 0$, insert it in to (1.18) and obtain

$$E_{S}(\mathbf{r},t) = -\frac{iV_{\nu}}{8\pi^{3}}e^{-i\omega_{0}t}\sum_{j=1}^{N}\int d\mathbf{k}\,\mathcal{E}_{k}g_{k}\int_{0}^{t}dt'e^{-i(\omega_{k}-\omega_{0})t'}e^{i\mathbf{k}\cdot(\mathbf{r}-\mathbf{r}_{j})}\beta_{j}(t-t').$$
(1.19)

Taking $d\mathbf{k} = k^2 dk \sin\theta d\phi d\theta$ and performing the integrations over θ and ϕ , (1.19) becomes

$$E_S(\mathbf{r},t) = -\frac{id\omega_0}{4\pi^2\epsilon_0}e^{-i\omega_0t}\sum_{j=1}^N\int_0^t dt'\beta_j(t-t')\int_0^\infty dk\,k\,\frac{\sin(k|\mathbf{r}-\mathbf{r}_j|)}{|\mathbf{r}-\mathbf{r}_j|}e^{-i(\omega_k-\omega_0)t'}$$
(1.20)

The scattered intensity will be centered about the incidence laser frequency ω_0 . The quantity $\omega_k = ck$ varies little around $\omega_k = \omega_0$ for which the time integral in (1.20) is not negligible. We can therefore replace k by ω_0/c and the lower limit in the k integration by $-\infty$. The integral

$$\int_{-\infty}^{\infty} dk \sin(kR) e^{-ic(k-k_0)t'} = \frac{\pi}{ic} \left[e^{ik_0 R} \delta(t-R/c) - e^{-ik_0 R} \delta(t+R/c) \right]$$
(1.21)

One-dimensional photonic band gaps in optical lattices

yields

$$E_S(\mathbf{r},t) = -\frac{dk_0^2}{4\pi\epsilon_0} \sum_{j=1}^N \frac{e^{ik_0|\mathbf{r}-\mathbf{r}_j|}}{|\mathbf{r}-\mathbf{r}_j|} \beta_j(t-|\mathbf{r}-\mathbf{r}_j|/c).$$
(1.22)

The delay in β_j can be neglected, as previously assumed in (1.10), if the photon time-of-flight $|\mathbf{r} - \mathbf{r}_j|/c$ is much smaller than the characteristic time during which the atomic variables change appreciably, obtaining

$$E_S(\mathbf{r},t) = -\frac{dk_0^2}{4\pi\epsilon_0} \sum_{j=1}^N \frac{e^{ik_0|\mathbf{r}-\mathbf{r}_j|}}{|\mathbf{r}-\mathbf{r}_j|} \beta_j(t).$$
(1.23)

At distances r much larger than the cloud, a far-field expression can be derived. Using $|\mathbf{r} - \mathbf{r}_j| \approx r - \hat{\mathbf{n}} \cdot \mathbf{r}_j$, where $\hat{\mathbf{n}} = \mathbf{r}/r$, (1.23) turns into

$$E_S(\mathbf{k}) \approx -\frac{dk_0^2}{4\pi\epsilon_0} \frac{e^{ik_0r}}{r} \sum_{j=1}^N \beta_j(t) e^{-i\mathbf{k}\cdot\mathbf{r}_j},\qquad(1.24)$$

where $\mathbf{k} = k_0 \hat{\mathbf{n}}$. Notice that with (1.23) we recover the field present in (1.17) apart from the self-contribution of *j*th-atom to the field. Thus, the excitation of the atomic dipole in (1.17) can be rewritten as

$$\beta_j = \frac{d}{\hbar \left(\Delta_0 + \Gamma/2\right)} E_{tot\backslash j}(\mathbf{r}_j), \qquad (1.25)$$

where $E_{tot\backslash j}(\mathbf{r}_j) = (E_0/2) \exp(i\mathbf{k}_0 \cdot \mathbf{r}_j) + E_S(\mathbf{r}_j) - E_{self}(\mathbf{r}_j)$ is the total electric field minus the atom self-contribution. Hence, (1.17) describes the light response of a set of point-like dielectric particles (in the linear optics regime), where the self-field contributions are treated separately to avoid singularities.

Measurements of the atomic density distribution after a time of free expansion yield the atomic momentum distribution after interaction with the light beam. The momentum distribution, which is modified by the recoil imparted to the atomic cloud due to the radiation pressure force, contains important information on the light scattering in disordered systems [Courteille et al. (2010); Bachelard et al. (2012a); Piovella et al. (2013)]. The strong relationships between scattered light and radiation pressure force are a manifestation of collective effects induced by scattered photons and described in the present model [Bienaimé et al. (2013)]. Until now we assumed free atoms. However, in optical lattices the laser beams used to create the lattice potential also generate a strong confining force localizing the atoms within a regime smaller than the Lamb Dicke limit. so that the simple link between scattered light and radiation pressure force is broken. Other effects which take into account the atomic trapping need to be included in order to access more exhaustive information revealed by time-of-flight measurements.

Paper-mh

My Book Title

Bragg scattering

Bragg scattering is the reflection of light by a periodic structure due to constructive interference. This phenomenon has been used to probe atomic structure [Wollan (1932)], molecular dynamics [Doucet and Benoit (1987)], optical lattices [Birkl *et al.* (1995); Weidemüller *et al.* (1995, 1998)], and photonic bands in photonic crystals [Koenderink and Vos (2003)].

In a one-dimensional structure of period d a wave with wavelength λ_0 is reflected provided its incident angle θ_0 (with respect to the structure axis) satisfies the Bragg condition:

$$\sin \theta_0 = n \frac{\lambda_0}{2d},\tag{1.26}$$

where n is an integer. The condition (1.26) is actually that required to provide constructive interference from the scatterers. In the case of an atomic lattice dense enough to modify the phase of the propagating wave, a correction is necessary to account for the medium effective index m [Deutsch *et al.* (1995)], and the Bragg condition turns into

$$\sin \theta_0 = n \frac{\lambda_0}{2md}.\tag{1.27}$$

Model (1.17) naturally describes Bragg scattering since the radiation field (1.22) takes into account the interference from the atomic scatterers. Yet the collective term in (1.17) which describes the radiation from the neighbour is unimportant as far as Bragg scattering is concerned. Let us consider the radiation of the lattice without the collectivity. Then (1.17)simply turns into

$$\beta_j = \Omega_0 \frac{e^{i\mathbf{k}_0 \cdot \mathbf{r}_j}}{(2\Delta_0 + i\Gamma)}.$$
(1.28)

Each atomic dipole follows the laser only, from which follows a state that is similar to the mean-field ansatz called the timed Dicke-state [Scully *et al.* (2006)]. The many-body aspect of the problem only appears as the field radiated by the atoms interferes in (1.22), but this interference is sufficient to modify the structure factor of the atomic lattice and reflect the incoming wave. This effect is described by Rayleigh scattering, where the synchronization of the atomic dipoles by coherent light results in longrange correlations and in an emission of light in the Bragg directions for large clouds. However, here the lattice period constrains the light emission to specific directions.

Bragg scattering of a Gaussian beam using models (1.28) and (1.17) is displayed in Fig. 1.2. Despite both describing the light reflection, model

World Scientific Book - $9in \times 6in$

Paper-mh

19

One-dimensional photonic band gaps in optical lattices

(1.28) does not conserve energy, nor is it able to account for the progressive attenuation of the light in the lattice as the photons are reflected. This highlights the necessity of including collective terms, such as those in (1.17), when investigating PBGs. We point out that the study of photonic band




Fig. 1.2 Bragg reflection of a Gaussian beam by a one-dimensional optical lattice for models (1.28) neglecting collective (multiple) scattering [top] and (1.17) including collective scattering [bottom]. The laser beam has a waist $4.5\lambda_0$, a detuning $\Delta_0 = \Gamma$, and arrives from the top left with angle $\theta_0 = 0.2$ rad. The lattice is composed of 5000 atoms spread over 60 disks of radius $9\lambda_0$ and thickness $0.04\lambda_0$, separated by $d = \lambda_0/2\cos\theta_0$. The rectangles mark the limit of the atomic lattice.

gaps on a finite cloud requires a finite beam, thus the Gaussian beam used above. Indeed if one considers the scattering of a plane-wave on a finite lattice, the effect of the cloud in the far-field limit becomes negligible because of diffraction.

Paper-mh

My Book Title

One-dimensional photonic band gap

A PBG corresponds to a range of light frequencies which cannot propagate through the medium, and instead are reflected. As discussed above accounting for multiple scattering, and thus collective effects, is necessary to describe the progressive reflection of the light as it penetrates the lattice. We here show how one-dimensional PBGs are described by the collective model (1.17).

As shown in Fig. 1.3, the microscopic model (1.17) provides a good description of the reflection of the light. Aside from a spontaneous emission contribution that is emitted in all directions, most of the light is reflected and almost no light propagates in transmission behind the lattice. Note that only an infinite lattice is actually able to completely stop the light.

The capacity for a given wavelength to propagate in a medium is quantified by the local density of states (LDOS). In the case of one-dimensional systems the LDOS at the center of the lattice can be conveniently calculated using the complex reflection coefficients $r_{-,+}$ corresponding to the reflection from the two halves of the lattice, i.e., from the lattice beginning to the center, and from the center to the end [Boedecker and Henkel (2003)]:

$$LDOS = \operatorname{Re}\left(\frac{2+r_{-}+r_{+}}{1-r_{-}r_{+}}-1\right).$$
(1.29)


The complex reflection coefficient $r_{-} = \sqrt{R_{-}}e^{i\phi}$ is computed numerically using the reflectivity R_{-} of the first semi-lattice, i.e., the ratio of the reflected to the incident power, and the phase ϕ of the wave reflected at the origin of the lattice.

The quantity of light reflected by the lattice naturally increases with the number of atomic layers, as can be observed in Fig. 1.4. Furthermore, the larger the detuning the weaker the interaction between the light and each layer, and therefore the longer the lattice needs be to efficiently reflect the incoming light. In particular, larger detunings allow for a reduced spontaneous emission since the imaginary part of the atomic polarizability is reduced, yet this requires lattice lengths beyond what can be feasibly be simulated numerically using the model (1.17). Consequently in what follows we focus on resonant or near-resonant light.

Finally, by tuning the laser wavelength around the atomic transition and calculating the one-dimensional LDOS (1.29), the opening of forbidden bands in the one-dimensional lattice is observed in Fig. 1.5. In fact, we observe a lowering of the LDOS to values of ~ 0.1 over a range of a few Γ s.

World Scientific Book - 9in x 6in

Paper-mh




One-dimensional photonic band gaps in optical lattices

Fig. 1.3 Top: Intensity of the light in the y = 0 plane as it enters a one-dimensional optical lattice and its reflection. The rectangle marks the limit of the atomic structure. Bottom: Zoom of the leftmost region of the atomic lattice. The luminous grains correspond to the strong field radiated by the atoms close to the y = 0 plane. The simulations are realized for N = 9000 atoms randomly distributed over $N_d = 100$ layers of thickness $a = 0.06\lambda_0$ and radius $R = 9\lambda_0$, the distance between the atomic disks is $d = 0.508\lambda_0$ with λ_0 being the resonance wavelength. The input Gaussian beam has waist $4.5\lambda_0$ and power 100 mW with a detuning $\Delta_0 = \Gamma$ and is incident at an angle $\theta_0 = 0.2$ rad with respect to the lattice axis.

This value of the LDOS is in agreement with the measurements of Schilke and collaborators [Schilke *et al.* (2011)], although in that experiment the band was a dozen Γ s large due to the larger length of the lattice. Again, only an infinite lattice is able to support a perfect band gap with D = 0.

In conclusion, the microscopic model (1.17) captures well onedimensional forbidden bands: it describes the finite penetration in the lattice, the reflection of the wave and the vanishing LDOS as the lattice size increases.

Paper-mh



My Book Title

Fig. 1.4 Reflection coefficient R (left) and LDOS (right) as a function of the number of atoms in the lattice, for detunings $\delta = 0, 2, 4, 6$ and 8. The one-dimensional lattice is composed of N_d disks with 60 atoms each, with spacing $d = \lambda_0/2$, disk radius $9\lambda_0$ and thickness $0.04\lambda_0$. The laser waist is $4.5\lambda_0$.

1.2.2 Transfer matrix formalism


It is customary to study the photonic properties of one-dimensional optical lattices using the transfer matrix (TM) formalism [Deutsch *et al.* (1995)]. One of the foundation hypotheses of the TM theory is that the atomic cloud is assimilated in to a dielectric medium. Moreover, each lattice site is assumed to be a radially unlimited slice called a *slab*. This allows one to reduce the light scattering to the one-dimensional propagation of a plane wave, where the coupled evolution of the forward and backward waves is captured in an iterative 2×2 matrix problem.

In this section we establish a formal link between the microscopic model (1.16) and the TM approach, providing more details of the derivation presented in [Samoylova *et al.* (2013)]. The TM results will also be used as a benchmark to understand the signatures of finite-size effects and atomic disorder in later sections.

Introducing the normalized detuning $\delta = \Delta_0 / \Gamma$, ((1.16)) can be rewritten as:

$$\frac{\Omega_0}{\Gamma}e^{i\mathbf{k}_0\mathbf{r}_j} = (2\delta + i)\beta_j + i\sum_{k\neq j} \frac{\exp(ik_0|\mathbf{r}_j - \mathbf{r}_k|)}{ik_0|\mathbf{r}_j - \mathbf{r}_k|}\beta_k.$$
(1.30)

One can then adopt a fluid description of the atomic cloud, i.e., the coarse-grained field $\beta(\mathbf{r})$ describing the local atomic dipole moment field



One-dimensional photonic band gaps in optical lattices

Fig. 1.5 Reflection coefficient R (left) and LDOS (right) as a function of the detuning δ for atom numbers $N = 5.10^3$, 10^4 and 2.10^4 . The one-dimensional lattice is composed of $N_d = 100$ disks, with spacing $d = \lambda_0/2$, disk radius $9\lambda_0$ and thickness $0.04\lambda_0$. The laser beam waist is $4.5\lambda_0$.

is introduced. This approach has been particularly useful in studying the superradiant and subradiant radiation modes [Svidzinsky *et al.* (2010)], non-local effects [Svidzinsky (2012)], scattering from clouds with arbitrary spherical densities [Piovella and Bachelard (2012); Piovella *et al.* (2013)] and the coherent nature of this scattering [Prasad and Glauber (2000); Glauber and Prasad (2000); Prasad and Glauber (2010, 2011)]. The coarse-grained equation for the cloud excitation in the steady state reads:

$$\frac{\Omega_0}{\Gamma} e^{i\mathbf{k}_0\mathbf{r}} = (2\delta + i)\overline{\beta}(\mathbf{r}) + i\int d\mathbf{r}'\rho(\mathbf{r}')\frac{\exp(ik_0|\mathbf{r} - \mathbf{r}'|)}{ik_0|\mathbf{r} - \mathbf{r}'|}\overline{\beta}(\mathbf{r}'), \qquad (1.31)$$

where $\rho(\mathbf{r})$ is the atomic density. In the case of extended slab planes in the transverse directions with density $\rho(z)$ depending only on the longitudinal coordinate, we can assume $\overline{\beta}(\mathbf{r}) = \overline{\beta}(z)e^{ik_{0y}y}$ where $k_0^2 = k_{0y}^2 + k_{0z}^2$, so that (1.31) can be written as:

$$\frac{\Omega_0}{\Gamma} e^{ik_{0z}z} = (2\delta + i)\overline{\beta}(z) + \int dz' \rho(z')\overline{\beta}(z') \\
\times \int dx' \int dy' \frac{\exp(ik_0|\mathbf{r} - \mathbf{r}'|)}{k_0|\mathbf{r} - \mathbf{r}'|} e^{ik_{0y}(y'-y)}.$$
(1.32)

By introducing $s = \sqrt{(x'-x)^2 - (y'-y)^2}$ and $s\sin\phi = y' - y$, the

24

$My \ Book \ Title$

integral over transverse coordinates x' and y' becomes [Manassah (2012b)]:

$$\int dx' \int dy' \frac{\exp(ik_0 |\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|} e^{ik_{0y}(y'-y)} = \int_0^\infty \frac{dss}{\sqrt{s^2 + (z-z')^2}} e^{ik_0\sqrt{s^2 + (z-z')^2}} \times \int_0^{2\pi} d\phi e^{ik_{0y}s\sin\phi}.$$
(1.33)

The integration over ϕ and s gives:

$$\int dx' \int dy' \frac{\exp(ik_0 |\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|} e^{ik_{0y}(y'-y)} = 2\pi \int_0^\infty \frac{dss J_0(k_{0y}s)}{\sqrt{s^2 + (z-z')^2}} e^{ik_0\sqrt{s^2 + (z-z')^2}} = \frac{2\pi i}{k_{0z}} e^{ik_{0z}|z-z'|}.$$
(1.34)

From Eqs.(1.32) and (1.34), the equation for the 1D scattering problem is:

$$\frac{\Omega_0}{\Gamma}e^{ik_{0z}z} = (2\delta+i)\overline{\beta}(z) + \frac{2\pi i}{k_{0z}k_0} \int dz' \rho(z')\overline{\beta}(z')e^{ik_{0z}|z-z'|}.$$
 (1.35)

Then, for an optical lattice consisting of N_d parallel slabs of uniform density ρ_0 and having thickness a, separated by a free space drift d and bounded by the planes z = 0 and $z = N_d a + (N_d - 1)d$, (1.35) reduces to N_d equations, one for each slab:

$$\frac{\Omega_0}{\Gamma} e^{ik_{0z}z} = (2\delta + i)\overline{\beta}_n(z) + \frac{2\pi i\rho_0}{k_{0z}k_0} \sum_{m=1}^N \int_{z_m}^{z_m+a} dz' e^{ik_{0z}|z-z'|}\overline{\beta}_m(z'), \quad (1.36)$$

for $z_n < z < z_n + a$ (n = 1, ...N), where $z_n = (n - 1)(a + d)$ and $z_n + a$ are the slab edges, while $\overline{\beta}_n(z)$ refers to the value of $\overline{\beta}(z)$ inside the slab. In (1.36) $k_{0z} = \sqrt{k_0^2 - k_{0y}^2} = k_0 \cos \theta_0$ with θ_0 being the angle of incidence of the laser with respect to the lattice axis z.

While the one-dimensional problem is usually solved for the electric field by calculating its propagation layer by layer (see, e.g., Ref. [Deutsch *et al.* (1995)]), ((1.36)) explicitly shows the dependence of the field on the radiation of each dipole in the system. The collective nature of the scattering process is thus more evident in our formalism and, although the derivation is less straightforward, it is formally equivalent to the standard approach.

Let us now use the fact that the kernel of (1.36) is the Green function for the 1D Helmholtz equation

$$\left(\partial_z^2 + k_{0z}^2\right)e^{ik_{0z}|z-z'|} = 2ik_{0z}\delta(z-z').$$
(1.37)

One-dimensional photonic band gaps in optical lattices

Hence, applying the operator $(\partial_z^2 + k_{0z}^2)$ to ((1.36)) we obtain the Helmholtz equation which takes the following form inside the *n*th slab:

$$\frac{\partial^2 \overline{\beta}_n(z)}{\partial z^2} + k_z^2 \overline{\beta}_n(z) = 0, \qquad (1.38)$$

where $k_z^2 = k_0^2(m_0^2 - \sin^2 \theta_0)$ and m_0 is the refractive index of the atomic layer, which can be expressed as

$$m_0^2 = 1 - \frac{4\pi\rho_0}{k_0^3(2\delta + i)}.$$
(1.39)

Notice that, according to the Snell-Descartes law $(\sin \theta_0 = m_0 \sin \theta)$, the z component of the wave vector *inside* the medium can be written as $k_z = k_0 m_0 \cos \theta$, where θ is the angle of refraction. For a homogeneous slab of refractive index m_0 only two modes are allowed (forward and backward), so the general solution of ((1.38)) for the *n*th slice is given by

$$\overline{\beta}_n(z) = \frac{1}{2\delta + i} \left[x_n e^{ik_z(z - z_n)} + y_n e^{-ik_z(z - z_n)} \right].$$
(1.40)

The coefficients x_n and y_n are obtained by substituting (1.40) into (1.36) evaluated at the boundary conditions $z = z_n$ and $z = z_n + a$. The long but straightforward calculation is detailed in Appendix (.6).

The recurrence relation for x_n and y_n is given by [Boedecker and Henkel (2003)]

$$\begin{bmatrix} x_{n+1} \\ y_{n+1} \end{bmatrix} = T \begin{bmatrix} x_n \\ y_n \end{bmatrix}, \tag{1.41}$$

where

$$T = \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$$
(1.42)

is the transfer matrix with the elements

$$T_{11,22} = \left[\cos(k_{0z}d) \pm i\frac{k_{0z}^2 + k_z^2}{2k_{0z}k_z}\sin(k_{0z}d)\right]e^{\pm ik_z a},$$

$$T_{12,21} = \pm i\frac{k_{0z}^2 - k_z^2}{2k_{0z}k_z}\sin(k_{0z}d)e^{\mp ik_z a}.$$
(1.43)

Since det(T) = 1, its eigenvalues can be written in the form $\lambda_{\pm} = e^{\pm i\phi}$, where $\cos \phi = \text{Tr}(T)/2$, or explicitly

$$\cos\phi = \left[\cos(k_{0z}d)\cos(k_{z}a) - \frac{k_{0z}^2 + k_z^2}{2k_{0z}k_z}\sin(k_{0z}d)\sin(k_za)\right].$$
 (1.44)

In the case of real refractive index m_0 and if $|\cos \phi| < 1$, the eigenvalues λ_{\pm} are on the unit circle (real ϕ) and correspond to propagating (extended)

Paper-mh

My Book Title

Bloch modes, whereas if $|\cos \phi| > 1$, the eigenvalues are real (imaginary ϕ) and can be found in band gaps. In atomic clouds, the index m_0 is always complex and so are the eigenvalues. It is then necessary to resort to the Local Density of States to properly characterize the band gaps (see later discussion).

The scattered field in the smooth density limit is readily obtained from ((1.23)) and reads [Bachelard *et al.* (2011)]:

$$E_S = -\frac{dk_0^2}{4\pi\varepsilon_0} \int d\mathbf{r}' \rho(\mathbf{r}') \overline{\beta}(\mathbf{r}') \frac{\exp(ik_0|\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|}.$$
 (1.45)

Using (1.34), we obtain the following expression for the total electric field:

$$E(z) = \frac{E_0}{2} \left[e^{ik_{0z}z} + i\frac{k_z^2 - k_{0z}^2}{2k_{0z}}(2\delta + i) \right] \times \sum_{n=1}^N \int_{z_n}^{z_n + a} dz' e^{ik_{0z}|z - z'|} \overline{\beta}_n(z') , \qquad (1.46)$$

where E_0 is the incident laser field amplitude. (1.46) allows extraction of the reflection and transmission coefficients r_N and t_N , respectively:

$$r_{N} = \sum_{n=1}^{N} e^{ik_{0z}z_{n}} \left\{ \frac{k_{z} - k_{0z}}{2k_{0z}} \left[e^{i(k_{0z} + k_{z})a} - 1 \right] x_{n} - \frac{k_{z} + k_{0z}}{2k_{0z}} \left[e^{i(k_{0z} - k_{z})a} - 1 \right] y_{n} \right\},$$
(1.47)

$$t_N = 1 + \sum_{n=1}^{N} e^{-ik_{0z}z_n} \left\{ \frac{k_z + k_{0z}}{2k_{0z}} \left[e^{i(k_z - k_{0z})a} - 1 \right] x_n - \frac{k_z - k_{0z}}{2k_{0z}} \left[e^{-i(k_z + k_{0z})a} - 1 \right] y_n \right\}.$$
 (1.48)

Iterating (1.41) we can write

$$\begin{bmatrix} x_{n+1} \\ y_{n+1} \end{bmatrix} = T^n \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}.$$
 (1.49)

The matrix T^n can be easily calculated in the Bloch basis [Boedecker and Henkel (2003)]:

$$T^{n} = M \begin{bmatrix} e^{in\phi} & 0\\ 0 & e^{-in\phi} \end{bmatrix} M^{-1}, \qquad (1.50)$$

with

$$M = \begin{bmatrix} N_{+} & N_{-} \\ N_{+}c_{+} & N_{-}c_{-} \end{bmatrix},$$
 (1.51)

World Scientific Book - 9in x 6in

One-dimensional photonic band gaps in optical lattices

where $c_{\pm} = (\lambda_{\pm} - T_{11})/T_{12}$ and N_{\pm} are normalization constants chosen for the Bloch eigenstates. The explicitly written elements of the matrix T^n have the following form:

$$T_{11,22}^{n} = \frac{1}{\sin\phi} \left[T_{11,22} \sin(n\phi) - \sin[(n-1)\phi] \right],$$

$$T_{12,21}^{n} = \frac{1}{\sin\phi} T_{12,21} \sin(n\phi).$$
 (1.52)

The values x_1 and y_1 appearing in (1.49) can be determined explicitly from (1.36) by considering the first slab n = 1 and the last slab n = Nand (1.49) at n = N - 1 (see the Appendix), which allows rewriting the expressions (1.47) and (1.48) for the reflection and transmission coefficients r_N and t_N in terms of the reflection and transmission coefficient amplitudes r and t for a single slab:

$$t_N = \frac{t\sin\phi}{\sin N\phi - t\sin(N-1)\phi},\tag{1.53}$$

$$r_N = t_N \frac{r}{t} \frac{\sin N\phi}{\sin \phi},\tag{1.54}$$

where

$$r = \frac{k_{0z} - k_z}{k_{0z} + k_z} \frac{1 - e^{2ik_z a}}{1 - \left(\frac{k_{0z} - k_z}{k_{0z} + k_z}\right)^2 e^{2ik_z a}}$$
(1.55)

$$t = e^{i(k_{0z}d + k_z a)} \frac{1 - \left(\frac{k_{0z} - k_z}{k_{0z} + k_z}\right)^2}{1 - \left(\frac{k_{0z} - k_z}{k_{0z} + k_z}\right)^2 e^{2ik_z a}}.$$
 (1.56)

1.2.3 Vectorial microscopic theory

1

1.2.3.1 Discrete model

Up to now we have considered a scalar cooperative scattering theory, where the vectorial nature of the electromagnetic field is disregarded. The scalar theory remains valid for s-polarized light incident on a 1D periodic stack of planes and is partially satisfactory for disordered systems, where polarization effects have a minor role due to the random orientation of the atomic dipoles. However, the scalar theory does not accurately describe scattering by periodic systems of higher dimensions, where a vectorial description is required. The vectorial model of light scattering can be derived from a quantum field theory (see for instance Ref.[Manassah (2012b)]) considering

Paper-mh

My Book Title

electric dipole transitions between a singlet ground state and a degenerate triplet excited state (as for instance in a $J = 0 \rightarrow J = 1$ transition). Identifying z as a quantization axis, the equations for the three components of the complex polarization vector of the *j*th atom are given by (see Appendix .7):

$$\dot{\beta}_{j}^{(\alpha)} = \left(i\Delta_{0} - \frac{\Gamma}{3}\right)\beta_{j}^{(\alpha)} - i\frac{dE_{0}}{2\hbar}\hat{e}_{0\alpha}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}} - \frac{\Gamma}{2}\sum_{\beta}\sum_{m\neq j}G_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)},\tag{1.57}$$

where $\alpha = (x, y, z), \mathbf{r}_{jm} = \mathbf{r}_j - \mathbf{r}_m$,

$$G_{\alpha,\beta}(\mathbf{r}) = \frac{e^{ik_0r}}{ik_0r} \left\{ \left[\delta_{\alpha,\beta} - \hat{r}_{\alpha}\hat{r}_{\beta} \right] + \left[\delta_{\alpha,\beta} - 3\hat{r}_{\alpha}\hat{r}_{\beta} \right] \left[\frac{i}{k_0r} - \frac{1}{(k_0r)^2} \right] \right\}$$
(1.58)

and $\hat{\mathbf{r}} = \mathbf{r}/r$. The incident laser beam has electric field $\mathbf{E}_{in}(\mathbf{r},t) = E_0 \hat{\mathbf{e}}_0 \cos(i\mathbf{k}_0 \cdot \mathbf{r} - i\omega_0 t)$ with polarization unit vector $\hat{\mathbf{e}}_0$, wave vector \mathbf{k}_0 and frequency $\omega_0 = ck_0$ near the atomic transition frequency ω_a . The vectorial Green function $G_{\alpha,\beta}(\mathbf{r})$ can be written as a function of the scalar Green function $G(r) = \exp(ik_0 r)/(ik_0 r)$ in the following form:

$$G_{\alpha,\beta}(\mathbf{r}) = \left[\delta_{\alpha,\beta} + \frac{1}{k_0^2} \frac{\partial^2}{\partial x_\alpha \partial x_\beta}\right] G(r) = \mathcal{D}_{\alpha,\beta} G(r), \qquad (1.59)$$

where $x_{\alpha} = (\mathbf{r})_{\alpha}$. The steady-state problem reduces to solving the coupled equations:

$$\frac{dE_0}{\hbar\Gamma}\hat{e}_{0\alpha}e^{i\mathbf{k}_0\cdot\mathbf{r}_j} = \left(2\delta + \frac{2}{3}i\right)\beta_j^{(\alpha)} + i\sum_{\beta}\sum_{m\neq j}G_{\alpha,\beta}(\mathbf{r}_{jm})\beta_m^{(\beta)}.$$
 (1.60)

This equation is similar to (1.17). The only major difference is the (2/3) coefficient appearing in the self-decay term due to single-atom spontaneous emission isotropy.

1.2.3.2 Continuous model

Neglecting granularity, (1.60) can be converted into a continuous integral equation for the vectorial field $\overline{\beta}(\mathbf{r})$ with the components $(\overline{\beta})_{\alpha} = \overline{\beta}_{\alpha}$:

$$\frac{d\mathbf{E}_{0}}{\hbar\Gamma}e^{i\mathbf{k}_{0}\cdot\mathbf{r}} = (2\delta + 2i/3)\overline{\boldsymbol{\beta}} + i\int d\mathbf{r}'\rho(\mathbf{r}')\left\{1 + \frac{1}{k_{0}^{2}}\nabla_{\mathbf{r}}\nabla_{\mathbf{r}}\right\}G(|\mathbf{r} - \mathbf{r}'|)\overline{\boldsymbol{\beta}}(\mathbf{r}'), \quad (1.61)$$

World Scientific Book - 9in x 6in

One-dimensional photonic band gaps in optical lattices

where $\mathbf{E}_0 = E_0 \hat{\mathbf{e}}_0$ and $\rho(\mathbf{r})$ is the atomic density. Since $\nabla_{\mathbf{r}} \nabla_{\mathbf{r}} G(|\mathbf{r} - \mathbf{r}'|) = \nabla_{\mathbf{r}'} \nabla_{\mathbf{r}'} G(|\mathbf{r} - \mathbf{r}'|)$, by integrating by parts the last term of (1.61) we obtain:

$$\frac{d\mathbf{E}_{0}}{\hbar\Gamma}e^{i\mathbf{k}_{0}\cdot\mathbf{r}} = (2\delta + 2i/3)\overline{\beta} + i\int d\mathbf{r}'\rho(\mathbf{r}')G(|\mathbf{r} - \mathbf{r}'|)\left\{\overline{\beta}(\mathbf{r}') + \frac{1}{k_{0}^{2}}\nabla_{\mathbf{r}'}\left[\nabla_{\mathbf{r}'}\cdot\overline{\beta}(\mathbf{r}')\right]\right\}(1.62)$$

Since

$$(\nabla^2 + k_0^2)G(|\mathbf{r} - \mathbf{r}'|) = \frac{4\pi i}{k_0}\delta(\mathbf{r} - \mathbf{r}')$$
(1.63)

and $(\nabla^2 + k_0^2) \exp(i\mathbf{k}_0 \cdot \mathbf{r}) = 0$, applying the operator $(\nabla^2 + k_0^2)$ to both sides of (1.62), we obtain the following differential equation:

$$\left[\nabla^2 + k_0^2 m^2(\mathbf{r})\right] \overline{\beta}(\mathbf{r}) = \left[1 - m^2(\mathbf{r})\right] \nabla \left[\nabla \cdot \overline{\beta}(\mathbf{r})\right], \qquad (1.64)$$

where $m(\mathbf{r})$ is the cloud refractive index given by

$$m^{2}(\mathbf{r}) = 1 - \frac{2\pi\rho(\mathbf{r})}{k_{0}^{3}(\delta + i/3)}.$$
(1.65)

We notice that in the scalar radiation theory (1.64) reduces to $[\nabla^2 + k_0^2 m^2(\mathbf{r})]\overline{\beta}_{\alpha}(\mathbf{r}) = 0$ for each α -components. Hence, in the scalar approximation $\overline{\beta}$ is a purely transverse field that satisfies $\nabla \cdot \overline{\beta} = 0$.

1.2.3.3 Scattered field

The electric field of the scattered radiation can be evaluated directly from the macroscopic Maxwell equations with polarization $\mathbf{P} = -d \sum_{j=1}^{N} \beta_j \delta(\mathbf{r} - \mathbf{r}_j)$. The result, as demonstrated in Appendix .8, is

$$E_{\alpha}(\mathbf{r}) = -i \frac{dk_0^3}{4\pi\epsilon_0} \sum_{\beta} \sum_{j=1}^N G_{\alpha,\beta}(\mathbf{r} - \mathbf{r}_j) \beta_j^{(\beta)}.$$
 (1.66)

Considering $G_{\alpha,\beta}$ as a spatial component of the symmetric tensor **G**, (1.66) can be written in the vectorial form:

$$\mathbf{E}(\mathbf{r}) = -i \frac{dk_0^3}{4\pi\epsilon_0} \sum_{j=1}^N \mathbf{G}(\mathbf{r} - \mathbf{r}_j) \cdot \boldsymbol{\beta}_j.$$
(1.67)

Combining (1.66) and (1.60) we find:

$$\boldsymbol{\beta}_{j} = \frac{d}{\hbar \left(\Delta_{0} + i\Gamma/3 \right)} \left[\mathbf{E}_{in}(\mathbf{r}_{j}) + \mathbf{E}(\mathbf{r}_{j}) \right], \tag{1.68}$$

where $\mathbf{E}_{in}(\mathbf{r}) = (\mathbf{E}_0/2)e^{i\mathbf{k}_0\cdot\mathbf{r}}$ is the incident field and $j = 1, \ldots, N$. As expected, the electric dipole moment of the single atom is proportional to the sum of the incident field and the field scattered by all the other atoms.

Paper-mh

My Book Title

1.2.3.4 Single-scattering contribution

Neglecting the contribution due to the internal field $\mathbf{E}(\mathbf{r}_j)$ in (1.68) (so, the effect of multiple scattering is not taken into account) from (1.67) and (1.68) we obtain:

$$\mathbf{E}^{(1)}(\mathbf{r}) = \kappa(\delta) \sum_{j} \mathbf{G}(\mathbf{r} - \mathbf{r}_{j}) \cdot \mathbf{E}_{in}(\mathbf{r}_{j}), \qquad (1.69)$$

where $\kappa(\delta) = 1/(2i\delta - 2/3)$. Far away from the scatters we can approximate $|\mathbf{r} - \mathbf{r}_j| \approx r - \mathbf{\hat{n}} \cdot \mathbf{r}_j$ in the exponent factor of (1.58), where $\mathbf{\hat{n}}$ is a unit vector in the direction of the observation and r is the distance to the system of scatters. Retaining only the terms decreasing as $1/|\mathbf{r} - \mathbf{r}_j| \approx 1/r$, (1.58) is approximated by

$$G_{\alpha,\beta}(\mathbf{r} - \mathbf{r}_j) \approx \frac{e^{ik_0 r}}{ik_0 r} \left(\delta_{\alpha,\beta} - \hat{n}_\alpha \hat{n}_\beta\right) e^{-i\mathbf{k}\cdot\mathbf{r}_j},\tag{1.70}$$

where $\mathbf{k} = k_0 \hat{\mathbf{n}}$. Inserting (1.70) in (1.69) we obtain:

$$\mathbf{E}^{(1)}(\mathbf{r}) \approx i\kappa(\delta) \frac{e^{ik_0 r}}{2k_0 r} E_0[\hat{\mathbf{n}} \times (\hat{\mathbf{n}} \times \hat{\mathbf{e}}_0)] \sum_{j=1}^N e^{i(\mathbf{k}_0 - \mathbf{k}) \cdot \mathbf{r}_j}, \qquad (1.71)$$

where $\hat{\mathbf{e}}_0$ is the incident polarization vector. Equation (1.71) coincides with the well-known expression for the radiation field emitted by a collection of point-like scatters, known as Rayleigh scattering [Jackson (1999)]. The scattered field results from a coherent superposition of the field amplitudes generated by each atom and is proportional to the structure factor $\mathcal{F}(\mathbf{q}) = \sum_j \exp(i\mathbf{q}\cdot\mathbf{r}_j)$, where $\mathbf{q} = \mathbf{k}_0 - \mathbf{k}$ is the transferred momentum. Equation (1.71) also shows that the electric dipole moment of each atom is $\mathbf{p} = \alpha\epsilon_0 \mathbf{E}_{in}$ with polarizability $\alpha(\delta) = 4\pi i \kappa(\delta)/k_0^3$.

1.2.4 Photonic band gaps for three-level atoms

The microscopic model of cooperative scattering and photonic band gaps, so far developed only for two-level atoms, can be extended to three-level atoms in Λ or cascade configurations. These systems offer the advantage of further manipulations and control of the photonic band gaps, exploiting the electromagnetically induced transparency (EIT) properties [Harris (1997)]. EIT is a quantum interference effect characterized by the presence of a frequency region where absorption is greatly reduced, accompanied by steep dispersion [Fleischhauer *et al.* (2005)]. Both the transparency bandwidth

One-dimensional photonic band gaps in optical lattices

and the steep dispersion near the EIT resonance are controlled by the corresponding driving field [Petrosyan (2007)]. The possible advantages and limitations of using the photonic band gaps near the EIT transparency bandwidth has been investigated in a recent experiment [Schilke *et al.* (2012b)]. Here we develop a microscopic description of the scattering by three-level atoms, showing how the TM formalism can take into account EIT, simply by modifying the atomic medium refractive index.

1.2.4.1 Single-particle dynamics

We consider a cascade configuration between the three states $|g\rangle$, $|e\rangle$ and $|m\rangle$ with energy differences $\hbar\omega_{eg} = E_e - E_g$ and $\hbar\omega_{me} = E_m - E_e$ and decay rates Γ_{eg} and Γ_{me} , respectively. For instance, for ⁸⁸Sr atoms the three states may be $|g\rangle = (5s^2)^1 S_0$, $|e\rangle = (5s5p)^3 P_1$ and $|m\rangle = (5s5d)^3 D_1$. The transition $|m\rangle \rightarrow |g\rangle$ is forbidden and the two transitions $|g\rangle \rightarrow |e\rangle$ and $|e\rangle \rightarrow |m\rangle$ are driven by two external fields with Rabi frequencies $\Omega_1 = d_{ge} E_1/\hbar$, $\Omega_2 = d_{em} E_2/\hbar$ and frequencies $\omega_{1,2}$, where d_{ab} is the dipole matrix elements for the generic transition $|a\rangle \rightarrow |b\rangle$. The single-particle Hamiltonian is

$$H = \hbar \omega_{eg} |e\rangle \langle e| + \hbar (\omega_{eg} + \omega_{me}) |m\rangle \langle m| - \hbar \left\{ \Omega_1 e^{-i\omega_1 t} |e\rangle \langle g| + \Omega_2 e^{-i\omega_2 t} |m\rangle \langle e| + \text{h.c.} \right\}.$$
(1.72)

Assuming $|\psi\rangle = c_g(t)|g\rangle + c_e(t)|e\rangle + c_m(t)|m\rangle$, from the Schrödinger equation, $i\hbar\partial_t|\psi\rangle = H|\psi\rangle$, we obtain:

$$\dot{c}_g = i\Omega_1^* e^{i\omega_1 t} c_e, \tag{1.73}$$

$$\dot{c}_e = -i\omega_{eg}c_e + i\Omega_1 e^{-i\omega_1 t}c_g + i\Omega_2^* e^{i\omega_2 t}c_m, \qquad (1.74)$$

$$\dot{c}_m = -i(\omega_{eg} + \omega_{me})c_m + i\Omega_2 e^{-i\omega_2 t}c_e.$$
(1.75)

Introducing the coherences $\rho_{ge} = c_g^* c_e \exp(i\omega_1 t)$, $\rho_{gm} = c_g^* c_m \exp[i(\omega_1 + \omega_2)t]$ and $\rho_{em} = c_e^* c_m \exp(i\omega_2 t)$, we write:

$$\dot{\rho}_{ge} = [i\Delta_1 - \Gamma_{eg}/2]\rho_{ge} + i\Omega_1(\rho_{gg} - \rho_{ee}) + i\Omega_2^*\rho_{gm}, \qquad (1.76)$$

$$\dot{\rho}_{gm} = [i(\Delta_1 + \Delta_2) - \Gamma_{me}/2]\rho_{gm} - i\Omega_1\rho_{em} + i\Omega_2\rho_{ge}, \qquad (1.77)$$

$$\dot{\rho}_{em} = [i\Delta_2 - (\Gamma_{eg} + \Gamma_{me})/2]\rho_{em} + i\Omega_2(\rho_{ee} - \rho_{mm}) - i\Omega_1^*\rho_{gm}, (1.78)$$

where $\rho_{ii} = |c_i|^2$, $\Delta_1 = \omega_1 - \omega_{eg}$ and $\Delta_2 = \omega_2 - \omega_{me}$, and we add the decay terms with rates Γ_{eg} and Γ_{me} . We assume the field Ω_1 to be so weak that $\rho_{gg} \approx 1$, $\rho_{ee} \approx 0$ and $\rho_{mm} \approx 0$. With these approximations (1.78) yields at steady-state

$$\rho_{em} \approx \frac{\Omega_1^* \rho_{gm}}{\Delta_2 + i(\Gamma_{eg} + \Gamma_{me})/2},$$

Paper-mh

My Book Title

so that ρ_{em} can be neglected in (1.77). The latter then yields

$$\rho_{gm} \approx -\frac{\Omega_2 \rho_{ge}}{(\Delta_1 + \Delta_2) + i\Gamma_{me}/2}$$

and can be inserted into (1.76) to obtain

f

$$\rho_{ge} = -\frac{\Omega_1}{\left[\Delta_1 + i\Gamma_{eg}/2 - \frac{|\Omega_2|^2}{(\Delta_1 + \Delta_2) + i\Gamma_{me}/2}\right]}.$$
(1.79)

Defining the polarization $P = \rho d_{ge} \rho_{ge} = \epsilon_0 \chi E_1$, where ρ is the atomic density, we obtain the following expression:

$$\chi = -\frac{\chi_0}{\left[2\delta + i - \frac{a^2}{2\delta + i\gamma}\right]},\tag{1.80}$$

where $\chi_0 = (2nd_{ge}^2)/(\epsilon_0\hbar\Gamma_{eg})$, $\delta = \Delta_1/\Gamma_{eg}$, $\gamma = \Gamma_{me}/\Gamma_{eg}$, $a = 2\Omega_2/\Gamma_{eg}$ and we assumed $\Delta_2 = 0$. Separating the real and imaginary parts, we write:

$$\operatorname{Re}\left(\frac{\chi}{\chi_0}\right) = -2\delta \frac{4\delta^2 - \gamma^2 - a^2}{(4\delta^2 - \gamma - a^2)^2 + 4\delta^2(1+\gamma)^2},\tag{1.81}$$

$$\operatorname{Im}\left(\frac{\chi}{\chi_0}\right) = \frac{\gamma(\gamma + a^2) + 4\delta^2}{(4\delta^2 - \gamma - a^2)^2 + 4\delta^2(1+\gamma)^2}.$$
 (1.82)

For instance, for ⁸⁸Sr, $\Gamma_{eg} = (2\pi)7.6$ kHz and $\Gamma_{me} = (2\pi)90.3$ kHz, so that $\gamma = 11.8$. In Fig. 1.6 the real and imaginary parts of χ/χ_0 are represented as a function of δ for two- and three-level atoms. For three-level atoms the imaginary part of χ has a minimum near $\delta = 0$ provided $a^2 > \gamma^3/(1+2\gamma)$, which for $\gamma = 11.8$ gives $\Omega_2 > 4.1\Gamma_{eg}$.

1.2.4.2 Cooperative emission

Once the single-particle dynamics have been determined it is straightforward to obtain the cooperative scattering model for the three-level cascade configuration. Considering a collection of such N three-level atoms, the Hamiltonian of the system in the scalar field theory, including the coupling with the vacuum radiation modes, is given by

$$H = \hbar \sum_{j=1}^{N} \left\{ \omega_{eg} |e_{j}\rangle \langle e_{j}| + (\omega_{eg} + \omega_{me}) |m_{j}\rangle \langle m_{j}| \right\}$$
$$- \hbar \sum_{j=1}^{N} \left\{ \Omega_{1} e^{-i\omega_{1}t + i\mathbf{k}_{1}\cdot\mathbf{r}_{j}} |e_{j}\rangle \langle g_{j}| + \Omega_{2} e^{-i\omega_{2}t + i\mathbf{k}_{2}\cdot\mathbf{r}_{j}} |m_{j}\rangle \langle e_{j}| + \text{h.c.} \right\}$$
$$+ \hbar \sum_{j=1}^{N} \sum_{\mathbf{k}} g_{\mathbf{k}} \{ a_{\mathbf{k}} e^{-i\omega_{k}t + i\mathbf{k}\cdot\mathbf{r}_{j}} + a_{\mathbf{k}}^{\dagger} e^{i\omega_{k}t - i\mathbf{k}\cdot\mathbf{r}_{j}} \} \{ |e_{j}\rangle \langle g_{j}| + |g_{j}\rangle \langle e_{j}|)$$
(1.83)

World Scientific Book - 9in × 6in

One-dimensional photonic band gaps in optical lattices

Paper-mh




Fig. 1.6 Real (blue) and imaginary (red) parts of the normalized susceptibility for two-(dashed) and three-level (solid) atoms. For three-level atoms the following parameters are used: a = 20 and $\gamma = 11.8$.

We assume that the cooperative effects manifest themselves only in the lower transition $|e\rangle \rightarrow |g\rangle$, as described by the last term in the Hamiltonian (1.83). The single-atom spontaneous emission from the upper level $|m\rangle$ is described simply by adding a damping term, as performed in the previous section. Assuming at most one atom in the excited state $|e\rangle$, the complete system is described by the following state:

$$\begin{split} |\psi\rangle &= c_g(t)|g_1, \dots, g_N\rangle|0\rangle_{\mathbf{k}} + \sum_{j=1}^N c_{ej}(t)|g_1, \dots, e_j, \dots, g_N\rangle|0\rangle_{\mathbf{k}} \\ &+ \sum_{j=1}^N c_{mj}(t)|g_1, \dots, m_j, \dots, g_N\rangle|0\rangle_{\mathbf{k}} \\ &+ \sum_{\mathbf{k}} \gamma_{\mathbf{k}}(t)|g_1, \dots, g_N\rangle|1\rangle_{\mathbf{k}} \\ &+ \sum_{\mathbf{k}} \sum_{j\neq j'} \epsilon_{j,j',\mathbf{k}}(t)|g_1, \dots, e_j, \dots, e_{j'}, \dots, g_N\rangle|1\rangle_{\mathbf{k}}. \end{split}$$
(1.84)

World Scientific Book - $9 \text{in} \times 6 \text{in}$

Paper-mh

My Book Title

The evolution of the probability amplitudes is governed by:

$$\dot{c}_g(t) = i\Omega_1^* \sum_{j=1}^N c_{ej}(t) e^{i\omega_1 t - i\mathbf{k}_1 \cdot \mathbf{r}_j},$$
(1.85)

$$\dot{c}_{ej}(t) = -i\omega_{eg}c_e(t) + i\Omega_1 c_g(t)e^{-i\omega_1 t + i\mathbf{k}_1 \cdot \mathbf{r}_j} + i\Omega_2^* c_m(t)e^{i\omega_2 t - i\mathbf{k}_2 \cdot \mathbf{r}_j} - i\sum g_{\mathbf{k}} \left[\gamma_{\mathbf{k}}(t) + \sum \epsilon_{j,j',\mathbf{k}}(t) \right] e^{-i\omega_k t + i\mathbf{k} \cdot \mathbf{r}_j},$$
(1.86)

$$-i\sum_{\mathbf{k}}g_{\mathbf{k}}\left[\gamma_{\mathbf{k}}(t)+\sum_{j'\neq j}\epsilon_{j,j',\mathbf{k}}(t)\right]e^{-i\omega_{k}t+i\mathbf{k}\cdot\mathbf{r}_{j}},\qquad(1.86)$$

$$\dot{c}_{mj}(t) = -i(\omega_{eg} + \omega_{me})c_{mj}(t) + i\Omega_2 c_{ej}(t)e^{-i\omega_2 t + i\mathbf{k}_2 \cdot \mathbf{r}_j}, \qquad (1.87)$$

$$\dot{\gamma}_{\mathbf{k}}(t) = -ig_{\mathbf{k}} \sum_{j=1}^{N} c_{ej}(t) e^{i\omega_{k}t - i\mathbf{k}\cdot\mathbf{r}_{j}}$$
(1.88)

$$\dot{\epsilon}_{j,j'\mathbf{k}}(t) = -ig_{\mathbf{k}}e^{i\omega_k t} \left[e^{-i\mathbf{k}\cdot\mathbf{r}_j}c_{ej'}(t) + e^{-i\mathbf{k}\cdot\mathbf{r}_{j'}}c_{ej}(t) \right].$$
(1.89)

Introducing again the coherences and assuming the field Ω_1 to be weak and the ground state undepleted, we obtain:

$$\dot{\rho}_{ge}^{(j)}(t) = i\Delta_1 \rho_{ge}^{(j)}(t) + i\Omega_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_j} + i\Omega_2^* e^{-i\mathbf{k}_2 \cdot \mathbf{r}_j} \rho_{gm}^{(j)}(t) - i\sum_{\mathbf{k}} g_{\mathbf{k}} \left[\gamma_{g\mathbf{k}}(t) + \sum_{j' \neq j} \epsilon_{g,\mathbf{k}}^{(j,j')}(t) \right] e^{-i(\omega_k - \omega_1)t + i\mathbf{k} \cdot \mathbf{r}_j}, \quad (1.90)$$

$$\dot{\rho}_{gm}^{(j)}(t) = [i(\Delta_1 + \Delta_2) - \Gamma_{me}/2]\rho_{gm}^{(j)}(t) + i\Omega_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_j} \rho_{ge}^{(j)}(t), \quad (1.91)$$

$$\dot{\gamma}_{g\mathbf{k}}(t) = -ig_{\mathbf{k}} \sum_{j=1}^{N} \rho_{ge}^{(j)}(t) e^{i(\omega_{k}-\omega_{1})t-i\mathbf{k}\cdot\mathbf{r}_{j}}, \qquad (1.92)$$

$$\dot{\epsilon}_{g\mathbf{k}}^{(j,j')}(t) = -ig_{\mathbf{k}}e^{i(\omega_{k}-\omega_{1})t} \left[e^{-i\mathbf{k}\cdot\mathbf{r}_{j}}\rho_{ge}^{(j')}(t) + e^{-i\mathbf{k}\cdot\mathbf{r}_{j'}}\rho_{ge}^{(j)}(t) \right], \quad (1.93)$$

where $\gamma_{g\mathbf{k}} = c_g^* \gamma_{\mathbf{k}}$ and $\epsilon_{g\mathbf{k}}^{(j,j')} = c_g^* \epsilon_{j,j',\mathbf{k}}$. As we see, the only difference with respect to the two-level atom case is the third term on the right hand side of (1.90) coupled to the upper transition by the field Ω_2 . By eliminating the variables $\gamma_{g\mathbf{k}}$ and $\epsilon_{g\mathbf{k}}^{(j,j')}$ in the usual Markov approximation, (1.90) becomes

$$\dot{\rho}_{ge}^{(j)}(t) = i\Delta_1 \rho_{ge}^{(j)}(t) + i\Omega_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_j} + i\Omega_2^* e^{-i\mathbf{k}_2 \cdot \mathbf{r}_j} \rho_{gm}^{(j)}(t) - \frac{\Gamma_{eg}}{2} \sum_{m=1}^N \frac{\exp[ik_1 |\mathbf{r}_j - \mathbf{r}_m|]}{ik_1 |\mathbf{r}_j - \mathbf{r}_m|} \rho_{ge}^{(m)}(t).$$
(1.94)

Neglecting granularity and assuming a continuous density distribution $\rho(\mathbf{r})$, (1.91) at steady-state reads

$$\rho_{gm}(\mathbf{r}) = -\frac{\Omega_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_j}}{(\Delta_1 + \Delta_2) + i\Gamma_{me}/2} \rho_{ge}(\mathbf{r}).$$

One-dimensional photonic band gaps in optical lattices

Inserted in to (1.94), this gives

$$-\Omega_{1}e^{i\mathbf{k}_{1}\cdot\mathbf{r}} = \left[\Delta_{1} + i\frac{\Gamma_{eg}}{2} - \frac{|\Omega_{2}|^{2}}{\Delta_{1} + \Delta_{2} + i\Gamma_{me}/2}\right]\rho_{ge}(\mathbf{r}) + \frac{\Gamma_{eg}}{2}\int d\mathbf{r}'\rho(\mathbf{r}')\frac{\exp(ik_{1}|\mathbf{r} - \mathbf{r}'|)}{k_{1}|\mathbf{r} - \mathbf{r}'|}\rho_{ge}(\mathbf{r}').$$
(1.95)

Assuming for simplicity $\Delta_2 = 0$ and defining $\delta = \Delta_1 / \Gamma_{eg}$, $a = 2|\Omega_2| / \Gamma_{eg}$, $\gamma = \Gamma_{me} / \Gamma_{eg}$ and $\overline{\beta}(\mathbf{r}) = -\rho_{ge}(\mathbf{r})$, (1.95) becomes

$$\frac{\Omega_1}{\Gamma_{eg}} e^{i\mathbf{k}_1 \cdot \mathbf{r}} = \left[2\delta + i - \frac{a^2}{2\delta + i\gamma} \right] \overline{\beta}(\mathbf{r})
+ \int d\mathbf{r}' \rho(\mathbf{r}') \frac{\exp(ik_1 |\mathbf{r} - \mathbf{r}'|)}{k_1 |\mathbf{r} - \mathbf{r}'|} \overline{\beta}(\mathbf{r}').$$
(1.96)

Equation (1.96) generalizes the coarse-grained (1.31) for the case of threelevel atoms. The only difference is the extra term in the squared parenthesis, proportional to the driving intensity. Moreover, the TM formalism remains unchanged, only the refractive index in (1.39) is replaced by the expression

$$m_0^2 = 1 - \frac{4\pi\rho_0}{k_1^3 \left(2\delta + i - \frac{a^2}{2\delta + i\gamma}\right)}.$$
 (1.97)

In (1.97), the possibility to manipulate the cloud's refractive index using a third level becomes clear. An illustration is shown in Fig.1.7, where the spectrum in reflection and transmission of a lattice are plotted for two- and three-level atoms.

1.3 Measurement of a photonic band gap

One-dimensional photonic band gaps in optical lattices have been observed in a recent experiment [Schilke *et al.* (2011)]. The basic idea of this experiment was to create a pile of pancake-shaped atomic clouds, to irradiate them with a probe laser tuned close to an atomic resonance at an incident angle satisfying the Bragg condition (1.26). The reflected light intensity is then monitored while the probe laser frequency is scanned across resonance to look for signatures of the presence of a band gap in the reflection spectrum.




Fig. 1.7 Reflection R and transmission T coefficients for two- and three-level atom lattices. The lattice is composed of 8000 atoms spread over 200 disks of thickness $0.04\lambda_0$ and radius 4.5 λ_0 . For the three-level atoms we choose a = 4 and $\gamma = 11.8$.

1.3.1Description of the experiment

Technically, a cloud of rubidium atoms was trapped and cooled in a standard magneto-optical trap and then transferred into an optical dipole trap formed by a retroreflected laser beam with 1.3 W of power focused into a 220 μ m waist and red detuned from the ⁸⁷Rb D2-line at $\lambda_0 = 780.24$ nm. Inside the standing wave formed by the laser beam the atoms arranged themselves at the antinodes and formed a lattice adopting the shape of a pile of pancakes aligned along the optical axis. About 7700 pancakes were filled, each one with on average 6500 atoms. At temperatures around 100 μ K the pancakes represented Gaussian density distributions with radii of $\sigma_r = 60 \ \mu m$ and thicknesses of $\sigma_z = 47 \ nm$. The optical layout of the experiment is shown in Fig. 1.1.

Reflection spectra were obtained by irradiating a probe laser (wavevector $k_{pr} = 2\pi/\lambda_{pr}$ under an angle of $\theta = 2^{\circ}$ focused down to 35 μ m into the optical lattice. The Bragg condition is satisfied when the difference between the incident and reflected wavevectors matches the lattice wave vector, $2nk_0\cos\theta = 4\pi/\lambda_{lat}$, where the real part of the average refractive index $n = n(\delta)$ depends on the probe laser detuning from resonance. However, it is experimentally more convenient to tune the Bragg condition varying the wavelength of the lattice laser, instead of varying the angle of incidence. Introducing the lattice laser detuning from the geometrically

One-dimensional photonic band gaps in optical lattices

ideal (n = 1) Bragg condition, $\Delta \lambda_{lat} \equiv \lambda_{lat} - \lambda_0 / \cos \theta$, the Bragg condition can be written as $n(\delta) - 1 = -\Delta \lambda_{lat} / \lambda_{lat}$.

For a chosen lattice wavelength, the probe laser was scanned across the $F = 2 \rightarrow F' = 3$ transition of the rubidium D2-resonance, while recording the reflected and transmitted light intensity. As explained in Ref. [Slama *et al.* (2005a)], when the atoms are axially strongly localized in the standing light wave, the atoms scattering into the Bragg angle do not receive recoil, hence the reflected light is not Doppler-shifted, but scattered elastically. As a consequence, the atoms are not heated by the probe beam and the reflected light is not Doppler-broadened. At low atomic densities, the spectral linewidth observed in the reflection spectrum was equal to the natural linewidth of $\Gamma/2\pi = 6$ MHz. In contrast, at high atomic densities, when the Bragg condition was met a very efficient reflection of up to 80% was observed as well as a dramatic and asymmetric spectral broadening.

1.3.2 Reaching the thick grating regime

The experimental challenge in reaching the regime where the lattice responds to incident light by the formation of photonic bands arises from the necessity of realizing a *thick grating*, which means that the light is Braggreflected back and forth many times between subsequent atomic layers.

This requires, first of all, that the lattice's optical density is high enough for multiple scattering. Second, the atoms need to be well arranged in axial direction, ideally forming two-dimensional sheets exactly located at the antinodes of the optical lattice. Any atom dislocated from this plane reduces specular reflection through diffuse scattering into arbitrary solid angles. Furthermore, the incident light gets absorbed before it has a chance of penetrating deep into the lattice and to probe many atomic layers. The diffuse scattering problem can be partially circumvented by detuning the probe laser slightly from resonance. The main reason for axial disorder is the axial Gaussian distribution of the atoms within a pancake at finite temperatures whose *rms*-width is $\sigma_z \propto \sqrt{T}$. One could think of lowering the temperature by some means, but this also reduces the radial extent of the pancakes, $\sigma_r \propto \sqrt{T}$, which introduces another problem.

As discussed in Section 1.4.1, due to the non-zero angle of incidence, the finite radial extent of the pancakes causes the light to walk off the optical axis after a certain number of reflections [Slama *et al.* (2005b)], which also limits the number of pancakes available to multiple reflections. Choosing a small angle of incidence reduces this problem, but still walk-off losses

Paper-mh

My Book Title

represent a serious limitation in some circumstances [Schilke et al. (2012a)].

1.3.3 Interpretation of the observations

In the low optical density limit (thin grating limit), the light reflection results from Bragg scattering, which is understood as constructive interference of the Rayleigh-scattered radiation patterns emitted by the individual periodically arranged atoms. In this regime the reflection coefficient of the lattice turns out to be nearly real, phase shifts are negligible, and the spectral lineshape, which is determined by the natural decay rate Γ , is Lorentzian. Consequently, Bragg scattering in this regime is cooperative but not superradiant.

When the density of the atomic gas, or the optical thickness of the lattice, is increased, the probability that photons are reflected multiple times between adjacent layers also increases (thick grating regime). The interference between the light reflected from or transmitted through the layers gives rise to stopping bands for certain light frequencies or irradiation angles. In this regime absorption can generally be neglected, but large phase shifts occur and the spectral lineshape dramatically broadens under the influence of superradiance and becomes asymmetric (see Fig. 1.8). In the extreme limit of very high densities the broadened reflection spectrum can be interpreted as a frequency band in which the propagation of light traversing the optical lattice under the Bragg angle is forbidden. This frequency band is known as "photonic band gap".

To support this interpretation of the observations, Schilke *et al.* compared the observed reflection spectra to the prediction of the TM formalism detailed in Sec. 1.2.2 and found excellent agreement for all density regimes. The model also allows calculation of the LDOS. For the set of parameters which best fit the reflection spectra a considerable reduction of the LDOS below the free space value of 1 was found.

1.4 Finite size effects

The TM formalism is broadly used to describe photonic band gaps, yet it may present severe limitations in the case of cold atoms in relatively small optical lattices (as compared to dielectric photonic crystals). Typical threedimensional lattices may only contain $10^5 - 10^6$ atoms, which corresponds to several dozens of periods in each direction only. Hence, finite-size ef-

One-dimensional photonic band gaps in optical lattices




Fig. 1.8 Bragg reflection (left) and transmission (right) spectra for lattices with various numbers of layers N_d . Here we assumed an atomic density $5 \times 10^{11} cm^3$. The lattice is slightly detuned from the Bragg condition $\lambda_{dip} = 1nm$, so that the line shapes become asymmetric.

fects which can not be accounted for by the TM approach are expected to come into play. Furthermore, while some of the experimental peculiarities and imperfections can be incorporated into a *generalized transfer matrix approach* [Slama *et al.* (2006)] other effects are intrinsically beyond it.

1.4.1 Applicability of the TM formalism to experiment

Let us now discuss several limitations of the TM formalism with respect to the experimental situation:

1. If a probe beam enters an optical lattice under a finite angle the beam walks out of the stack of slabs after a finite amount of reflections due to the finite radial distribution. This is similar to the well-known situation in unstable optical cavities. This effect limits the effective number of available slabs contributing to multiple scattering. The effect is particularly pronounced for large angles of incidence or small radial extension of the slabs. While the finite angle of incidence can be incorporated into the formalism via a modified quasi-momentum, the beam walk-off can not because the TM formalism, being purely one-dimensional, assumes not only radially infinite atomic layers, but also a radially homogeneous density distribution. In reality the radial density distribution is rather Gaussian which implying a variation of the penetration depth with the distance from the optical axis. The experimentally observed reflection spectra thus represent an average of reflection spectra taken at *different* optical densities.

Paper-mh

My Book Title

2. The finite radial extent has another important impact on the reflection angle. Although the pancake's aspect ratio is smaller than $\sigma_z/\sigma_r \approx 10^{-3}$ with respect to Bragg scattering, it tends to behave like a chain of pointlike scatterers rather than a dielectric mirror [Slama *et al.* (2005b)]. This means that if the lattice constant is detuned from the Bragg condition, the reflection angle tends to self-adjust in order to fulfil the Bragg condition rather than to be equal to the angle of incidence as assumed in the TM formalism. This self-adjustment of the Bragg condition impedes a controlled tuning of quasi-momentum.

3. Atoms distributed over the lattice potential, the depth of which is on the order of $U_0 = h \cdot 7$ MHz, experience individual dynamical Stark shifts of their resonances which vary with the atomic location. This effect shifts and inhomogeneously broadens the Bragg spectra [Slama *et al.* (2005a)]. The axial modulation of the Stark shift can be included into the TM formalism by dividing every atomic layer into a number of thin sublayers for which the transfer matrices are evaluated based on the local density and detuning of the probe light. However, the radial variation of the Stark shift can not be described.

4. When the probe laser is detuned from the resonance, the average refractive index of the cloud differs from 1. This means that the incident light is deflected by the refraction when it enters the optically thick atomic cloud. This slightly modifies the angle of incidence, and thus its deviation from the Bragg angle can be as much as 0.1° depending on the probe beam detuning [Slama *et al.* (2006)]. Moreover, the optically thick cloud focuses or defocuses the incident beam depending on its detuning.

5. A good collimation of the probe beam is important. A divergent probe beam may be expanded into several plane waves, and each of them might have a slightly different angle of incidence with respect to the Bragg condition. On the other hand, the probe beam needs to have a waist smaller than the radial extent of the atomic cloud in order to yield high reflection coefficients.

6. Collective effects arising from the disordered part of the atomic cloud (forward scattering, Mie scattering, superradiance, etc.) can play a role even at moderate atomic densities. However, local disorder is disregarded in the TM formalism. Only disorder along the optical axis of the 1D lattice (finite Debye-Waller factor) could be introduced into the TM formalism.

7. As already mentioned, the atoms are axially strongly localized by the lattice potential, so that the Bragg-reflected light is scattered elastically due to the Lamb-Dicke effect. This does not hold for diffuse scattering into

October 8, 2013 0:27

World Scientific Book - 9in x 6in

Paper-mh

One-dimensional photonic band gaps in optical lattices

other solid angles, which is inelastic as it involves the transfer of photonic recoil to atoms in radial direction. The approach of dividing the atomic cloud into a perfectly ordered optical lattice and a separate homogeneous cloud is reminiscent to an ansatz frequently made in describing Bragg scattering of X-rays on crystals [Coley (1995); Weidemüller *et al.* (1998)], where it is found that disorder and impurities do not broaden the angular distribution of the reflected radiation, but increase the background of isotropically distributed diffuse scattering.

8. The microscopic model can obviously be extended to arbitrary lattice geometries including 3D lattices of any crystalline structure. This is simply due to the fact that before performing a numerical simulation the position of all atoms must be defined, as well as the angle of incidence and the polarization of the probe beam. It makes no difference to the algorithm whether the positions are chosen randomly (to account for disorder) or periodically (to ascribe long-range-order).

All these effects and experimental constraints are beyond (or have some aspects beyond) the TM formalism, but *all effects* are naturally included in the cooperative scattering model [Courteille *et al.* (2010); Bachelard *et al.* (2011, 2012b)]. Effect 7., for example, is confirmed by the appearance of a background of speckle-like scattering well distinct from the solid angle into which the reflected light is scattered, as shown in Fig. 1.9.




Fig. 1.9 Far-field radiation from a one-dimensional optical lattice illuminated with an incident beam inclined by $\theta_0 = 20^{\circ}$ with respect to the z-axis. The lattice is composed of 100 disks of radius $15\lambda_0$ and thickness $0.04\lambda_0$, spaced by $\lambda_0/2/\cos\theta_0$, with the system being composed of N = 8000 atoms overall. The incident Gaussian beam has waist $9\lambda_0$ at the lattice center.

Paper-mh

My Book Title

1.4.2 Suppression of absorption

The hallmark of a photonic band is the suppression of spontaneous emission. Models describing the propagation of light inside photonic crystals assign this suppression to a reduction of the density of optical modes available for spontaneous decay [John and Stephen (1983)], as shown in Fig. 1.10. The suppression of spontaneous emission in a one-dimensional lattice may seem surprising since scattered photons always have the option to escape sideways. However, the reduction of absorption inside a 1D band gap can be understood in classical terms [Slama *et al.* (2006)]: the standing wave formed by the incident probe beam and the Bragg-reflected light adjusts its phase so that its *intensity nodes coincide with the atomic layers*. In that way absorption is minimized. If the length of the lattice is finite, the contrast of the standing wave is smaller than 1, meaning that the probe light intensity at the locations of the atomic layers does not vanish. Hence, a finite absorption persists even for a perfect but finite lattice with fulfilled Bragg condition, and the photonic band gap cannot be completely dark.




Fig. 1.10 Local density of state (circles) and spontaneous emission (dashed line) for a lattice consisting of 10000 atoms spread over 200 disks spaced by $\lambda_0/2$, having thickness $0.04\lambda_0$ and radius $15\lambda_0$. The spontaneous emission of the lattice is normalized by that of a single-atom.

October 8, 2013 0:27

World Scientific Book - 9in x 6in

One-dimensional photonic band gaps in optical lattices

1.4.3 Modelling finite-size effects

To illustrate the superiority of the microscopic model under experimentally realistic conditions we now study the deflection of light reflection in onedimensional lattices. As observed in [Slama *et al.* (2005b)], due to the finite lattice size, any deviation of the lattice constant from the Bragg condition for the lattice period leads to an extra inclination of the scattered beam. Fig. 1.11 shows the position where the reflected beam impinges on a CCD camera for various lattice constants chosen to satisfy or not the Bragg condition for a given angle of incidence.




Fig. 1.11 Variation of the reflection angle with the lattice period when the probe laser is at resonance ($\Delta_0 = 0$). For the chosen angle of incidence, $\theta_0 = 15.9^{\circ}$, the Bragg condition is fulfilled for the lattice laser wavelength of $\lambda_{lat} = 811.07$ nm. The data is recorded using the experimental set-up in Ref. [Slama *et al.* (2005b)].

According to [Slama *et al.* (2005b)], the reflection angle is $\theta_s = \theta_0$ in the case of atomic layers extended to infinity, but should be given by

$$\cos\theta_0 + \cos\theta_s = 2\frac{\lambda_0}{\lambda_{lat}} \tag{1.98}$$

in the case of a *one-dimensional line* of point-like scatterers. This prediction (blue solid line in Fig. 1.12) is confirmed by simulations based on our microscopic model (1.17).

In the intermediate regime, where the slabs are pancake-shaped but have finite radial extension, any deviation of the lattice wavelength from the infinite-slab-Bragg condition $\cos \theta_0 = \cos \theta_s = \frac{\lambda_0}{\lambda_{lat}}$ leads to a deviation of

My Book Title

Paper-mh




Fig. 1.12 Deviation of the scattering angle θ_s from the infinite-slab-Bragg reflection angle $-\theta_0$. The blue solid line represents a theoretical expectation (1.98), the black circles are obtained from simulations of the microscopic model (1.17). The simulations are realized for a lattice of 8000 atoms spread over 400 small disks of thickness $0.04\lambda_0$, radius $0.001\lambda_0$ and spaced by λ_{lat} . The incident laser is inclinated by $\theta_0 = 15.9^{\circ}$.


the scattered beam, $\theta_s \neq \theta_0$ depending on the aspect ratio of the pancakes. In this regime the Bragg condition (1.98) has to be generalized, as shown in Ref. [Slama *et al.* (2005b)]. On the other hand, as shown above, this physics is contained in our microscopic simulations based on (1.17).

Another example demonstrates the utility of the microscopic model. While the TM formalism describes the scattering of a plane-wave from a lattice of infinite transverse size, in reality the probe laser has a Gaussian intensity profile and a finite divergence. In order to study the influence of a finite sized probe laser beam on the PBG, we perform numerical simulations for disks of radius $R_d = 9\lambda_0$. Fig. 1.13 shows that the reflectivity reaches a maximum value and the LDOS is minimal for a laser waist in the range of $3 - 6\lambda_0$. The presence of this maximum is rather intuitive, as if the waist is larger than radial size of the lattice, the laser beam will not be fully intercepted by the lattice, while if it is too small, its divergence is so high that the beam encounters a lower number of disks. In both cases, the contrast of the PBG is reduced.

As a final remark, we note that finite-size effects of the optical lattice have been investigated for a 3D diamond configuration [Antezza and Castin

World Scientific Book - 9in x 6in

Paper-mh



One-dimensional photonic band gaps in optical lattices

Fig. 1.13 (color online) Reflection coefficient (dark circles) and LDOS (blue crosses) derived from the scalar model as the laser waist is tuned. The simulations are realized for N = 6000 atoms distributed over $N_d = 60$ disks of radius $R_0 = 9\lambda_0$ and thickness $a = 0.05\lambda_0$. The laser is incident under the angle $\theta_0 = 2^\circ$ and is resonant, i.e., $\Delta_0 = 0$. Picture taken from [Samoylova *et al.* (2013)].

(2013)]. In this work, the three-dimensional band gap, and more specifically the LDOS, is shown to be strongly affected by either vacancies in the lattice sites or by the finite size.

1.5 Conclusions and Perspectives

The opening of a photonic band gap in an optical lattice is just another manifestation of cooperative scattering. The entire physics of photonics bands is therefore contained within a microscopic collective scattering model that was first described by R. Dicke [Dicke (1954)] and earned increasing attention in recent years [Friedberg *et al.* (1973); Scully *et al.* (2006); Courteille *et al.* (2010)]. In comparison to the single-photon superradiance, Bragg and Mie scattering, photonic bands occur in an optically dense regime where photons are scattered by different atoms multiple times. Moreover, in contrast to the most frequently studied situation of disordered atomic clouds, photonic bands are a consequence of the order imposed onto the scatterers by the presence of a lattice.

In this paper we aimed at supporting the above assertion by applying

Paper-mh

My Book Title

the microscopic cooperative scattering model to the only physical system that has been studied in experiment: a one-dimensional optical lattice, producing a stack of pancake-shaped atomic clouds. We proceeded with two approaches. First, the classical TM formalism approach, used in the past to treat 1D optical lattices, has been derived from our cooperative scattering model under certain idealized assumptions. This was followed by numerical simulations of the cooperative scattering model which are compared to theoretical predictions of the TM formalism. However, the cooperative scattering model allows working beyond the TM formalism and the incorporation of a variety of experimental constraints, some of which can have a huge impact on observations. To grasp the full range of phenomena expected in light scattering from optically dense lattices, a vectorial treatment of the light is necessary. Nevertheless, many features can be determined from a simplified scalar model, which turns out to be an excellent approximation in the case of 1D lattices.

The localization of light and the suppression of spontaneous emission, which represent the holy grail of photonic band gaps, can only be expected for omnidirectional band gaps, which in turn are only supported in threedimensional lattices. In fact, only few lattice geometries are expected to sustain complete omnidirectional band gaps, for example, diamond-shaped lattices [Antezza and Castin (2009)] and cubic lattices of driven three-level atoms [Yu (2011)]. Such a lattice would reflect light incident from any direction. Additionally, an excited atom inside the lattice (one could imagine, for example, that the excited state of the atom was reached via spontaneous emission from an upper state) can not emit a photon because the lattice does not offer electromagnetic modes the photon could use. Hence, spontaneous emission is inhibited and the light is trapped in a localized spot. In this respect, there is an interesting analogy to Anderson localization except that in the case of PBGs it is not disorder, but order causing localization.

It is important to stress that the microscopic approach not only is perfectly suited in describing any type of 3D optical lattices, but it naturally offers the possibility to *incorporate disorder and finite-size effects*. It will be the task of upcoming studies to identify the most suitable lattice geometries and other parameters for the detection of omnidirectional band gaps under experimentally realistic conditions. Only then will it be possible to evaluate the potential technological power of optical lattices in molding the flow of light and to compare it with alternative techniques. Indeed, photonic crystals made from dielectric materials are very advanced and powerful devices [Ishizaki *et al.* (2013)]. However, these are technically limited as photonic October 8, 2013 0:27

World Scientific Book - 9in x 6in

Paper-mh

One-dimensional photonic band gaps in optical lattices

crystals have issues maintaining their periodic order over long distances due to the fabrication process. Here optical lattices are superior because the periodicity is imposed by the lattice lasers and is inherently perfect. Another decisive advantage is the dynamic control possible over optical lattices, which may be ramped up and down or even reshaped in real time.

Acknowledgements

This work has been supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and the Research Executive Agency (Program COSCALI, Grant No. PIRSES-GA-2010-268717).

.6 Appendix: Derivation of the Transfer Matrix T

Setting $z = z_n$ and $z = z_n + a$ in (1.36) we obtain, respectively

$$\frac{\Omega_0}{\Gamma} e^{ik_{0z}z_n} = (2\delta + i)\overline{\beta}_n(z_n) + i\alpha e^{-ik_{0z}z_n} \sum_{m \ge n} \int_{z_m}^{z_m + a} dz' e^{ik_{0z}z'} \overline{\beta}_m(z')
+ i\alpha e^{ik_{0z}z_n} \sum_{m < n} \int_{z_m}^{z_m + a} dz' e^{-ik_{0z}z'} \overline{\beta}_m(z'),$$
(.99)

and

S

$$\frac{Q_0}{\Gamma} e^{ik_{0z}(z_n+a)} = (2\delta+i)\overline{\beta}_n(z_n+a)
+ i\alpha e^{ik_{0z}(z_n+a)} \sum_{m \le n} \int_{z_m}^{z_m+a} dz' e^{-ik_{0z}z'}\overline{\beta}_m(z')
+ i\alpha e^{-ik_{0z}(z_n+a)} \sum_{m>n} \int_{z_m}^{z_m+a} dz' e^{ik_{0z}z'}\overline{\beta}_m(z'), \quad (.100)$$

where $\alpha = 2\pi\rho_0/(k_0k_{0z})$. Using (1.40), the integrals give:

$$i\alpha \int_{z_m}^{z_m+a} dz' e^{ik_{0z}z'} \overline{\beta}_m(z') = [Ax_m + By_m] e^{ik_{0z}z_m},$$

$$i\alpha \int_{z_m}^{z_m+a} dz' e^{-ik_{0z}z'} \overline{\beta}_m(z') = \left[Be^{i(k_z - k_{0z})a} x_m + Ae^{i(k_z + k_{0z})a} y_m \right] e^{-ik_{0z}z_m},$$

where

$$A = \frac{k_z - k_{0z}}{2k_{0z}} \left[1 - e^{i(k_z + k_{0z})a} \right], \tag{.101}$$

$$B = \frac{k_z + k_{0z}}{2k_{0z}} \left[1 - e^{-i(k_z - k_{0z})a} \right], \qquad (.102)$$

Paper-mh

My Book Title

and (1.39) and $k_0^2(m_0^2-1) = k_z^2 - k_{0z}^2$ were used. By inserting the above equations into Eqs.(.99) and (.100), we obtain:

$$\frac{\Omega_0}{\Gamma} = \left[(1+A)x_n + (1+B)y_n \right] e^{-ik_{0z}z_n}
+ \sum_{m < n} \left(Be^{ik_z a}x_m + Ae^{-ik_z a}y_m \right) e^{-ik_{0z}(z_m+a)}
+ e^{-2ik_{0z}z_n} \sum_{m > n} \left(Ax_m + By_m \right) e^{ik_{0z}z_m}, \quad (.103)$$

$$\frac{\Omega_0}{\Gamma} = \left[(1+B)e^{ik_z a}x_n + (1+A)e^{-ik_z a}y_n \right] e^{-ik_{0z}(z_n+a)}
+ \sum \left(Be^{ik_z a}x_m + Ae^{-ik_z a}y_m \right) e^{-ik_{0z}(z_m+a)}$$

$$= \sum_{m < n}^{m < n} + e^{-2ik_{0z}(z_n + a)} \sum_{m > n} (Ax_m + By_m) e^{ik_{0z}z_m}.$$
 (.104)

Subtraction (.104) from (.103) gives:

$$0 = e^{ik_{0z}z_n} \left[(1+A) - (1+B)e^{i(k_z - k_{0z})a} \right] x_n + e^{ik_{0z}z_n} \left[(1+B) - (1+A)e^{-i(k_z + k_{0z})a} \right] y_n + \sum_{m=n+1}^N \left(1 - e^{-2ik_{0z}a} \right) e^{ik_{0z}z_m} \left(Ax_m + By_m \right),$$
(.105)

where n = 1, ..., N - 1. Multiplying (.104) by $\exp(2ik_{0z}a)$ and subtracting it from (.103) we obtain:

$$\frac{\Omega_0}{\Gamma} \left(1 - e^{2ik_{0z}a} \right) = e^{-ik_{0z}z_n} \left[(1+A) - (1+B)e^{i(k_z+k_{0z})a} \right] x_n
+ e^{-ik_{0z}z_n} \left[(1+B) - (1+A)e^{-i(k_z-k_{0z})a} \right] y_n
+ \sum_{m=1}^{n-1} \left(1 - e^{2ik_{0z}a} \right) e^{-ik_{0z}z_m}
\times \left[Be^{i(k_z-k_{0z})a}x_m + Ae^{-i(k_z+k_{0z})a}y_m, \right]$$
(.106)

where n = 2, ..., N. In order to find the solution for N slabs the complete system provided by Eqs. (.105) and (.106) for the coefficients x_n and y_n should be solved. The most convenient way to do that is the iterative method. We introduce

$$K_n = e^{ik_{0z}z_n} \left\{ \left[(1+A) - (1+B)e^{i(k_z - k_{0z})a} \right] x_n + \left[(1+B) - (1+A)e^{-i(k_z + k_{0z})a} \right] y_n \right\}$$
(.107)

One-dimensional photonic band gaps in optical lattices

and

$$L_n = \left(1 - e^{-2ik_{0z}a}\right) e^{ik_{0z}z_n} \left(Ax_n + By_n\right), \qquad (.108)$$

so that (.105) is equivalent to

$$K_n + \sum_{m=n+1}^{N} L_m = 0, \quad n = 1, 2, ..., N - 1$$
 (.109)

or

$$K_{n+1} - L_{n+1} = K_n, \quad n = 1, 2, ..., N - 1.$$
 (.110)

Analogously, (.106) can be written in the following form:

$$M_n + \sum_{m=1}^{n-1} N_m = \frac{\Omega_0}{\Gamma} \left(1 - e^{2ik_{0z}a} \right), \quad n = 2, ..., N,$$
(.111)

or

$$M_{n+1} = M_n - N_n, \quad n = 1, 2, ..., N - 1,$$
 (.112)

where

$$M_n = e^{-ik_{0z}z_n} \left\{ \left[(1+A) - (1+B)e^{i(k_z+k_{0z})a} \right] x_n + \left[(1+B) - (1+A)e^{-i(k_z-k_{0z})a} \right] y_n \right\}$$
(.113)

and

$$N_n = \left(1 - e^{2ik_{0z}a}\right)e^{-ik_{0z}z_n} \left\{Be^{i(k_z - k_{0z})a}x_n + Ae^{-i(k_z + k_{0z})a}y_n\right\}.$$
 (.114)

From Eqs.(.110) and (.112) it is possible to obtain recurrence equations connecting the coefficients (x_{n+1}, y_{n+1}) with (x_n, y_n) in a similar fashion described in [Boedecker and Henkel (2003)]. Using Eqs.(.101) and (.102), a straightforward calculation allows to write Eqs.(.110) and (.112) in the matrix form:

$$R\begin{bmatrix} x_{n+1}\\ y_{n+1} \end{bmatrix} = S\begin{bmatrix} x_n\\ y_n \end{bmatrix}$$
(.115)

where

$$R = \begin{bmatrix} k_{0z} + k_z & k_{0z} - k_z \\ k_{0z} - k_z & k_{0z} + k_z \end{bmatrix}$$
(.116)

and

$$S = \begin{bmatrix} (k_{0z} + k_z)e^{i(k_{0z}d + k_z a)} & (k_{0z} - k_z)e^{i(k_{0z}d - k_z a)} \\ (k_{0z} - k_z)e^{-i(k_{0z}d - k_z a)} & (k_{0z} + k_z)e^{-i(k_{0z}d + k_z a)} \end{bmatrix}.$$
 (.117)

50

My Book Title

Multiplying both sides of (.115) by the inverse of the left matrix, we obtain:

$$\begin{bmatrix} x_{n+1} \\ y_{n+1} \end{bmatrix} = T \begin{bmatrix} x_n \\ y_n \end{bmatrix}$$
(.118)

where

$$T = \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$$
(.119)

is the transfer matrix with the elements

01

$$T_{11,22} = \left[\cos(k_{0z}d) \pm i\frac{k_{0z}^2 + k_z^2}{2k_{0z}k_z}\sin(k_{0z}d)\right]e^{\pm ik_z a},$$

$$T_{12,21} = \pm i\frac{k_{0z}^2 - k_z^2}{2k_{0z}k_z}\sin(k_{0z}d)e^{\mp ik_z a}.$$
(.120)

We notice that $M_1 = \frac{\Omega_0}{\Gamma} (1 - e^{2ik_{0z}a})$ and $K_N = 0$. In addition to these, the iteration equation (1.49) at n = N - 1 provides a link between the coefficients (x_N, y_N) for the Nth slab with those written for the first slice (x_1, y_1) :

$$x_N = T_{11}^{N-1} x_1 + T_{12}^{N-1} y_1,$$

$$y_N = T_{21}^{N-1} x_1 + T_{22}^{N-1} y_1,$$
(.121)

so that the coefficients (x_1, y_1) can be derived explicitly. The long but straightforward calculations yield:

$$x_1 = \frac{2\kappa_{0z}}{(k_{0z} + k_z)^2 - (k_{0z} - k_z)^2 e^{2ik_z a}}$$
(.122)

$$\times \frac{[k_{0z} + k_z - (k_{0z} - k_z)re^{2ik_{0z}d}]\sin(N-1)\phi - t(k_{0z} + k_z)\sin(N-2)\phi}{(1 - r^2e^{2ik_{0z}d})\sin(N-1)\phi - t\sin(N-2)\phi},$$

$$y_{1} = \frac{2\kappa_{0z}}{(k_{0z} - k_{z})^{2} - (k_{0z} + k_{z})^{2} e^{-2ik_{z}a}}$$
(.123)

$$\times \frac{[k_{0z} - k_{z} - (k_{0z} + k_{z})re^{2ik_{0z}d}]\sin(N-1)\phi - t(k_{0z} - k_{z})\sin(N-2)\phi}{(1 - r^{2}e^{2ik_{0z}d})\sin(N-1)\phi - t\sin(N-2)\phi},$$

where r and t are the reflection and transmission coefficients of a single slab given by Eqs.(1.55) and (1.56), respectively.

.7 Appendix: Vectorial model

In this appendix we summarize the derivation of equations (1.57) for the vectorial scattering model [Manassah (2012b)]. We consider N two-level

World Scientific Book - 9in x 6in

One-dimensional photonic band gaps in optical lattices

atoms, where the ground and excited states are connected by an electric dipole transition. We assume that for the *j*th atom the ground state $|g_j\rangle$ is a singlet and the excited state $|e_j^{(\alpha)}\rangle$ is a degenerate triplet, with $\alpha = x, y, z$. For the *j*th atom we define the lowering operators $\hat{\sigma}_j^{(\alpha)} = |g_j\rangle\langle e_j^{(\alpha)}|$ and the electric dipole operator $\hat{\mathbf{d}}_j$ with components $\hat{d}_j^{(\alpha)} = d\hat{\sigma}_j^{(\alpha)} \exp(-i\omega_a t) + c.c.$, where *d* is the matrix element equal for the three transitions. The interaction Hamiltonian includes three contributions:

$$H_I = H_L + H_{dd} + H_{rad}, (.124)$$

where

$$H_L = \frac{d}{2} \sum_{\alpha} E_{0\alpha} \sum_{j=1}^{N} \left[\hat{\sigma}_j^{(\alpha)} e^{i(\Delta_0 t - \mathbf{k}_0 \cdot \mathbf{r}_j)} + \text{h.c.} \right]$$
(.125)

is the interaction between the atoms and the incident classical field with frequency $\omega_0 = ck_0$ and wave vector \mathbf{k}_0 , where $E_{0\alpha}$ are the electric field components and $\Delta_0 = \omega_0 - \omega_a$,

$$H_{dd} = \frac{1}{4\pi\epsilon_0} \sum_{i < j} \frac{1}{r_{ij}^3} \left[\hat{\mathbf{d}}_i \cdot \hat{\mathbf{d}}_j - 3(\hat{\mathbf{d}}_i \cdot \hat{\mathbf{r}}_{ij})(\hat{\mathbf{d}}_j \cdot \hat{\mathbf{r}}_{ij}) \right]$$
(.126)

is the instantaneous Coulomb interaction between pairs of atoms, where $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j = r_{jm} \hat{\mathbf{r}}_{jm}$, and

$$H_{rad} = \hbar \sum_{j=1}^{N} \sum_{\alpha} \sum_{\mathbf{k}, \hat{\epsilon}} g_k \hat{\epsilon}_{\alpha} \left(\hat{\sigma}_j^{(\alpha)} e^{-i\omega_a t} + \text{h.c.} \right) \left[\hat{a}_{\mathbf{k}, \hat{\epsilon}} e^{-i(\omega_k t - \mathbf{k} \cdot \mathbf{r}_j)} + \text{h.c.} \right]$$
(127)

is the interaction between atoms and the vacuum radiation field, $g_k = d(\omega_a^2/2\hbar\epsilon_0 V_{ph}\omega_k)^{1/2}$ is the single-photon Rabi frequency, V_{ph} is the photon volume and $\hat{\epsilon}$ is the polarization unit vector. The dipole-dipole interaction (.126) can be written in a more compact form, neglecting the rapidly time varying terms, as

$$H_{dd} = \frac{1}{8\pi\epsilon_0} \sum_{j \neq m} \sum_{\alpha,\beta} \left[\hat{\sigma}_j^{(\alpha)} \hat{\sigma}_m^{\dagger(\beta)} + \hat{\sigma}_j^{\dagger(\alpha)} \hat{\sigma}_m^{(\beta)} \right] W_{\alpha,\beta}(\mathbf{r}_{jm}), \qquad (.128)$$

where

$$W_{\alpha,\beta}(\mathbf{r}) = \frac{d^2}{r^3} (\delta_{\alpha,\beta} - 3\hat{r}_{\alpha}\hat{r}_{\beta}).$$
(.129)

For a weak incident beam we restrict our analysis to the case where only one photon is absorbed by the atoms. We also consider the non-rotatingwave terms in the Hamiltonian (.127), so the system atoms+photons has

Paper-mh

My Book Title

the form:

$$\begin{split} |\Psi\rangle &= \alpha(t)|g_1\dots g_N\rangle|0\rangle_{\mathbf{k}} + e^{-i\Delta_0 t}\sum_{j=1}^N\sum_{\alpha}\beta_j^{(\alpha)}(t)|g_1\dots e_j^{(\alpha)},\dots g_N\rangle|0\rangle_{\mathbf{k}} \\ &+ \sum_{\mathbf{k},\hat{\epsilon}}\gamma_{\mathbf{k},\hat{\epsilon}}(t)|g_1\dots g_N\rangle|1\rangle_{\mathbf{k},\hat{\epsilon}} \\ &+ \sum_{\mathbf{k},\hat{\epsilon}}\sum_{m\neq n}\sum_{\alpha,\beta}\epsilon_{m,n,\mathbf{k},\hat{\epsilon}}^{(\alpha,\beta)}(t)|g_1\dots e_m^{(\alpha)}\dots e_n^{(\beta)}\dots g_N\rangle|1\rangle_{\mathbf{k},\hat{\epsilon}}. \end{split}$$
(.130)

The first term in (.130) corresponds to the initial ground state without photons, the second term is the sum over the states where a single atom has been excited by the classical field. The third term corresponds to the atoms that returned to the ground state having emitted a photon in the mode \mathbf{k} with polarization $\hat{\epsilon}$, whereas the last term characterizes the presence of two excited atoms and one virtual photon with 'negative' energy. It is due to the counter-rotating terms in the Hamiltonian (.127) and disappears when the rotating wave approximation is made.

With the ansatz (.130) the Schrödinger's equation reduces to the following system of coupled equations:

$$\dot{\alpha} = -i\frac{d}{2\hbar}\sum_{j,\alpha} E_{0\alpha}\beta_{j}^{(\alpha)}e^{-i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}, \qquad (.131)$$

$$\dot{\beta}_{j}^{(\alpha)} = i\Delta_{0}\beta_{j}^{(\alpha)} - \frac{i}{\hbar}\sum_{m,\beta} W_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)} - i\frac{dE_{0\alpha}}{2\hbar}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}\alpha$$

$$-i\sum_{\mathbf{k},\hat{\epsilon}}g_{k}\hat{\epsilon}_{\alpha}e^{-i(\omega_{k}-\omega_{0})t+i\mathbf{k}\cdot\mathbf{r}_{j}}\gamma_{\mathbf{k},\hat{\epsilon}}$$

$$-i\sum_{\mathbf{k},\hat{\epsilon}}g_{k}\sum_{m\neq j,\beta}\hat{\epsilon}_{\beta}e^{-i(\omega_{a}+\omega_{k}-\Delta_{0})t+i\mathbf{k}\cdot\mathbf{r}_{m}}\epsilon_{j,m,\mathbf{k},\hat{\epsilon}}^{(\alpha,\beta)}, \qquad (.132)$$

$$\dot{\gamma}_{\mathbf{k},\hat{\epsilon}} = -ig_k e^{i(\omega_k - \omega_0)t} \sum_j \sum_{\alpha} \hat{\epsilon}_{\alpha} e^{-i\mathbf{k}\cdot\mathbf{r}_j} \beta_j^{(\alpha)}, \qquad (.133)$$

$$\dot{\epsilon}^{(\alpha,\beta)}_{j,m,\mathbf{k},\hat{\epsilon}} = -ig_k e^{i(\omega_k + \omega_a - \Delta_0)t} \hat{\epsilon}_{\alpha} \left[e^{-i\mathbf{k}\cdot\mathbf{r}_j} \beta_m^{(\beta)} + e^{-i\mathbf{k}\cdot\mathbf{r}_m} \beta_j^{(\beta)} \right]. \quad (.134)$$

By integrating Eqs.(.133) and (.134) with $\gamma_{\mathbf{k},\hat{\epsilon}}(0) = 0$, $\epsilon_{j,m,\mathbf{k},\hat{\epsilon}}^{(\alpha,\beta)}(0) = 0$ and

October 8, 2013 0:27

One-dimensional photonic band gaps in optical lattices

substituting them into (.132), assuming $\alpha \approx 1$ (linear regime), we obtain:

$$\dot{\beta}_{j}^{(\alpha)} = i\Delta_{0}\beta_{j}^{(\alpha)} - \frac{i}{\hbar}\sum_{m,\beta} W_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)} - i\frac{dE_{0\alpha}}{2\hbar}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}$$

$$-\sum_{\mathbf{k},\hat{\epsilon}}g_{k}^{2}\sum_{m,\beta}\hat{\epsilon}_{\alpha}\hat{\epsilon}_{\beta}e^{i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}\int_{0}^{t}dt'\beta_{m}^{(\beta)}(t')e^{-i(\omega_{k}-\omega_{0})(t-t')}$$

$$-\sum_{\mathbf{k},\hat{\epsilon}}g_{k}^{2}\sum_{\beta}\hat{\epsilon}_{\alpha}\hat{\epsilon}_{\beta}\int_{0}^{t}dt'e^{i(\Delta_{0}-\omega_{k}-\omega_{a})(t-t')}$$

$$\times\left[\sum_{m\neq j}e^{-i\mathbf{k}\cdot(\mathbf{r}_{j}-\mathbf{r}_{m})}\beta_{m}^{(\beta)}(t') + (N-1)\beta_{j}^{(\beta)}(t')\right]. \quad (.135)$$

The interaction with the vacuum field yields diagonal terms with m = j, whose real part corresponds to the single-atom decay term and imaginary part to the self-energy shift, and off-diagonal term with $m \neq j$, which is related to the atom-atom interaction mediated by the photon. By separating the two contributions, we write:

$$\dot{\beta}_{j}^{(\alpha)} = i\Delta_{0}\beta_{j}^{(\alpha)} - \frac{i}{\hbar}\sum_{m,\beta} W_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)} - i\frac{dE_{0\alpha}}{2\hbar}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}$$
$$-\sum_{\beta}\int_{0}^{t}d\tau\Lambda_{\alpha,\beta}(\tau)\beta_{j}^{(\beta)}(t-\tau)$$
$$-\sum_{m\neq j}\sum_{\beta}\int_{0}^{t}d\tau K_{\alpha,\beta}(\mathbf{r}_{jm},\tau)\beta_{m}^{(\beta)}(t-\tau), \qquad (.136)$$

where

$$\Lambda_{\alpha,\beta}(\tau) = \sum_{\mathbf{k},\hat{\epsilon}} g_k^2 \sum_{\beta} \hat{\epsilon}_{\alpha} \hat{\epsilon}_{\beta} \left[e^{-i(\omega_k - \omega_0)\tau} + (N-1)e^{i(\Delta_0 - \omega_k - \omega_a)\tau} \right]$$
(.137)

is the self-interaction term and

$$K_{\alpha,\beta}(\mathbf{r}_{jm},\tau) = e^{i\Delta_0\tau} \sum_{\mathbf{k},\hat{\epsilon}} g_k^2 \hat{\epsilon}_{\alpha} \hat{\epsilon}_{\beta} e^{-i\omega_k\tau} \left\{ e^{i\mathbf{k}\cdot\mathbf{r}_{jm}} e^{i\omega_a\tau} + e^{-i\omega_a\tau} e^{-i\mathbf{k}\cdot\mathbf{r}_{jm}} \right\}$$
(.138)

is the inter-atom interaction term. The photon polarization factor in Eqs.(.137) and (.138) can be written as

$$\sum_{\hat{\epsilon}} \hat{\epsilon}_{\alpha} \hat{\epsilon}_{\beta} = \delta_{\alpha,\beta} - \hat{k}_{\alpha} \hat{k}_{\beta}$$

Paper-mh

My Book Title

and the summation over \mathbf{k} can be replaced by integration:

$$\int d\mathbf{k} \to \frac{V_{ph}}{8\pi^3} \int dk k^2 \int_0^{2\pi} d\phi \int_0^{\pi} d\theta \sin \theta.$$

The self-interaction term (.137) varies on a time scale much faster than the atomic response, and we can set $\beta_j^{(\beta)}(t-\tau) \approx \beta_j^{(\beta)}(t)$ and $t \to \infty$ in the integral with $\Lambda_{\alpha,\beta}$ in (.136). Then, the real part of this integral yields:

$$\int_{0}^{\infty} d\tau \operatorname{Re}\Lambda_{\alpha,\beta}(\tau) = \frac{V_{ph}}{8\pi^{3}} \int_{0}^{\infty} dkk^{2}g_{k}^{2} \sum_{\beta} \int_{0}^{2\pi} d\phi \int_{0}^{\pi} d\theta \sin\theta \left(\delta_{\alpha,\beta} - \hat{k}_{\alpha}\hat{k}_{\beta}\right) \\ \times \pi \left[\delta(\omega_{k} - \omega_{0}) + (N - 1)\delta(\omega_{k} + \omega_{a})\right], \qquad (.139)$$

where $\omega_k = ck$. The last term, coming from the counter-rotating wave terms of the Hamiltonian (.127), does not contribute since it corresponds to a negative photon energy $\omega_k = -\omega_a$, so that

$$\int_{0}^{\infty} d\tau \operatorname{Re}\Lambda_{\alpha,\beta}(\tau) = \frac{\Gamma}{8\pi} \sum_{\beta} \int_{0}^{2\pi} d\phi \int_{0}^{\pi} d\theta \sin\theta \left(\delta_{\alpha,\beta} - \hat{k}_{\alpha}\hat{k}_{\beta}\right), (.140)$$

where $\Gamma = d^2 k_0^3 / 2\pi \epsilon_0 \hbar$. The remaining angular integration gives

$$\int_0^{2\pi} d\phi \int_0^{\pi} d\theta \sin \theta \left(\delta_{\alpha,\beta} - \hat{k}_{\alpha} \hat{k}_{\beta} \right) = \frac{8\pi}{3} \delta_{\alpha,\beta}.$$

Hence, the energy decay rate of the isolated atom is $(2/3)\Gamma$. Notice that it differs by a factor 2/3 from the scalar decay rate Γ , which does not include polarization effects.

The imaginary part of the time integral of $\Lambda_{\alpha,\beta}$ consists of the self-energy shift of the atom in the excited state plus the self-energy contribution of the atom in the ground state. Its effect is an adjustment to the transition frequency ω_a , which we assume to be already introduced. Indeed, it cannot be directly treated in our model that treats the atoms as point particles, so it is disregarded in the present approach. With these results, (.136) becomes:

$$\dot{\beta}_{j}^{(\alpha)} = \left(i\Delta_{0} - \frac{\Gamma}{3}\right)\beta_{j}^{(\alpha)} + \frac{1}{i\hbar}\sum_{m,\beta}W_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)} - i\frac{dE_{0\alpha}}{2\hbar}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}}$$
$$-\sum_{m\neq j}\sum_{\beta}\int_{0}^{t}d\tau K_{\alpha,\beta}(\mathbf{r}_{jm},\tau)\beta_{m}^{(\beta)}(t-\tau).$$
(.141)

Now we face the evaluation of the time-dependent kernel $K_{\alpha,\beta}(\mathbf{r}_{jm},\tau)$ defined by (.138). Summing over polarization and replacing **k** by integration,

One-dimensional photonic band gaps in optical lattices

we obtain:

$$K_{\alpha,\beta}(\mathbf{r}_{jm},\tau) = \frac{V_{ph}}{(2\pi)^3} e^{i\Delta_0\tau} \int_0^\infty dk k^2 g_k^2 e^{-i\omega_k\tau} \int_0^{2\pi} d\phi \int_0^\pi d\theta \sin\theta \\ \times \left(\delta_{\alpha,\beta} - \hat{k}_\alpha \hat{k}_\beta\right) \left\{ e^{ikr_{jm}\cos\theta + i\omega_a\tau} + e^{-ikr_{jm}\cos\theta - i\omega_a\tau} \right\},$$
(.142)

where $r_{jm} = |\mathbf{r}_{jm}|$. Notice that the polar angle θ depends on the orientation of the vector \mathbf{r}_{jm} , but for simplicity of notation we have omitted the subscript (j, m). The integration over the azimuth angle ϕ yields:

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi \left(\delta_{\alpha,\beta} - \hat{k}_\alpha \hat{k}_\beta \right) = \frac{1}{2} (1 + \cos^2 \theta) \delta_{\alpha,\beta} + \frac{1}{2} (1 - 3\cos^2 \theta) \hat{r}_\alpha \hat{r}_\beta.$$
(.143)

Using (.143) in (.142) and integrating over θ ,

$$K_{\alpha,\beta}(\mathbf{r},\tau) = \frac{\Gamma c^2}{\pi \omega_a} e^{i\Delta_0 \tau} \cos(\omega_a \tau) \int_0^\infty dk k e^{-ick\tau} \left\{ \left(\delta_{\alpha,\beta} - \hat{r}_\alpha \hat{r}_\beta\right) \frac{\sin(kr)}{kr} + \left(\delta_{\alpha,\beta} - 3\hat{r}_\alpha \hat{r}_\beta\right) \left(\frac{\cos(kr)}{(kr)^2} - \frac{\sin(kr)}{(kr)^3}\right) \right\},$$
(.144)

where we used the definitions of g_k^2 and Γ . Then, by integrating by parts the last term in the braces of (.144) and assuming $\exp(i\Delta_0\tau)\cos(\omega_a\tau) \approx \cos(\omega_0\tau)$ since $\Delta_0 \ll \omega_0$ and $\omega_a \sim \omega_0$, we obtain:

$$K_{\alpha,\beta}(\mathbf{r},\tau) = \frac{\Gamma c^2}{\pi\omega_0} \cos(\omega_0 \tau) \left\{ \frac{1}{r} \left(\delta_{\alpha,\beta} - \hat{r}_{\alpha} \hat{r}_{\beta} \right) \int_0^\infty dk e^{-ick\tau} \sin(kr) \right. \\ \left. + \frac{1}{r^3} \left(\delta_{\alpha,\beta} - 3\hat{r}_{\alpha} \hat{r}_{\beta} \right) \left(ic\tau \int_0^\infty dk e^{-ick\tau} \frac{\sin(kr)}{k} - r \right) \right\} (.145)$$

Equation (.145) contains the following integrals:

$$\int_0^\infty dk e^{-ick\tau} \sin(kr) = \frac{1}{2} \lim_{\epsilon \to 0^+} \left[\frac{1}{r - c\tau + i\epsilon} + \frac{1}{r + c\tau - i\epsilon} \right], \quad (.146)$$

$$\int_0^\infty dk e^{-ick\tau} \frac{\sin(kr)}{k} = \frac{1}{2i} \left[\ln \left| \frac{\tau + r/c}{\tau - r/c} \right| + i\pi\theta(r - c\tau) \right], \qquad (.147)$$

where $\theta(x) = 1$ for x > 0 and $\theta(x) = 0$ for x < 0. We observe that $K_{\alpha,\beta}(\mathbf{r},\tau)$ depends on $\tau \pm r/c$. Assuming that $\beta_j^{(\alpha)}$ varies on a characteristic time much longer than r_{jm}/c between any two atoms j, m in the sample, $\beta_j^{(\beta)}(t-\tau)$ can be replaced by $\beta_j^{(\beta)}(t)$ during the integration time and the

World Scientific Book - $9 \text{in} \times 6 \text{in}$

Paper-mh

My Book Title

upper integration limit can be extended to infinity. The integration over τ of $K_{\alpha,\beta}(\mathbf{r},\tau)$ requires the further evaluation of the integrals:

$$\int_0^\infty d\tau \cos(\omega_0 \tau) \left[\frac{1}{r - c\tau + i\epsilon} + \frac{1}{r + c\tau - i\epsilon} \right] = \int_{-\infty}^\infty d\tau \frac{\cos(\omega_0 \tau)}{r - c\tau + i\epsilon} = \frac{\pi}{ic} e^{ik_0 r}$$
(.148)

and

$$\int_{0}^{\infty} d\tau \cos(\omega_{0}\tau) \left\{ \frac{c\tau}{2} \left[\ln \left| \frac{\tau + r/c}{\tau - r/c} \right| + i\pi\theta(r - c\tau) \right] - r \right\}$$
$$= \frac{i\pi}{2ck_{0}^{2}} \left[(1 - ik_{0}r)e^{ik_{0}r} - 1 \right].$$
(.149)

Using Eqs.(.146)-(.149), we obtain:

$$K_{\alpha,\beta}(\mathbf{r}) = \frac{\Gamma}{2} \left\{ \frac{1}{i(k_0 r)} \left(\delta_{\alpha,\beta} - \hat{r}_{\alpha} \hat{r}_{\beta} \right) e^{ik_0 r} + \frac{i}{(k_0 r)^3} \left(\delta_{\alpha,\beta} - 3\hat{r}_{\alpha} \hat{r}_{\beta} \right) \left[(1 - ik_0 r) e^{ik_0 r} - 1 \right] \right\}.$$
 (.150)

The contribution from the non-oscillating term in (.150) cancels the electrostatic term $-(i/\hbar)W_{\alpha,\beta}$ in (.141). Reintroducing the subscript i, j, the equation for $\beta_j^{(\alpha)}$ is finally obtained:

$$\dot{\beta}_{j}^{(\alpha)} = \left(i\Delta_{0} - \frac{\Gamma}{3}\right)\beta_{j}^{(\alpha)} - i\frac{dE_{0\alpha}}{2\hbar}e^{i\mathbf{k}_{0}\cdot\mathbf{r}_{j}} - \frac{\Gamma}{2}\sum_{m\neq j}\sum_{\beta}G_{\alpha,\beta}(\mathbf{r}_{jm})\beta_{m}^{(\beta)},$$
(.151)

where

$$G_{\alpha,\beta}(\mathbf{r}) = \frac{e^{ik_0r}}{ik_0r} \left\{ \left[\delta_{\alpha,\beta} - (\hat{r}_{\alpha}\hat{r}_{\beta}] + \left[\delta_{\alpha,\beta} - 3\hat{r}_{\alpha}\hat{r}_{\beta} \right] \left[\frac{i}{k_0r} - \frac{1}{(k_0r)^2} \right] \right\}.$$
(.152)

.8 Appendix: Vectorial field

The expression for the radiation field can be obtained from the Maxwell equations in the presence of a polarization \mathbf{P} :

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},\tag{153}$$

$$\nabla \times \mathbf{B} = \mu_0 \frac{\partial \mathbf{D}}{\partial t},\tag{154}$$

$$\nabla \cdot \mathbf{D} = 0, \tag{.155}$$

October 8, 2013 0:27

One-dimensional photonic band gaps in optical lattices

where $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$. From Eqs.(.153) and (.154) we obtain:

$$\nabla \times \nabla \times \mathbf{E} = -\frac{\partial}{\partial t} (\nabla \times \mathbf{B}) = -\mu_0 \frac{\partial^2}{\partial t^2} (\epsilon_0 \mathbf{E} + \mathbf{P}), \quad (.156)$$

$$\nabla \cdot \mathbf{E} = -\frac{1}{\epsilon_0} \nabla \cdot \mathbf{P}. \tag{.157}$$

Since $\nabla \times \nabla \times \mathbf{E} = \nabla (\nabla \cdot \mathbf{E}) - \nabla^2 \mathbf{E}$ and $\epsilon_0 \mu_0 = c^{-2}$, then

$$\nabla^{2}\mathbf{E} - \frac{1}{c^{2}}\frac{\partial^{2}\mathbf{E}}{\partial t^{2}} = \frac{1}{\epsilon_{0}c^{2}}\frac{\partial^{2}\mathbf{P}}{\partial t^{2}} - \frac{1}{\epsilon_{0}}\nabla(\nabla\cdot\mathbf{P}).$$
 (.158)

For a monochromatic field with frequency $\omega_0 = ck_0$ we have:

$$\left(\nabla^2 + k_0^2\right) \mathbf{E} = -\frac{k_0^2}{\epsilon_0} \left[\mathbf{P} + \frac{1}{k_0^2} \nabla (\nabla \cdot \mathbf{P}) \right].$$
(.159)

Using (1.63), the solution of (.159) is

$$\mathbf{E}(\mathbf{r}) = i \frac{k_0^3}{4\pi\epsilon_0} \int d\mathbf{r}' G(|\mathbf{r} - \mathbf{r}'|) \left[\mathbf{P}(\mathbf{r}') + \frac{1}{k_0^2} \nabla(\nabla \cdot \mathbf{P}(\mathbf{r}')) \right]$$
(.160)

or for each component:

$$E_{\alpha}(\mathbf{r}) = i \frac{k_0^3}{4\pi\epsilon_0} \sum_{\beta} \int d\mathbf{r}' G(|\mathbf{r} - \mathbf{r}'|) \left[\delta_{\alpha,\beta} + \frac{1}{k_0^2} \frac{\partial^2}{\partial x'_{\alpha} \partial x'_{\beta}} \right] P_{\beta}(\mathbf{r}'). \quad (.161)$$

By integrating by parts and using (1.59), we obtain:

$$E_{\alpha}(\mathbf{r}) = i \frac{k_0^3}{4\pi\epsilon_0} \sum_{\beta} \int d\mathbf{r}' G_{\alpha,\beta}(\mathbf{r} - \mathbf{r}') P_{\beta}(\mathbf{r}').$$
(.162)

Taking $P_{\alpha}(\mathbf{r}) = -d\rho\beta^{(\alpha)}(\mathbf{r})$ and returning to a discrete distribution of atoms with position \mathbf{r}_{j} , the scattered field components at the position \mathbf{r} are

$$E_{\alpha}(\mathbf{r}) = -i\frac{dk_0^3}{4\pi\epsilon_0} \sum_{\beta} \sum_m G_{\alpha,\beta}(\mathbf{r} - \mathbf{r}_m)\beta_m^{(\beta)}.$$
 (.163)

Combining Eqs.(.163) and (.151), we find that the stationary atomic polarization components are given by:

$$\beta_j^{(\alpha)} = \frac{d}{\hbar \left(\Delta_0 + i\Gamma/3\right)} \left[\frac{E_{0\alpha}}{2} e^{i\mathbf{k}_0 \cdot \mathbf{r}_j} + E_\alpha(\mathbf{r}_j)\right].$$
 (.164)

58

 $My \ Book \ Title$

Bibliography

- Antezza, M. and Castin, Y. (2009). Fano-hopfield model and photonic band gaps for an arbitrary atomic lattice, *Phys. Rev. A* 80, p. 013816.
- Antezza, M. and Castin, Y. (2013). Photonic band-gap in a realistic atomic diamond lattice: penetration depth, finite-size and vacancy effects, arXiv:1304.7188.
- Bachelard, R., Bender, H., Courteille, P. W., Piovella, N., Stehle, C., Zimmermann, C. and Slama, S. (2012a). Role of mie scattering in the seeding of matter-wave superradiance, *Phys. Rev. A* 86, p. 043605.
- Bachelard, R., Courteille, P., Kaiser, R. and Piovella, N. (2012b). Resonances in mie scattering by an inhomogeneous atomic cloud, *Europhys. Lett.* 97, p. 14004.
- Bachelard, R., Piovella, N. and Courteille, P. W. (2011). Cooperative scattering and radiation pressure force in dense atomic clouds, *Phys. Rev. A* 84, p. 013821.
- Bender, H., Stehle, C., Slama, S., Kaiser, R., Piovella, N., Zimmermann, C. and Courteille, P. W. (2010). Observation of cooperative mie scattering from an ultracold atomic cloud, *Phys. Rev. A* 82, p. 011404.
- Bienaimé, T., Bachelard, R., Chabé, J., Rouabah, M., Bellando, L., Courteille, P., Piovella, N. and Kaiser, R. (2013). Interplay between radiation pressure force and scattered light intensity in the cooperative scattering by cold atoms, *submitted*.
- Bienaimé, T., Bux, S., Lucioni, E., Courteille, P., Piovella, N. and Kaiser, R. (2010). Observation of a cooperative radiation force in the presence of disorder, *Phys. Rev. Lett.* **104**, p. 183602.
- Bienaime, T., Petruzzo, M., Bigerni, D., Piovella, N. and Kaiser, R. (2011). Atom and photon measurement in cooperative scattering by cold atoms, J. Mod. Opt. 58, p. 1942.
- Birkl, G., Gatzke, M., Deutsch, I. H., Rolston, S. L. and Phillips, W. D. (1995). *Phys. Rev. Lett.* **75**, p. 2823.
- Boedecker, G. and Henkel, C. (2003). All-frequency effective medium theory of a photonic crystal, *Opt. Express* **11**, p. 1590.
- Bux, S., Lucioni, E., Bender, H., Bienaim, T., Lauber, K., Stehle, C., Zimmer-

My Book Title

mann, C., Slama, S., Courteille, P., Piovella, N. and Kaiser, R. (2010). Cooperative scattering by cold atoms, *J. of Mod. Opt.* **57**, pp. 1841–1848.

- Coley, J. M. (1995). Diffraction Physics (North-Holland, Amsterdam).
- Courteille, P. W., Bux, S., Lucioni, E., Lauber, K., Bienaim, T., Kaiser, R. and Piovella, N. (2010). Modification of radiation pressure due to cooperative scattering of light, *Euro. Phys. J. D* 58, p. 69.
- Deutsch, I. H., Spreeuw, R. J. C., Rolston, S. L. and Phillips, W. D. (1995). Photonic band gaps in optical lattices, *Phys. Rev. A* 52, p. 1394.
- Dicke, R. H. (1954). Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, p. 99.
- Doucet, J. and Benoit, J. P. (1987). Molecular dynamics studied by analysis of the x-ray diffusion scattering by isozyme crystals, *Nature (London)* **325**, p. 644.
- Fleischhauer, M., Imamoglu, A. and Marangos, J. P. (2005). Rev. Mod. Phys. 77, p. 633.
- Friedberg, R., Hartmann, S. R. and Manassah, J. T. (1973). Frequency shifts in emission and absorption by resonant systems of two-level atoms, *Phys. Rep. C* 7, p. 101.
- Glauber, R. J. and Prasad, S. (2000). Polarium model: Reflection and transmission of coherent radiation, *Phys. Rev. A* 61, p. 063815.
- Harris, S. E. (1997). Electromagnetically induced transparency, *Phys. Today* **50**, p. 36.
- Ishizaki, K., Koumura, M., Suzuki, K., Gondaira, K. and Noda, S. (2013). Realization of three-dimensional guiding of photons in photonic crystals, *Nature Phot.* 7, p. 133.
- Jackson, J. D. (1999). Classical electrodynamics, 3rd edn. (Wiley, New York, NY), ISBN 9780471309321.
- John, S. and Stephen, M. J. (1983). Wave propagation and localization in a longrange correlated random potential, *Phys. Rev. B* 28, p. 6358.
- Keaveney, J., Sargsyan, A., Krohn, U., Hughes, I. G., Sarkisyan, D. and Adams, C. S. (2012). Cooperative lamb shift in an atomic vapor layer of nanometer thickness, *Phys. Rev. Lett.* **108**, p. 173601.
- Khaykovich, L. and Davidson, N. (1999). Compression of a cold atomic cloud by on-resonance laser light, J. Opt. Soc. Am. B 16, p. 702.
- Koenderink, A. F. and Vos, W. L. (2003). Light exiting from real photonic band gap crystals is diffuse and strongly directional, *Phys. Rev. Lett.* **91**, p. 213902.
- Manassah, J. T. (2012a). Comparison of the cooperative emission profile from a spherical distribution of two-level atoms resulting from the choice of the interaction kernel, *Phys. Rev. A* **85**, p. 015801.
- Manassah, J. T. (2012b). Cooperative radiation from atoms in different geometries: decay rate and frequency shift, Advances in Optics and Photonics 4, pp. 108–156.
- Mazets, I. E. and Kurizki, G. (2007). Multiatom cooperative emission following single-photon absorption: Dicke-state dynamics, J. Phys. B 40, p. F105.
- Petrosyan, D. (2007). Phys. Rev. A 76, p. 053823.

World Scientific Book - 9in x 6in

Bibliography

- Piovella, N. and Bachelard, R. (2012). Cooperative light scattering by a cold atomic fluid, AIP Conf. Proc. 1421, p. 73.
- Piovella, N., Bachelard, R. and Courteille, P. (2013). Fluid description of the cooperative scattering of light by spherical atomic clouds, J. of Plasma Physics .
- Prasad, S. and Glauber, R. (2010). Coherent radiation by a spherical medium of resonant atoms, *Phys. Rev. A* 82, p. 063805.
- Prasad, S. and Glauber, R. J. (2000). Polarium model: Coherent radiation by a resonant medium, *Phys. Rev. A* **61**, p. 063814.
- Prasad, S. and Glauber, R. J. (2011). Coherent scattering by a spherical medium of resonant atoms, *Phys. Rev. A* 83, p. 063821.
- Röhlsberger, R., Schlage, K., Sahoo, B., Couet, S. and Rüffer, R. (2010). Collective lamb shift in single-photon superradiance, *Science* **328**, p. 1239.
- Samoylova, M., Piovella, N., Bachelard, R. and Courteille, P. (2013). Microscopic theory of photonic band gaps in optical lattices, *submitted*.
- Schilke, A., Zimmermann, C., Courteille, P. and Guerin, W. (2011). Photonic band gaps in one-dimensionally ordered cold atomic vapors, *Phys. Rev. Lett.* **106**, p. 223903.
- Schilke, A., Zimmermann, C., Courteille, P. and Guerin, W. (2012a). Optical parametric oscillation with distributed feedback in cold atoms, *Nature Phot.* 6, p. 101.
- Schilke, A., Zimmermann, C. and Guerin, W. (2012b). Photonic properties of one-dimensionally-ordered cold atomic vapors under conditions of electromagnetically induced transparency, *Phys. Rev. A* 86, p. 023809.
- Scully, M. O., Fry, E., Ooi, C. H. R. and Wodkiewicz, K. (2006). Directed spontaneous emission from an extended ensemble of n atoms: Timing is everything, *Phys. Rev. Lett.* **96**, p. 010501.
- Scully, M. O. and Svidzinsky, A. A. (2009). The effects of the n atom collective lamb shift on single photon superradiance, *Phys. Lett. A* 373, p. 1283.
- Scully, M. O. and Svidzinsky, A. A. (2010). The lamb shiftyesterday, today, and tomorrow, *Science* **328**, p. 1239.
- Sesko, D. W., Walker, T. G. and Wieman, C. E. (1991). Behavior of neutral atoms in a spontaneous force trap, J. Opt. Soc. Am. B 8, p. 946.
- Skipetrov, S. E. and Goetchy, A. (2011). J. Phys. A 44, p. 065102.
- Slama, S., Bux, S., Krenz, G., Zimmermann, C. and Courteille, P. W. (2006). Multiple reflections and diffuse scattering in bragg scattering at optical lattices, *Phys. Rev. A* 73, p. 023424.
- Slama, S., von Cube, C., Deh, B., Ludewig, A., Zimmermann, C., and Courteille, P. W. (2005a). Phase-sensitive detection of bragg scattering at 1d optical lattices, *Phys. Rev. Lett.* **94**, p. 193901.
- Slama, S., von Cube, C., Ludewig, A., Kohler, M., Zimmermann, C. and Courteille, P. W. (2005b). Dimensional crossover in bragg scattering from an optical lattice, *Phys. Rev. A* 72, p. 031402(R).
- Svidzinsky, A. A. (2012). Nonlocal effects in single-photon superradiance, *Phys. Rev. A* 85, p. 013821.
- Svidzinsky, A. A., Chang, J. and Scully, M. O. (2008). Dynamical evolution of

Paper-mh

My Book Title

correlated spontaneous emission of a single photon from a uniformly excited cloud of n atoms, *Phys. Rev. Lett.* **100**, p. 160504.

Svidzinsky, A. A., Chang, J. and Scully, M. O. (2010). Cooperative spontaneous emission of n atoms: Many-body eigenstates, the effect of virtual lamb shift processes, and analogy with radiation of n classical oscillators, *Phys. Rev.* A **81**, p. 053821.

van Coevorden, D. V., Sprik, R., Tip, A. and Lagendijk, A. (1996). Photonic band structure of atomic lattices, *Phys. Rev. Lett.* **77**, p. 2412.

- Walker, T., Sesko, D. and Wieman, C. (1990). Collective behavior of optically trapped neutral atoms, *Phys. Rev. Lett.* 64, p. 408.
- Weidemüller, M., Görlitz, A., Hänsch, T. W. and Hemmerich, A. (1998). *Phys. Rev. A* 58, p. 4647.
- Weidemüller, M., Hemmerich, A., Görlitz, A., Esslinger, T. and Hänsch, T. W. (1995). Phys. Rev. Lett. 75, p. 4583.
- Wollan, E. O. (1932). X-ray scattering and atomic structure, aRev. Mod. Phys. 4, pp. 205–258.
- Yu, D. (2011). Photonic band structure of the three-dimensional ⁸⁸sr atomic lattice, *Phys. Rev. A* 84, p. 043833.