

Original Study

Open Access

Stefano De Benedetti, Giorgio Lucchini, Alessandro Marocchi, Silvana Penco,
Christian Lunetta, Stefania Iametti*, Elisabetta Gianazza, Francesco Bonomi

Serum metal evaluation in a small cohort of Amyotrophic Lateral Sclerosis patients reveals high levels of thiophylic species

DOI 10.1515/ped-2015-0004

Received July 28, 2015; accepted October 16, 2015

Abstract: Amyotrophic Lateral Sclerosis (ALS) has often been associated with improper/changed metal metabolism. Analysis of thiophylic metals in serum from a small and geographically restricted cohort of ALS patients indicates contents of Pb and Ni much higher in patients than in controls (Ni, 5-fold; Pb, 2-fold). Se levels are also higher in the patients' group, which has instead lower As levels than controls. Thiophylic metals may impair biogenesis of FeS clusters or substitute for iron, even in folded proteins; Se may non-functionally replace S. Thus, improper assembly/function of FeS proteins could represent another possible issue to be considered in ALS pathogenesis.

Keywords: Amyotrophic Lateral Sclerosis, Serum metals, ICP-MS, Iron Sulfur Clusters

*Corresponding author **Stefania Iametti:** DeFENS, University of Milan, Via Celoria 2, 20133, Milan, Italy, E-mail: stefania.iametti@unimi.it

Stefano De Benedetti, Francesco Bonomi, Department of Food, Environmental and Nutritional Sciences (DEFENS), University of Milan, 20133 Milan, Italy

Giorgio Lucchini, Department of Agricultural and Environmental Sciences (DiSAA), University of Milan, 20133 Milan, Italy.

Stefano De Benedetti, Alessandro Marocchi, Silvana Penco, Medical Genetics Unit, Department of Laboratory Medicine, Niguarda Ca' Granda Hospital, 20169, Milan, Italy

Christian Lunetta, NEuroMuscular Omnicentre (NEMO), Fondazione Serena Onlus, Niguarda Ca' Granda Hospital, 20169, Milan, Italy

Elisabetta Gianazza, Proteomics and Protein Structure Study Group, Division of Pharmacological Sciences, Department of Pharmacological and Biomolecular Sciences (DiSFeB), University of Milan, 20133 Milano, Italy

1 Introduction

Amyotrophic Lateral Sclerosis (ALS) is a rare progressive neurodegenerative disorder with an incidence of 2.1 per 100 000 person-years and an estimated prevalence of 5.4 cases per 100 000 population [1]. This progressive disease is characterized by a selective degeneration of both upper and lower motor neurons in the brain, brainstem, and spinal cord resulting in paralysis due to muscle weakness and atrophy. ALS usually leads to death in 3 to 5 years upon the onset of symptoms [2]. Two forms of ALS can be recognized: the sporadic (SALS) accounts for 90–95% of the cases, while the familial form (FALS) accounts for the remaining 5–10% [3]. Both forms show similar clinical features suggesting a common pathogenic pathway. ALS is the most common disorder among motor neuron diseases with adult onset. Despite extensive research, its etiology is still unknown. Mutations in several genes, including SOD1, FUS and TARDBP, have been identified as responsible for the disease in 25–35% of cases of the familial forms [4–6]. Recently, the pathogenic expansion of a non-coding hexanucleotide repeat sequence (GGGGCC) in the C9ORF72 gene was reported in familial and sporadic forms of ALS [7,8].

Other proposed pathogenic mechanisms involve protein aggregation, oxidative stress, impairment of mitochondrial function, transcription dysfunctions, alterations in the proteasome pathway, inflammation, excitotoxicity, and environmental influences [9]. Among the environmental factors, metals have been hypothesized to play a – still much un-clarified – role in ALS. Indeed, through the years many neurodegenerative disorders, such as Alzheimer Disease, Parkinson Disease and ALS, have been linked to iron and metal metabolism in a number of studies [10,11].

Transition metal-induced toxicity also has been proposed to be involved in the pathology [12] and different authors reported increased concentrations of metals,

as well as of proteins regulating their homeostasis, in ALS patients [13-15]. Metals are essential for different physiological processes and their concentrations must be strictly regulated since imbalance can lead to toxic effects. Environmental, occupational and nutritional levels of metals could influence their body concentrations and result in overload.

In this work, we report on a small cohort of subjects affected by the sporadic form of ALS, originating from a restricted geographical area with a high prevalence of the disease. We evaluated concentrations of different metals in serum trying to establish a correlation with ALS, assuming that the limited geographical area would allow to assess the role played by specific chemical species in the environment and in the diet of the involved subjects.

2 Methods

The study enrolled seven patients (4 men and 3 women) and five controls (2 men and 3 women, age matched – 69 years vs. 64 years – and living in the same geographical area) (Table 1 and Table 2). All patients and controls were recruited by Medical Genetics Unit of Niguarda Ca' Granda Hospital and signed an informed consent approved by local ethics committee. Controls were represented by caregivers of the patients, known to be free from neurological disorders and living in the same geographic area. Blood was collected from all subjects. Serum was obtained by centrifugation after clotting and stored at -80°C.

ALS diagnosis was according El Escorial criteria [9] with clinically defined sporadic cases; one presenting bulbar form at the onset, while the six with spinal form. All subjects were under Riluzole treatment; two of them underwent Percutaneous Endoscopic Gastrostomy (PEG).

For metal quantitation, samples were analyzed by ICP-MS (Bruker AURORA M90 ICP-MS). Serum was diluted 1:20 with 0.05% Triton X-100 in MilliQ water. Seronom™ Trace Elements Serum L-1 was used to build appropriate calibration curves. An aliquot of a 2 mg L⁻¹ of an internal standard solution (⁴⁵Sc, ⁸⁹Y, ¹⁵⁹Tb) was added to both samples and standards to give a final concentration of 20 µg L⁻¹ of each element. Typical analysis interferences were removed by using Collision-Reaction-Interface (CRI) with an H₂ flow of 70 mL min⁻¹ through the skimmer cone.

Statistical analyses were carried out with Student's t-Test and through multivariate statistical analyses tools, including Principal Component Analysis (PCA) and Auto Contractive Map algorithm (Auto-CM), a type of Artificial Neural Network (ANN) able to define the strength of the associations of each variable with all the others and to visually show the map of the main connections; this approach has already been applied to studies on the ALS disease [16].

3 Results

Serum concentrations of Cr, Fe, Ni, Cu, Zn, As, Se, Sr, Cd, Pb are reported in Table 3. A statistically significant difference between patients and controls was found for Ni

Table 1: Minimum clinical data for ALS patients. LL: lower limbs, UL: upper limbs. Duration of disease is calculated as time passed from symptoms onset to blood sampling or PEG start.

ID	Gender	Age at onset (y)	Site at onset	Duration of disease (y)	PEG	BMI at onset (m ² Kg ⁻¹)	Occupation	Smoke
1 SU	M	48	LL	2.1	N	29.4	Shopkeeper	N
2 SU	M	57	Bulbar	3.0	Y	23.0	Manager	Y
3 SU	M	65	UL	3.2	N	28.1	Workman	N
4 SU	F	60	LL	19.1	N	22.9	Seamstress	N
5 SU	F	71	LL	12.1	N	27.9	Farmer	N
6 SU	M	69	LL	9.0	N	23.7	Manager	N
7 SU	F	59	LL	4.2	Y	29.3	Teacher	EX

Table 2: Data for control subjects.

ID	Gender	Current age (y)	BMI (m ² Kg ⁻¹)	Occupation	Smoke
244 N	F	55	20.8	Odontologist	N
245 N	F	66	22.3	Teacher	Y
246 N	F	65	25.7	Housewife	EX
247 N	M	72	28.1	Professor	N
248 N	M	61	31.1	Manager	EX

($p = 0.0001$), Pb ($p = 0.01$) and As ($p = 0.05$). The median Ni concentration was $9.44 \mu\text{g L}^{-1}$ for the ALS patients, about five times higher than in controls ($2.14 \mu\text{g L}^{-1}$). The median Pb concentration in the patients' group was almost twice that of controls (2.16 vs $1.26 \mu\text{g L}^{-1}$). Conversely, the median As concentration was lower in the patients' group than in controls' ($0.51 \mu\text{g L}^{-1}$ vs $0.73 \mu\text{g L}^{-1}$). In both groups, Ni and Pb levels were higher than the values reported for the general Italian population. Indeed, according to data reported by the Italian Superior Health Institute [17], Ni concentrations should be in the $0.26\text{--}0.75 \mu\text{g L}^{-1}$ range, and Pb concentrations in the $0.20\text{--}0.98 \mu\text{g L}^{-1}$ range.

Principal Component Analysis (PCA) was able to discriminate between the patients and the controls. The

most prominent feature of the control group was a high concentration of As and a low concentrations of all the other metals analyzed. The semantic connectivity map generated with the Auto-CM system (Figure 1) confirmed this observation. This algorithm is able to visually show the connections between variables and to describe the strength of the connections with a numerical value from 0 to 1. Results clearly discriminate the two groups and define an ALS area characterized by high levels of metals, except for As. Among the three metals that were significantly different, none individually emerged as more relevant than the others in the discrimination of the controls' group from the patients' group. The Auto-CM analysis also indicated that high levels of Se were closely

Table 3: Averages of the measures of metals concentrations in sera. *: p -value ≤ 0.05 , NA: Not Available.

Element	Average Patients \pm SD ($\mu\text{g/L}$)	Average Controls \pm SD ($\mu\text{g/L}$)	Reference Values ($\mu\text{g/L}$) [17]
Cr	1.57 ± 0.12	1.54 ± 0.06	0.07-0.28
Fe	1261.28 ± 429.00	1225.94 ± 160.00	648-1301
Ni	$9.44 \pm 1.02^*$	$2.10 \pm 0.92^*$	0.26-0.75
Cu	1130.24 ± 157.00	1141.55 ± 108.00	648-1301
Zn	811.03 ± 114.00	835.88 ± 72.40	597-1028
As	$0.51 \pm 0.14^*$	$0.73 \pm 0.18^*$	NA
Se	97.71 ± 10.20	89.54 ± 6.32	56-105
Sr	39.73 ± 12.50	34.54 ± 5.71	23-61.5
Cd	0.08 ± 0.03	0.06 ± 0.01	0.03-0.2
Pb	$2.16 \pm 0.72^*$	$1.26 \pm 0.29^*$	0.2-0.98




Figure 1: Semantic connectivity map obtained with Auto-Cm System. The Minimum Spanning Tree (MST) shows the connections between the variables. Values on the arches of the graph refer to the strength of the association between two adjacent nodes. The range of this value is

1 linked to the ALS group, whereas low levels of Se were
2 connected with the controls. Even if Se levels did not
3 reach statistical significance with common statistic tools
4 ($p = 0.12$), artificial neural network multivariate statistical
5 analyses point to a possible relevance of the observed
6 differences (Figure 1).

7 8 **4 Discussion**

9
10 In the last decades, Amyotrophic Lateral Sclerosis has
11 been the focus of intense research. Despite this, the
12 etiology of the disease still has to be clarified. Genetic
13 traits only partly explain this disorder [18] and, at
14 present, fail to assess a common and clear molecular
15 pattern that leads to the disease. Among subjects
16 bearing a pathological mutation in one of the causative
17 gene, a wide phenotypical heterogeneity is present [19],
18 and this is true also in the sporadic form of the disease
19 [20]. Clearly, some other factors must be involved in the
20 pathogenesis of the disease; some of the triggers could be
21 environmental influences [18]. We decided to focus our
22 investigations on metals' concentrations in a small group
23 of subjects originating from a restricted geographical
24 region due to the higher prevalence of the disease in this
25 area, in comparison with literature data. In our intentions,
26 limiting the investigation to a restricted area could help
27 in pinpointing environmental or dietary contributions to
28 specific metal species.

29
30 We did not performed any study on environmental
31 pollution, but we are aware of the presence of a steel plant
32 within this area. Furthermore, there is a Cu and Fe mine,
33 closed in 1962, located 8 km behind the inhabited area,
34 within the basin of a creek, whose waters are reported to
35 be strongly polluted due to Acid Mine Drainage. Waste-
36 rock dumps can store significant amounts of potentially
37 deleterious metals that can be released to solutions
38 during transformation processes, induced by variations in
39 the physicochemical parameters or by ageing. This could
40 lead to the release, to the circulating solutions, of high
41 amounts of potentially toxic metals during weathering of
42 sulfide mineralization (Fe, Cu, Zn, etc.) and host rocks (Cr,
43 Ni, V, etc.) [21]. Among the metals analyzed, Pb resulted
44 higher in patients' than in controls' group. Much evidence
45 in the literature describes Pb as a neurotoxic agent and as
46 being related to ALS [22,23], since high levels of this metal
47 have been found both in blood and in cerebrospinal fluid
48 of ALS patients [24,25]. In our analyses, Pb levels were
49 twice higher in ALS subjects than in controls, supporting
50 the hypothesis of a possible involvement of this metal in
51 the etiology of the disease.

Pb²⁺ ions have a substantial affinity for protein cysteine
thiols, and could be involved in direct substitution of iron
in various types of FeS clusters, a property that has been
demonstrated *in vitro* for various thiophylic metals [26,27].
Rates and yields of cluster-metal substitution often depend
on the redox state of the cluster iron [27,28] and on the
accessibility of the ligands into the protein structure, as
dictated by folding/unfolding rates and equilibria [27,29].

Another metal ion able to substitute for iron (in
particular when ligands other than cysteine are involved)
is Ni²⁺. In this study, Ni concentration was found almost
five times higher in ALS patients than in controls. A
Ni overload could promote oxidative stress and the
production of Reactive Oxygen Species, as well as an
increase in inflammatory markers [30]. However, recent
evidence points to a direct role of Ni in the modulation
of the biosynthetic pathways of FeS proteins in neural
cells [31]. According to this report, high Ni²⁺ levels act by
repressing the expression of the mitochondrial scaffold
proteins IscU1 and IscU2, which are essential to FeS
biogenesis [32]. According to this study, this leads to a
cellular energy metabolism shift, likely as a result of an
alteration in the mitochondrial respiratory chain and
in the related energy fluxes [33]. The relevance of these
issues is currently being investigated.

Arsenic can form both inorganic and organic
compounds in the environment and in the human body,
as either arsenite (As³⁺) or arsenate (As⁵⁺). Since absorbed
arsenate is mostly reduced to arsenite in blood, the effect
of the two forms appears to be very similar [30]. Many
studies confirmed the generation of free radicals during
As metabolism in cells, and oxidative stress has been
linked to the development of As-related diseases [34]. In
the light of this, the result obtained is unexpected, as the
levels in our study controls were higher than in patients.
One possible interpretation is that low levels of As are
a feature of this peculiar group of ALS patients, without
hypothesizing that an As overload may exert a protective
role in control subjects. On the other hand, He and Ma
reported that, in mouse hepa1c1c7 cells, As³⁺ induces the
transcription of metallothionein I (MT1) through direct
interaction with metal-activated transcription factor 1
(MTF1) [35], and this could possibly result in a protective
effect, in case of non-toxic levels of As overload.

Auto-CM analysis also pointed to a possible
relationship between ALS and Se levels in serum,
although statistical significance was close to the limit of
acceptability and expanding the number of subjects is
still needed for verification. Abnormal levels of various
Se species have been found in cerebrospinal fluid of ALS
patients [36] and circulating levels of Se, usually bound to

serum albumin, could reflect the Se levels present in the Central Nervous System. Finally, we were not able to find a correlation between Fe levels and ALS in the group of patients considered in this study, although several reports previously assessed this relationship [14,37-39].

Despite these phenomenological reports, the role of these metals in the pathogenesis of ALS remains unclear. We plan to evaluate it in a wider cohort of subjects while investigating a broader panel of metals. Impaired metal homeostasis, hypothetically attributable to environmental exposure, in the end could lead to overload of different elements. Besides promoting oxidative stress, metals can compete for the binding sites of metals-containing proteins, such as those containing iron-sulfur clusters. This damaging effect could be exerted by all those metals having affinity for thiols, like mercury(II), silver(I), copper(I), lead(II), cadmium(II), nickel(II), zinc(II) and cobalt(II) whose toxicity in bacteria has been shown to be proportional to the affinity for sulfur [40].

The role of iron-sulfur clusters biosynthetic machinery in ALS is still poorly explored. Upregulation of Frataxin (Fxn) and of the human ortholog of the cluster-assembly scaffold protein (IscU), two of the proteins involved in carrying out or in regulating FeS biosynthesis, has been explained as a response of the cell to iron overload [38], whereas Ni overload results in the repression of IscU biosynthesis, as discussed above [31]. However, further investigation is required to assess a possible relationship between metal imbalance and the many biosynthetic steps leading to functional metalloproteins, as well as the molecular basis of mid- to long-term effects of metal imbalance on the energy status and the redox balance of neuronal cells. Effects on other proteins involved in metal homeostasis within the cytoplasm or in specific cellular compartments (such as metallothioneins or ferritin [35,38,41]) and on proteins involved in controlling the intracellular redox potential and/or the concentration of active chemical species (such as peroxides and superoxides [19]) also remain to be assessed.

Acknowledgments: We thank the “Comitato per la Fondazione Giovanni Raffo – Onlus” for support and partial funding of this work. We also thank the involved patients and their relatives, along with healthy volunteers, for their kind participation.

References

- [1] Chiò A., Logroscino G., Traynor B.J., Collins J., Simeone J.C., Goldstein L.A., White L.A., Global epidemiology of amyotrophic lateral sclerosis: a systematic review of the published literature, *Neuroepidemiology*, 2013, 41, 118-30.
- [2] Mitchell J.D., Borasio G.D., Amyotrophic lateral sclerosis, *Lancet*, 2007, 369, 2031-2041.
- [3] McLaughlin R.L., Vajda A., Hardiman O., Heritability of Amyotrophic Lateral Sclerosis: Insights From Disparate Numbers., *JAMA Neurol* Published online June 01, 2015. doi:10.1001/jamaneurol.2014.4049.
- [4] Rosen D.R., Siddique T., Patterson D., Figlewicz D.A., Sapp P., Hentati, A., Donaldson D., Goto J., O'Regan J.P., Deng H.X., et al., Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis, *Nature*, 1993, 362, 59–62.
- [5] Sreedharan J., Blair I.P., Tripathi V.B., Hu X., Vance C., Rogelj B., Ackerley S., Durnall J.C., Williams K.L., Buratti E., et al., TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis, *Science*, 2008, 319, 1668–1672.
- [6] Vance C., Rogelj B., Hortobagyi T., De Vos K.J., Nishimura A.L., Sreedharan J., Hu X., Smith B., Ruddy D., Wright P., et al., Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6, *Science*, 2009, 323, 1208–1211.
- [7] DeJesus-Hernandez M., Mackenzie I.R., Boeve B.F., Boxer A.L., Baker M., Rutherford N.J., Nicholson A.M., Finch N.A., Flynn H., Adamson J., et al., Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS, *Neuron*, 2011, 72, 245–456.
- [8] Renton A.E., Majounie E., Waite A., Simón-Sánchez J., Rollinson S., Gibbs J.R., Schymick J.C., Laaksovirta H., van Swieten J.C., Myllykangas L., et al., A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD, *Neuron*, 2011, 72, 257–268.
- [9] Brooks B.R., Miller R.G., Swash M., Munsat T.L., World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis, *Amyotroph Lateral Scler Other Motor Neuron Disord*, 2000, 1(5), 293-9.
- [10] Crichton R.R., Ward R.J., Metal-based neurodegeneration. From molecular mechanisms to therapeutic strategies, *John Wiley and Sons*, pp 227, 2006.
- [11] Hadzhieva M., Kirches E., Mawrin C., Review: iron metabolism and the role of iron in neurodegenerative disorders, *Neuropathol Appl Neurobiol*, 2014, 40(3), 240-57.
- [12] Carrí M.T., Ferri A., Cozzolino M., Calabrese L., Rotilio G., Neurodegeneration in amyotrophic lateral sclerosis: the role of oxidative stress and altered homeostasis of metals, *Brain Res Bull*, 2003, 61(4), 365-74.
- [13] Goodall E.F., Haque M.S., Morrison K.E., Increased serum ferritin levels in amyotrophic lateral sclerosis (ALS) patients, *J Neurol*, 2008, 255(11),1652-6.
- [14] Nadjar Y., Gordon P., Corcia P., Bensimon G., Pieroni L., Meininger V., Salachas F., Elevated serum ferritin is associated with reduced survival in amyotrophic lateral sclerosis, *PLoS One*, 2012, 7(9), e45034.
- [15] Roos P.M., Lierhagen S., Flaten T.P., Syversen T., Vesterberg O., Nordberg M., Manganese in cerebrospinal fluid and blood plasma of patients with amyotrophic lateral sclerosis, *Exp Biol Med (Maywood)*, 2012, 237(7), 803-10.
- [16] Buscema M., Penco S., Grossi E., A Novel Mathematical Approach to Define the Genes/SNPs Conferring Risk or

[1] Chiò A., Logroscino G., Traynor B.J., Collins J., Simeone J.C., Goldstein L.A., White L.A., Global epidemiology of amyotrophic

- 1 Protection in Sporadic Amyotrophic Lateral Sclerosis Based
 2 on Auto Contractive Map Neural Networks and Graph Theory,
 3 *Neurol Res Int.*, 2012, 478560.
- 4 [17] ISTISAN, Alimonti A., Bocca B., Mattei D., Pino A., 2010.
 5 Report 10/22: Biomonitoraggio della popolazione
 6 italiana per l'esposizione ai metalli: valori di riferimento
 7 1990–2009 [Biomonitoring of Italian population for metals
 8 exposure: reference values 1990–2009]. ISSN: 1123-3117.
 9 Istituto Superiore della Sanità [Italian Superior Health
 10 Institute], Available at: [http://www.iss.it/binary/publ/](http://www.iss.it/binary/publ/cont/10ventidueWEB.pdf)
 11 [cont/10ventidueWEB.pdf](http://www.iss.it/binary/publ/cont/10ventidueWEB.pdf). Accessed May 2015.
- 12 [18] Ingre C., Roos P.M., Piehl F., Kamel F., Fang F., Risk factors for
 13 amyotrophic lateral sclerosis, *Clin Epidemiol*, 2015, 7, 181-93.
- 14 [19] Penco S., Lunetta C., Mosca L., Maestri E., Avemaria F.,
 15 Tarlarini C., Patrosso M.C., Marocchi A., Corbo M., Phenotypic
 16 Heterogeneity in a SOD1 G93D Italian ALS Family: An Example
 17 of Human Model to Study a Complex Disease, *J Mol Neurosci*,
 18 2011, 44, 25–30.
- 19 [20] Chio` A., Calvo A., Moglia C., Mazzini L., Mora G., PARALS
 20 study group, Phenotypic heterogeneity of amyotrophic lateral
 21 sclerosis: a population based study, *J Neurol Neurosurg*
 22 *Psychiatry*, 2011, 82, 740-746.
- 23 [21] Marescotti P., Azzali E., Servida D., Carbone C., Grieco G., de
 24 Capitani L., Lucchetti G., Mineralogical and geochemical spatial
 25 analyses of a waste-rock dump at the Libiola Fe-Cu sulphide
 26 mine (Eastern Liguria, Italy), *Environmental Earth Sciences*,
 27 2010, 61 (1), 187-199.
- 28 [22] Oh S.S., Kim E.A., Lee S.W., Kim M.K., Kang S.K., A case of
 29 amyotrophic lateral sclerosis in electronic parts manufacturing
 30 worker exposed to lead, *Neurotoxicology*, 2007, 28, 324–327.
- 31 [23] Fang F., Kwee L.C., Allen K.D., Umbach D.M., Ye W., Watson
 32 M., Keller J., Oddone E.Z., Sandler D.P., Schmidt S., et al.,
 33 Association between blood lead and the risk of amyotrophic
 34 lateral sclerosis, *Am. J. Epidemiol*, 2010, 171, 1126–1133.
- 35 [24] Vinceti M., Guidetti D., Bergomi M., Caselgrandi E., Vivoli R.,
 36 Olmi M., Rinaldi L., Rovesti S., Solime F., Lead, cadmium, and
 37 selenium in the blood of patients with sporadic amyotrophic
 38 lateral sclerosis, *Ital. J. Neurol. Sci.*, 1997, 18, 87–92.
- 39 [25] Roos P.M., Vesterberg O., Syversen T., Peder Flaten T., Nordberg
 40 M., Metal concentrations in cerebrospinal fluid and blood
 41 plasma from patients with amyotrophic lateral sclerosis, *Biol.*
 42 *Trace Elem. Res*, 2013, 151, 159–170.
- 43 [26] Iametti S., Uhlmann H., Sala N., Bernhardt R., Ragg E.M.,
 44 Bonomi F., Reversible, non-denaturing metal substitution in
 45 bovine adrenodoxin and spinach ferredoxin and the different
 46 reactivity of [2Fe-2S]-cluster-containing proteins, *Eur. J.*
 47 *Biochem.*, 1996, 239, 818-826.
- 48 [27] Iametti S., Uhlmann H., Ragg E.M., Sala N., Grinberg A., Beckert
 49 V., Bernhardt R., Bonomi F., Cluster iron substitution is related
 50 to structural and functional features of adrenodoxin mutants
 51 and to their redox state, *Eur. J. Biochem.*, 1998, 251, 673-681
- [28] Lorusso M., Cocco R., Sardanelli A.M., Minuto M., Bonomi F.,
 Papa S., Interaction of zinc ions with the bovine mitochondrial
 b-c1 complex, *Eur. J. Biochem.*, 1991, 197, 555-561.
- [29] Bonomi F., Iametti S., Kurtz D.M., Ragg E.M., Richie K.A., Direct
 metal ion substitution at the [M(SCys)₄]²⁻ site of rubredoxin, *J.*
Biol. Inorg. Chem., 1998, 3, 595-605.
- [30] Valko M., Morris H., Cronin M.T.D., Metals, Toxicity and
 Oxidative Stress, *Current Medicinal Chemistry*, 2005, 12,
 1161-1208.
- [31] He M., Lu Y., Xu S., Mao L., Zhang L., Duan W., Liu C., Pi H.,
 Zhang Y., Zhong M., Yu Z., Zhou Z., MiRNA-210 modulates a
 nickel-induced cellular energy metabolism shift by repressing
 the iron-sulfur cluster assembly proteins ISCU1/2 in Neuro-2a
 cells, *Cell Death Dis.*, 2014, 5, e1090
- [32] Pierik A.J., Netz D.J., Lill R., Analysis of iron-sulfur protein
 maturation in eukaryotes. *Nat. Protoc.*, 2009, 4, 753-766.
- [33] Lill R., Dutkiewicz R., Elsässer H.P., Hausmann A., Netz D.J.,
 Pierik A.J., Stehling O., Urzica E., Mühlenhoff U., Mechanisms
 of iron-sulfur protein maturation in mitochondria, cytosol and
 nucleus of eukaryotes, *Biochim. Biophys. Acta*, 2006, 1763,
 652-667.
- [34] Shi H., Shi X., Liu K.J., Oxidative mechanism of arsenic toxicity
 and carcinogenesis, *Mol Cell Biochem*, 2004, 255(1-2), 67-78.
- [35] He X. and Ma Q., Induction of Metallothionein I by Arsenic via
 Metal-activated Transcription Factor 1, *Jour of Biol Chem*, 2009,
 284(19), 12609-12621.
- [36] Vinceti M., Solovyev N., Mandrioli J., Crespi C.M., Bonvicini
 F., Arcolin E., Georgouloupoulou E., Michalke B., Cerebrospinal
 fluid of newly diagnosed amyotrophic lateral sclerosis patients
 exhibits abnormal levels of selenium species including
 elevated selenite, *NeuroToxicology*, 2013, 38, 25–32.
- [37] Hozumi I., Hasegawa T., Honda A., Ozawa K., Hayashi Y.,
 Hashimoto K., Yamada M., Koumura A., Sakurai T., Kimura
 A., et al., Patterns of levels of biological metals in CSF differ
 among neurodegenerative diseases, *J Neurol Sci*, 2011, 15,
 303(1-2):95-9.
- [38] Hadzhieva M., Kirches E., Wilisch-Neumann A., Pachow D.,
 Wallesch M., Schoenfeld P., Paegle I., Vielhaber S., Petri S.,
 Keilhoff G., et al., Dysregulation of iron protein expression in
 the G93A model of amyotrophic lateral sclerosis, *Neuroscience*,
 2013, 29;230, 94-101.
- [39] Veyrat-Durebex C., Corcia P., Mucha A., Benzimra S., Mallet C.,
 Gendrot C., Moreau C., Devos D., Piver E., Pagès J.C., et al., Iron
 metabolism disturbance in a French cohort of ALS patients,
Biomed Res Int, 2014, 2014:485723.
- [40] Nies D.H., Efflux-mediated heavy metal resistance in
 prokaryotes, *FEMS Microbiol.*, 2003, Rev. 27, 313–339.
- [41] Hashimoto K., Hayashi Y., Watabe K., Inuzuka T., Hozumi I.,
 Metallothionein-III prevents neuronal death and prolongs
 life span in Amyotrophic Lateral Sclerosis model mice,
Neuroscience, 2011, 189, 293-298.