

IS A BIRTH-MONTH-DEPENDENCE OF HUMAN LONGEVITY INFLUENCED BY HALF-YEARLY CHANGES IN GEOMAGNETICS?

Germaine Cornelissen, Cristina Maggioni*, Franz Halberg

Halberg Chronobiology Center, University of Minnesota, Minneapolis, MN, USA (e-mail: halbe001@umn.edu and corne001@umn.edu) *University of Milan, Milan, Italy

Introduction

About-daily and about-yearly variations in organisms are commonly viewed as evolutionary adaptations to changes in the proximate environmental temperature and illumination. There is now ample evidence that a much broader time structure (chronome), long known to characterize the environment, is built into biological variables, from the level of an ecological niche revealed by demographic statistics to that of molecular genetics. While a molecular basis has been established for circadian rhythms, indirect evidence for some endogenicity had long been available by the persistence of rhythms with a period differing from the environmental cycle under constant conditions (free-running) (1, 2). Natural environmental factors have also been shown to play a critical role, notably in terms of synchronizing built-in rhythms (1-3). As reviewed elsewhere (4), non-photic as well as photic effects of the sun may play a role in shaping the element of multifrequency rhythms currently, and may have done so in the past, resulting in even broader chronomes with added elements of chaos and trends (4, 5). Non-photic signatures include, with the biological week (5), the about 10.5-year solar activity cycle, the about 21.0-year Hale bipolarity variation, and a prominent about half-year rhythm peaking at the equinoxes. This natural physical half-year characterizes various indices of geomagnetic activity (6-15) and may relate to the tilt angle of the earth's dipole axis toward and away from the sun, which reportedly is not constant according to Robert L. McPherron. It is particularly prominent when analyzed in K_p (5, 16, 17) by the population-mean cosinor method (18, 19).

Methods

In response to a recent article reviewing non-photic solar influences on biota, in the context of intersecting antiaging and chronoastrobiological issues (4), researchers at the Center on Aging at the University of Chicago kindly brought to our attention their work on the effect of birth month on human longevity (20). In checking for confounding factors in studies on possible trade-offs between human longevity and biological fertility, Gavrilov and Gavrilova unexpectedly found that the month of birth is a predictor for the life expectancy of adult women (30 years and older): women born in May and December tend to live about 3 years longer on the average by comparison with women born in August. Their data are expressed as a difference from the August value, used as reference. They are based on the point estimates of the differential intercept coefficients obtained after adjusting for other variables (such as calendar year of birth, maternal and paternal lifespan, maternal and paternal age at birth, birth order, nationality, cause of death, and loss of mother, father, or both before age 20).

These data were taken off the published graph (20), to be re-analyzed by cosinor (18, 19) in the light of prior association of geomagnetic activity with anthropometry notably at birth (21-24) and at the time of conscription (25, 26) and thus with growth, previously related to light acting via melatonin (26) and/or vasopressin (27). Trial periods of one year and harmonics thereof were considered. More specifically, cosine curves were fitted by least-squares to the data, and the hypothesis of a zero amplitude (no-rhythm) was tested. Components in non-harmonic relation with the year, or any infra-annual component were not investigated because the data, representing cohorts born in 1800-1880, had been stacked over an idealized 1-year span. Data on the antipodal index of geomagnetic activity, aa, available only since 1868 were retrieved from the internet for the span 1868-1880 and similarly stacked as monthly means over an idealized one-year span. A phase difference between aa and lifespan was examined by parameter test (28).

Results

Both geomagnetic activity and human longevity are characterized by a highly statistically significant (P<0.001) halfyear component accounting for over 80% of the variance, as illustrated in Figure 1. The lifespan data lag behind the antipodal geomagnetic aa index by 2 months, as shown by cross-correlation analysis, Figure 2. The phase difference between lifespan and aa is statistically validated (P<0.001). Half-year changes are also documented for length at birth of babies from South Africa (21), Germany (22), Spain (23), and Denmark (24). In the latter data and similar time series from Minnesota and Russia available longitudinally over spans covering up to 112 years, the demonstration of about 10.5 and/or about 21.0-year components (4, 25), signatures of the Schwabe and Hale solar activity cycles, lends further evidence for an influence of non-photic physical environmental (solar?) factors.

A similarity of periodicities between the natural environmental (aa) and the biological (longevity) variables constitutes no more than a hint of putative associations. The result is strengthened by similar findings in other studies, involving longitudinal data series lending themselves better to additional analyses by superposed epochs, cross-spectral






Figure 1. Human longevity studied by Gavrilov and Gavrilova (20) in adult women (30 years and older), in cohorts born in 1800-1880, expressed as a difference for the August value, used as reference, are plotted vs. the month of birth. For comparison, data from the geomagnetic index aa (1868-1880) are plotted after stacking over an idealized 1-year span.

coherence, and remove-andreplace approaches (25, 29). In particular, body length at birth in Russia is crossspectrally coherent with the geomagnetic activity index K_p. Cross-spectral coherence with K_p is also found at a frequency of about 2 cycles per year in our re-analyses of data on adult height from Weber et al. (26), who examined 507,125 male Austrian recruits at 18 years of age, and which show a dependence on the month of birth. Weber et al. postulated effect of seasonably an changing light, accounted for by photic effects upon melatonin. without commenting on a 10-year cyclic aspect of their data that they reported.

Discussion

Putative hypothalamic-pineal mechanisms come to mind in accounting for these associations. A half-year component was documented to characterize vasopressin (27). The pineal, in turn, is reportedly capable of responding to variations of the order of a few nT induced by the solar wind via the magnetosphere (30). Of particular importance is the report by Lerchl et al. (31) that both the synthesis and the release of pineal melatonin in vitro are significantly reduced (P<0.001) by exposure to extremely low frequency Ca⁺⁺-cyclotron resonance, presumably caused by the enzyme N-acetyltransferase. Melatonin, in turn, is implicated in growth. Serum growth hormone concentrations have been reported to increase fivefold after melatonin administration in clinically healthy men (32). Melatonin secretion is also known to increase during pregnancy (33), and is thought to be implicated with cortisol in mechanisms underlying intrauterine growth retardation (34). Whereas no differences in circulating maternal melatonin were found in terms of circadian rhythm characteristics, a half-year component was present only for IUGR. Circulating estriol in maternal blood, which has a fetal origin, shows a marked difference between healthy and IUGR groups at a half-year, as well as along the 24-hour scale. Circulating melatonin itself has been shown to be latitude dependent (35). Furthermore, it undergoes a half-year variation by night occasionally at middle latitudes, and even at noon at higher latitudes (36-38), where magnetic disturbances induced by the solar wind tend to be more pronounced (39). A reduced nocturnal excretion of a melatonin metabolite (6-hydroxymelatonin sulfate) has also been observed in humans exposed to fluctuating magnetic fields (40-42).








Figure 2. Cross-correlation of human longevity vs. the geomagnetic activity index aa shows a maximum at a lag of 2 months, in keeping with the possibility, quantified below, that the maximum of human longevity follows the maximum in aa two months later. A half-year periodicity characterizes the cross-correlation function, which results in maxima occurring at lags -10, -4, 2 and 8 months, and in minima at lags -7, -1, 5 and 11 months. Whereas the results indicate that human longevity and aa share a common period of six months, longevity peaking 2 months after the peak in aa, it cannot be said without further work whether the two variables are causally related. If they were, directly or by mediating variables, then it cannot be said whether the association is positive or negative. It is also not clear at which moment of the child's (embryo's or fetus's) life the effect of geomagnetics operates, if at all. With a possible causal relation and a 2-month lag, the effect may take place when the child is born. Alternative possibilities include a negative association with zero lag if the effect occurs during the second or 8^{th} month of gestation (or a positive association taking place in the 5^{th} gestational month). Should the original data become available in toto (instead of only as 12 monthly means), dividing the data into subsamples of varying aa exposures would allow the testing of a graded response which, should it validate an association, would lend support to geomagnetic influences on human longevity.

A melatonin-dependent circannual variability of the quality of the oocyte has been postulated by Smits et al. (43) to account for an association of fecundity with the month of birth. These authors studied a cohort of 1526 women who married between 1802 and 1931, using only women whose first marriage occurred before the age of 35 years. Childless women, as compared with fecunds, reportedly showed a birth distribution that was represented by a bimodal curve peaking in January and July (43). The question will have to be elucidated whether we deal with a vacation or a geomagnetic effect, which latter should show a latitude dependence (36-38) or probably with both, including perhaps other social effects as well. Effects of melatonin supplementation on longevity, investigated in the experimental laboratory are inconclusive (44-46). Physiological effects of exposure to geomagnetic fields appear complex (47). A systematic recording of biological signals complementing the data bases physicists are assembling on natural environmental factors would gain from analyses aimed at assessing broad interacting time structures (chronomes) (4, 5, 29).

In summary, chronobiologic aging research documented earlier that circadian rhythms (e.g., in core temperature) can involve a greater extent of change than does aging over several decades (48). Hence, if unassessed, these rhythms constitute a great confounder of aging trends. The same warning of a potential confounder applies to an about half-year change. The Gavrilovs' meta-analyzed discovery (20) strengthens the case of heliogeophysical associations with research on aging (25, 49). The similarity of periods, in us and around us, constitutes a hint, but no proof. Latitude-dependence would strengthen the case further, as in a decadal aspect of human motivation (50). The in-phase-ness of

G. Cornelissen, C. Maggioni, and F. Halberg

the cycles in the northern and southern hemispheres would support the global effect again, as shown earlier (25). Crossspectral coherence studies remain to be performed in conjunction with remove-and-replace approaches wherein the sun rather than a surgeon may do the removal and replacement of one of its spectral components (51, 52). With the accumulating evidence, the story of built-in circadians, now extensively covered in journals such as Science and Nature, repeats itself in biological populations and individual components with longer than circadian periods, associated with non-photic corpuscular radiation directly or indirectly affecting the embryo and fetus, and thus influencing longevity.

Acknowledgements. This work was supported by the U.S. Public Health Service (GM-13981); University of Minnesota Supercomputing Institute; Lynn Peterson (United Business Machines, Fridley, Minnesota)

References

- 1. Halberg F. Temporal coordination of physiologic function. Cold Spr Harb Symp quant Biol 1960; 25: 289-310.
- 2. Ehret CF. The sense of time: evidence for its molecular basis in the eukaryotic gene-action system. Advances in Biological and Medical Physics 1974; 15 (0): 47-77.
- 3. Pauly JE. Chronobiology: anatomy in time. Am J Anat 1983; 168: 365-388.
- 4. Halberg F, Cornelissen G, Watanabe Y, et al. Near 10-year and longer periods modulate circadians: intersecting anti-aging and chronoastrobiological research. J Gerontol A Biol Sci Med Sci 2001; 56: M304-M324.
- Halberg F, Cornelissen G, Bingham C, Hillman D, Katinas G, Sampson M, Revilla M, Prikryl P Sr, Prikryl P Jr, Sanchez de la Pena S, Gonzalez C, Amory-Mazaudier C, Bouvet J, Barnwell F, Maggioni C, Sothern RB, Wang ZR, Schwartzkopff O, Bakken E. Season's Appreciations 2001. Neuroendocrinol Lett 2002; 23: 170-187.
- 6. Grafe A. Einige charakterische Besonderheiten des geomagnetischen Sonneneruptionseffektes. Geofisica Pura e Applicata 1958; 40: 172-179.
- 7. Russell CT, McPherron RL. Semiannual variation of geomagnetic activity. J Geophys Res 1973; 78: 92-108.
- Chernosky EJ. Double sunspot-cycle variation in terrestrial magnetic activity, 1884-1963. J Geophys Res 1966; 71: 965-974.
- 9. Lincoln JV, ed. Geomagnetic and solar data. J Geophys Res 1966; 71: 975-977.
- Crooker N, Cliver EW, Tsurutani BT. The semiannual variation of great geomagnetic storms and the postshock Russell-McPherron effect preceding coronal mass ejection. Geophys Res Ltrs 1992; 19: 429-432.
- Gonzalez WD, Clua de Gonzalez AL, Tsurutani BT. Comment on "The semiannual variation of great geomagnetic storms and the postshock Russell-McPherron effect preceding coronal mass ejecta" by N Crooker, EW Cliver, and BT Tsurutani. Geophys Res Ltrs 1993; 20: 1659-1660.
- Crooker NU, Cliver EW. Reply to "Comment on 'The semiannual variation of great geomagnetic storms and the postshock Russell-McPherron effect preceding coronal mass ejecta' by WD Gonzalez, AL Clua de Gonzalez, and BT Tsurutani". Geophys Res Ltrs 1993; 20: 1661-1662.
- 13. McPherron RL, O'Brien TP. Solar wind coupling to the ring current as a function of season and solar cycle (Abstract #SM21D-03). Eos, Trans Am Geophys Union 1999; 80: S274.
- 14. McPherron RL. Diurnal and seasonal effects observed in the Dst index. Abstract #S1-09, 1st S-Ramp Conference, Book of Abstracts, SCOSTEP, Sapporo, Japan, 2000, 13 pp.
- 15. Cliver EW, Kamide Y, Ling AG. Mountains versus valleys: Semiannual variation of geomagnetic activity. J Geophys Res 2000; 105 (A2): 2413-2424.
- Cornelissen G, Halberg F, Obridko VN, Breus TK. Quasi-11-year modulation of global and spectral features of geomagnetic disturbance. Biofizika 1998; 43: 677-680.
- 17. Cornelissen G, Halberg F, Sothern RB, Nikityuk BA, Garcia Alonso L, Syutkina EV, Grafe A, Bingham C. Toward a chronoastrobiology: sunspot cycles and geomagnetism as well as sunshine may modulate human morphology. Russian Morphological Newsletter 1998; [v. 5] N. 3(4): 133-137.
- 18. Halberg F. Chronobiology. Ann Rev Physiol 1969; 31: 675-725.
- 19. Cornelissen G, Halberg F. Chronomedicine. In: Armitage P, Colton T, editors-in-chief. Encyclopedia of Biostatistics, v. 1. Chichester, UK: John Wiley & Sons Ltd., 1998: 642-649.
- 20. Gavrilov LA, Gavrilova NS. Season of birth and human longevity. J Anti-Aging Med 1999; 2 (4): 365-366.
- 21. Henneberg M, Louw GJ. Further studies on the month-of-birth effect on body size: rural schoolchildren and an animal model. Am J Physical Anthropology 1993; 91: 235-244.
- 22. Otto W, Reissig G. Zur Anthropologie der Neugeborenen. 4. Mitteilung. Lange und Gewicht der Neugeborenen in den verschiedenen Monaten. Monatsberichte der Deutschen Akademie der Wissenschaften zu Berlin 1963; 5: 549-559.
- 23. Garcia Alonso L, Hillman D, Cornelissen G, Garcia Penalta X, Wang ZR, Halberg F. Nature, not solely nurture: chronome as well as season governs growth patterns of infants. In: Otsuka K, Cornelissen G, Halberg F, eds. Chronocardiology and Chronomedicine: Humans in Time and Cosmos. Tokyo: Life Science Publishing, 1993: 71-75.
- 24. Wohlfahrt J, Melbye M, Christens P, Andersen A-MN, Hjalgrim H. Secular and seasonal variation of length and weight at birth. The Lancet 1998; 352 (Dec 19/26): 1990.
- 25. Halberg F, Cornelissen G, Otsuka K, et al. Cross-spectrally coherent ~10.5- and 21-year biological and physical cycles, magnetic storms and myocardial infarctions. Neuroendocrinol Lett 2000; 21: 233-258.

Is a birth-month-dependence of human longevity influenced by half-yearly changes in geomagnetics?

- 26. Weber GW, Prossinger H, Seidler H. Height depends on month of birth. Nature 1998; 391: 754-755.
- 27. Portela A, Cornelissen G, Halberg Franz, et al. Metachronanalysis of circannual and circasemiannual characteristics of human suprachiasmatic vasopressin-containing neurons. In vivo 1995; 9: 347-358.
- Bingham C, Arbogast B, Cornelissen Guillaume G, Lee JK, Halberg F. Inferential statistical methods for estimating and comparing cosinor parameters. Chronobiologia 1982; 9: 397-439.
- Cornelissen G, Halberg F, Schwartzkopff O, et al. Chronomes, time structures, for chronobioengineering for "a full life". Biomedical Instrumentation & Technology 1999; 33: 152-187.
- Demaine C, Semm P. Magnetic fields abolish nycthemeral rhythmicity of responses of Purkinje cells to the pineal hormone melatonin in the pigeon's cerebellum. Neuroscience Letters 1986; 72: 158-162.
- 31. Lerchl A, Reiter RJ, Howes KA, Nonaka KO, Stokkan KA. Evidence that extremely low-frequency Ca⁺⁺-cyclotron resonance depresses pineal melatonin synthesis in vitro. Neuroscience Letters 1991; 124: 213-215.
- 32. Coiro V, Vescovi PP. Alcoholism abolishes the effects of melatonin on growth hormone secretion in humans. Neuropeptides 1998; 32: 211-214.
- 33. Kivela A. Serum melatonin during human pregnancy. Acta endocrinologica 1991; 124: 233-237.
- 34. Maggioni C, Cornelissen G, Antinozzi R, Ferrario M, Grafe A, Halberg F. A half-yearly aspect of circulating melatonin in pregnancies complicated by intrauterine growth retardation. Neuroendocrinology Letters 1999; 20: 55-68.
- 35. Wetterberg L, Bratlid T, Knorring L v, Eberhard G, Yuwiler A. A multinational study of the relationships between nighttime urinary melatonin production, age, gender, body size, and latitude. Eur Arch Psychiatr Neurosci 1999; 249: 256-262.
- 36. Tarquini B, Cornelissen G, Perfetto F, Tarquini R, Halberg F. Chronome assessment of circulating melatonin in humans. In vivo 1997; 11: 473-484.
- Martikainen H, Tapanainen J, Vakkuri O, Leppaluoto J, Huhtaniemi I. Circannual concentrations of melatonin, gonadotrophins, prolactin and gonadal steroids in males in a geographical area with a large annual variation in daylight. Acta endocrinol (Copenhagen) 1985; 109: 446-450.
- Bergiannaki J, Paparrigopoulos TJ, Stefanis CN. Seasonal pattern of melatonin excretion in humans: relationship to daylength variation rate and geomagnetic field fluctuations. Experientia 1996; 52: 253-258.
- 39. Randall W. The solar wind and human birth rate: a possible relationship due to magnetic disturbances. Int J Biometeorol 1990; 34: 42-48.
- 40. Burch JB, Reif JB, Yost MG, Keefe TJ, Pitrat CA. Reduced excretion of a melatonin metabolite in workers exposed to 60 Hz magnetic fields. Am J Epidemiol 1999; 150: 27-36.
- 41. Burch JB, Reif JB, Yost MG. Geomagnetic disturbances are associated with reduced nocturnal secretion of a melatonin metabolite in humans. Neurosci Lett 1999; 266: 209-212.
- 42. Weydahl A, Sothern RB, Cornelissen G, Wetterberg L. Geomagnetic activity influences the melatonin secretion at latitude 70?N. Biomedicine and Pharmacotherapy 2001; 55: 57-62.
- 43. Smits LJ, Van Poppel FW, Verduin JA, Jongbloet PH, Straatman H, Zielhuis GA. Is fecundability associated with month of birth? An analysis of 19th and early 20th century family reconstitution data from The Netherlands. Human Reproduction 1997; 12: 2572-2578.
- 44. Anisimov VN, Zavarzina NI, Zabezhinskii MA, et al. Vliianie melatonina na pokazateli biologicheskogo vozrasta, prodolzhitel'nost' zhizni i razvitie spontannykh opukholei u myshei [The effect of melatonin on the indices of biological age, on longevity and on the development of spontaneous tumors in mice]. [Russian] Voprosy Onkologii 2000; 46(3):311-319.
- 45. Bulian D, Pierpaoli W. The pineal gland and cancer. I. Pinealectomy corrects congenital hormonal dysfunctions and prolongs life of cancer-prone C3H/He mice. Journal of Neuroimmunology 2000; 108(1-2): 131-135.
- 46. Natelson BH, Ottenweller JE, Tapp WN, Heung S, Beldowicz D. The pineal affects life span in hamsters with heart disease. Physiology & Behavior 1997; 62 (5): 1059-1064.
- 47. Bellossi A, Rocher C, Ruelloux M. Effect of a pulsed magnetic field and of first cold-pressure sunflower oil on mice. Zeitschrift fur Naturforschung. Section C. Journal of Biosciences 2000; 55 (3-4): 267-270.
- 48. Halberg E, Halberg J, Halberg Francine, Sothern RB, Levine H, Halberg F. Familial and individualized longitudinal autorhythmometry for 5 to 12 years and human age effects. J Gerontol 1981; 36: 31-33.
- 49. Cornelissen G, Halberg F, Gheonjian L, Paatashvili T, Faraone P, Watanabe Y, Otsuka K, Sothern RB, Breus T, Baevsky R, Engebretson M, Schroder W. Schwabe's ~10.5- and Hale's ~21-year cycles in human pathology and physiology. In: Schroder W, editor. Long- and Short-Term Variability in Sun's History and Global Change. Bremen: Science Edition, 2000: 79-88.
- 50. Starbuck S, Cornelissen G, Halberg F. Is motivation influenced by geomagnetic activity? Scripta medica (Brno), in press.
- Cornelissen G, Halberg F. Introduction to Chronobiology. Medtronic Chronobiology Seminar #7, April 1994, 52 pp. (Library of Congress Catalog Card #94-060580; URL [accessed 1994] http://revilla.mac.cie.uva.es/chrono or http://www.msi.umn.edu/~halberg/)
- 52. Cornelissen G, Halberg F, Wendt HW, Bingham C, Sothern RB, Haus E, Kleitman E, Kleitman N, Revilla MA, Revilla M Jr, Breus TK, Pimenov K, Grigoriev AE, Mitish MD, Yatsyk GV, Syutkina EV. Resonance of about-weekly human heart rate rhythm with solar activity change. Biologia (Bratislava) 1996; 51: 749-756.