



Endocytic control of signaling at the plasma membrane

Elisa Barbieri¹, Pier Paolo Di Fiore^{1,2,3} and Sara Sigismund¹

Signaling is regulated by endocytosis at multiple levels along endocytic routes. Endocytic control of signaling starts already at the plasma membrane, where cells employ different mechanisms to finely tune the type and strength of signals emanating from the cell surface. Here, we will review some of the most recently described endocytic mechanisms controlling signaling at the plasma membrane, through the regulation of internalization dynamics and through the integration of different internalization pathways triggered by canonical chemical stimuli or physical forces.

Addresses

¹ IFOM, Fondazione Istituto FIRC di Oncologia Molecolare, Via Adamello 16, 20139 Milan, Italy

² Dipartimento di Scienze della Salute, Università degli Studi di Milano, Via di Rudini 8, 20122 Milan, Italy

³ Istituto Europeo di Oncologia, Via Ripamonti 435, 20141 Milan, Italy

Corresponding authors: Di Fiore, Pier Paolo (pierpaolo.difiore@ifom.eu) and Sigismund, Sara (sara.sigismund@ifom.eu)

Current Opinion in Cell Biology 2016, 39:21–27

This review comes from a themed issue on **Cell regulation**

Edited by **Manuela Baccarini** and **Ivan Dikic**

<http://dx.doi.org/10.1016/j.ceb.2016.01.012>

0955-0674/© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Signal location is a critical parameter that cells use to decode complex signaling circuitries and to compute specific biological responses. While signaling initiates at the plasma membrane (PM), it requires membrane dynamics for its sustainment/extinction and, more importantly, for its deconvolution. Endocytosis provides this dimension through numerous mechanisms (for a review see [1]) enacted in different subcellular compartments. For instance, it provides spatial constraints to biomembrane-associated signaling molecules (e.g., PM versus endosomes) and dictates differential access to signaling effectors. Moreover, endocytosis regulates the internalization and fate (i.e., recycling versus degradation) of signaling molecules through distinct endocytic pathways and/or via endosomal sorting. Finally, endocytosis is critical in the control of membrane turnover and plasticity in fundamental cellular programs, such as mitosis, adhesion and migration, as well as in the relocalization of signaling/adhesion molecules to PM ‘competent’ regions

[1]. Through these integrated functions, endocytosis determines signal strength and diversification of biological outputs.

In this section, we will highlight recent evidence demonstrating how endocytosis controls signaling at the PM level (for signaling control at the endosomal level see Chapter 7 of this issue), through the regulation of internalization dynamics and the integration of different internalization pathways triggered by canonical chemical stimuli or physical forces.


Clathrin-coated pit dynamics regulate signaling from the PM

Regulation of receptor levels at the PM, receptor availability, and ligand accessibility are established mechanisms affecting the timing and strength of signaling responses [1]. Another critical parameter in signal regulation is clathrin-coated pit (CCP) dynamics. Recent advances in live imaging and data analysis, which allow the large-scale simultaneous detection of the endocytic machinery and cargoes, have helped to establish that CCP dynamics is directly controlled by the cargo [2^{••},3]. While cargoes have been long regarded as ‘passengers’ in internalization structures, it is now clear that — at least in some instances — they directly influence the formation and maturation of CCPs [4,5] by locally communicating with the endocytic machinery [6[•]]. Failure to recruit cargo generates short-lived, abortive CCPs [7–9], revealing the existence of an early checkpoint required to monitor the fidelity of CCP formation, which depends on cargo, dynamin and AP2 [10[•]]. For signaling receptors, this might prolong the time available for clustering at the PM and for the initiation of productive signaling.

Interestingly, different ligands can induce clustering of the same cargo receptor in dynamically distinct CCPs. This is the case of the cannabinoid receptor 1 (CB1R) that, depending on the type of agonist, is clustered into CCPs with different dwell times (i.e., the time required for the receptor to be clustered in CCPs together with the adaptor β -arrestin), which in turn affects the signaling output (Figure 1a). Indeed, pits with long dwell times elicit productive and robust β -arrestin-dependent ERK1/2 activation (Figure 1a, right), while pits with short dwell times generate scarce β -arrestin signaling (Figure 1a, left) [11^{••}].

Similar scenarios have been uncovered for receptor tyrosine kinases (RTKs), such as the RET (REarranged during Transfection) isoforms, which are internalized in CCPs displaying different kinetics [12]. Interestingly, the different RET isoforms assemble specific signaling

Fig. 1



Endocytosis controls PM signaling through different mechanisms. **(a)** CCP dynamics control signaling. Left, the agonist WIN 55,212-2 induces recruitment and internalization of the cannabinoid receptor (CB1R) through CCPs with fast dwell times, which results in trimeric G-protein (α , β , γ) signaling at the PM, but fails to generate productive β -arrestin-dependent signaling. Right, the agonist 2-AG recruits CB1R into CCPs with slow dwell times allowing for both G-protein and productive β -arrestin-dependent signaling, which leads to phosphorylation of Erk1/2 and cell proliferation. **(b)** Left, different agonists differentially regulate signaling by activating distinct internalization pathways. WNT3a binding to its receptor Frizzled (Fz) induces the formation of signalosomes at the PM that include LRP6, Dishevelled (Dvl) and Rab8, and which mediate internalization via caveolae. This allows phosphorylation of LRP6 by CK1 γ kinase and β -catenin accumulation, leading to cell proliferation. In contrast, the Wnt pathway antagonist, DKK1, recruits LRP6 to clathrin-mediated endocytosis (CME), which ultimately leads to β -catenin degradation and signal extinction. Right, signal strength (extracellular ligand concentration) controls EGFR signaling by activating distinct endocytic pathways: CME and non-clathrin endocytosis (NCE). At low doses of EGF, ~90% of the EGFR is internalized via CME, which primarily leads to recycling and sustainment of signaling, resulting in cell proliferation. At high doses of EGF, the EGFR becomes significantly ubiquitinated, concomitantly with the activation of NCE (note, however, that CME persists). NCE targets EGFRs mainly to degradation in the lysosome causing long-term attenuation of signaling. **(c)** Physical and mechanical stimuli, such as temperature (left) or PM tension (right), control endocytic routes and signaling. Left, in *Drosophila* at low temperatures, Notch is constitutively internalized mainly via Deltex-mediated endocytosis, which does not require cholesterol and allows signaling activation. At higher temperatures, Sudex-mediated endocytosis is enhanced, which is dependent on cholesterol and leads to Notch degradation. This pathway acts to compensate for the increased ligand-dependent signaling occurring at high temperatures. This dual mechanism ensures thermal balance of Notch signaling during development. Right, under low PM tension (top), assembly of the clathrin coat is sufficient to deform the PM and CME does not require the actin cytoskeleton. Bottom, under high PM tension (i.e., on the apical side of polarized cells, or in cells subjected to mechanical stretching), clathrin assembly is unable to counteract the tension force, and invaginations stall; actin polymerization then provides the energy needed to complete membrane bending. Hip1R is the link between the assembling clathrin coat and actin polymers. CCPs in this latter case have an extended lifetime, which is necessary to allow Hip1R to be recruited and actin polymerization to occur.

complexes and are connected to distinct developmental roles, raising the possibility that internalization kinetics might influence RTK biological outputs.

Endocytic routes and receptor fate

Several signaling receptors can be internalized through multiple endocytic routes, the activation of which depends on the cellular context and environmental conditions. Notably, these different endocytic routes are coupled to distinct receptor fates [1,13].

The Wnt signaling pathway is critical for development throughout evolution and is controlled by distinct endocytic pathways [14]. When Wnt binds to its receptor, Frizzled (Fz), it induces clustering of Fz and of the co-receptor LRP6 (low density lipoprotein receptor-related protein 6), followed by recruitment of Dishevelled (Dvl) and formation of signalosomes, which include scaffolding proteins (Axin1) and kinases (GSK3 β and CK1 γ kinases) [15]. In mammalian cells, different entry routes are used by the LRP6-Fz complex and are associated with distinct outputs (Figure 1b, left); specifically, clathrin-mediated endocytosis (CME) and caveolar endocytosis, which lead to receptor degradation and signaling, respectively [16].

The molecular mechanisms underlying the different endocytic routes are starting to be dissected. Upon WNT3a stimulation and Fz-LRP6 activation, Rab8 is recruited by Dvl to the PM, where it interacts with LRP6 and engages its guanine exchange factor (GEF) RABIN8, promoting caveolar endocytosis of the LRP6-signalosome complex (Figure 1b, left) [15,17]. Internalization of the complex promotes sequestration of GSK3 β into multivesicular bodies (MVBs) and inactivation of the β -catenin destruction complex, inducing β -catenin stabilization and signaling via CK1 γ [16,18^{*},19]. LRP6 can also bind the WNT3a antagonist, Dickkopf (DKK), which removes LRP6 from lipid rafts (where CK1 γ is localized) and diverts its endocytosis from the caveolar to the CME pathway, ultimately resulting in enhanced β -catenin degradation and signal extinction (Figure 1b, left) [16,19]. There are additional layers of complexity, though, as suggested by findings that, depending on the level of the adaptor protein Dab2 in the cell, WNT3a can also direct the Fz-LRP6 complex into the CME pathway, thereby leading to signal attenuation [20^{*}].

The above findings highlight the relevance of the cellular context in the integration of endocytosis and signaling, a notion further supported by studies in lower organisms. Here, the differential impact of endocytic pathways on Wnt signaling is less clearly defined [21,22]; however, differently from mammalian cells, CME appears to exert a positive role in the regulation of both the non-canonical (β -catenin-independent) [23–25] and canonical (β -catenin-dependent) Wnt signaling pathways [26].

An additional level of control, conserved in evolution, is exerted by the ZNFR3/RNF43 transmembrane Ub ligases [27]. These enzymes promote the continuous ubiquitination, constitutive endocytosis and lysosomal degradation of Fz, thus regulating receptor availability at the PM [28,29]. ZNFR3 and RNF43 are also direct transcriptional targets of β -catenin in the WNT3a signaling pathway, thus constituting a negative-feedback loop [30,31^{*}]. This loop can, in turn, be inhibited by secreted R-Spondins [32] that antagonize ZNFR3/RNF43, stabilizing Fz receptors and increasing Wnt signaling strength [30,31^{*}]. In summary, the emerging picture is that Wnt signaling is finely tuned by endocytosis, through mechanisms regulated by the cellular context and environmental cues.

In the case of the transforming growth factor receptor β (TGF β R) and epidermal growth factor receptor (EGFR), CME and various forms of non-clathrin endocytosis (NCE) are associated with distinct receptor fates, although with opposite outcomes as compared with LRP6-Fz in mammalian cells. For these receptors, CME is predominantly associated with receptor recycling and sustainment of signaling, while NCE mainly directs receptors to degradation and signal extinction [33,34]. For the EGFR, NCE occurs through different mechanisms [35–37]. In all cases, however, NCE is activated only when the ligand is present at high, nearly saturating, doses (>10 ng/ml) (Figure 1b, right). One mechanism involves the conversion of a linear gradient of EGF into an almost all-or-nothing EGFR ubiquitination (Ub) response, which in turn leads to the sharp activation of NCE and to receptor degradation [38]. Thus, NCE likely protects cells from overstimulation under conditions of excess ligand. A combination of mathematical modeling and wet-lab experiments revealed that EGFR ubiquitination — and its recruitment into NCE — is critically controlled by EGFR levels [39^{*}]. Indeed, in cells displaying high EGFR levels — a hallmark of some cancer cells — the receptor is inefficiently ubiquitinated while being highly phosphorylated/activated [39^{*}]. The prediction here is that, under these conditions, EGFR would escape internalization through NCE and the ensuing signal extinction, thereby providing a proliferative advantage to cancer cells.

Of note, a second level of analogical-to-digital control of EGFR signaling intensity occurs in the endosomal compartment, where increasing EGF concentrations induce a proportional increment in the number of endosomes, so that the number of active EGFRs/endosome remains constant. A linear EGF gradient is thus converted into ‘quanta’ of signaling receptor. In addition to the EGFR, other receptors, such as hepatocyte growth factor receptor (HGFR) or nerve growth factor receptor (NGFR), can induce ‘quanta’ of different magnitudes, which correlate with distinct biochemical and biological outputs in

specific cellular context ([40**], see also Chapter 7 of this issue).

Similarly, endocytosis-controlled modalities of analogical-to-digital conversion of signals have been described in the establishment/decoding of morphogen gradients during development, as well as in the control of collective cellular motility in physiology and in cancer [41]. For instance, it has recently been shown how malignant lymphocytes, traditionally regarded as individual movers, have an intrinsic tendency to gather into clusters that display unique migratory and chemotactic properties [42]. These collective entities, at variance with single cells, display chemotactic prowess in shallow chemokine gradients and are resistant to the chemorepulsion that normally results from increases in gradient steepness. Not surprisingly, endocytic dynamics are at the core of these processes, although the precise molecular mechanisms remain to be elucidated [42].

Physical stimuli and regulation of endocytic pathways

Physical stimuli, such as temperature or mechanotension, are emerging as important and specific activators/regulators of endocytic routes. Notch signaling during *Drosophila* development is maintained across a wide range of temperatures through a compensation mechanism relying on distinct internalization pathways [43**]. Ligand-independent endocytosis of Notch occurs through two routes with distinct lipid and temperature requirements (Figure 1c, left). At low temperatures, Notch is mainly internalized via Deltex-mediated endocytosis, which occurs through glycosylphosphatidylinositol (GPI)-negative endosomes in a cholesterol-independent manner, and leads to Notch signaling activation. At higher temperatures, Sudex-mediated endocytosis is activated, characterized by GPI-positive endosomes and cholesterol sensitivity. This pathway dampens signaling by targeting receptors for lysosomal degradation, thereby counterbalancing the increased ligand-receptor binding kinetics occurring at high temperature [43**]. The net result is thermal robustness of Notch signaling. Moreover, temperature might control the cholesterol-enriched pathway by directly influencing membrane fluidity and tension (see also below). These regulatory mechanisms are clearly relevant in ectothermic organisms; it will be interesting to investigate whether they apply also to endothermic organisms, to specific organs/tissues or under inflammatory conditions.

Mammalian cells are constantly subjected to environmental mechanical forces that regulate PM tension [44]. A tight bidirectional regulation exists between membrane tension, the extracellular matrix (ECM) and endocytosis (see Section 3 of this issue), which can be harnessed by cancer cells to establish invasive programs [45]. Through

membrane recycling and turnover, endocytosis can influence the physical properties of the PM. In addition, through trafficking of adhesion molecules, endocytosis regulates cellular communication with the ECM [46]. In turn, endocytosis is regulated by external forces through the activation of specific internalization pathways [44], as exemplified by the endocytic dynamics of integrins. Integrins display different endocytic responses to mechanical forces, which influence their signaling [47]. Endocytosis of integrin-beta3 is controlled by ECM-originated cues: RGD (Arg-Gly-Asp) ligands immobilized on supported lipid membranes cannot generate traction on the engaged integrin nor promote its clustering and removal from the PM via CME. Conversely, the increase in force, obtained by using rigid RGD ligands immobilized on glass, induces the reinforcement of the actomyosin network and the recruitment of focal adhesion adaptors, such as talin, resulting in CME inhibition and focal adhesion formation [48*]. Thus, mechanical forces control the balance of adhesion signaling versus integrin turnover.

The cortical actin cytoskeleton functions both as a sensor and as a transducer of membrane tension. During phagocytosis, a bidirectional crosstalk between membrane tension and the actin cytoskeleton allows for pseudopodia extension and particle engulfment. Upon particle engagement, Rac1 is activated and the actin cytoskeleton pushes the PM forward [49*]. As a consequence of PM stretching, membrane tension increases, leading to inactivation of Rac1 and to its redistribution from the pseudopodia to the cell center, thereby promoting actin reorganization in the pseudopodia and induction of actin-mediated exocytic events at the site of engagement. The newly exocytic-delivered membranes relieve membrane tension and allow wrapping of the particle [49*]. Thus, continuous communication exists between the PM, deformed by particle engagement, and the actin machinery required for phagocytosis to progress.

Also in CME, the action and recruitment of the actin cytoskeleton is regulated by membrane tension (Figure 1c, right) [50**]. At variance with yeast, in mammalian cells CME proceeds also in absence of actin polymerization [51]. However, the scenario changes under conditions of high membrane tension in which actin recruitment via the clathrin-light-chain adaptor Hip1R becomes essential. Mechanistically, membrane tension opposes clathrin polymerization and hinders the closure of CCPs by varying the membrane budding energy [52*]. This provides enough time for assembly of actin filaments via Hip1R, rescuing the stalled coat (Figure 1c, right) [50**]. Since actin-dependent and -independent CCPs have distinct lifetimes [50**], it is tempting to speculate that they might influence the retention time of signaling molecules and receptors, thus regulating signaling outputs.

Outlook: mechanotransduction, cell context and endocytosis at the system level

Membrane tension is a critical stimulus to which cells packed in a tissue, or a culture dish, are subjected [53]. Recent studies showed that endocytic pathways are directly controlled by local cell density in culture [54**]. This is achieved through a PM-based mechanism, centered on focal adhesion kinase (FAK), which senses local crowding and responds by controlling membrane lipid composition, via a feedback loop that does not require the exchange of a 'chemical signal' between cells [55]. These data show that local crowding (and, most likely, mechanical forces) contributes to the generation of single-cell heterogeneity of endocytic pathways [55].

Single-cell heterogeneity has been previously described for different cellular processes (e.g., gene transcription [56]) and cell signaling pathways influenced by endocytosis, including calcium signaling [57], and NF- κ B [58] and Erk signal transduction pathways [59]. Interestingly, when single-cell variability is taken into account in the analysis of endocytosis data from perturbation screenings, it improves the statistical significance of the results [60,61,62*,63].

Overall, these studies highlight the importance of monitoring endocytic and signaling events at the system level, coupling single cell measurements to quantitative computational analysis of large datasets and mathematical modeling, to untangle the impact of the endocytic machinery on cell regulation.

Acknowledgements

We thank Rosalind Gunby for reviewing the manuscript. Work in the authors' lab is supported by grants from: the Associazione Italiana per la Ricerca sul Cancro (IG 10349 and 14404 and MCO 10.000), MIUR (the Italian Ministry of University and Scientific Research), the Italian Ministry of Health, the European Research Council (Mammastem Project), the Monzino Foundation and the European Community (Network of Excellence FP6, 100601–201012).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Sigismund S, Confalonieri S, Ciliberto A, Polo S, Scita G, Di Fiore PP: **Endocytosis and signaling: cell logistics shape the eukaryotic cell plan.** *Physiol Rev* 2012, **92**:273-366.
 2. Li D, Shao L, Chen BC, Zhang X, Zhang M, Moses B, Milkie DE, Beach JR, Hammer JA 3rd, Pasham M *et al.*: **ADVANCED IMAGING. Extended-resolution structured illumination imaging of endocytic and cytoskeletal dynamics.** *Science* 2015, **349**:aab3500.
- This paper describes the most recent advances in live-imaging structured illumination microscopy (SIM).
3. Cocucci E, Aguet F, Boulant S, Kirchhausen T: **The first five seconds in the life of a clathrin-coated pit.** *Cell* 2012, **150**:495-507.
 4. Mettlen M, Stoeber M, Loerke D, Antonescu CN, Danuser G, Schmid SL: **Endocytic accessory proteins are functionally distinguished by their differential effects on the maturation of clathrin-coated pits.** *Mol Biol Cell* 2009, **20**:3251-3260.
 5. Puthenveedu MA, von Zastrow M: **Cargo regulates clathrin-coated pit dynamics.** *Cell* 2006, **127**:113-124.
 6. Henry AG, Hislop JN, Grove J, Thorn K, Marsh M, von Zastrow M: **Regulation of endocytic clathrin dynamics by cargo ubiquitination.** *Dev Cell* 2012, **23**:519-532.
- Evidence is provided showing that ubiquitination of the cargo (in this case a GPCR) directly controls CCP maturation and release.
7. Ehrlich M, Boll W, Van Oijen A, Hariharan R, Chandran K, Nibert ML, Kirchhausen T: **Endocytosis by random initiation and stabilization of clathrin-coated pits.** *Cell* 2004, **118**:591-605.
 8. Loerke D, Mettlen M, Yarar D, Jaqaman K, Jaqaman H, Danuser G, Schmid SL: **Cargo and dynamin regulate clathrin-coated pit maturation.** *PLoS Biol* 2009, **7**:e57.
 9. Taylor MJ, Lampe M, Merrifield CJ: **A feedback loop between dynamin and actin recruitment during clathrin-mediated endocytosis.** *PLoS Biol* 2012, **10**:e1001302.
 10. Aguet F, Antonescu CN, Mettlen M, Schmid SL, Danuser G: **Advances in analysis of low signal-to-noise images link dynamin and AP2 to the functions of an endocytic checkpoint.** *Dev Cell* 2013, **26**:279-291.
- This paper shows the existence of a molecular endocytic checkpoint for the correct maturation of CCPs, which requires dynamin and the AP2 appendage domain.
11. Flores-Otero J, Ahn KH, Delgado-Peraza F, Mackie K, Kendall DA, Yudowski GA: **Ligand-specific endocytic dwell times control functional selectivity of the cannabinoid receptor 1.** *Nat Commun* 2014, **5**:4589.
- This paper provides evidence demonstrating that different agonists induce CCPs with distinct kinetics differentially regulating β -arrestin-dependent signaling.
12. Crupi MJ, Yoganathan P, Bone LN, Lian E, Fetz A, Antonescu CN, Mulligan LM: **Distinct temporal regulation of RET isoform internalization: roles of clathrin and AP2.** *Traffic* 2015, **16**:1155-1173.
 13. Johannes L, Parton RG, Bassereau P, Mayor S: **Building endocytic pits without clathrin.** *Nat Rev Mol Cell Biol* 2015, **16**:311-321.
 14. Clevers H, Nusse R: **Wnt/beta-catenin signaling and disease.** *Cell* 2012, **149**:1192-1205.
 15. Bilic J, Huang YL, Davidson G, Zimmermann T, Cruciat CM, Bienz M, Niehrs C: **Wnt induces LRP6 signalosomes and promotes dishevelled-dependent LRP6 phosphorylation.** *Science* 2007, **316**:1619-1622.
 16. Yamamoto H, Komekado H, Kikuchi A: **Caveolin is necessary for Wnt-3a-dependent internalization of LRP6 and accumulation of beta-catenin.** *Dev Cell* 2006, **11**:213-223.
 17. Demir K, Kirsch N, Beretta CA, Erdmann G, Ingelfinger D, Moro E, Argenton F, Carl M, Niehrs C, Boutros M: **RAB8B is required for activity and caveolar endocytosis of LRP6.** *Cell Rep* 2013, **4**:1224-1234.
 18. Taelman VF, Dobrowolski R, Plouhinec JL, Fuentealba LC, Vorwald PP, Gumper I, Sabatini DD, De Robertis EM: **Wnt signaling requires sequestration of glycogen synthase kinase 3 inside multivesicular endosomes.** *Cell* 2010, **143**:1136-1148.
- This paper dissects the mechanism through which GSK3 inhibition by Wnt signaling is achieved, which involves the sequestration of GSK3 from the cytosol into multivesicular bodies, so that it is inaccessible to its substrates.
19. Yamamoto H, Sakane H, Michiue T, Kikuchi A: **Wnt3a and Dkk1 regulate distinct internalization pathways of LRP6 to tune the activation of beta-catenin signaling.** *Dev Cell* 2008, **15**:37-48.
 20. Jiang Y, He X, Howe PH: **Disabled-2 (Dab2) inhibits Wnt/beta-catenin signalling by binding LRP6 and promoting its internalization through clathrin.** *EMBO J* 2012, **31**:2336-2349.
- This paper describes how the increase in Dab2 cellular levels diverts LRP6 from caveolae to clathrin-mediated endocytosis upon WNT3a stimulation, thereby inhibiting Wnt/beta-catenin signaling.
21. Niehrs C: **The complex world of WNT receptor signalling.** *Nat Rev Mol Cell Biol* 2012, **13**:767-779.

22. Feng Q, Gao N: **Keeping Wnt signalosome in check by vesicular traffic.** *J Cell Physiol* 2015, **230**:1170-1180.
23. Chen W, ten Berge D, Brown J, Ahn S, Hu LA, Miller WE, Caron MG, Barak LS, Nusse R, Lefkowitz RJ: **Dishevelled 2 recruits beta-arrestin 2 to mediate Wnt5A-stimulated endocytosis of Frizzled 4.** *Science* 2003, **301**:1391-1394.
24. Yu A, Rual JF, Tamai K, Harada Y, Vidal M, He X, Kirchhausen T: **Association of Dishevelled with the clathrin AP-2 adaptor is required for Frizzled endocytosis and planar cell polarity signaling.** *Dev Cell* 2007, **12**:129-141.
25. Sato A, Yamamoto H, Sakane H, Koyama H, Kikuchi A: **Wnt5a regulates distinct signalling pathways by binding to Frizzled2.** *EMBO J* 2010, **29**:41-54.
26. Hagemann AI, Kurz J, Kauffeld S, Chen Q, Reeves PM, Weber S, Schindler S, Davidson G, Kirchhausen T, Scholpp S: **In vivo analysis of formation and endocytosis of the Wnt/beta-catenin signaling complex in zebrafish embryos.** *J Cell Sci* 2014, **127**:3970-3982.
27. Malinauskas T, Jones EY: **Extracellular modulators of Wnt signalling.** *Curr Opin Struct Biol* 2014, **29**:77-84.
28. Jiang X, Charlat O, Zamponi R, Yang Y, Cong F: **Dishevelled promotes Wnt receptor degradation through recruitment of ZNRF3/RNF43 E3 ubiquitin ligases.** *Mol Cell* 2015, **58**:522-533.
29. Moffat LL, Robinson RE, Bakoulis A, Clark SG: **The conserved transmembrane RING finger protein PLR-1 downregulates Wnt signaling by reducing Frizzled, Ror and Ryk cell-surface levels in *C. elegans*.** *Development* 2014, **141**:617-628.
30. Hao HX, Xie Y, Zhang Y, Charlat O, Oster E, Avello M, Lei H, Mickanin C, Liu D, Ruffner H *et al.*: **ZNRF3 promotes Wnt receptor turnover in an R-spondin-sensitive manner.** *Nature* 2012, **485**:195-200.
31. Koo BK, Spit M, Jordens I, Low TY, Stange DE, van de Wetering M, van Es JH, Mohammed S, Heck AJ, Maurice MM *et al.*: **Tumour suppressor RNF43 is a stem-cell E3 ligase that induces endocytosis of Wnt receptors.** *Nature* 2012, **488**:665-669.
- This paper (together with Ref.30 and Ref.28) shows how the availability of Wnt receptors at the cell surface is constantly regulated through ubiquitination, mediated by the transmembrane ligases, RNF43/ZNRF3.
32. Sato T, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M, Barker N, Shroyer NF, van de Wetering M, Clevers H: **Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts.** *Nature* 2011, **469**:415-418.
33. Di Guglielmo GM, Le Roy C, Goodfellow AF, Wrana JL: **Distinct endocytic pathways regulate TGF-beta receptor signalling and turnover.** *Nat Cell Biol* 2003, **5**:410-421.
34. Sigismund S, Argenzio E, Tosoni D, Cavallaro E, Polo S, Di Fiore PP: **Clathrin-mediated internalization is essential for sustained EGFR signaling but dispensable for degradation.** *Dev Cell* 2008, **15**:209-219.
35. Boucrot E, Ferreira AP, Almeida-Souza L, Debard S, Vallis Y, Howard G, Bertot L, Sauvonnnet N, McMahon HT: **Endophilin marks and controls a clathrin-independent endocytic pathway.** *Nature* 2015, **517**:460-465.
36. Orth JD, Krueger EW, Weller SG, McNiven MA: **A novel endocytic mechanism of epidermal growth factor receptor sequestration and internalization.** *Cancer Res* 2006, **66**:3603-3610.
37. Sigismund S, Woelk T, Puri C, Maspero E, Tacchetti C, Transidico P, Di Fiore PP, Polo S: **Clathrin-independent endocytosis of ubiquitinated cargos.** *Proc Natl Acad Sci U S A* 2005, **102**:2760-2765.
38. Sigismund S, Algisi V, Nappo G, Conte A, Pascolutti R, Cuomo A, Bonaldi T, Argenzio E, Verhoef LG, Maspero E *et al.*: **Threshold-controlled ubiquitination of the EGFR directs receptor fate.** *EMBO J* 2013, **32**:2140-2157.
39. Capuani F, Conte A, Argenzio E, Marchetti L, Priami C, Polo S, Di Fiore PP, Sigismund S, Ciliberto A: **Quantitative analysis reveals how EGFR activation and downregulation are coupled in normal but not in cancer cells.** *Nat Commun* 2015, **6**:7999.

Through a combination of modeling and experiments, this paper (together with Ref. [38]) describes how cells are able to convert a linear EGF gradient into a threshold response of EGFR ubiquitination, regulating receptor fate.

40. Villasenor R, Nonaka H, Del Conte-Zerial P, Kalaidzidis Y, Zerial M: **Regulation of EGFR signal transduction by analogue-to-digital conversion in endosomes.** *eLife* 2015:4.
- In this paper, the mechanism regulating the number of endosomes by signaling receptors is dissected through a combination of mathematical modeling and experiments.
41. Gonzalez-Gaitan M, Julicher F: **The role of endocytosis during morphogenetic signaling.** *Cold Spring Harbor Perspect Biol* 2014, **6**:a016881.
42. Malet-Engra G, Yu W, Oldani A, Rey-Barroso J, Gov NS, Scita G, Dupre L: **Collective cell motility promotes chemotactic prowess and resistance to chemorepulsion.** *Curr Biol* 2015, **25**:242-250.
43. Shimizu H, Woodcock SA, Wilkin MB, Trubenova B, Monk NA, Baron M: **Compensatory flux changes within an endocytic trafficking network maintain thermal robustness of Notch signaling.** *Cell* 2014, **157**:1160-1174.
- This paper shows how thermal robustness of Notch signaling in *Drosophila* is achieved through the activation of distinct internalization pathways directly regulated by external temperature.
44. Gauthier NC, Masters TA, Sheetz MP: **Mechanical feedback between membrane tension and dynamics.** *Trends Cell Biol* 2012, **22**:527-535.
45. Frittoli E, Palamidessi A, Marighetti P, Confalonieri S, Bianchi F, Malinverno C, Mazzarol G, Viale G, Martin-Padura I, Garre M *et al.*: **A RAB5/RAB4 recycling circuitry induces a proteolytic invasive program and promotes tumor dissemination.** *J Cell Biol* 2014, **206**:307-328.
46. Ezratty EJ, Bertaux C, Marcantonio EE, Gundersen GG: **Clathrin mediates integrin endocytosis for focal adhesion disassembly in migrating cells.** *J Cell Biol* 2009, **187**:733-747.
47. Schiller HB, Fassler R: **Mechanosensitivity and compositional dynamics of cell-matrix adhesions.** *EMBO Rep* 2013, **14**:509-519.
48. Yu CH, Rafiq NB, Cao F, Zhou Y, Krishnasamy A, Biswas KH, Ravasio A, Chen Z, Wang YH, Kawachi K *et al.*: **Integrin-beta3 clusters recruit clathrin-mediated endocytic machinery in the absence of traction force.** *Nat Commun* 2015, **6**:8672.
- This paper shows how rigidity of the extracellular matrix regulates recruitment of adaptor/adhesion molecules, directly controlling endocytosis of Integrin-beta3.
49. Masters TA, Pontes B, Viasnoff V, Li Y, Gauthier NC: **Plasma membrane tension orchestrates membrane trafficking, cytoskeletal remodeling, and biochemical signaling during phagocytosis.** *Proc Natl Acad Sci U S A* 2013, **110**:11875-11880.
- Evidence is provided demonstrating that, during phagocytosis, a peak in membrane tension inactivates Rac1, allowing for completion of phagocytosis.
50. Boulant S, Kural C, Zeeh JC, Ubelmann F, Kirchhausen T: **Actin dynamics counteract membrane tension during clathrin-mediated endocytosis.** *Nat Cell Biol* 2011, **13**:1124-1131.
- This paper demonstrates how elevated membrane tension (e.g. in the apical surfaces of polarized cells, upon osmotic swelling or mechanical stretching) determines the actin dependence of CCP assembly in mammalian cells.
51. Aghamohammadzadeh S, Ayscough KR: **Differential requirements for actin during yeast and mammalian endocytosis.** *Nat Cell Biol* 2009, **11**:1039-1042.
52. Saleem M, Morlot S, Hohendahl A, Manzi J, Lenz M, Roux A: **A balance between membrane elasticity and polymerization energy sets the shape of spherical clathrin coats.** *Nat Commun* 2015, **6**:249.
- In this paper, clathrin budding was reconstituted *in vitro*, allowing the measurement of changes in membrane tension during clathrin polymerization and the estimation of membrane budding energy.
53. Iskratsch T, Wolfenson H, Sheetz MP: **Appreciating force and shape—the rise of mechanotransduction in cell biology.** *Nat Rev Mol Cell Biol* 2014, **15**:825-833.

54. Snijder B, Sacher R, Ramo P, Damm EM, Liberali P, Pelkmans L:
 ●● **Population context determines cell-to-cell variability in endocytosis and virus infection.** *Nature* 2009, **461**:520-523.
 Quantitative high-throughput analysis is provided dissecting the mechanism by which cells are able to sense local crowding within a population, resulting in adaptation of genes controlling lipid membrane homeostasis.
55. Frechin M, Stoeger T, Daetwyler S, Gehin C, Battich N, Damm EM, Stergiou L, Riezman H, Pelkmans L: **Cell-intrinsic adaptation of lipid composition to local crowding drives social behaviour.** *Nature* 2015, **523**:88-91.
56. Elowitz MB, Levine AJ, Siggia ED, Swain PS: **Stochastic gene expression in a single cell.** *Science* 2002, **297**:1183-1186.
57. Thurley K, Tovey SC, Moenke G, Prince VL, Meena A, Thomas AP, Skupin A, Taylor CW, Falcke M: **Reliable encoding of stimulus intensities within random sequences of intracellular Ca²⁺ spikes.** *Sci Signal* 2014, **7**:ra59.
58. Lee RE, Walker SR, Savery K, Frank DA, Gaudet S: **Fold change of nuclear NF-kappaB determines TNF-induced transcription in single cells.** *Mol Cell* 2014, **53**:867-879.
59. Cohen-Saidon C, Cohen AA, Sigal A, Liron Y, Alon U: **Dynamics and variability of ERK2 response to EGF in individual living cells.** *Mol Cell* 2009, **36**:885-893.
60. Dey G, Gupta GD, Ramalingam B, Sathe M, Mayor S, Thattai M: **Exploiting cell-to-cell variability to detect cellular perturbations.** *PLOS ONE* 2014, **9**:e90540.
61. Gupta GD, Dey G, Swetha MG, Ramalingam B, Shameer K, Thottacherry JJ, Kalappurakkal JM, Howes MT, Chandran R, Das A *et al.*: **Population distribution analyses reveal a hierarchy of molecular players underlying parallel endocytic pathways.** *PLOS ONE* 2014, **9**:e100554.
62. Liberali P, Snijder B, Pelkmans L: **A hierarchical map of regulatory genetic interactions in membrane trafficking.** *Cell* 2014, **157**:1473-1487.
 RNAi screening of endocytic and trafficking genes, combining high-throughput image analysis of different cell perturbations and cell-to-cell variability.
63. Sachs K, Perez O, Pe'er D, Lauffenburger DA, Nolan GP: **Causal protein-signaling networks derived from multiparameter single-cell data.** *Science* 2005, **308**:523-529.