

Draft Genome Sequence of *Rhodotorula mucilaginosa*, an Emergent Opportunistic Pathogen

Massimo Deligios,^a Cristina Fraumene,^b Marcello Abbondio,^b Ilaria Mannazzu,^c Alessandro Tanca,^b Maria Filippa Addis,^b Sergio Uzzau^{a,b}

Dipartimento di Scienze Biomediche, Università di Sassari, Sassari, Italy^a; Porto Conte Ricerche, Parco Scientifico e Tecnologico della Sardegna, Alghero, Italy^b; Dipartimento di Agraria, Università di Sassari, Sassari, Italy^c

***Rhodotorula mucilaginosa*, a yeast with valuable biotechnological features, has also been recorded as an emergent opportunistic pathogen that might cause disease in both immunocompetent and immunocompromised individuals. Here, we report the draft genome sequence of *R. mucilaginosa* strain C2.5t1, which was isolated from cacao seeds in Cameroon.**

Received 18 February 2015 Accepted 3 March 2015 Published 9 April 2015

Citation Deligios M, Fraumene C, Abbondio M, Mannazzu I, Tanca A, Addis MF, Uzzau S. 2015. Draft genome sequence of *Rhodotorula mucilaginosa*, an emergent opportunistic pathogen. *Genome Announc* 3(2):e00201-15. doi:10.1128/genomeA.00201-15.

Copyright © 2015 Deligios et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Sergio Uzzau, uzzau@portocontericerche.it.

Rhodotorula species are ubiquitous saprophytic yeasts that can be recovered from many environmental sources, including humans, animals, and a large variety of foods and beverages (1). Among these species, *Rhodotorula mucilaginosa* has been receiving increasing attention because it can be isolated not only from naturally fermented milk (2) and other food matrices (3, 4), but also from different and extreme ecosystems, including the complex core gut microbiota of carnivore wild fish (5), marine shores, glacial core cold environments (6), and hydrocarbon-contaminated soil (7).

In keeping with its capability to survive and grow in many unfavorable conditions, *R. mucilaginosa* possesses valuable biotechnological features, including copper biotransformation (8), the production of biosurfactants (9), and high yields of unsaturated fatty acids (6) and carotenoids (10). Moreover, yeasts ascribed to this species have a potential use in the biocontrol of postharvest fungal spoilage on fruit (11, 12).

Indeed, *R. mucilaginosa* also represents an emergent opportunistic pathogen for humans and animals, causing skin infections, otitis, lung infections, endocarditis, fungemia, and keratitis in both immunocompetent and immunocompromised individuals (1, 13–18).

Altogether, this information indicates that knowledge of the genomic potential of *R. mucilaginosa* should be extended to the exploration of the cellular physiology and biochemical and biotechnological potentials of this peculiar yeast species to facilitate research for both human and animal health and bioindustry.

R. mucilaginosa strain C2.5t1 was isolated from cacao seeds (*Theobroma cacao* L) in Cameroon. This isolate was previously assigned as *Rhodotorula glutinis* and fully characterized for its capability to produce high yields of biomass and carotenoids when grown in glycerol-containing media (19, 20). A DNA library was prepared using Nextera XT and sequenced using the Illumina HiScanSQ DNA sequencing platform. Sequence data generated about 57 million 2 × 93 bp paired reads that were assembled using Velvet v1/2/10 in 1,034 scaffolds containing 1,254 contigs longer

than 500 bp (G+C content of 60.5%). Strain C2.5t1 assignment was confirmed by 18S rRNA gene identity of 100% to *R. mucilaginosa* strain MAFF237983 (accession no. AB042787) and identity of 99% to *R. glutinis* (accession no. HQ420261). A total of 6,413 protein-coding genes were automatically predicted using the Web server Augustus (<http://bioinf.uni-greifswald.de/webaugustus>) trained using *Ustilago maydis* parameters. Among these genes, 65% were assigned to protein functions using an identity ≥50% to a database containing *Sporidiobolales* (11,439 entries). As expected, among all predicted genes and protein functions, the genome of strain C2.5t1 showed a number of genes that might account for the ability of *R. mucilaginosa* to exert biotechnological features, including copper binding and biotransformation and pigment biosynthesis.

Nucleotide sequence accession numbers. The genome sequence of *R. mucilaginosa* strain C2.5t1 has been deposited at DDBJ/EMBL/GenBank under the accession no. [JWTJ000000000](https://www.ncbi.nlm.nih.gov/nuccore/JWTJ000000000). The version described in this paper is version [JWTJ010000000](https://www.ncbi.nlm.nih.gov/nuccore/JWTJ010000000).

ACKNOWLEDGMENTS

This work was supported by the Porto Conte Ricerche art.26_37/89_2012 Program “Tecnologie Omiche,” funded by Sardegna Ricerche—Science and Technology Park of Sardinia, and partially granted by Regione Autonoma della Sardegna (LR/07 -2010, to I.M.).

REFERENCES

1. Wirth F, Goldani LZ. 2012. Epidemiology of *Rhodotorula*: an emerging pathogen. *Interdiscip Perspect Infect Dis* 2012:465717. <http://dx.doi.org/10.1155/2012/465717>.
2. Bai M, Qing M, Guo Z, Zhang Y, Chen X, Bao Q, Zhang H, Sun TS. 2010. Occurrence and dominance of yeast species in naturally fermented milk from the Tibetan Plateau of China. *Can J Microbiol* 56:707–714. <http://dx.doi.org/10.1139/w10-056>.
3. Sabate J, Cano J, Esteve-Zarzo B, Guillamón JM. 2002. Isolation and identification of yeasts associated with vineyard and winery by RFLP analysis of ribosomal genes and mitochondrial DNA. *Microbiol Res* 157:267–274. <http://dx.doi.org/10.1078/0944-5013-00163>.
4. Lucena-Padrós H, Caballero-Guerrero B, Maldonado-Barragán A,

- Ruiz-Barba JL. 2014. Microbial diversity and dynamics of Spanish-style green table-olive fermentations in large manufacturing companies through culture-dependent techniques. *Food Microbiol* 42:154–165. <http://dx.doi.org/10.1016/j.fm.2014.03.020>.
5. Raggi P, Lopez P, Diaz A, Carrasco D, Silva A, Velez A, Opazo R, Magne F, Navarrete PA. 2014. *Debaryomyces hansenii* and *Rhodotorula mucilaginosa* comprised the yeast core gut microbiota of wild and reared carnivorous salmonids, croaker and yellowtail. *Environ Microbiol* 16: 2791–2803. <http://dx.doi.org/10.1111/1462-2920.12397>.
 6. Singh P, Tsuji M, Singh SM, Roy U, Hoshino T. 2013. Taxonomic characterization, adaptation strategies and biotechnological potential of cryophilic yeasts from ice cores of Midre Lovénbreen glacier, Svalbard, Arctic. *Cryobiology* 66:167–175. <http://dx.doi.org/10.1016/j.cryobiol.2013.01.002>.
 7. Chandran P, Das N. 2012. Role of plasmid in diesel oil degradation by yeast species isolated from petroleum hydrocarbon-contaminated soil. *Environ Technol* 33:645–652. <http://dx.doi.org/10.1080/09593330.2011.587024>.
 8. Salvadori MR, Ando RA, Oller do Nascimento CA, Corrêa B. 2014. Intracellular biosynthesis and removal of copper nanoparticles by dead biomass of yeast isolated from the wastewater of a mine in the Brazilian Amazonia. *PLoS One* 9:e87968. <http://dx.doi.org/10.1371/journal.pone.0087968>.
 9. Kawahara H, Hirai A, Minabe T, Obata H. 2013. Stabilization of astaxanthin by a novel biosurfactant produced by *Rhodotorula mucilaginosa* KUGPP-1. *Biocontrol Sci* 18:21–28. <http://dx.doi.org/10.4265/bio.18.21>.
 10. Aksu Z, Eren AT. 2005. Carotenoids production by the yeast *Rhodotorula mucilaginosa*: use of agricultural wastes as a carbon source. *Proc Biochem* 40:2985–2991. <http://dx.doi.org/10.1016/j.procbio.2005.01.011>.
 11. Li R, Zhang H, Liu W, Zheng X. 2011. Biocontrol of postharvest gray and blue mold decay of apples with *Rhodotorula mucilaginosa* and possible mechanisms of action. *Int J Food Microbiol* 146:151–156. <http://dx.doi.org/10.1016/j.ijfoodmicro.2011.02.015>.
 12. Zhang H, Ge L, Chen K, Zhao L, Zhang X. 2014. Enhanced biocontrol activity of *Rhodotorula mucilaginosa* cultured in media containing chitosan against postharvest diseases in strawberries: possible mechanisms underlying the effect. *J Agric Food Chem* 62:4214–4224. <http://dx.doi.org/10.1021/jf500065n>.
 13. Capoor MR, Aggarwal S, Raghvan C, Gupta DK, Jain AK, Chaudhary R. 2014. Clinical and microbiological characteristics of *Rhodotorula mucilaginosa* infections in a tertiary-care facility. *Indian J Med Microbiol* 32:304–309. <http://dx.doi.org/10.4103/0255-0857.136576>.
 14. Tsiodras S, Papageorgiou S, Meletiadis J, Tofas P, Pappa V, Panayiotides J, Karakitsos P, Armaganidis A, Petrikos G. 2014. *Rhodotorula mucilaginosa* associated meningitis: a subacute entity with high mortality. Case report and review. *Med Mycol Case Rep* 6:46–50. <http://dx.doi.org/10.1016/j.mmcr.2014.08.006>.
 15. Coppola R, Zanframundo S, Rinati MV, Carbotti M, Graziano A, Galati G, De Florio L, Panasiti V. 14 March 2014. *Rhodotorula mucilaginosa* skin infection in a patient treated with sorafenib. *J Eur Acad Dermatol Venerol*. <http://dx.doi.org/10.1111/jdv.12455>.
 16. Villar JM, Delgado JDJ, Velasco CG. 2012. Fungemia due to *Rhodotorula mucilaginosa* in an immunocompetent, critically ill patient. *J Infect Chemother* 18:581–583. <http://dx.doi.org/10.1007/s10156-011-0347-6>.
 17. Kim HA, Hyun M, Ryu S. 2013. Catheter-associated *Rhodotorula mucilaginosa* fungemia in an immunocompetent host. *J Infect Chemother* 45: 339–342. <http://dx.doi.org/10.3947/ic.2013.45.3.339>.
 18. Simon MS, Somersan S, Singh HK, Hartman B, Wickes BL, Jenkins SG, Walsh TJ, Schuetz AN. 2014. Endocarditis caused by *Rhodotorula* infection. *J Clin Microbiol* 52:374–378. <http://dx.doi.org/10.1128/JCM.01950-13>.
 19. Taccari M, Canonico L, Comitini F, Mannazzu I, Ciani M. 2012. Screening of yeasts for growth on crude glycerol and optimization of biomass production. *Bioresour Technol* 110:488–495. <http://dx.doi.org/10.1016/j.biortech.2012.01.109>.
 20. Cutzu R, Coi A, Rosso F, Bardi L, Ciani M, Budroni M, Zara G, Zara S, Mannazzu I. 2013. From crude glycerol to carotenoids by using a *Rhodotorula glutinis* mutant. *World J Microbiol Biotechnol* 29: 1009–1017. <http://dx.doi.org/10.1007/s11274-013-1264-x>.