

Perilla Derived Compounds Mediate Human TRPA1 Channel Activity

Alberto Maria Cattaneo^{1,2*}, Yuriy V Bobkov², Elizabeth A Corey², Gigliola Borgonovo¹ and Angela Bassoli¹

¹DeFENS, Department of Food, Environmental and Nutritional Sciences, Università degli Studi di Milano, Milan (MI), Italy

²Whitney Laboratory for Marine Bioscience, Center for Smell and Taste, and McKnight Brain Institute University of Florida, Gainesville, FL, USA

Abstract

Compounds from food plants affecting the somatosensory system, like *Perilla frutescens* (L.), are well known for their flavoring, pharmacological and medical properties. Yet the exact mechanisms underlying their activity are still poorly understood. Transient Receptor Potential (TRP) channels involved in chemesthetic sensations likely represent some of the primary targets for these compounds. Using a heterologous expression system and calcium imaging we show that a number of *Perilla* derived compounds (S-(-)-1,8-*p*-menthadiene-7-al (perillaldehyde, PA); 3-(4-methyl-1-oxopentyl)furan (perillaketone, PK); 1,2,4-trimethoxy-5-[(E)-prop-1-enyl]benzene (α -asarone, ASA)) and synthetic compounds derivative from *Perilla* (3-(4-methoxy-phenyl)-1-furan-2-yl-propenone (PK-16) and 3-(4-chloro-phenyl)-1-furan-2-yl-propenone (PK-18)) are capable of activating the human TRP Ankyrin family channel (h-TRPA1). The compounds tested appear to be partial agonists of the channel with the potency sequence (EC50, μ M): PK-16(107.7)>PA (160.5)>ASA(210.9)>PK(350). Our findings provide important insight into the functional properties of the compounds derived from *P. frutescens* and reveal new perspectives for the design of tools for pharmaceutical, agricultural and food industry applications.

Keywords: *Perilla frutescens*; Somato-sensory compounds; Transient receptor potential; TRPA1 ion channel; Calcium imaging

Abbreviations: MO: Mustard Oil, 3-Isothiocyanato-1-Propene; PA: Perillaldehyde; S-(-)-1,8-P-Menthadiene-7-Al; PK: Perillaketone; 3-(4-Methyl-1-Oxopentyl) Furan; IK: Isoegomaketone; 3-(4-Methyl-1-Oxo-2-Pentenyl) Furan; ASAA: Asarone; 1,2,4-Trimethoxy-5-[(E)-Prop-1-Enyl]Benzene; PK-16: 3-(4-Methoxy-Phenyl)-1-Furan-2-yl-Propenone; PK-18: 3-(4-Chloro-Phenyl)-1-Furan-2-yl-Propenone.

Introduction

Transient Receptor Potential (TRP) channels are an evolutionarily ancient and diverse superfamily of cation channels [1,2]. Many TRP channels are involved in the transduction of stimuli of different sensory modalities: photo-, thermoreception, chemosensory reception [3] and perhaps, mechanoreception [4,5] and the activity of several TRP channels appear to be directly mediated by osmolarity and/or humidity [6]. Like many other species across different phyla, mammals use TRP channels to detect a variety of physical and chemical stimuli including compounds derived from food plants and spices [3,7-9]. An interesting feature of some TRP channel family members is their ability to detect and integrate qualitatively different stimuli. For example, the temperature sensitive mammalian TRPV1 (sensitive to high temperature) and TRPM8 (sensitive to low temperature) channels were also demonstrated to interact with chemical compounds associated with high or low temperature perception, such as chili pepper derived capsaicinoids [7] and menthol from peppermint [9] respectively.

Mammalian TRPA1 channels are abundantly expressed in the somatosensory system, including in trigeminal neurons. They were initially suggested to be involved in noxious cold-sensing, potentially perceived as a burning pain [8,10-13], but in parallel were identified as receptors for mustard oil (MO) derived from *Sinapis* ssp. (Brassicales: Brassicaceae) (allyl-isothiocyanates) and phytocannabinoids from *Cannabis sativa* L. (Rosales: Cannabaceae) (tetra-hydro-cannabinol) [8]. Like MO and phytocannabinoids, several additional ligands derived from food plants and spices can activate TRPA1 channels. For instance, allylic and diallyl disulfide from garlic [11] cinnamaldehyde from essential oils of cinnamon [14], and several other natural ligands, e.g. Artepillin C, ligustilide, unsaturated dialdehyde terpenes,

stilbenoids and phenol derivatives [15,16] have been reported to cause TRPA1 channel mediated Ca⁺⁺ influx.

The Asian plant *Perilla frutescens* L. (Lamiales: Lamiaceae), known as *kaennip* in Korea and as *shiso* in Japan, is commonly used in traditional Eastern cuisine. Applications of extracts of this plant also have been used in traditional Chinese medicine for the treatment of atopic dermatitis as well as other inflammatory and allergy-related symptoms [17], resulting in a renewed interest in the economic importance and cultivation of different varieties of this species [18]. There are several varieties of *P. frutescens* characterized by different chemical composition of the essential oils extracted from their leaves, they are referred to as chemotypes and are named for the primary component of the oil. Among these, some have been identified as monoterpenes like S-(-)-1,8-*p*-menthadiene-7-al (perillaldehyde, PA), which is the main component in the PA-type varieties used primarily for cooking in Japan and 3-(4-methyl-1-oxopentyl) furan (perillaketone, PK) the main component in the PK-type varieties very diffused in Korea. Despite the large diffusion of *Perilla* in Asian food and the absence of epidemiological evidences seems to indicate that there is not a relevant safety problem for humans, the plant is suspected to arise some safety issues. PK resulted toxic to some animals being associated to the insorgence of pulmonary disease in equine and ovine [19,20]; PA has been long used as a natural aroma compound for food ingredients, but a very recent report by the European Food Safety Authority (EFSA) concluded that PA is genotoxic *in-vivo* and that, accordingly, there is a safety concern for its use as flavouring substance [21].

***Corresponding author:** Alberto Maria Cattaneo, Department of Food, Environmental and Nutritional Sciences, Università degli Studi di Milano, Via Celoria, 2 20133, Milan (MI) - Italy, Tel. +39 392 2939008; E-mail: albertomaria.cattaneo@gmail.com

Received January 20, 2017; **Accepted** February 01, 2017; **Published** February 06, 2017

Citation: Cattaneo AM, Bobkov YV, Corey EA, Borgonovo G, Bassoli A (2017) *Perilla* Derived Compounds Mediate Human TRPA1 Channel Activity. Med Aromat Plants (Los Angel) 6: 283. doi: [10.4172/2167-0412.1000283](https://doi.org/10.4172/2167-0412.1000283)

Copyright: © 2017 Cattaneo AM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

At the molecular level, the properties of *P. frutescens* derived compounds and their mechanisms of action are still poorly understood. PA and PK were reported to interact with rat TRPA1 (r-TRPA1) and TRPM8 channels [22]. Further experiments revealed that synthetic derivatives of PK are more potent than the natural variant [23].

Here we show that the human TRPA1 (h-TRPA1) channel orthologue is sensitive to *Perilla* derived compounds. Among the varieties of *P. frutescens*, we screened essential oils derived from the PA and PK chemotypes, as well as from the PKIK chemotype which contains varying amounts of 3-(4-methyl-1-oxo-2-pentenyl)furan (isogomaketone, IK) in combination with PK [24], for their ability to activate h-TRPA1. In our screen, we also included two PK-derivatives recently reported to activate r-TRPA1: 3-(4-methoxy-phenyl)-1-furan-2-yl-propenone (PK-16) and 3-(4-chloro-phenyl)-1-furan-2-yl-propenone (PK-18) [23]. Additional tests were conducted with the somatosensory phenylpropanoid 1,2,4-trimethoxy-5-[(E)-prop-1-enyl]benzene, identified in the essential oil content of *P. frutescens* var. *acutifolia*, a cultivar mainly used in traditional medicine [25]. The compounds tested appear to be partial agonists of the h-TRPA1 channel with the potency sequence (EC₅₀, μM): PK-16(107.7)>PA (160.5)>ASA (210.9)>PK (350). Our findings provide important insight into the functional properties of the compounds derived from *P. frutescens*.

Material and Methods


Preparations of essential oils and compounds

The *P. frutescens* plants used in the present study were cultivated in an open field. The samples with the highest concentration of perillaldehyde (in PA-type plants), perillaketone (in PK-type plants) and perillaketone with isogomaketone (in PKIK-type plants) were collected from May to October, which covers the entire period of the year during which the plants produce these compounds at the altitude and under the climatic conditions in which the plants were cultivated (Fondazione Minoprio, province of Como, Italy; 380 AMSL). Essential oils from PA- PK- and PKIK-type *P. frutescens* were prepared as previously reported [22,26]. PK derivatives 3-(4-methoxy-phenyl)-1-furan-2-yl-propenone (PK-16) and 3-(4-chloro-phenyl)-1-furan-2-yl-propenone (PK-18) were synthesized according with Bassoli et al. [23]. α-Asarone was isolated as the main component of the essential oil obtained from dry leaves of *P. frutescens* var. *acutifolia* (*soyeop*, Bassoli and Borgonovo, unpublished) bought at the medicinal plants market in Seoul, Korea. The essential oil has an average yield of 0.25 g/100 g dry leaves. The chemical structure of this metabolite was determined by NMR and GC-MS investigations and was consistent with those described in the literature for α-asarone [24] and with an authentic sample. For the *in-vitro* assays a commercial sample of asarone (Sigma Aldrich, St. Louis, MO, USA, >97% GC) was used. Structures of *Perilla* derived compounds are reported in Figure 1.

Heterologous expression and transient transfection

HEK293T cells (ATCC, Manassas, VA, USA) were grown in HEK293T cell media [Dulbecco's modified Eagle's medium (DMEM) enriched with 10% heat-inactivated fetal bovine serum (MP Biomedicals, Solon, OH, USA), 2 mM L-glutamine, and 100 μg/mL penicillin/streptomycin (Invitrogen, Life Technologies, Grand Island, NY, USA) at 37°C with 5% CO₂.

For transient expression, semi-confluent HEK293T cells were grown in 35-mm cell culture-treated dishes and transfected with pcDNA5-FRT carrying the entire protein-coding region for h-TRPA1



described by Doerner et al. [27] and generously provided by Dr. Gunter Gisselmann and Dr. Hanns Hatt. For parallel control of the channel expression in HEK293T cells, 0.67 μg of a separate plasmid [pEBFP2-Nuc a gift from Robert Campbell (Addgene plasmid # 14893, [28])] carrying the coding sequence of a nucleus-targeted blue fluorescent protein (BFP) under the regulation of the same CMV promoter used for expression of h-TRPA1 was co-transfected with calyculin following the manufacturer's recommended protocol (SigmaGen Laboratories, Rockville, MD, USA). At 10 to 18 hours post-transfection, the transfection mix was replaced with fresh HEK293T cell media and the cells were incubated for eight additional hours. Cells were then split in 35-mm cell culture treated dishes and allowed to recover for up to 4 hours prior calcium imaging experiments. Measurements were performed within 44 to 72 hours post-transfection. To estimate transfection efficiency, a parallel transfection was conducted using the positive control vector pcDNA5/TO/LACZ (Invitrogen) and staining with 0.1% 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (XGal, Thermo Scientific, Wilmington, DE, USA). The predominant majority of LACZ transfected cells demonstrated a strong staining suggesting well optimized transfection protocol/transfection efficiency.

Calcium imaging

HEK293T cells co-transfected with the h-TRPA1 and pEBFP2-Nuc plasmids, were incubated for 30m-2h at room temperature in 0.5-1.0 mL HEK cell solution (mM: 140 NaCl, 5 KCl, 1.0 CaCl₂, 1.0 MgCl₂, 10 HEPES, 10 Glucose, pH 7.5) including the fluorescent calcium indicator Fluo-4AM (Invitrogen) at 5.0-15 μM prepared with 0.2-0.06% Pluronic F-127 (Invitrogen). After incubation, the buffer was removed and cells were rinsed with 4.0 mL HEK Ca⁺⁺ solution (mM: 140 NaCl, 2.0 CaCl₂, 10 HEPES, pH 7.5). Cells were placed on the stage of an inverted microscope (Olympus IX-71) equipped with a cooled CCD camera (ORCA R2, Hamamatsu, Hamamatsu City, Japan). A standard FITC or Fura-2 filter set (excitation at 510 nm, emission at 530 nm and excitation at 340 nm or 380 nm, emission at 510 nm) were used to estimate Fluo-4 and BFP related fluorescence intensity. Two gravity fed perfusion contours were used. The first contour continuously washed the cells with HEK Ca⁺⁺ solution (~250 μL/min). The second was used for stimulation and/or application of the compounds tested. Switches between the perfusion channels and regulation of pulse duration were controlled by a multi-channel rapid solution changer (RSC-160, Bio-Logic) and Clampex 9 software (Molecular Devices, Sunnyvale, CA,

USA). Stimulus duration was 10 seconds when cells were stimulated with MO and 20 seconds when stimulated with other compounds. More complex stimulation protocols are specified in the Results.

Calcium imaging experiments were carried out under the control of Imaging Workbench 6 software (INDEC BioSystems, Santa Clara, CA, USA). Stored time series image stacks were analyzed off-line using Imaging Workbench 6, Clampfit 10.5, SigmaPlot 11 or exported as TIFF files into ImageJ 1.42 (available from public domain at <http://rsbweb.nih.gov/ij/index.html>). Continuous traces of multiple responses were compensated for slow drift of the baseline fluorescence when necessary. All recordings were performed at room temperature (22°C to 25°C).

Stimulus

Aliquots of stimulus solutions were prepared in different volumes of HEK Ca⁺⁺ solution depending on experimental needs, diluting stock solutions of our samples previously dissolved in dimethyl sulfoxide (DMSO, Sigma Aldrich).

MO (Sigma Aldrich), PK, PA, ASA, PK-16 and PK-18 samples were dissolved in DMSO at millimolar (mM) concentrations and diluted in HEK Ca⁺⁺ solution to prepare stimuli at micromolar (μM) concentrations. PKIK was dissolved in DMSO at milligram/mL (mg/mL) to prepare stimulus in HEK Ca⁺⁺ solution at microgram/mL (μg/mL). For the latter, concentrations of PK and IK compounds were calculated according with their respective percentage content in the essential oil sample. Concentrations of stimulus solutions and their sequences of application varied between different experiments.

According to HPLC based estimations, PKIK-type essential oil from *P. frutescens* contains a mixture of PK as the major component and IK as the secondary metabolite; the relative content of the two compounds changes during growing of the plant (Bassoli and Borgonovo, unpublished). For experiments reported here, we used a mixture of samples collected throughout the entire harvesting season. The amount of PK and IK in the sample determined by HPLC is 88% and 11% respectively. In our experiments, we used *P. frutescens* samples diluted to a final PK dose of ~535 μM (saturating PK concentration). Thus, the PK dose in the PKIK-type essential oil used in this study contained ~67 μM IK.

Generation of dose/response relationships

Dose/response relationships were generated for pure compounds (ASA, PK-16, PK-18) and *P. frutescens* essential oils predominantly containing only one compound, as estimated by HPLC [PK-type (PK > 98%) and PA-type (PA > 98%)]. Different concentration ranges were chosen for different compounds/samples, depending on compound concentration eliciting the minimal detectable response and the stimulus saturating concentration (Table 1). To estimate and if necessary correct the system sensitivity to excitation light intensity and to the mechanical disturbance of the cells caused by perfusion system we performed control tests before each experiment using HEK

Sample	Human TRPA1 (μM)	Rat TRPA1 (μM)
MO	85.25 ± 14.99	2.5 ± 0.7
PK-16	107.66 ± 10.71	20.9 ± 2.27
PA	160.47 ± 9.12	40.7 ± 7.63
ASA	210.92 ± 36.43	Inactive
PK	349.92 ± 53.01	21.9 ± 1.93

Table 1: Comparison of the potencies (EC50) (Potencies were calculated for *P. frutescens* samples comparing human and rat TRPA1, according with our findings and Bassoli et al. in 2013. MO: Mustard oil, as a reference).

Ca⁺⁺ solution without stimulus. MO (200 μM) was used as a positive control and as a reference for normalization. To minimize rundown of the calcium responses especially to saturating stimuli the consecutive stimuli were applied every 45-60 min and multiple 35-mm dishes were used for different doses. To estimate EC50s, the normalized dose/response data were approximated by a modified Hill equation.

Results

To confirm functional expression of h-TRPA1, we began by applying MO, a known agonist. Many of the HEK293T cells transfected with h-TRPA1 generated transient calcium signal upon application of MO. The amplitude of the responses depended on MO concentration, and the overall sensitivity of the cells to the agonist correlated with BFP expression. The parameters of concentration dependence for the MO were estimated to be EC50~ 85.3 ± 15 μM with Hill coefficient $h \sim 2.9 \pm 1.3$ μM, consistent with previous estimates [14,29]. Untransfected cells were not responsive to MO when tested under the same conditions. The responses to MO at close to saturating concentration (200 μM) were used for normalization to allow comparison among the agonists tested in further experiments.

We then screened a panel of *P. frutescens*-derived compounds, including those previously found to be r-TRPA1 channel modulators [22,23]. The main components of *P. frutescens* essential oils (PA, PK, and ASA) and the PK synthetic derivative PK-16 were able to activate h-TRPA1 channel mediated calcium responses. However, all of the compounds tested evoked substantially lower calcium responses in comparison to MO (saturating concentrations of agonists, Figure 2). The dose response relationship of the h-TRPA1 channel response to PA yielded an EC50 of ~160.5 ± 9.1 μM (n = 76-164) with a normalized peak amplitude of 0.33 ± 0.04. Similarly, PK activated a much less robust h-TRPA1 channel dependent calcium response in comparison to MO, with normalized peak amplitude of 0.44 ± 0.03. The concentration-dependence yielded an EC50 of ~349.9 ± 53 μM (n = 78-151). The concentration dependences for ASA and PK-16 yielded comparable EC50s of ~210.9 ± 36.4 μM and ~107.66 ± 10.71 μM and normalized amplitudes of 0.28 ± 0.04 (n = 52-139) and 0.47 ± 0.05 (n = 16-114), respectively. HEK293T cells not expressing h-TRPA1 were not appreciably responsive to any of the compounds tested. Overall, the potency sequence of the compounds was PK-16>PA>ASA>PK, with the synthetic derivative PK-16 and its natural analog PK acting as the most and the least potent ligands, respectively.

Unlike the other agonists tested, synthetic PK-derivative, PK-18, resulted in relatively slow, incoherent and mostly irreversible responses in h-TRPA1 transfected cells (Figure 3). These properties as well as low solubility of PK-18 [1.2E-4 mol/L, Log P=3.48, Scifinder, 2015; Chemical Abstracts Service: Columbus, OH, 2015; CAS Registry Number 111042-59-2 (accessed Nov 12, 2015); calculated using ACD/Labs software, version 11.02; ACD/Labs 1994-2015] made the compound impractical in terms of obtaining experimental data and interpretation.

The relatively low efficacy of the compounds tested in comparison to MO suggests that they are partial agonists of the h-TRPA1 channel. To probe whether they may promote the h-TRPA1 channel desensitisation [23] and/or inhibit channel activity by competing with MO, we used following experimental paradigm: the h-TRPA1 channel-mediated calcium signal was initially activated by exposure of the cells to MO (40 μM, 20 seconds), followed by application of a close to saturating concentration of an agonist (PA or PK) and then by a combined application of MO and an agonist (PA or PK). Each




Figure 2: Effects of *Perilla* derived compounds on human TRPA1 channel activity. The HEK293T cells expressing h-TRPA1 channels generate calcium signals in response to application of the compounds (insets). The effects were concentration dependent and reversible. Graphs show the concentration dependences of the compounds expressed as a function of normalized mean peak fluorescence intensity change (ΔF) vs drug concentration. In all cases, the responses were individually normalized to the saturating concentration of MO (200 μM). Error bars represent standard error of mean. Data were fit with the Hill equation (solid lines). Insets represent average calcium response elicited by a drug application obtained in a single experiment. Grey color depicts standard deviation. Top right panel, summary plot of the concentration dependences.


compound was tested in an individual series of experiments. The amplitudes of the calcium responses were (mean \pm SEM): MO, 40 μM 19.48 ± 8.30 ; PA, 300 μM 13.73 ± 9.10 ; PA, 300 μM + MO, 40 μM 28.33 ± 15.22 ; MO, 40 μM 32.73 ± 14.82 ; PK, 400 μM 32.08 ± 12.12 ; PK, 400 μM + MO, 40 μM 54.82 ± 16.41 (Figure 4). In each of the series, the combined application of the ligands (MO + PA or PK) generated the strongest responses.

Discussion

Here we found that a number of compounds from the Asian food plant *P. frutescens* can activate the h-TRPA1 channel. The potency sequence of h-TRPA1 activation derived for the compounds, PK-16>PA>ASA>PK, differs from that previously derived for r-TRPA1 activation, PK-16>PK>PA>ASA (Table 1). Notably, r-TRPA1 is almost equally responsive to PK-16 and PK, while h-TRPA1 is more sensitive to PK-16 than PK. Furthermore, tests conducted on r-TRPA1 verified that it does not respond to ASA (Dr. Luciano De Petrocellis,

personal communication; Table 1), which is a compound that we have found to activate h-TRPA1 ($\text{EC}_{50} \sim 210.9 \pm 36.4 \mu\text{M}$). Consistent with previous data for r-TRPA1, the synthetic derivative of PK, PK-16 ($\text{EC}_{50} \sim 107.7 \pm 10.7 \mu\text{M}$), is the most potent of the non-MO h-TRPA1 agonists tested. The previous findings [23] also suggested that some *Perilla*-derived compounds promoted r-TRPA1 channel desensitization and/or inhibited the channel activity, possibly by competing with MO for receptor binding. In direct contrast, we found that the strongest of the antagonists reported for r-TRPA1, PA and PK did not reduce the h-TRPA1 calcium signal activated by MO. Instead, the h-TRPA1 channel mediated calcium signal was proportionally augmented in the presence of close to saturating concentrations of these compounds. Further structure-function relationship analysis is necessary to understand what structural features of these compounds and the TRPA1 orthologues account for the different potencies and the different kinetic parameters of the responses.

The significant differences in overall channel properties of r-TRPA1



and h-TRPA1 could be explained by either the different experimental approaches used in the two studies or, most likely, by species specific heterogeneity in their functional properties. The full length (untrimmed) protein sequences of the two orthologues differ in amino acid identity by approximately 20%, with differences throughout including within transmembrane domains 5 and 6 which have been found to determine the specificity of proton sensitivity of TRPA1. More targeted structure - function correlation analyses of these TRPA1 channel orthologues, similar to those of previous studies [30-32], will be required to identify the differences in the critical functional domains/residues that result in the biophysical and pharmacological heterogeneity of the responses to the compounds tested here.

The agonists of TRPA1 channels are usually divided into two main categories based on their mechanism of activation: electrophiles and non-electrophiles. Electrophiles such as allyl-isothiocyanates [8] and α - β unsaturated aldehydes [33,34], activate TRPA1 channels through covalent modifications, condensing α - β -unsaturated bonds with nucleophilic mercapto-groups of cysteine residues of the channels in a Michael addition. Non-electrophiles, such as carvacrol from oregano [35], activate TRPA1 channels without causing covalent modifications. According to the molecular features of the ligands tested here (Figure 1), ASA lacks of α - β -unsaturated bonds, indicate that this compound may be involved in non-covalent interactions with the channel. In

contrast, the electrophilic carbonyl of PA as well as those of ketones PK and its synthetic analogs PK-16 and PK-18 may serve as electrophilic targets in the reversible attack of nucleophilic residues of N-terminal cysteines in a Michael addition.

One of the compounds that we identified as a partial agonist of h-TRPA1 is ASA ($\text{EC}_{50} \sim 210.9 \pm 36.4$ μM), a phenylpropanoid normally present in essential oils from ginger species within *Asarum* (Piperales: Aristolochiaceae), but also found in *P. frutescens* var. *acutifolia* (Bassoli and Borgonuovo, unpublished data). Contrary to our findings, tests conducted on r-TRPA1 found no response to ASA (Table 1), which is consistent with significant differences in the overall channel properties between r-TRPA1 and h-TRPA1. ASA and other phenylpropanoids (allyltetramethoxybenzene, caffeic acid, dillapiole, elemicin, myristicin, nothoapiole, rosmarinic acid and methyl ester) contained in *P. frutescens* extracts, were recently found to inhibit proinflammatory cytokine generation in lipopolysaccharide-treated murine lung cells [36]. Conversely, monoterpenes from *P. frutescens* such as PK have long been known to induce pulmonary toxicity in mammals such as horses and sheep [19,20]. While the molecular targets for these compounds in these processes were not identified, there is evidence that implicates TRP channels, including TRPA1, in the physiological and pathophysiological reactions associated with respiratory depression and neurogenic lung inflammation [37,38].




Figure 4: Effects of PA and PK on the MO activated h-TRPA1 channel. The h-TRPA1 channel mediated calcium signal was initially activated by MO (40 μM , left panels), then by high responsive doses of an agonist (PA: 300 μM , $n=111$, or PK: 400 μM , $n=86$, middle panels) followed by combined application of MO and an agonist (PA or PK, right panels). Amplitudes of the calcium responses were (mean \pm standard error of mean): MO, 40 μM 19.48 ± 8.30 ; PA, 300 μM 13.73 ± 9.10 ; PA, 300 μM +MO, 40 μM 28.33 ± 15.22 ; MO, 40 μM 32.73 ± 14.82 ; PK, 400 μM 32.08 ± 12.12 ; PK, 400 μM +MO, 40 μM 54.82 ± 16.41 . Note, in both experimental sets the combination of ligands (MO+PA or PK) generated the highest responses. Horizontal black bars below traces mark timing of the drug pulse application (20 s).




Figure 5: Effects of PK and PKIK on the h-TRPA1 channel. The h-TRPA1 channel mediated calcium signal was initially activated by PK (535 μM , $n=84$, left panel) and then by a PKIK dose containing the same concentration of PK (535 μM PK+67 μM IK, right panel). The amplitudes of the calcium responses were (mean \pm standard error of mean) PK, 535 μM 27.27 ± 13.30 ; PK, 535 μM +IK, 67 μM 21.36 ± 10.86 . Horizontal black bars below traces mark timing of the drug pulse application (20 s).

Our findings indicate that these compounds can activate h-TRPA1, suggesting a potential molecular mechanism for their role in human inflammatory processes in general, and in pulmonary inflammation in particular.

The major components of the essential oil derived from the PKIK chemotype of *P. frutescens* [24] are PK and IK [3-(4-methyl-1-oxo-2-pentenyl)furan]. IK is an unsaturated form of PK and a *P. frutescens* monoterpen known for antitumor properties [39]. Although its molecular targets have not been identified *in-vivo*, IK was recently reported to activate r-TRPA1 channels ($\text{EC}_{50} = 7.6 \pm 0.2 \mu\text{M}$), even more potently than *P. frutescens* derived natural ligands (PA: 40.7 ± 7.63 ; PK: 21.9 ± 1.93 ; Table 1) [23]. Although it has not been tested with r-TRPA1, the h-TRPA1-based calcium signal activated by the essential oil derived from the PKIK chemotype was lower (~78%) than that activated by a similar concentration of PK alone (Figure 5). While

speculative in the context of the current study given the molecular complexity of essential oils, the lower response to PKIK suggests that PK and IK may be competitive agonists of h-TRPA1 channel.

Conclusion

In conclusion, the chemotypic variations of *P. frutescens* and resulting differing physiological effects underlie the distinct uses of specific varieties of this plant species in food [18] and medicine [17]. Here we report that two natural terpenoids, PA and PK, found in *P. frutescens* essential oils and PK-16, a synthetic derivative of PK, can activate the h-TRPA1 channel. We use dose response experiments to demonstrate that the channel is more sensitive to PK-16 and PA and find that the ability of these compounds to activate TRPA1 differs between the human and rat orthologues. Notably, we also report that h-TRPA1 but not r-TRPA1 responds to a phenylpropanoid ASA

which has a possible inhibitory role of inflammatory responses and is a compound known to be synthesized by *Asarum* plants that was recently identified in the essential oil content of a novel chemotype of *P. frutescens* (var. *acutifolia*). Contrary to previous findings with r-TRPA1, we find no detectable inhibitory activity of PK and PA towards the well-known TRPA1 agonist MO. We rather demonstrate a potential inhibitory interaction between two monoterpenes, PK and IK, emitted by the PKIK chemotype of this plant. The varying levels of TRPA1 activation in response to the compounds derived from the essential oils of different *P. frutescens* chemotypes, likely resulting from differences in their modes of interaction with the channels, highlight the importance of the chemotype mapping of this food and medicinal plant [24]. Further, they draw attention to the potential importance of this plant as a source for naturally derived drugs that may be used to target the TRPA1 channel.

Acknowledgments

We thank Dr. Mark Q. Martindale and Dr. Barry Ache from the Whitney Laboratory for Marine Bioscience (USA) and Dr. Gianfranco Anfora from Fondazione Edmund Mach (Italy) for stimulating discussion and general support. We thank Dr. Luciano De Petrocellis from the Institute of Biomolecular Chemistry (Italy) for sharing part of his results. This research was supported by the administration of the PhD course in Chemistry, Biochemistry and Ecology of Pesticides (CBEA) of Università degli Studi di Milano, DeFENS—Department of Food, Nutritional and Environmental Sciences and by the Italian Ministry of Education, Universities and Research (MIUR) [project "Analisi integrate dei processi molecolari e cellulari responsabili dell'elaborazione di segnali sensoriali in condizioni normali e patologiche" (no. 2010599KBR_005, PRIN 2011)]. We also wanted to acknowledge Fondazione Edmund Mach, Università degli Studi di Cagliari and the government of Regione Autonoma della Sardegna (Italy) for further financial support while performing part of the experiments and during the manuscript preparation phase.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interests.

References

- Clapham DE (2003) TRP channels as cellular sensors. *Nature* 426: 517-524.
- Wu LJ, Sweet TB, Clapham DE (2010) International Union of Basic and Clinical Pharmacology. LXXVI. Current progress in the mammalian TRP ion channel family. *Pharmacol Rev* 62: 381-404.
- Liedtke W, Heller S (2007) TRP ion channel function in sensory transduction and cellular signaling cascades. *Frontiers in Neuroscience*, Taylor & Francis, CRC Press. ISBN-10: 0-8493-4048-9 ISBN-13: 978-0-8493-4048-2, Boca Raton, FL.
- Walker RG, Willingham AT, Zuker CS (2000) A *Drosophila* mechano-sensory transduction channel. *Science* 287: 2229-2234.
- Sun Y, Liu L, Ben-Shahar Y, Jacobs JS, Eberl DF, et al. (2009) TRPA channels distinguish gravity sensing from hearing in Johnston's organ. *Proc Natl Acad Sci USA* 106: 13606-13611.
- Liu L, Li Y, Wang R, Yin C, Dong Q, et al. (2007) *Drosophila* hygro-sensation requires the TRP channels water witch and nanchung. *Nature*, 450: 294-298.
- Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, et al. (1997) The capsaicin receptor: A heat-activated ion channel in the pain pathway. *Nature* 389: 816-824.
- Jordt SE, Bautista DM, Chuang HH, McKemy DD, Zygmunt PM, et al. (2004) Mustard oils and cannabinoids excite sensory nerve fibres through the TRP channel ANKTM1. *Nature* 427: 260-265.
- Bautista DM, Siemens J, Glazer JM, Tsuruda PR, Basbaum AI, et al. (2007) The menthol receptor TRPM8 is the principal detector of environmental cold. *Nature* 448: 204-208.
- Story GM, Peier AM, Reeve AJ, Eid SR, Mosbacher J, et al. (2003) ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures. *Cell* 112: 819-829.
- Bandell M, Story GM, Hwang SW, Viswanath V, Eid SR, et al. (2004) Noxious cold ion channel TRPA1 is activated by pungent compounds and bradykinin. *Neuron* 41: 849-857.
- Moparathi L, Survery S, Kreir M, Simonsen C, Kjellbom P, et al. (2014) Human TRPA1 is intrinsically cold- and chemosensitive with and without its N-terminal ankyrin repeat domain. *Proc Natl Acad Sci USA* 111: 16901-16906.
- Jabba S, Goyal R, Sosa-Pagan JO, Moldenhauer H, Wu J, et al. (2014) Directionality of temperature activation in mouse TRPA1 ion channel can be inverted by single-point mutations in ankyrin repeat six. *Neuron* 82: 1017-1031.
- Hinman A, Chuang HH, Bautista DM, Julius D (2006) TRP channel activation by reversible covalent modification. *Proc Natl Acad Sci USA* 103: 19564-19568.
- Hata T, Tazawa S, Ohta S, Rhyu M, Misaka T (2012). Artepillin C, A major ingredient of Brazilian propolis, induces a pungent taste by activating TRPA1 channels. *PLOS ONE* 7: e48072.
- Nilius B, Flockerzi V (2014) Mammalian transient receptor potential (TRP) cation channels. New York, NY: Handbook of experimental pharmacology, Volume II, Springer Cham Heidelberg.
- Yu HC, Kosuna K, Haga M (1997) *Perilla*, the Genus *Perilla*. Taylor & Francis.
- Ito M (2008) Studies on *Perilla*, agar-wood, and cinnamon through a combination of fieldwork and laboratory work. *J Nat Med* 62: 387-395.
- Breeze RG, Laegreid WW, Bayly WW, Wilson BJ (1984) *Perilla* ketone toxicity: A chemical model for the study of equine restrictive lung disease. *Equine Vet J*, 16: 180-184.
- Abernathy VJ, Roselli RJ, Parker PE, Pou NA (1992) Effects of *Perilla* ketone on the in situ sheep lung. *J Appl Physiol*, 72: 505-514.
- EFSA (2015) Panel on food contact materials, enzymes, flavourings and processing aids. Scientific opinion on flavouring group evaluation 208 Revision 1 (FGE.208Rev1): Consideration of genotoxicity data on representatives for 10 alicyclic aldehydes with the α,β -unsaturation in ring / side-chain and precursors from chemical subgroup 2.2 of FGE.19. *EFSA Journal* 2015, 13: 4173.
- Bassoli A, Borgonovo G, Caimi S, Scaglioni L, Morini G, et al. (2009) Taste-guided identification of high potency TRPA1 agonists from *Perilla frutescens*. *Bioorg Med Chem* 17: 1636-1639.
- Bassoli A, Borgonovo G, Morini G, De Petrocellis L, Schiano Moriello A, et al. (2013) Analogues of *perilla* ketone has highly potent agonists of TRPA1 channel. *Food Chem* 141: 2044-2051.
- Nitta M, Kobayashi H, Ohnishi-Kameyama M, Nagamine T (2006) Essential oil variation of cultivated and wild *Perilla* analyzed by GC/MS. *Biochem Syst Ecol* 34: 25-37.
- Wang Z, Levinson SR, Sun L, Heinbockel T (2014) Identification of both GABAA receptors and voltage-activated Na⁺ channels as molecular targets of anticonvulsant α -asarone. *Front Pharmacol* 5: 40.
- Cattaneo AM, Bengtsson JB, Borgonovo G, Bassoli A, Anfora G (2014) Response of the European grapevine moth *Lobesia botrana* to somatosensory-active volatiles emitted by the non-host plant *Perilla frutescens*. *Physiol Entomol* 39: 229-236.
- Doerner JF, Gisselmann G, Hatt H, Wetzel CH (2007) Transient receptor potential channel A1 is directly gated by calcium ions. *J Biol Chem* 282: 13180-13189.
- Ai HW, Shaner NC, Cheng Z, Tsieng RY, Campbell RE (2007) Exploration of new chromophore structures leads to the identification of improved blue fluorescent proteins. *Biochemistry* 46: 5904-5910.
- Klionsky L, Tamir R, Gao B, Wang W, Immke DC, et al. (2007) Species-specific pharmacology of trichloro(sulfanyl)ethyl benzamides as transient receptor potential ankyrin 1 (TRPA1) antagonists. *Mol Pain* 3:39.
- Samad A, Sura L, Benedikt J, Ettrich R, Minofar B, et al. (2011) The C-terminal basic residues contribute to the chemical- and voltage-dependent activation of TRPA1. *Biochem J* 433: 197-204.
- Nakatsuka K, Gupta R, Saito S, Banzawa N, Takahashi K, et al. (2013) Identification of molecular determinants for a potent mammalian TRPA1 antagonist by utilizing species differences. *J Mol Neurosci* 51: 754-762.
- De La Roche J, Eberhardt MJ, Klinger AB, Stanslowsky N, Wegner F, et al. (2013) The molecular basis for species-specific activation of human TRPA1

- protein by protons involves poorly conserved residues within trans-membrane domains 5 and 6. *J Biol Chem* 288: 20280-20292.
33. Macpherson LJ, Dubin AE, Evans MJ, Marr F, Schultz PG, et al. (2007) Noxious compounds activate TRPA1 ion channels through covalent modification of cysteines. *Nature Letters* 445: 541-545.
34. Trevisani M, Siemens J, Materazzi S, Bautista DM, Nassini R, et al. (2007) 4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1. *Proc Nat Acad Sci USA* 104: 13519-13524.
35. Xu H, Delling M, Jun JC, Clapham DE (2006) Oregano, thyme and clove-derived flavors and skin sensitizers activate specific TRP channels. *Nat Neurosci* 9: 628-635.
36. Lim HJ, Woo KW, Lee KR, Lee SK, Kim HP (2014) Inhibition of pro-inflammatory cytokine generation in lung inflammation by the leaves of *Perilla frutescens* and its constituents. *Biomol Ther (Seoul)* 22: 62-67.
37. Bessac F, Jordt S (2008) Breathtaking TRP channels: TRPA1 and TRPV1 in airway chemosensation and reflex control. *Physiology (Bethesda)* 23: 360-370.
38. Lin AH, Liu MH, Ko HK, Perng DW, Lee TS, et al. (2015) Lung epithelial TRPA1 transduces the extracellular ROS into transcriptional regulation of lung inflammation induced by cigarette smoke: the role of influxed Ca²⁺. *Mediators Inflamm* 148367.
39. Cho BO, Jin CH, Park YD, Ryu HW, Byun MW, et al. (2011) Isoeugenol induces apoptosis through caspase-dependent and caspase-independent pathways in human DLD1 cells. *Biosci Biotechnol Biochem* 75: 1306-1311.

Citation: Cattaneo AM, Bobkov YV, Corey EA, Borgonovo G, Bassoli A (2017) *Perilla* Derived Compounds Mediate Human TRPA1 Channel Activity. *Med Aromat Plants* (Los Angel) 6: 283. doi: [10.4172/2167-0412.1000283](https://doi.org/10.4172/2167-0412.1000283)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>