

The DNA methylation pattern of prepubertal and pubertal Alpine goats

Stefano Frattini¹, Emanuele Capra², Barbara Lazzari^{2,3}, Beatrice Coizet¹, Debora Groppetti¹, Pietro Riccaboni¹, Alessandro Pecile¹, Stefania Chessa², Bianca Castiglioni², Andrea Talenti¹, John L. Williams⁴, Stephanie McKay⁵, Paola Crepaldi¹, Alessandra Stella², Giulio Pagnacco¹

¹ Department of Veterinary Science and Public Health University of Milan, Milan, Italy

² Institute of Agricultural Biology and Biotechnology CNR UOS of Lodi, Lodi, Italy

³ PTP Science Park Lodi Italy

⁴ School of Animal and Veterinary Sciences University of Adelaide, Roseworthy, Australia

⁵ Department of Animal & Veterinary Sciences University of Vermont, Burlington, Vermont



INTRODUCTION

- Puberty timing is controlled by many genes and the elements coordinating this process have not yet been completely identified.
- Hypothalamus is a pivotal organ in the control of sexual maturation¹.
- There is evidence that epigenetic modifications, such as DNA methylation, play a key role in the process.
- GNRHR and Cbx7 are two key genes in the initiation of mammalian puberty^{2,3}.



OBJECTIVES

Highlight epigenetic differences at DNA methylation level that occur when reaching puberty in goat

METHODS

- The methylome of the hypothalamus of 10 Alpine goats, 5 at a prepubertal stage and 5 at their pubertal stage was analysed.




Figure 1 - MethylMiner™ Methylated DNA Enrichment Kit.

- Methylated DNA Binding Domain sequencing (MBD-seq) (Figure 1) and NGS (Hiseq2000 Illumina) were performed.
- An average of 24.13 million of reads (range 18.00 and 30.11 million of reads) were produced per sample.

RESULTS

- The analysis highlighted an increase in methylation in the prepubertal goats. The extent of methylation had a median value (\pm IQR) of 12.32 \pm 10.21 Mbp of the genome for prepubertal goats, compared with 8.18 \pm 9.71 Mbp for pubertal goats.
- Significantly increased methylation was seen on 18 chromosomes in prepubertal goats. Chromosomes 2, 4 and 7 were the most highly significant differentially methylated ones (Figure 2).




Figure 2 - Number of methylation peaks per chromosome corrected for chromosome length in the hypothalamus of prepubertal and pubertal goats.

- A higher methylation level ($P < 0.001$) was observed in a small fragment of the terminal part of Cbx7 gene and of the body of the gene of GNRHR (coding for GNRH receptor) in prepubertal goats (Table 1 and Figure 3).

Gene	Fragment size (bp)	Coverage of methylation \pm DS		P-Value
		Prepubertal	Pubertal	
GNRHR	115	820.5 \pm 72.9	751.0 \pm 63.4	<0.001*
Cbx7	390	62.8 \pm 15.6	47.55 \pm 12.7	<0.001*

Table 1 – Coverage of methylation in a part of GNRHR and Cbx7 genes.




Figure 3 – Methylation profile of Cbx7 in the hypothalamus of prepubertal and pubertal goats (IGV software). In the red box the methylated fragments as in Table 1.

CONCLUSIONS

- These results confirm that female puberty in goats is associated with a modification of the DNA methylation pattern in the hypothalamus

REFERENCES

- 1 - Ojeda, S.R., Skinner, M.K., 2006. Puberty in the rat. In: Neill, J.D. (Ed.), The Physiology of Reproduction, third ed. Academic Press/Elsevier, San Diego, pp. 2061-2126.
- 2 - Lomniczi, A., Loche, A., Castellano, J.M., Ronnekleiv, O.K., Bosh, M., Kaidar, G., Knoll, J.G., Wright, H., Pfeifer, G.P., Ojeda, S.R., 2013. Epigenetic control of female puberty. Nat. Neurosci. 16, 281-289.
- 3 - Lomniczi, A., Wright, H., Ojeda, S.R., 2015. Epigenetic regulation of female puberty. Front. Endocrinol. 36, 90-107.

ACKNOWLEDGMENTS



funded by



Ministero dell'Istruzione, dell'Università e della Ricerca