

VALIDITY AND RELIABILITY OF SUBMAXIMAL FITNESS TESTS BASED ON PERCEPTUAL VARIABLES

Matteo Crotti¹, Andrea Bosio², Pietro L. Invernizzi³

¹School of Exercise and Sports Sciences, Università degli Studi di Milano, Milan, Italy

²Human Performance Laboratory, Mapeisport Research Centre, Varese, Italy

³Department of Biomedical Sciences for Health, Università degli studi di Milano, Italy

Corresponding author:

Prof. Pietro Luigi Invernizzi

Department of Biomedical Sciences for Health

Università degli Studi di Milano, Milan, Italy

Address: Via Kramer 4/A, 20129, Milan, Italy

Email: pietro.invernizzi1@unimi.it

Phone: +39 02 503 15163

Fax: +39 02 503 15163

Other authors:

Matteo Crotti; Email: crotti.mc@gmail.com; Phone: +39 3202318944

Andrea Bosio; Email: and.bos@tiscali.it; Phone: +39 3392731823

Running title: Submaximal tests based on perception

ABSTRACT

BACKGROUND: Few studies examined the validity of fitness tests based on perception scales. The field-based fitness tests batteries are composed by maximal tests and currently, a battery of submaximal tests does not exist.

METHODS: Seventeen physically active male participants (age $40,5 \pm 10,7$ years) performed the Multistage fitness test, the Crunch test, the Push-up test and a Trunk flexion test with a submaximal protocol based on perceptual variables that were defined as non-validated submaximal. The mentioned tests were repeated three times to assess their reliability. The participants also performed four validated maximal tests comprising a velocity dependent ramp test on the treadmill, the Crunch test, the Push-up test, and Sit-and-reach test. The submaximal tests were correlated to the maximal ones to assess their criterion validity.

RESULTS: All the submaximal tests showed an ICC superior to 0.8 apart from the submaximal Crunch test. The correlation between maximal treadmill test and submaximal Multistage fitness test was 0.77. The correlation between Sit-and-reach test and submaximal Trunk flexion test was 0.71. Correlations equal to 0.66 and 0.60 were found between the maximal and submaximal Crunch test and push-up test respectively.

CONCLUSIONS: The use of submaximal tests based on a fixed value of perception can represent a suitable tool to assess cardiorespiratory fitness and flexibility because of their correlation with the corresponding maximal evaluation tests and because of their reliability. On the contrary, muscular endurance capacity seems not to be assessable in the same manner.

Keywords: Perception, Physical Fitness, Exercise Test, Reliability

INTRODUCTION

It is well documented by scientific literature that fit people present lower mortality and morbidity rates together with better physical and physiological health status compared to unfit people ⁽¹⁻⁶⁾. Given the importance of fitness level for health it is fundamental to create tools that allow to monitor it and its different components comprising endurance, strength and flexibility. The most accurate and reliable methods to evaluate physical fitness require sophisticated instruments such as spirometers, metabolimeter, accelerometers, ergometers, dynamometers and force platforms ⁽²⁾. Unfortunately, most of the mentioned tools present some characteristics that might limit their use in field-based research and in testing a great number of participants. In fact, most of them are expensive and, as a consequence, it might be difficult for several public or private organizations to afford their costs. Furthermore, as this tools are used in specialized scientific laboratories, it could be difficult to move them from a place to another. Lastly, they do not allow to evaluate several participants simultaneously or in a short amount of time. As an alternative to laboratory testing, several field-based batteries of tests have been validated such as the Health-Related Fitness Test Battery for Adults, the AAHPERD, the Fitness Canada and the Eurofit for adults ⁽⁷⁻¹¹⁾. These type of tests batteries allow to assess a great amount of participants in a relatively short amount of time without the need of expensive equipment. Moreover, they are suitable to evaluate the different components of the physical fitness. Currently, all the field-based fitness tests batteries are composed by maximal tests which are commonly considered safe for people's health ⁽¹²⁻¹⁴⁾. However, the safety of maximal tests could be argued as previous studies analyzing it involved highly qualified experts in a controlled environment where the health story, and the possible risks for the participant's health were well known differently from what might happen in real life settings ⁽¹²⁻¹⁴⁾. Otherwise, submaximal testing could be considered as the most appropriate way to monitor physical fitness when some variables concerning participants health are not known, or when the participants do not tolerate strenuous efforts or when the tests are not administered by experts ⁽¹⁵⁾. It should be

noted that a field-based alternative to maximal tests, such as a battery of submaximal tests, does not exist. During everyday-sessions of sport and physical activity, where little equipment is available and only few information are available about people's health risks, the employment of submaximal tests based on hearth rate, blood pressure, rate of perceived exertion and other types of perceptual variables could provide valid and helpful information regarding people's capacity to tolerate physical activity⁽¹⁵⁾. A growing interest in quantifying perceptual variables such as exertion, pain, comfort or temperature led to the validation of several perception scales. However, the studies examining the validity of fitness tests based on perception only regard the evaluation of cardiorespiratory fitness^(2, 16). Indeed, it is possible to evaluate cardiorespiratory fitness using submaximal testing thanks to the relation between rate of perceived exertion and workload⁽¹⁶⁻¹⁸⁾. On the contrary, submaximal tests to evaluate strength and flexibility do not exist even if perception scales of strength performance and stretching intensity have been validated. On the basis of what previously said, this study aims to validate a battery of submaximal field-based fitness tests that takes in consideration perceptual variables such as rate of perceived exertion and stretching intensity to evaluate endurance, strength and flexibility in adults⁽¹⁹⁻²²⁾. It is hypothesized that the submaximal tests represent a valid and reliable tool to assess an individual's physical fitness without the exertion of maximal strain.

METHODS

Participants

Seventeen male participants practicing running regularly at competitive or non-competitive level were recruited in this study (Table I). The participants were instructed to follow their normal training activities and to refrain from participating in competition or strenuous maximal tests during the period of participation to the study. Moreover, they were required not to drink alcohol *or* caffeine during the testing days. All the participants signed an informed consent after a detailed

2
3
4
5
6
7
8
9 explanation of the procedures and possible risks and after being informed that they could interrupt
10 the participation to the study at any time. The study protocol was approved by the Ethics Committee
11 of our Institution and was in accordance with the Declaration of Helsinki for the Humans Rights.
12
13
14
15
16
17
18
19
20
21

22 **Fitness testing**

23 *Maximal cardiorespiratory fitness test*

24
25
26 The maximum oxygen uptake (VO₂max) was assessed using a velocity dependent ramp test
27 protocol performed on a treadmill as it has been demonstrated that this type of protocol is
28 appropriate for cardiorespiratory testing⁽²³⁾. A commercially available metabolimeter (Fitmate
29 PRO, Cosmed, Roma, Italy), calibrated prior to each test according to the manual, was used to
30 monitor continuously the oxygen uptake during the test. The protocol of the test involved 5 minutes
31 at a velocity of 8 km/h followed by an increase in speed of 1 km/h every minute⁽²⁴⁾. The test
32 terminated when a person could no longer maintain the speed.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 *Maximal Muscular endurance*

54
55
56 The Crunch test and the Push-up test described below were used to assess the muscular endurance
57 of Rectus abdominis and upper limbs muscles respectively as they are considered valid, reliable
58 tests to assess this construct^(22, 25, 26). The participants were asked to lie in a supine position and had
59 to flex their trunk, reaching their hands forward. A metronome was set at 40 beats per minute At the
60 first beep, the subject began the curl-up, reaching the top position at the second beep and returning
61 to the starting position at the third. Repetitions were counted each time the subject reached the
62 bottom position. The test was concluded either when the subject reached 75 curl-ups, or the cadence
63 was broken.
64
65
66
67
68
69
70
71
72
73
74
75

76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95

During the test, the participants had to raise the body by straightening the arms and to return to the down position after that, until their chin touched the floor. The maximal number of push-ups performed consecutively without rest was counted as the score.

Maximal Flexibility

The Sit-and-reach test was used to assess the flexibility as it is considered a valid and reliable test to assess the flexibility of the posterior chain muscles^(27, 28). This test consists in a trunk flexion from a seated position. The distance in centimeters between the top of the middle fingers and the axis passing between the extremities of the heels was recorded.

Submaximal cardiorespiratory fitness test

The Multistage fitness test, also called shuttle run test, is represented by a shuttle running between two markers placed 20 meters apart at a progressively increasing speed. This test is widely used to evaluate the cardiorespiratory fitness as it was demonstrated to be a valid tool to predict $VO_2\max$ ⁽²⁹⁾. Moreover, the Multistage fitness test was used in previous studies to predict $VO_2\max$ from submaximal performances by means of the existent relation between workload and perceived exertion^(16, 30, 31). The participants were asked to perform the Multistage fitness test and to interrupt the performance when they perceived an exertion equal to 5 (strong) on the CR-10 Borg's Scale. The distance covered during the whole test was recorded. Previous studies suggested not to exceed an RPE equal to 15 (hard) on the 6-20 Borg's Scale to reduce the risk of cardiovascular disease in clinical populations⁽³⁰⁾. As an RPE equal to 5 on the CR-10 Borg's Scale corresponds to an RPE of 15 on the 6-20 Borg's Scale we decided to use it as an endpoint for the submaximal testing^(32, 33). The same criterion was applied to the submaximal Crunch test and to the Submaximal Push-up test presented in this study. We decided to use the CR-10 Borg's Scale instead of 6-20 Borg's Scale because it is suitable for the assessment of the RPE both during endurance and during resistance exercise^(34, 35).

Submaximal muscular endurance tests

The same protocol of the Crunch test was used for the submaximal testing of the abdominal muscular endurance defined submaximal Crunch test, however, the participants were required to stop the test when they perceived an exertion equal to 5 (strong) on the CR-10 Borg's Scale⁽³³⁾. The muscular endurance of the upper limbs was tested using a modified version of the Push-up test defined submaximal Push-up test where the participants were asked to execute the push-ups at a pace of 40 beats per minutes and to interrupt the performance when they perceived an exertion equal to 5 (strong) on the CR-10 Borg's Scale.

Submaximal flexibility tests

An exercise of trunk flexion from a standing position defined submaximal Trunk flexion test was used to assess the flexibility of the posterior chain in a submaximal way. The participants were asked to stand on a 40 cm high stair and to perform a trunk flexion maintaining the leg in a straight position. In particular, the participants were instructed to reach the highest level of trunk flexion without feeling pain, according to the stretching intensity scale (SIS)⁽³⁶⁾, and the distance between the extremity of the fingers and the plane of the stair where the participants were standing was assessed.

Protocol

The participants took part in one testing session per week for five consecutive weeks at the same time and during the same day of the week. During the first day, the participants did a familiarization to the tests' protocols and to the perceptual scales. During the other four days, the participants performed all the maximal tests or all the submaximal tests selected in this study in a specific order (Figure 1).

****Figure 1 about here****

Three testing days concerned submaximal tests while only one testing day concerned maximal tests, therefore, the participants performed the submaximal tests three times and the maximal tests one time. The order of the testing days was randomized (Figure 2).

****Figure 2 about here****

All the submaximal tests selected for the validation in this study were chosen because they satisfy the assumption of logic validity. Indeed, all the submaximal tests of endurance and muscular endurance used in this study present the protocol of previously validated tests such as the Multistage fitness test, the Crunch test and the Push-up test^(18, 22, 25, 26, 29, 37, 38). Differently, the logic validity of the submaximal test of flexibility is guaranteed by the fact that it involves the same movements assessed by the validated test used as a standard in this study and in other studies such as trunk flexion and hip flexion^(22, 39, 40).

Statistical analysis

The statistical analysis was performed using the software “SPSS statistics 20”. The relative reliability of the submaximal tests, that consists in the degree to which individuals maintain their position in a sample with repeated measurement, was assessed calculating the Intraclass Correlation Coefficient (ICC)⁽⁴¹⁾. The absolute reliability of the submaximal tests, that consists in the degree to which repeated measurements vary from individuals, was evaluated by means of the standard error of the measurement (SEM)⁽⁴¹⁾. The Shapiro-Wilk test was used to assess the normality assumption of the data distribution. The Pearson and Spearman tests were used to evaluate the correlation between submaximal tests and maximal tests evaluating the same construct in order to assess the

construct validity of the submaximal tests. The Pearson correlation was used when the data distribution satisfied the assumption of normality, on the contrary, when the assumption was violated the Spearman correlation was employed. An ICC equal or superior to 0.8 was the criterion to accept the reliability of the tests while a correlation equal or superior to 0.7 was the criterion to accept the construct validity of the submaximal tests ⁽⁴¹⁾.

RESULTS

Most of the data resulted normally distributed apart from the results of the submaximal Push-up test and the maximal Crunch test. The results of the maximal and submaximal tests are presented in Table II and Table III.

**** Table II, Table III about here****

The ICC analysis showed significant results (all p-values lower than 0.05) with all the submaximal tests showing an ICC superior to 0.8 apart from the submaximal Crunch test (Table IV).

**** Table IV about here****

The SEM values were heterogeneous as the submaximal tests presented in this study concerned different fitness variables and involved different unit of measurement apart from the submaximal Crunch test and the submaximal Push-up test which both comprised arbitrary units (a.u.). In particular, the multistage fitness test showed a SEM of 102 m, the submaximal Crunch test showed a SEM of 4.08 a.u., the submaximal Push-up showed a SEM of 2.02 a.u. and the Submaximal trunk

flexion test showed a SEM of 1.85 cm. Interestingly, the SEM of the submaximal Crunch test was more than twice the SEM of the submaximal Push-up test.

**** Table V about here****

A relation between the submaximal and the maximal tests evaluating the same construct was found in all of the test with moderate to low correlations (Figure 3). The correlation between maximal treadmill test and submaximal Multistage fitness test was the highest found in this study with a correlation coefficient equal to 0.77. Similarly, a significant correlation was found between Sit-and-reach test and submaximal Trunk flexion with a coefficient equal to 0.71. Significant moderate correlations were found between maximal and submaximal test of muscular endurance performance, indeed correlations equal to 0.66 and 0.60 were found between the maximal and submaximal Crunch test and push-up test respectively (Figure 3).

****Figure 3 about here****

DISCUSSION

The submaximal Multistage fitness test and Trunk flexion test are the only two tests that satisfied the assumption of criterion validity obtaining a correlation with their reference standard higher than 0.7. The submaximal Multistage fitness test is the one that showed the highest correlation ($r = 0.77$) with the respective reference standard represented by the VO₂max measured during an incremental running test to exhaustion performed on a treadmill. This is in line with previous studies where the submaximal rate of perceived exertion was used to estimate the VO₂max^(30, 31). A relation was

found between maximal Crunch test and submaximal Crunch tests ($\rho = 0.60$). A similar relation was found between submaximal and maximal Push-up tests with a correlation coefficient equal to 0.66. Unfortunately, these relations does not satisfy the standards for the assumption of criterion validity ⁽⁴¹⁾. The difference between the protocol of the maximal tests and the protocol of the submaximal tests could be the cause of the low correlations. Indeed, previous studies showed a low correlation (< 0.50) between fitness tests designed to assess abdominal muscular endurance which presented slightly different protocols ⁽⁴²⁾. A further explanation to the low correlations could be attributed to the people's neuromuscular characteristics. In particular, the participants with a higher percentage of type I fibers could be able to execute a relatively higher number of repetitions after reaching a perceived exertion equal to 5 on the Borg's scale compared to the other participants ⁽⁴³⁾. In addition, a recent study of Grosicki et al. reported a high between-subject variability in muscular endurance performances ⁽⁴⁴⁾. In the mentioned study the participants performed a bicep curl exercise at 60% of their Maximal Voluntary Contraction (MVC) until exhaustion and, as a result, a great between-subject variability in the number of repetition was reported ⁽⁴⁴⁾. Knowing that people report equal RPE scores at the same MCV% it is possible to say that there is a substantial variability in the number of repetitions that people can perform during a muscular endurance exercise after the onset of the fatigue ^(34, 45). As regards the flexibility tests, a correlation equal to 0.71 was found between the submaximal Trunk flexion test and the Sit-and-reach test. Notably, this correlation is higher than the ones recorded in previous studies where the Sit-and-reach test was used as a reference standard for criterion validity to validate the back saver Sit-and-reach test and correlations between 0.45 and 0.65 were found ⁽³⁹⁾. Therefore, the submaximal Trunk flexion test is a better estimator of posterior chain muscles flexibility than other previously published test such as the sit and reach back saver test ⁽³⁹⁾.

Three of the submaximal tests satisfied the reliability criteria by achieving an ICC superior to 0.80 while only the submaximal Crunch test did not satisfy this criterion. Particularly, the submaximal Multistage fitness test and Trunk flexion test presented the highest values of reliability with an ICC

equal to 0.94 and 0.96 respectively. These results are in line with the reliability coefficients found in previous studies concerning the Multistage fitness test and the Sit-and-reach test where ICCs between 0,92 and 0,96 were found for the first test and 0,94 for the second ^(7, 30). The submaximal Push-up test also presented reliable results showing an ICC equal to 0.85. Notably, the submaximal Push-up test showed a higher reliability than the Push-up test performed in previous studies (ICC = 0.73) ⁽⁴⁶⁾. Only the submaximal Crunch test presented a low reliability with an ICC equal to 0.76. This results is slightly lower than the one obtained by adults performing the sit-up test of the Eurofit battery of adults (ICC = 0.83), that evaluates muscular endurance of the Rectus abdominis muscles ⁽⁷⁾. However, it should be noted that the Crunch test and the sit-up test present different protocols ⁽⁷⁾. The SEM was also evaluated in this study (Table V) as it is considered a parameter of absolute reliability that consists in the degree to which repeated measures vary for individuals ⁽⁴¹⁾. The SEM is necessary to evaluate the capacity of a test to monitor the changes over time. In other words, SEM allows to assess to which extent a test provides accurate scores. Furthermore, it could represent a practical indication for the people who use the tests to understand whether the change in the measurement over time is due to measurement errors or to other factors such as the effect of a training. Unfortunately, the lack of data in the current literature concerning the SEM in fitness test similar to the ones exposed in this study make it difficult to interpret this results. However, a method to interpret the SEM was proposed by clinimetrics, that is a methodological discipline which investigates the quality of medical research and clinical practice ⁽⁴⁷⁾. Clinimetrics established that the amount of absolute error of the measurement (e.g. SEM) should be lower than the minimal amount of change that is considered to be relevant, also called Minimal Important Change (MIC) ⁽⁴⁸⁾. Hopkins was the first who proposed to apply this concept to the field of sport performance, and he suggested that the MIC related to performance variables and the absolute error of the measurement of exercise tests should be calculated to assess whether a test is appropriate or not for a specific purpose ⁽⁴⁹⁾. Impellizzeri and colleagues applied this method in a study concerning the evaluation of the validity of a repeated sprint ability test in soccer players ⁽⁵⁰⁾.

The main limit of this study is the fact that the participants are not representative of the general population as the sample was composed by people who regularly participate in physical activity and therefore presents a good or high fitness level. In fact, a great part of the population is represented by people who are unfit, obese or overweight and sedentary. It is possible that people with low level of fitness and experience in physical activity might have difficulties in performing submaximal tests based on their perception given their lack of practice in the evaluation of their self-perception during physical efforts. Therefore, It is fundamental to make people familiarize and practice with perceptual scales, such as the Borg scale and the SIS scale, according to the indication presented by the respective authors of the scales^(33, 36).

CONCLUSIONS

The submaximal Multistage fitness test and the submaximal Trunk flexion test satisfied the standard for logic validity, criterion validity and reliability and therefore can be considered as valid tools to assess cardiorespiratory fitness and flexibility respectively. The submaximal Crunch test did not show an adequate reliability and correlation with the maximal Crunch test while the submaximal Push-up test did not satisfy the standard of criterion validity, therefore, both the tests cannot be considered a valid tool to predict maximal muscular endurance capacities. Consequently, submaximal tests based on perception seem to be a suitable method to evaluate cardiorespiratory fitness and flexibility while they seem to be inadequate to assess muscular endurance capacities. On the basis of the finding of this study future research in the field of fitness and wellness should consider the opportunity to implement and validate submaximal fitness test based on perception to evaluate cardiorespiratory fitness and flexibility. Future studies should clarify the relation between rate of perceived exertion (internal load), load (%MVC) and number of repetitions, and the factors that might influence this relation should be identified. The practitioners that work in the field of health promotion could consider new ways of training and testing based on the perceptions or

internal load rather than on external load. This could help to avoid problems like injuries and overtraining and it could educate the people to new perspectives of self-awareness and knowledge. This might help fostering health and wellness in the population.

ACKNOWLEDGMENTS

We would like to thank all the participants in this research. No conflict of interest does exist between the authors.

REFERENCES

1. Bemben DA, Bemben MG. Dose-response effect of 40 weeks of resistance training on bone mineral density in older adults. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2011 Jan;22(1):179-86.
2. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Medicine and science in sports and exercise* 2011 Jul;43(7):1334-59.
3. Johnson JL, Slentz CA, Houmard JA, Samsa GP, Duscha BD, Aiken LB, McCartney JS, Tanner CJ, Kraus WE. Exercise training amount and intensity effects on metabolic syndrome (from Studies of a Targeted Risk Reduction Intervention through Defined Exercise). *The American journal of cardiology* 2007 Dec 15;100(12):1759-66.

4. Lee DC, Sui X, Ortega FB, Kim YS, Church TS, Winett RA, Ekelund U, Katzmarzyk PT, Blair SN. Comparisons of leisure-time physical activity and cardiorespiratory fitness as predictors of all-cause mortality in men and women. *British journal of sports medicine* 2011 May;45(6):504-10.
5. Ruiz JR, Sui X, Lobelo F, Morrow JR, Jr., Jackson AW, Sjostrom M, Blair SN. Association between muscular strength and mortality in men: prospective cohort study. *BMJ (Clinical research ed)* 2008;337:a439.
6. Sillanpaa E, Laaksonen DE, Hakkinen A, Karavirta L, Jensen B, Kraemer WJ, Nyman K, Hakkinen K. Body composition, fitness, and metabolic health during strength and endurance training and their combination in middle-aged and older women. *European journal of applied physiology* 2009 May;106(2):285-96.
7. Tsigilis N, Douda H, Tokmakidis SP. Test-retest reliability of the Eurofit test battery administered to university students. *Perceptual and motor skills* 2002 Dec;95(3 Pt 2):1295-300.
8. Suni JH, Miilunpalo SI, Asikainen TM, Laukkanen RT, Oja P, Pasanen ME, Bos K, Vuori IM. Safety and feasibility of a health-related fitness test battery for adults. *Physical therapy* 1998 Feb;78(2):134-48.
9. Suni JH, Oja P, Laukkanen RT, Miilunpalo SI, Pasanen ME, Vuori IM, Vartiainen TM, Bos K. Health-related fitness test battery for adults: aspects of reliability. *Archives of physical medicine and rehabilitation* 1996 Apr;77(4):399-405.
10. Oja P, Tuxworth B, Council of E, Committee for the Development of S, Research UKKIfHP. Eurofit for adults : assessment of health-related fitness. Strasbourg; Tampere, Finland; Croton-on-Hudson, N.Y.: Council of Europe, Committee for the Development of Sport ; UKK Institute for Health Promotion Research ; Manhattan Pub. Co. [distributor]; 1995.
11. Canadian Association of Sports S, Fitness Appraisal C, Accreditation P, Fitness C. Canadian standardized test of fitness (CSTF) operations manual : (for 15 to 69 years of age). Ottawa: Government of Canada, Fitness and Amateur Sport; 1987.

12. Ferraz RB, Gualano B, Filho CM, Almeida MG, Perandini LA, Dassouki T, Sa-Pinto AL, Lima FR, Roschel H. Safety and feasibility of maximal physical testing in rheumatic diseases: a cross-sectional study with 5,910 assessments. *Rheumatology international* 2015 Jun;35(6):1027-36.
13. Scardovi AB, Coletta C, De Maria R, Perna S, Aspromonte N, Feola M, Rosso G, Greggi M, Ceci V. The cardiopulmonary exercise test is safe and reliable in elderly patients with chronic heart failure. *Journal of cardiovascular medicine (Hagerstown, Md)* 2007 Aug;8(8):608-12.
14. Gibbons L, Blair SN, Kohl HW, Cooper K. The safety of maximal exercise testing. *Circulation* 1989 Oct;80(4):846-52.
15. Pescatello LS, American College of Sports M. ACSM's guidelines for exercise testing and prescription. Philadelphia: Wolters Kluwer/Lippincott Williams & Wilkins Health; 2014.
16. Coquart J, Tabben M, Farooq A, Tourny C, Eston R. Submaximal, Perceptually Regulated Exercise Testing Predicts Maximal Oxygen Uptake: A Meta-Analysis Study. *Sports medicine (Auckland, NZ)* 2016 Jan 20.
17. Coquart JB, Garcin M, Parfitt G, Tourny-Chollet C, Eston RG. Prediction of maximal or peak oxygen uptake from ratings of perceived exertion. *Sports medicine (Auckland, NZ)* 2014 May;44(5):563-78.
18. Davies RC, Rowlands AV, Eston RG. The prediction of maximal oxygen uptake from submaximal ratings of perceived exertion elicited during the multistage fitness test. *British journal of sports medicine* 2008 Dec;42(12):1006-10.
19. Lagally KM, Costigan EM. Anchoring procedures in reliability of ratings of perceived exertion during resistance exercise. *Perceptual and motor skills* 2004 Jun;98(3 Pt 2):1285-95.
20. Buckley JP, Borg GA. Borg's scales in strength training; from theory to practice in young and older adults. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme* 2011 Oct;36(5):682-92.

21. Robertson RJ, Goss FL, Rutkowski J, Lenz B, Dixon C, Timmer J, Frazee K, Dube J, Andreacci J. Concurrent validation of the OMNI perceived exertion scale for resistance exercise. *Medicine and science in sports and exercise* 2003 Feb;35(2):333-41.
22. USA Yot. YMCA fitness testing and assessment manual. [Champaign, Ill.]: Published for the YMCA of the USA by Human Kinetics; 2000.
23. Sperlich PF, Holmberg HC, Reed JL, Zinner C, Mester J, Sperlich B. Individual versus Standardized Running Protocols in the Determination of VO₂max. *J Sports Sci Med* 2015 Jun;14:386-93.
24. Kasch FW, Wallace JP, Huhn RR, Krogh LA, Hurl PM. VO₂max during horizontal and inclined treadmill running. *Journal of applied physiology* 1976 Jun;40(6):982-3.
25. Canadian Society for Exercise P. Canadian physical activity, fitness & lifestyle approach : CSEP - health & fitness program's health-related appraisal & counselling strategy. Ottawa, Ont.: Canadian Society for Exercise Physiology; 2004.
26. Blair SN, American College of Sports M. Resource manual for Guidelines for exercise testing and prescription. Philadelphia: Lea & Febiger; 1988.
27. Ayala F, Sainz de Baranda P, De Ste Croix M, Santonja F. Reproducibility and criterion-related validity of the sit and reach test and toe touch test for estimating hamstring flexibility in recreationally active young adults. *Phys Ther Sport* 2012 Nov;13:219-26.
28. Ayala F, Sainz de Baranda P, De Ste Croix M, Santonja F. Criterion-related validity of four clinical tests used to measure hamstring flexibility in professional futsal players. *Phys Ther Sport* 2011 Nov;12:175-81.
29. Ramsbottom R, Brewer J, Williams C. A progressive shuttle run test to estimate maximal oxygen uptake. *Br J Sports Med* 1988 Dec;22:141-4.
30. Coquart JB, Garcin M, Parfitt G, Tourny-Chollet C, Eston RG. Prediction of maximal or peak oxygen uptake from ratings of perceived exertion. *Sports Med* 2014 May;44:563-78.

- 2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
31. Davies RC, Rowlands AV, Eston RG. The prediction of maximal oxygen uptake from submaximal ratings of perceived exertion elicited during the multistage fitness test. *Br J Sports Med*2008 Dec;42:1006-10.
 32. Borg G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med*1970;2:92-8.
 33. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*1982;14:377-81.
 34. Lagally KM, Costigan EM. Anchoring procedures in reliability of ratings of perceived exertion during resistance exercise. *Percept Mot Skills*2004 Jun;98:1285-95.
 35. Buckley JP, Borg GA. Borg's scales in strength training; from theory to practice in young and older adults. *Appl Physiol Nutr Metab*2011 Oct;36:682-92.
 36. Freitas SR, Vaz JR, Gomes L, Silvestre R, Hilario E, Cordeiro N, Carnide F, Pezarat-Correia P, Mil-Homens P. A New Tool to Assess the Perception of Stretching Intensity. *J Strength Cond Res*2015 Sep;29:2666-78.
 37. Suni JH, Miilunpalo SI, Asikainen TM, Laukkanen RT, Oja P, Pasanen ME, Bos K, Vuori IM. Safety and feasibility of a health-related fitness test battery for adults. *Phys Ther*1998 Feb;78:134-48.
 38. Burden AM, Redmond CG. Abdominal and hip flexor muscle activity during 2 minutes of sit-ups and curl-ups. *J Strength Cond Res*2012 Aug;27:2119-28.
 39. Baltaci G, Un N, Tunay V, Besler A, Gerceker S. Comparison of three different sit and reach tests for measurement of hamstring flexibility in female university students. *Br J Sports Med*2003 Feb;37:59-61.
 40. Mayorga-Vega D, Merino-Marban R, Viciano J. Criterion-Related Validity of Sit-and-Reach Tests for Estimating Hamstring and Lumbar Extensibility: a Meta-Analysis. *J Sports Sci Med*2014 Jan;13:1-14.

41. Impellizzeri FM, Marcora SM. Test validation in sport physiology: lessons learned from clinimetrics. *Int J Sports Physiol Perform* 2009 Jun;4:269-77.
42. Sparling PB, Millard-Stafford M, Snow TK. Development of a cadence curl-up test for college students. *Res Q Exerc Sport* 1998 Dec;68:309-16.
43. Komi PV, Tesch P. EMG frequency spectrum, muscle structure, and fatigue during dynamic contractions in man. *Eur J Appl Physiol Occup Physiol* 1979 Sep;42:41-50.
44. Grosicki GJ, Miller ME, Marsh AP. Resistance exercise performance variability at submaximal intensities in older and younger adults. *Clinical interventions in aging* 2014;9:209-18.
45. Robertson RJ, Goss FL, Rutkowski J, Lenz B, Dixon C, Timmer J, Frazee K, Dube J, Andreacci J. Concurrent validation of the OMNI perceived exertion scale for resistance exercise. *Med Sci Sports Exerc* 2003 Feb;35:333-41.
46. Lubans DR, Smith JJ, Harries SK, Barnett LM, Faigenbaum AD. Development, test-retest reliability, and construct validity of the resistance training skills battery. *J Strength Cond Res* 2014 May;28:1373-80.
47. Fava GA, Tomba E, Sonino N. Clinimetrics: the science of clinical measurements. *Int J Clin Pract* 2011 Jan;66:11-5.
48. Terwee CB, Bot SD, de Boer MR, van der Windt DA, Knol DL, Dekker J, Bouter LM, de Vet HC. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 2006 Jan;60:34-42.
49. Hopkins WG, Hawley JA, Burke LM. Design and analysis of research on sport performance enhancement. *Med Sci Sports Exerc* 1999 Mar;31:472-85.
50. Impellizzeri FM, Rampinini E, Castagna C, Bishop D, Ferrari Bravo D, Tibaudi A, Wisloff U. Validity of a repeated-sprint test for football. *Int J Sports Med* 2008 Nov;29:899-905.

Table I. - Demographic information of the subjects.

Participants	Mean	Std. Deviation
Age (yrs)	40.5	10.7
Body mass (Kg)	69.3	4.7
Stature (cm)	174.6	6.2
Body Mass Index	22.7	1.1

PEER REVIEW COPY
The Journal of Sports Medicine and
Physical Fitness

Table II. – Descriptive statistics of the submaximal tests.

Submaximal test	Minimum	Maximum	Mean	Std. Deviation	Units
Submaximal Multistage fitness test	1133	2527	1656.9	413.6	m
Submaximal Crunch test	10.3	40	23.4	8.7	a.u.
Submaximal Push-up test	6	22	12.3	4.6	a.u.
Submaximal Trunk flexion test	-5	20	8.9	8.0	cm

a.u. = arbitrary units

PEER REVIEW COPY
The Journal of Sports Medicine and
Physical Fitness

Table III. - Descriptive statistics of the maximal tests.

Maximal tests	Minimum	Maximum	Mean	Std. Deviation	Units
Treadmill test	45.6	65.2	55.5	5.8	ml · kg ⁻¹ · min ⁻¹
Crunch test	15	75	35.7	18.1	a.u.
Push-up test	12	42	21.3	7.7	a.u.
Sit-and-reach	7	48	29.9	10.5	cm

a.u. = arbitrary units

PEER REVIEW COPY
The Journal of Sports Medicine and
Physical Fitness

Table IV. – Relative reliability of the submaximal tests.

Submaximal test	ICC	95% Confidence Interval		p-value
		Lower Bound	Upper Bound	
Submaximal Multistage fitness test	0.953	0.905	0.979	p < 0.001
Submaximal Crunch test	0.756	0.570	0.883	p < 0.001
Submaximal Push-up test	0.853	0.725	0.932	p < 0.001
Submaximal Trunk flexion test	0.946	0.892	0.976	p < 0.001

ICC: Intraclass Correlation.

PEER REVIEW COPY
The Journal of Sports Medicine and
Physical Fitness

Table V. – Absolute reliability of the submaximal tests.

Submaximal test	SEM	Units
Multistage fitness test	102	m
Submax Crunch test	4.08	a.u.
Submax Push-up test	2.02	a.u.
Submaximal Trunk flexion test	1.85	cm


a.u. : arbitrary units; SEM: Standard Error of the Measurement.

PEER REVIEW COPY
The Journal of Sports Medicine and
Physical Fitness

Test order

Maximal tests day	Submaximal tests day
<p>Tests order:</p> <ol style="list-style-type: none"> 1. Sit-and-reach test 2. Crunch test 3. Push-up test 4. Treadmill test 	<p>Tests order:</p> <ol style="list-style-type: none"> 1. Submaximal Trunk flexion test 2. Submaximal Crunch test 3. Submaximal Push-up test 4. Submaximal Multistage fitness test

Testing day order



Correlation between maximal and submaximal tests

