

# Thiol precursors in Catarratto Bianco Comune and Grillo grapes and effect of clarification conditions on the release of varietal thiols in wine

D. FRACASSETTI<sup>1</sup> , M. STUKNYTĖ<sup>1</sup>, C. LA ROSA<sup>2</sup>, M. GABRIELLI<sup>1</sup>, I. DE NONI<sup>1</sup> and A. TIRELLI<sup>1</sup>

<sup>1</sup> Department of Food, Environmental and Nutritional Sciences (DeFENS), Università degli Studi di Milano, 20133, Milan, Italy; <sup>2</sup> Department of Pharmaceutical Sciences, Section of General and Organic Chemistry 'A. Marchesini', Università degli Studi di Milano, 20133 Milan, Italy

Corresponding author: Professor Antonio Tirelli, email antonio.tirelli@unimi.it

## Abstract

**Background and Aims:** Varietal thiols characterise the typical aroma of several white wines, such as Sauvignon Blanc. Their presence and sensory perception were suggested in two Sicilian grape cultivars, Catarratto Bianco Comune (CBC) and Grillo, though it has not been analytically proven.

**Methods and Results:** Varietal thiol precursors and free varietal thiols were assessed in CBC and Grillo grapes, musts and wines by ultra-performance liquid chromatography/electrospray ionisation high-resolution MS. The isobaric compounds including *S*-3-(hexanal)-glutathione and *S*-3-(4-mercapto-4-methylpentan-2-one)-glutathione (GSH-4MMP) were distinguished by comparing their accurate masses and high-resolution-MS/MS spectra with those of their synthetic standards. *S*-3-(Hexanal)-glutathione, *S*-3-(hexan-1-ol)-glutathione and *S*-3-(hexan-1-ol)-cysteine were found in grape, must and wine, whereas GSH-4MMP and its hydrolysed forms were not found. Their concentration decreased during winemaking, mostly after grape pressing. We compared the effect of clarification conditions based on the exposure of must to either air or CO<sub>2</sub> on the concentration of thiol precursors and free thiols in wine; however, negligible differences were observed. The concentration of free thiols in the wines was found to be in the range 400–1100 ng/L and they were unaffected by the two clarification conditions tested.

**Conclusions:** The isobaric *S*-3-(hexanal)-glutathione and GSH-4MMP were clearly distinguished for the first time by ultra-performance liquid chromatography/high-resolution MS through their retention times and MS spectra. These varietal thiols were revealed in CBC and Grillo wines for the first time. The air-free and air-exposed clarification treatments had little effect on the concentration of the varietal thiols.

**Significance of the Study:** This study highlights the major impact of the varietal thiols (mainly 3-mercapto-hexan-1-ol and its acetate form) on the sensory properties of CBC and Grillo wines.

**Keywords:** *Catarratto Bianco Comune cultivar, Grillo cultivar, thiol precursors, UPLC/ESI-HRMS, varietal thiols*

## Introduction

Varietal thiols are grape-derived sulfur compounds contributing to the typical flavour of many white wines, such as Sauvignon Blanc; among them 3-mercaptohexan-1-ol (3MH), 3-mercaptohexylacetate (3MHA) and 4-mercapto-4-methylpentan-2-one (4MMP) are considered as the most important and pleasant compounds (Darriet et al. 1995, Tominaga et al. 1998b, 2000). 3-Mercaptohexan-1-ol and 3MHA have an olfactory perception threshold of 60 and 4 ng/L, respectively, and they are responsible for passion-fruit-like and grapefruit-like olfactory notes (Tominaga et al. 1998a). These compounds have also been found in red wines (Bouchilloux et al. 1998). 4-Mercapto-4-methylpentan-2-one has an olfactory perception threshold of 0.8 ng/L, and its aroma is described as box tree-like, black currant-like or even cat urine-like when occurring at high concentration (Tominaga et al. 1998a).

Varietal thiols occur as non-volatile precursors in the grape berry, where they share their sulfur atom with a cysteine residue. 3-Mercaptohexan-1-ol bound as a cysteinyl-conjugate [*S*-3-(hexan-1-ol)-cysteine (Cys-3MH)], a glutathionyl-conjugate [*S*-3-(hexan-1-ol)-glutathione (GSH-3MH)] and also a cysteinylglycine conjugate [*S*-3-(hexan-1-ol)-cysteinylglycine (CysGly-3MH)] have been reported (Coetzee

and Du Toit 2012, Peña-Gallego et al. 2012). Either (*E*)-2-hexenal or (*E*)-2-hexen-1-ol can act as a precursor of 3MH (Schneider et al. 2006, Harsch et al. 2013). (*E*)-2-Hexenal can be released from linolenic acid in the presence of oxygen by a lipoxygenase/lyase sequence, and then converted to GSH-3MH by coupling with glutathione (GSH) (Hatanaka et al. 1995, Allen et al. 2011, Harsch et al. 2013). Under aerobic conditions, *Saccharomyces* can oxidise (*E*)-2-hexen-1-ol into (*E*)-2-hexenal, and the reverse process can occur under anaerobic conditions (Harsch et al. 2013). (*E*)-2-Hexenal may act as a precursor when hydrogen sulfide is released in the early part of the fermentation (Schneider et al. 2006, Roland et al. 2010, Pinu et al. 2012). 4-Mercapto-4-methylpentan-2-one occurs in grape and must as the cysteine conjugate [*S*-3-(4-mercapto-4-methylpentan-2-one)-cysteine (Cys-4MMP)] and glutathione conjugate [*S*-3-(4-mercapto-4-methylpentan-2-one)-glutathione (GSH-4MMP)] (Fedrizzi et al. 2009, Roland et al. 2010). The free forms of varietal thiols are released along with the alcoholic fermentation by a *Saccharomyces cerevisiae* lyase (Tominaga et al. 1998b, Murat et al. 2001). Recently, Thibon et al. (2016) identified *S*-3-(hexan-1-ol)-glutathione (GSH-3MHA) in Sauvignon Blanc juice. This compound can be considered as a precursor of thiol aromas.

Varietal thiols have been found in some Italian grape cultivars, such as Verdicchio Bianco (also known as Trebbiano di Lugana) (Mattivi et al. 2012) and Arneis (Piano et al. 2014), and they were considered to occur in Grillo wine (Calò et al. 2006), which is used to produce Marsala wine together with the Catarratto Bianco Comune (CBC) cultivar. Both cultivars are important autochthonous grape cultivars of Sicily, the largest and the southern-most Italian wine region accounting for about 17.5% of the overall Italian wine production (De Lorenzis et al. 2014). Catarratto Bianco Comune is the most widespread grape cultivar in Sicily (Carimi et al. 2010) and the second most important white grape cultivar in Italy (Robinson et al. 2012). The Grillo grape was thought to be the offspring of CBC and Muscat of Alexandria (Robinson et al. 2012). Both its high vigour and sugar concentration make the Grillo grape more suitable than CBC for the production of Marsala wine (Calò et al. 2006). Even though the role of varietal thiols in the flavour of CBC and Grillo wines has been suggested (Corona 2010), it has not been analytically assessed to date.

This research aimed to characterise Grillo and CBC grapes, musts and wines based on the concentration of the varietal thiols and their precursors. The effect of must exposure to air on the formation of the thiol precursors was also evaluated, since it could increase the level of thiol precursors and, as a consequence, the concentration of varietal thiols in wine (Mattivi et al. 2012).

## Materials and methods

### Chemicals and reagents

Methanol, ethanol, dichloromethane (DCM), acetonitrile, formic acid (FA), anhydrous tetrahydrofuran, ammonium acetate, sodium fluoride, glutathione (GSH), (*E*)-2-hexenal, 2-hexyn-1-ol, thioacetic acid, activated manganese(IV) oxide, deuterium oxide, pyridine, Vitride [Red-Al sodium bis(2-methoxyethoxy) aluminium hydride], *p*-benzoquinone (*p*BQ), 3-mercaptopropanoic acid (3MPA), sodium chloride, sodium borohydride (NaBH<sub>4</sub>), ethanolamine (EA), *o*-phthalaldehyde (OPA), anhydrous sodium sulfate, calcium carbonate and boric acid were purchased from Sigma-Aldrich (St Louis, MO, USA); octadecyl-functionalised silica gel 60 RP-18 (40–63 mesh) and sodium hydroxide (NaOH) from Merck Millipore (Darmstadt, Germany); sodium metabisulfite from J. T. Baker (Deventer, The Netherlands); and polyvinylpyrrolidone (PVPP) resin from Dal Cin Gildo (Concorezzo, Italy). All chemicals were at least of analytical grade. Water of ultra-performance liquid chromatography (UPLC)-grade was obtained by a Milli-Q system (Merck Millipore).

### Grape sampling

Berry samples (500 g each) of the CBC and Grillo cultivars were collected during the ripening period until harvest in the 2014 vintage. Overall, four samplings of CBC and three of Grillo grapes were made. Berry samples were taken from the 5.5 ha vineyards Piana Regina and Sant' Anna at Tenuta Regaleali (Sicily, Italy, GPS HESI coordinates: 37.71°N, 13.85°E) by cutting the pedicel to prevent any berry damage. A maximum of three berries per sampled bunch was taken. The sampling position on the bunch (front, back, top and bottom) varied in and with each bunch sampled in order to collect a representative sample of grape berries. The samples were stored in aluminium containers at –18°C.

### Winemaking

Catarratto Bianco Comune grape bunches (81 000 kg) and Grillo grape bunches (52 000 kg) were hand harvested at ripening, and the winemaking followed the procedures usually adopted by the winery for both cultivars. In detail, during destemming and crushing, 10 g of pectolytic enzyme/1000 kg was added to the juices. Then, 20 000 and 14 000 kg of CBC and Grillo berries, respectively, were pressed at 60 kPa in a 120 min cycle in a closed-tank membrane press without air removal. The extracted juice received 40 mg/L SO<sub>2</sub> in the collection vessel and was cooled to 12°C in a heat exchanger. The two juices were directly transferred to stainless steel tanks. For each cultivar, the juice (12 hL from CBC and 80 hL from Grillo grapes) was pumped into two tanks through plastic pipes. One of the tanks was filled with air [samples named air-exposed must (AEM)], whereas the second tank was flushed and filled with CO<sub>2</sub> gas [samples named air-free must (AFM)] in order to strip oxygen. Both musts were cooled to 7°C in order to slow microbial growth and then submitted to pump-overs after 12 and 24 h (with air exposure for AEM and under CO<sub>2</sub> for AFM) before undergoing 12 h settling. The CBC musts were inoculated with *S. cerevisiae* 20 CRU611 strain (Ever Intec, Pramaggiore, Italy). The Grillo musts were inoculated with *S. cerevisiae* 25 NT116 strain (Vason, San Pietro in Cariano, Italy). Moreover, 15 g/hL of diammonium phosphate and 0.03 g/hL of thiamine were added, and the temperature was raised to 10–15°C. When about one-third of the sugar was fermented, the musts were further supplemented with 15 g/hL of diammonium phosphate and 0.03 g/hL of thiamine. The Grillo musts were also supplemented with 20 g/hL of yeast hulls. The density of the alcoholic fermentations was monitored daily with a hydrometer and the temperature was corrected. Once the alcoholic fermentation was completed, the wines were cooled to 10°C, racked off, supplemented with SO<sub>2</sub> up to 50 mg/L as the free form and bottled. These bottles were stored at 15°C until sensory evaluation. Winemaking for each cultivar was performed in duplicate.

### Must and wine sampling

The must samples were collected from the tank sampling valve into 500 mL plastic bottles after grape pressing, must clarification, yeast inoculation and at the end of fermentation. All these samples were collected in triplicate, filled with N<sub>2</sub> and immediately frozen at –18°C until analysis.

### Determination of GSH and grape reaction product

The concentration of GSH and grape reaction product (GRP) in grape, must and wine samples was assessed as described by Fracassetti and Tirelli (2015).

### Determination of thiol precursors


#### Synthesis of $\gamma$ -L-glutamyl-S-[(1R/S)-1-(2-hydroxyethyl)butyl]-L-cysteinyl-glycine (GSH-3MH).

The compound GSH-3MH was prepared according to the procedure reported by Grant-Preece et al. (2010) with a slight modification. (*E*)-2-Hexenal (0.16 g, 1.63 mmol) was added to a suspension of GSH (0.5 g, 1.63 mmol) in 50% aqueous acetonitrile (10 mL), following supplementation with pyridine (0.3 g, 3.8 mmol). The clear solution was stirred for 64 h at 25°C, diluted with water (10 mL) and washed with DCM (4 × 8 mL). The aqueous layer was concentrated under reduced pressure to obtain a yellowish

solid. A solution of this product in water (10 mL) was cooled to 0°C and treated with NaBH<sub>4</sub> (0.1 g, 2.6 mmol). The mixture was stirred at 0°C for 4 h before being quenched with 10% aqueous HCl to pH 3. After evaporation of the solvent, the resulting solid was purified by means of a C18 reversed-phase low-pressure column (LiChroprep RP-18, 40–63 μm, 3 g, 4 × 1.3 cm bed for 0.1 g of product, Merck, Darmstadt, Germany). The column was subsequently eluted with water (20 mL), 1% aqueous ethanol (20 mL), 5% aqueous ethanol (20 mL) and 15% aqueous ethanol (20 mL). Fractions of the last eluent were evaporated under reduced pressure to obtain GSH-3MH as a white solid (35%). It was a mixture of two epimers at C<sub>1</sub>, as evidenced from NMR spectrum; <sup>1</sup>H NMR spectra were recorded with a Varian-Gemini 200 MHz spectrometer (Agilent, Santa Clara, CA, USA). Chemical shifts (δ) are given in ppm in relation to tetramethylsilane. The MS spectra were recorded with a LCQ Advantage AP electrospray/ion trap equipped instrument (Thermo Fisher Scientific, San Jose, CA, USA) using a syringe pump device to directly inject sample solutions. The structure of GSH-3MH was confirmed by NMR spectrum.

<sup>1</sup>H NMR (δ, D<sub>2</sub>O): 0.73, 0.74 (3H, 2t, *J* = 7.0 Hz, H<sub>4</sub>); 1.20–1.80 (6H, m, H<sub>2',3',1''</sub>); 2.00 (2H, q, *J* = 7.3 Hz, H<sub>9</sub>); 2.38 (2H, 2t, *J* = 8.0 Hz, H<sub>8</sub>); 2.65–2.89 (2H, m, H<sub>1',12a</sub>); 2.92 (1H, dd, *J* = 13.9, 5.1 Hz, H<sub>12b</sub>); 3.52–3.62 (2H, m, H<sub>2''</sub>); 3.64 (1H, t, *J* = 6.2 Hz, H<sub>10</sub>); 3.79 (2H, s, H<sub>2</sub>); 4.39 (1H, m, H<sub>5</sub>) (Figure S1).

**Synthesis of γ-L-glutamyl-S-[(1*R*/S)-1-(2-hydroxyethyl)butyl-1*d*]-L-cysteinyl-glycine (d1-GSH-3MH).** Labelled compound d1-GSH-3MH (Figure 1a) was synthesised using (*E*)-2-hexenal-3-*d* prepared starting from the commercial 2-hexyn-1-ol as reported by Bennani et al. (2009). Glutathione (0.465 g, 1.515 mmol) was treated with (*E*)-2-hexenal-3-*d* (0.15 g, 1.515 mmol) and pyridine (0.285 g, 3.53 mmol) exactly as described by Bennani et al. (2009) for the synthesis of GSH-3MH. The subsequent reduction with NaBH<sub>4</sub> (0.093 g, 2.42 mmol) and purification by means of a C18 reversed-phase low-pressure column chromatography (LiChroprep RP-18, Merck) resulted in the compound d1-GSH-3MH (Figure 1a) as a white solid (32%). It was a mixture of two epimers at C<sub>1</sub>, as evidenced from the NMR spectrum.



**Figure 1.** Synthesis schemes of (a) γ-L-glutamyl-S-[(1*R*/S)-1-(2-hydroxyethyl)butyl-1*d*]-L-cysteinyl-glycine (d1-GSH-3MH) and (b) (*R*/S)-3-mercapto-1-hexan-3*d*-ol (d1-3MH).

<sup>1</sup>H NMR (δ, D<sub>2</sub>O): 0.73, 0.74 (3H, 2t, *J* = 7.0 Hz, H<sub>4</sub>); 1.20–1.75 (6H, m, H<sub>2',3',1''</sub>); 2.01 (2H, q, *J* = 7.0 Hz, H<sub>9</sub>); 2.38 (2H, 2t, *J* = 8.5 Hz, H<sub>8</sub>); 2.71 (1H, dd, *J* = 13.7, 8.7 Hz, H<sub>12a</sub>); 2.92 (1H, dd, *J* = 13.7, 5.1 Hz, H<sub>12b</sub>); 3.52–3.61 (2H, m, H<sub>2''</sub>); 3.66 (1H, t, *J* = 6.4 Hz, H<sub>10</sub>); 3.81 (2H, s, H<sub>2</sub>); 4.4 (1H, m, H<sub>5</sub>) (Figure S2).

#### Synthesis of γ-L-glutamyl-S-[(1*R*/S)-1-(2-oxoethyl)butyl]-L-cysteinyl-glycine (GSH-3MHAI).

Compound GSH-3MHAI was prepared as described above for the GSH-3MH except that the intermediate aldehyde was isolated. (*E*)-2-Hexenal (0.16 g, 1.63 mmol) was added to a suspension of GSH (0.5 g, 1.63 mmol) in 50% aqueous acetonitrile (10 mL), following supplementation with pyridine (0.3 g, 3.8 mmol). The clear solution was stirred for 64 h at 25°C, then diluted with water (10 mL) and washed with DCM (4 × 8 mL). The aqueous layer was concentrated under reduced pressure to obtain a yellowish solid (0.66 g) that was purified by means of a C18 reversed-phase, low-pressure column (3 g, 4 × 1.3 cm bed for 0.1 g of product). The column was subsequently eluted with water (20 mL), 5% aqueous ethanol (20 mL) and 15% aqueous ethanol (20 mL). Fractions of the last eluent were evaporated under reduced pressure to obtain GSH-3MHAI as a white solid (40%). The <sup>1</sup>H NMR spectrum was in agreement with what was previously reported (Thibon et al. 2016).

**Synthesis of γ-L-glutamyl-S-(1,1-dimethyl-3-oxo-butyl)-L-cysteinyl-glycine (GSH-4MMP).** The compound GSH-4MMP was prepared according to the procedure reported by Fedrizzi et al. (2009).

**Synthesis of S-[(1*R*,S)-1-(2-hydroxyethyl)butyl]-L-cysteine (Cys-3MH).** The compound Cys-3MH was prepared according to the procedure reported by Pardon et al. (2008) and the chloro-hydrate Cys-3MH was obtained.

#### Sample preparation for thiol precursor analysis

Grape juice, must and wine samples of both cultivars were analysed for thiol precursors. The grape juice was obtained from frozen grape samples, as described by Fracassetti and Tirelli (2015). The must and wine samples were collected directly from the tanks during and after fermentation.

All the samples were purified by solid-phase extraction (SPE) as described by Capone et al. (2010) with slight modification. The SPE cartridges (Strata X-polymeric sorbent 200 mg, Phenomenex, Torrance, CA, USA) were activated with 5 mL of methanol and 5 mL of Milli-Q-treated water. The sample (2 mL) was spiked with d1-GSH-3MH (500 μg/L; exact mass [M + H]<sup>+</sup>: 409.159) internal standard (IS), loaded onto the SPE column and eluted with 4 mL of methanol after a washing step with 2 mL water. The solvent was evaporated under N<sub>2</sub> flow to 400 μL, the sample was analysed by ultra-performance liquid chromatography coupled to electrospray ionisation high-resolution MS (UPLC/ESI-HRMS).

#### Ultra-performance liquid chromatography/electrospray ionisation high-resolution MS for the quantification of thiol precursors

The thiol precursors were analysed by UPLC/ESI-HRMS [Acquity UPLC separation module (Waters, Milford, MA, USA) coupled to a Q Exactive hybrid quadrupole-Orbitrap

mass spectrometer through an HESI-II probe for electrospray ionisation (Thermo Scientific, Waltham, MA, Stati Uniti). The HESI source parameters were optimised using the automated script in the Q Exactive acquisition software. To assess the thiol precursors, 4  $\mu\text{L}$  of SPE-purified wine extracts was separated on an Acquity UPLC BEH C18 column (50  $\times$  2.1 mm, 1.7  $\mu\text{m}$ , 130  $\text{\AA}$ ) (Waters, Milford, MA, USA) kept at 40°C, and using 0.1 mL/100 mL of FA in Milli-Q-treated water (solvent A) and 0.1 mL/100 mL of FA in acetonitrile (solvent B) as eluting solutions. For the UPLC separation, a linear elution gradient was applied (2–50% of solvent B in 5 min) at a flow rate of 0.5 mL/min. The eluate was analysed by MS using a full scan and data-dependent tandem MS analysis (ddMS<sup>2</sup>) of the nine most intense ions (Top9) from the inclusion list. The resolution was set at 70 000 and 17 500 for full MS and ddMS<sup>2</sup> scan types, respectively. The automatic gain control targets were  $5 \times 10^5$  and  $2 \times 10^5$ , and the maximum ion injection times were 100 and 60 ms for full MS and ddMS<sup>2</sup> scan types, respectively.

The MS data were processed with the Xcalibur 3.0 software (Thermo Scientific, Waltham, MA, Stati Uniti). Peak areas were calculated from extracted ion chromatograms of the thiol precursors with a 3 ppm mass tolerance. For quantification, a deuterated IS was used. Peak area ratios were compared with a five-point standard calibration curve obtained with synthetic thiol precursors.

#### Determination of varietal thiols

**Synthesis of (*R/S*)-3-mercapto-1-hexan-3 $\alpha$ -ol (*d1*-3MH).** Labelled *d1*-3MH (Figure 1b) was synthesised using (*E*)-2-hexenal-3-*d* prepared starting from the commercial 2-hexyn-1-ol as reported by Bennani et al. (2009). Thioacetic acid (0.73 mL, 10.2 mmol) was added to a solution of (*E*)-2-hexenal-3-*d* (1.0 g, 10.1 mmol) in DCM (20 mL) under N<sub>2</sub>. The mixture was stirred at 25°C for 60 h. The solvent was removed under reduced pressure and the pale yellow liquid obtained was used for the next step without further purification. It was dissolved in methanol (45 mL) and cooled on ice. After the addition of a solution of NaBH<sub>4</sub> (0.766 g, 10.1 mmol) in water (30 mL) dropwise to the stirred solution, the mixture was maintained for 1 h on ice. Subsequently, a solution of sodium hydroxide (0.310 g, 10.1 mmol) in water (12 mL) was added dropwise and the mixture was stirred for 2 h on ice. The pH was adjusted to 2 by adding 2 N sulfuric acid and the solution was extracted with DCM (4  $\times$  8 mL). The combined extracts were dried over anhydrous sodium sulfate and the solvent was evaporated under reduced pressure. Purification by column chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>) resulted in the compound *d1*-3MH (Figure 1b) as a colourless oil (45%). The compound structure was confirmed by NMR spectrum.

<sup>1</sup>H NMR ( $\delta$ , CDCl<sub>3</sub>): 0.92 (3H, t,  $J$  = 6.7 Hz, H<sub>6</sub>); 1.35–1.75 (5H, m, SH, H<sub>4</sub>, H<sub>5</sub>); 1.83 (1H, broad s, OH); 1.95 (2H, m, H<sub>2</sub>); 3.81 (2H, m, H<sub>1</sub>) (Figure S3).

#### Sample preparation for varietal thiol analysis.

The varietal thiols were assessed as described by Piano et al. (2015). Each wine sample was spiked with the *d1*-3MH (500  $\mu\text{g/L}$ ; exact mass OPA-derivatised *d1*-3MH [M + H]<sup>+</sup>: 295.159) prior to the sample preparation. Thiols were extracted by liquid/liquid extraction with DCM. After the solvent evaporation under vacuum and N<sub>2</sub> flow, the samples were derivatised with OPA and ethanolamine prior to the chromatographic separation.

**Quantification of varietal thiols with UPLC/ESI-HRMS.** The varietal thiols were determined with UPLC/ESI-HRMS as follows: 3  $\mu\text{L}$  of SPE-purified wine extracts (spiked with deuterated IS) were separated on a Kinetex phenyl-hexyl column (150  $\times$  2.1 mm, 2.6  $\mu\text{m}$ , 100  $\text{\AA}$ ) (Phenomenex, Torrance, CA, USA) kept at 28°C, and using 10 mmol ammonium acetate in MilliQ-treated water (solvent A) and methanol (solvent B) as eluting solvents. For UPLC separation, a linear elution gradient was applied (50–100% of solvent B in 14.5 min) at a flow rate of 0.25 mL/min. The LC eluate was analysed by MS using a ddMS<sup>2</sup> of six of the most intense ions (Top6) from the inclusion list. The MS parameters were as for thiol precursor analysis.

The MS data were processed using the Xcalibur 3.0 software. Peak areas were calculated from extracted ion chromatograms with a 3 ppm mass tolerance of OPA-derivatised thiols. For quantification, deuterated IS was used. Peak area ratios were compared with six-point standard calibration curves obtained using synthetic OPA-derivatised 3MH and 3-MHA.

#### Sensory analysis

A panel of 11 expert judges (three females and eight males) was enrolled for the sensory analysis of the bottled CBC and Grillo wines. The difference between the AEM wine and AFM wine for CBC and Grillo was evaluated with triangle tests. Each judge had a different randomised order of samples for both tests.

#### Statistical analysis

Statistica software (StatSoft, Tulsa, OK, USA) was used for all statistical analysis. The equations of the calibration curves were assessed by linear regression analysis. Differences were evaluated by the *t* test, and significance was set at a value of  $P < 0.05$ .

## Results and discussion

#### Identification of thiol precursors

The thiol precursors were identified in samples of SPE-purified CBC and Grillo grape juice, must and wine by UPLC/ESI-HRMS using synthetic standards as reference compounds, including GSH-3MH, Cys-3MH, GSH-4MMP and GSH-3MHA. Such an analytical approach for assessing these compounds was essential to properly identify them in grape juice, must and wine samples. These compounds were revealed based on their retention time in UPLC, exact mass calculated according to the chemical formula and MS/MS fragmentation spectrum (Table 1). Furthermore, the MS and MS/MS data were compared to those from the literature in order to confirm the assignment. In detail, two chromatographic peaks were assigned to GSH-3MH (not shown) both of them showing the same MS/MS fragmentation pattern. They corresponded to the (*S*)- and (*R*)-diastereomers of the thiol precursor, as reported elsewhere (Capone et al. 2010, Kobayashi et al. 2010, Roland et al. 2010). Moreover, the analysis of a standard mixture of (*R*)- and (*S*)-diastereomers of GSH-3MH resulted in the same chromatographic and MS/MS behaviour. Accurate mass and MS/MS fragmentation spectrum of GSH-4MMP were the same as those reported by other authors (Fedrizzi et al. 2009, Roland et al. 2010, Larcher et al. 2013a). The accurate mass of GSH-3MHA was in accordance with Thibon et al. (2016). The MS/MS spectra of GSH-4MMP and GSH-3MHA were similar in terms of accurate mass of the

**Table 1.** Identification of thiol precursors and *o*-phthaldialdehyde derivative varietal thiols by liquid chromatography coupled to electrospray ionisation-high resolution MS.

Compound	Formula	Retention time (min)	Exact mass [M + H] <sup>+</sup>	MS/MS fragments
Cys-3MH	C <sub>9</sub> H <sub>19</sub> NO <sub>3</sub> S	2.15 ( <i>S</i> diastereomer) 2.17 ( <i>R</i> diastereomer)	222.116	83.086, 101.096, 205.089, 176.074
CysGly-3MH	C <sub>11</sub> H <sub>22</sub> N <sub>2</sub> O <sub>4</sub> S	nd	279.137	nd
GluCys-3MH	C <sub>14</sub> H <sub>26</sub> N <sub>2</sub> O <sub>6</sub> S	nd	351.158	nd
GSH-3MH	C <sub>16</sub> H <sub>29</sub> N <sub>3</sub> O <sub>7</sub> S	2.58 ( <i>S</i> diastereomer) 2.62 ( <i>R</i> diastereomer)	408.180	83.086, 162.022, 262.111, 333.148
GSH-3MHAL	C <sub>16</sub> H <sub>27</sub> N <sub>3</sub> O <sub>7</sub> S	2.60	406.164	162.022, 174.095, 179.048, 259.111, 277.122, 331.132
Cys-4MMP	C <sub>9</sub> H <sub>17</sub> NO <sub>3</sub> S	nd	220.100	nd
CysGly-4MMP	C <sub>11</sub> H <sub>20</sub> N <sub>2</sub> O <sub>4</sub> S	nd	277.122	nd
GluCys-4MMP	C <sub>14</sub> H <sub>24</sub> N <sub>2</sub> O <sub>6</sub> S	nd	349.143	nd
GSH-4MMP	C <sub>16</sub> H <sub>27</sub> N <sub>3</sub> O <sub>7</sub> S	2.20 (nd)	406.164	162.022, 174.095, 179.048, 259.111, 277.122, 331.132
3MH-OPA	C <sub>16</sub> H <sub>23</sub> NO <sub>2</sub> S	7.95	294.151	83.086, 176.052, 194.063
3MHA-OPA	C <sub>18</sub> H <sub>25</sub> NO <sub>3</sub> S	10.15	336.162	83.086, 143.106, 194.063

nd, not detected; Cys-3MH, *S*-3-(hexan-1-ol)-cysteine; CysGly-3MH, *S*-3-(hexan-1-ol)-cysteinylglycine; GluCys-3MH, *S*-3-(hexan-1-ol)-glutamylcysteine; GSH-3MH, *S*-3-(hexan-1-ol)-glutathione; GSH-3MHAL, *S*-3-(hexanal)-glutathione; Cys-4MMP, *S*-3-(4-mercapto-4-methylpentan-2-one)-cysteine; CysGly-4MMP, *S*-3-(4-mercapto-4-methylpentan-2-one)-cysteinylglycine; GluCys-4MMP, *S*-3-(4-mercapto-4-methylpentan-2-one)-glutamylcysteine; GSH-4MMP, *S*-3-(4-mercapto-4-methylpentan-2-one)-glutathione; 3MH-OPA, OPA-derivatized 3-mercaptohexan-1-ol; 3MHA-OPA, OPA-derivatized 3-mercaptohexan-1-ol acetate.

fragments; however, the relative abundance of their fragments showed different patterns (Figure 2). The UPLC/ESI-HRMS of the synthetic GSH-4MMP and GSH-3MHAL permitted discrimination of the two isobaric compounds based on their retention time (Table 1).


#### Determination of thiol precursors and varietal thiols in grapes, musts and wines

The thiol precursors were quantified in juice, must and wine of CBC and Grillo grapes, for the first time, and the varietal thiols were investigated in the corresponding wine samples. Two must clarification treatments based on either exposure of must to air (AEM) or exclusion of air (AFM) were evaluated, because oxygen could increase the concentration of thiol precursors and, consequently, the concentration of varietal thiols in wine (Mattivi et al. 2012). Either an AEM or a CO<sub>2</sub>-must stored in a CO<sub>2</sub>-filled tank were obtained and vinified. Under such conditions, different exposure to oxygen can be obtained (Ribéreau-Gayon et al. 2006, Concejero et al. 2016). The amount of oxygen is difficult to quantify under commercial winemaking conditions due to the uneven oxygen concentration inside the air-exposed bulk, as well as to the oxygen consumption due to both the enzymatic and chemical oxidation phenomena occurring during the lengthy clarification process. The main chemical parameters of musts, obtained from both cultivars subjected to the two clarification treatments, showed negligible differences (Table 2). All alcoholic fermentations ran regularly to dryness though the AFM lasted 2 days longer than that of the AEM (data not shown). At the end of the alcoholic fermentation, negligible difference in the composition of the AEM and AFM wines of the two cultivars was observed, except for the pH and acidity of Grillo wines (Table 2).

As a high concentration of GSH had been reported to give rise to a significant concentration of thiol precursors (Roland et al. 2010), a GSH concentration of at least few tens of mg/L was expected in the grape juice. However, we detected only a trace of GSH (<1 mg/L) in all grape juice samples. Such quantity is among the lowest value reported in the literature, because several authors found a GSH concentration up to 200 mg/L in grape juice (Cheynier et al. 1989, Okuda and Yokotsuka 1999, Jančes et al. 2010, Fracassetti and Tirelli 2015). It is difficult to ascribe the low

GSH concentration to sample storage issues, because GRP concentration values lower than 4 mg/L (i.e. less than 2 mg/L of GSH) were found in the grape juices. A comparable concentration of GSH was detected in CBC and Grillo musts during fermentation, being in the range 8–9 mg/L in the final wines (Table 2). Similar data are reported in the literature (Cassol and Adams 1995, Du Toit et al. 2007, Kritzing et al. 2013).

The 4MMP precursors, as well as GluCys-3MH and CysGly-3MH, were not detected in the grape juice, must and wine samples of either cultivar. Meanwhile, Cys-3MH, GSH-3MH and GSH-3MHAL were identified in grape juice, must and wine samples of both cultivars (Table 3). The concentration of Cys-3MH and GSH-3MH in Grillo grape juice changed little during the monitored 3 weeks of ripening. Conversely, the concentration of the 3MH precursors in the CBC grape juice halved during the last 2 weeks of ripening, and their values were roughly half of the corresponding values found in Grillo grape juices. Ripening behaviour comparable to that of CBC grape was reported by Peyrot des Gachons et al. (2005) and Roland et al. (2010) for Sauvignon Blanc and Mellon B grape cultivars. The concentration of Cys-3MH and GSH-3MH matched those reported in previous research concerning Sauvignon Blanc and Pinot



**Figure 2.** High-resolution-MS/MS spectra of the synthetic standards: (a) (*S*)-3-(4-mercapto-4-methylpentan-2-one)-glutathione (GSH-4MMP) and (b) (*S*)-3-(hexan-1-ol)-glutathione (GSH-3MHAL).

**Table 2.** Composition of Catarratto Bianco Comune and Grillo musts and wines.

Cultivar	Clarification treatment	Sugars (g/L)	Total SO <sub>2</sub> (mg/L)	pH	TA (g/L)	Readily assimilable nitrogen (mg/L)			
Must									
Catarratto Bianco Comune	AEM	226 ± 2	42 ± 3	3.45 ± 0.03	6.15 ± 0.05	206 ± 8			
	AFM	226 ± 2	38 ± 2	3.45 ± 0.03	6.10 ± 0.05	210 ± 8			
Grillo	AEM	222 ± 3	36 ± 2	3.25 ± 0.03	6.05 ± 0.05	232 ± 9			
	AFM	222 ± 3	36 ± 2	3.25 ± 0.03	6.05 ± 0.05	232 ± 9			
	Clarification treatment	Sugars (g/L)	Free SO <sub>2</sub> (mg/L)	Total SO <sub>2</sub> (mg/L)	pH	TA (g/L)	Volatile acidity (g/L acetic acid)	Ethanol (% v/v)	Glutathione (mg/L)
Wine									
Catarratto Bianco Comune	AEM	1.1 ± 0.1	56 ± 4	80 ± 6	3.33 ± 0.01	6.35 ± 0.15	0.41 ± 0.02	13.30 ± 0.02	9.38 ± 0.47
	AFM	1.2 ± 0.3	50 ± 2	86 ± 4	3.51 ± 0.01	6.20 ± 0.05	0.42 ± 0.04	13.66 ± 0.06	7.85 ± 0.39
Grillo	AEM	2.9 ± 0.2	53 ± 3	83 ± 3	3.13 ± 0.01	7.28 ± 0.12	0.18 ± 0.01	12.81 ± 0.03	8.04 ± 0.40
	AFM	2.3 ± 0.6	35 ± 2	74 ± 3	3.14 ± 0.03	6.50 ± 0.15	0.13 ± 0.01	12.58 ± 0.33	8.24 ± 0.41

AEM, air-exposed must during clarification; AFM, air-free must during clarification.

**Table 3.** Concentration of thiol precursors in Catarratto Bianco Comune and Grillo grape juices during ripening in vintage 2014.

Grape cultivar	Sampling date	Concentration (µg/L)		
		Cys-3MH	GSH-3MH	GSH-3MHAl
Catarratto Bianco Comune	12.09	16.0 ± 4.0	445 ± 35	635 ± 51
	20.09	50.8 ± 4.1	475 ± 37	2679 ± 214
	29.09	15.5 ± 3.7	199 ± 16	10 672 ± 854
	08.10†	19.0 ± 4.8	216 ± 17	8096 ± 667
Grillo	21.08	25.5 ± 6.6	436 ± 34	7710 ± 632
	03.09	30.8 ± 2.4	479 ± 37	7294 ± 584
	10.09†	27.3 ± 2.2	460 ± 35	2644 ± 217

†Harvest date. Cys-3MH, *S*-3-(hexan-1-ol)-cysteine; GSH-3MH, *S*-3-(hexan-1-ol)-glutathione; GSH-3MHAl, *S*-3-(hexanal)-glutathione.


Gris grapes (Capone et al. 2010). The concentration of GSH-3MHAl decreased during the ripening of Grillo grape by 66%, while it greatly increased (more than 12-fold) in the CBC grape. Overall, when the two cultivars were compared at harvest, Grillo contained a higher concentration of both GSH-3MH and Cys-3MH, whereas CBC was higher in GSH-3MHAl (Table 3).

The concentration of thiol precursors dramatically decreased in all musts compared to that of the corresponding grape juices. About 50% of GSH-3MH and up to 98% of GSH-3MHAl were lost following the grape pressing (Figure 3a,b, Table 3). To the best of our knowledge, such a substantial loss of precursors under commercial pressing conditions is described for the first time. Capone et al. (2011) proved CysGly-3MH to be an intermediate of GSH-3MH degradation to Cys-3MH; however, this intermediate was not detected in the CBC and Grillo musts. Therefore, the loss of the glutathionyl- precursors cannot be ascribed to the hydrolysis of the GSH moiety of GSH-3MH. Moreover, the Cys-3MH concentration found in both the ripe grape juices matched the concentration found in the corresponding musts (Figure 3a,b, Table 3). Therefore, the degradation of GSH-3MH leading either to the release of its amino acid units or the oxidative phenomena involving the cysteinyl-3MH moiety is unlikely to have had a major role in the loss of the GSH-3MH, because thiol precursors are stable in oxic conditions due to their thioether bond (Roland et al. 2010). A similar behaviour was reported for

the loss of GSH following grape pressing under several commercial winemaking conditions (Fracassetti and Tirelli 2015). Therefore, factors affecting the GSH portion of the precursors, other than oxidation or proteolysis, could induce the loss of the thiol precursors.

The substantial loss of GSH-containing precursors that occurred following extraction and clarification of the must did not change the thiol precursor pattern observed in CBC and Grillo grape juices, because the higher concentration of GSH-3MH and Cys-3MH and lower concentration of GSH-3MHAl were still detected in Grillo must compared to that of CBC must, independent of the clarification treatment applied (Figure 3). Negligible differences were also observed between differently clarified Grillo musts, whereas AFM of CBC contained higher concentration of GSH-3MH and GSH-3MHAl compared to that of the corresponding AEM; however, only the concentration of GSH-3MHAl was statistically different. It is difficult to ascribe the observed high concentration of GSH-3MHAl to the applied anoxic conditions. The reducing conditions could hinder the lipoxygenase activity needed to produce additional 2-hexenal. Moreover, the negligible GSH concentration did not allow the formation of the precursor. More likely, the higher precursor concentration in the grape was retained in the must. The high concentration of GSH-3MHAl in the CBC AFM was expected to increase the formation of a higher concentration of GSH-3MH and then 3MH in wine due to the yeast activity. However, no significant increase in the






**Figure 3.** Effect of must treatment on the concentration of thiol precursors in must (■, ■) and wine (■, ■) produced from (a) Catarratto Bianco Comune and (b) Grillo grapes. The must was either air-exposed during clarification (■, ■) or air-free during clarification (■, ■). Different letters mean significant difference ( $P < 0.05$ ) between the two clarification treatments.

concentration of either GSH-3MH or 3MH in wine was detected, despite the GSH-3MHAl lost following vinification (Figures 3a,4). In all musts, the Cys-3MH concentration was lower than that of GSH-3MH and GSH-3MHAl, which is consistent with other published data (Capone et al. 2010, Thibon et al. 2016).

A high concentration of the thiol precursors still occurred in the wines even if at lower values in comparison to that of the corresponding musts (Figure 3). The alcoholic fermentation, however, had little effect on the concentration of GSH-3MH in CBC wine, likely due to the reduction of GSH-3MHAl by the yeast. An amount of GSH-3MH exceeding 50% was lost in Grillo musts following vinification. In spite of the large number of differences involved in the investigated vinification processes (grape cultivar, harvest date and processing conditions, fermenting yeast strains, thiol precursor concentration in the musts), the CBC and Grillo wines showed a similar concentration of each precursor; in addition a similar concentration of GSH-3MH and GSH-3MHAl was observed in each wine (Figure 3a,b). This behaviour is likely due to physical-chemical phenomena affecting solubility and/or adsorption on the lees rather than to microbial or chemical modifications, because the release of aromatic free thiols negligibly affected the loss of precursors. Once again, this suggests that thioether-bonded GSH derivatives can be lost in oenological matrices, although, explicit analytical data concerning their fate are needed to substantiate such a hypothesis.

In agreement with the previous research, no clear correlation between precursor concentration in the must and free thiol concentration in the wine was observed (Murat et al. 2001, Peyrot des Gachons et al. 2002, Dubourdieu et al. 2006, Subileau et al. 2008, Grant-Preece et al. 2010, Kobayashi et al. 2010, Roland et al. 2010, Winter et al. 2011, Concejero et al. 2016). Moreover, no significant difference was detected between AEM and AFM samples, though the AEM wines showed a slightly higher concentration of free thiols independent of the grape cultivar.



**Figure 4.** Effect of must treatment on the concentration of volatile thiols in Catarratto Bianco Comune (■, ■) and Grillo (■, ■) wines produced from air-exposed must during clarification (■, ■), and from air-free must during clarification (■, ■). Different letters mean significant difference ( $P < 0.05$ ) between the two clarification treatments.

The concentration of 3MH and 3MHA in CBC wine samples was consistently higher than that in Grillo wine (Figure 4) in spite of the lower concentration of GSH-3MH and Cys-3MH occurring in the former. Notably, CBC AEM and AFM contained more GSH-3MHAl (21 and 81%, respectively), which might have yielded the release of varietal thiols (Thibon et al. 2016). However, different yeast strains were employed, and their major role in yielding different thiol concentration cannot be excluded (Murat et al. 2001). The concentration of 3MH and 3MHA in Sauvignon Blanc wines from different countries has been reported to be in the range 688–18 681 and 10–2507 ng/L, respectively (Lund et al. 2009, Benkwitz et al. 2012, Piano et al. 2015). The concentration of varietal thiols found in CBC and Grillo wines was in the above-mentioned ranges and greatly exceeded their perception threshold. The calculated olfactory index of 3MH (expressed as a concentration detected to the perception threshold ratio) was 14.4 and 18.8 for AEM and AFM CBC wines, respectively, and 7.6 and 6.3 for AEM and AFM Grillo wines, respectively. The index of 3MHA was 17.2 and 15.5 for AEM and AFM CBC wines, respectively, and 3.8 and 1.6 for AEM and AFM Grillo wines, respectively. Such values highlight the major role of the varietal thiols in affecting the aromatic character of the wines made from CBC and Grillo.

Based on our data, a theoretical conversion yield of precursors into free thiols in the range 1–68% can be calculated (Table 4). However, the thiol formation arising from the Michael addition involving  $H_2S$  and 2-hexenal during the fermentation makes such yield values unreliable as it occurred in the AEM CBC wine where the calculated 68% yield is hardly plausible. The conversion yield of Cys-3MH into 3MH and 3MHA has been reported to be in the range 0.1–12%, while the conversion yield of GSH-3MH to 3MH and 3MHA is known to be less than 5% (Coetzee and Du Toit 2012, Concejero et al. 2016). Such a difference can be mainly ascribed to the yeast strains, as well as to their different kinetics of  $H_2S$  production (Harsch et al. 2013).

Finally, the sensory characteristics of AEM and AFM wines were compared with a triangle sensory test in order to confirm the analytical data for both CBC and Grillo wines. No significant difference was found between the CBC wines because only 5 out of 11 judges provided the correct response. The same result was obtained when the

**Table 4.** Theoretical conversion yield of thiol precursors into free thiols following vinification of Catarratto Bianco Comune and Grillo musts.

Wine	Clarification treatment	Loss of thiol precursors <sup>†</sup> in vinification (nmol/L)	Free thiols <sup>‡</sup> in wine (nmol/L)	Conversion yield (%)
Catarratto Bianco Comune	Air	10.07	6.82	67.73
	CO <sub>2</sub>	53.89	6.52	12.10
Grillo	Air	340.03	3.50	1.02
	CO <sub>2</sub>	194.54	2.87	1.48

<sup>†</sup>S-3-(Hexan-1-ol)-cysteine (Cys-3MH) + S-3-(hexan-1-ol)-glutathione (GSH-3MH). <sup>‡</sup>3-Mercaptohexan-1-ol (3MH) + 3-mercaptohexylacetate (3MHA).

AEM and AFM Grillo wines were tested. These findings suggest that the winemaking conditions adopted for these two Sicilian grape cultivars did not quantitatively affect the varietal thiols, and they even underline the main role of varietal thiols in the typical flavour of CBC and Grillo wines.

## Conclusions

We have, for the first time, clearly distinguished the isobaric GSH-4MMP and GSH-3MHA from each other, not only by their retention time in UPLC, but even by means of their HR-MS/MS spectra, and have demonstrated by analytical and sensory approaches the essential role of 3MH and 3MHA in CBC and Grillo grape flavours. As none of the 4MMP precursors were detected in both must and wine, a major role of this flavour compound in CBC and Grillo wines can be ruled out. The applied vinification conditions (air-free and air-exposed clarification) had little effect on the final aromatic thiol concentration in wine as was observed by Larcher et al. (2013b). The substantial loss of glutathionyl precursors that occurred during juice extraction may detrimentally affect the sensory properties of Grillo and CBC wine and should be better investigated.

## Acknowledgements

The authors thank the winery Conti Tasca d'Almerita, Valledlunga Pratameno, Italy for providing the grape and wine samples and Ms Kaitlyn Sulenski for her technical support. Special thanks to Dr Federico Guidugli (Thermo Fisher Scientific, Rodano, Italy) for his fruitful suggestions on UPLC/ESI-HRMS analysis.

## References

- Allen, T., Herbst-Johnstone, M., Girault, M., Butler, P., Logan, G., Jouanneau, S., Nicolau, L. and Kilmartin, P.A. (2011) Influence of grape-harvesting steps on varietal thiol aromas in Sauvignon blanc wines. *Journal of Agricultural and Food Chemistry* **59**, 10641–10650.
- Benkwitz, F., Tominaga, T., Kilmartin, P.A., Lund, C.M., Wohlers, M. and Nicolau, L. (2012) Identifying the chemical composition related to the distinct aroma characteristics of New Zealand Sauvignon blanc wines. *American Journal of Enology and Viticulture* **63**, 62–72.
- Bennani, Y.L., Maltais, F., Jung, Y.C., Chen, M., Tanoury, J., Perni, R.B., Mani, N., Laitinen, L., Huang, H., Liao, S., Gao, H., Tsao, H., Block, E., Ma, C., Shawgo, R.S., Town, C., Brummel, C. L., Howe, D., Pazhanisamy, S., Raybuck, S. and Namchuk, M. (2009) In vitro and in vivo isotope effects with hepatitis C protease inhibitors: enhanced plasma exposure of deuterated telaprevir versus telaprevir in rats. *Journal of Medicinal Chemistry* **52**, 7993–8001.
- Bouchilloux, P., Darriet, P., Henry, R., Lavigne-Cruège, V. and Dubourdieu, D. (1998) Identification of volatile and powerful odorous thiols in Bordeaux red wine varieties. *Journal of Agricultural and Food Chemistry* **46**, 3095–3099.
- Calò, A., Scienza, A. and Costacurta, A. (2006) Vitigni d'Italia (Calderini Edagricole: Milan, Italy).
- Capone, D.L., Sefton, M.A., Hayasaka, Y. and Jeffery, D.W. (2010) Analysis of precursors to wine odorant 3-mercaptohexan-1-ol using HPLC-MS/MS: resolution and quantification of diastereoisomers of 3-S-cysteinylhexan-1-ol and 3-S-glutathionylhexan-1-ol. *Journal of Agricultural and Food Chemistry* **58**, 1390–1395.
- Capone, D., Pardon, K.H., Cordente, A.G. and Jeffery, D.W. (2011) Identification and quantification of 3-S-cysteinylglycinehexan-1-ol (cysgly-3-MH) in Sauvignon blanc grape juice by HPLC-MS/MS. *Journal of Agricultural and Food Chemistry* **59**, 11204–11210.
- Carimi, F., Mercati, F., Abbate, L. and Sunseri, F. (2010) Microsatellite analyses for evaluation of genetic diversity among Sicilian grapevine cultivars. *Genetic Resources and Crop Evolution* **57**, 703–719.
- Cassol, T. and Adams, D.O. (1995) Detection of glutathione in white wines using an enzymatic analytical method. *American Journal of Enology and Viticulture* **46**, 410.
- Cheynier, V., Souquet, J.M. and Moutounet, M. (1989) Glutathione content and glutathione to hydroxycinnamic acid ratio in *Vitis vinifera* grapes and musts. *American Journal of Enology and Viticulture* **40**, 320–324.
- Coetzee, C. and Du Toit, W.J. (2012) A comprehensive review on Sauvignon blanc aroma with a focus on certain positive volatile thiols. *Food Research International* **45**, 287–298.
- Concejero, B., Hernandez-Orte, P., Astrain, J., Lacau, B., Baron, C. and Ferreira, V. (2016) Evolution of polyfunctional mercaptans and their precursors during Merlot alcoholic fermentation. *Food Science and Technology* **65**, 770–776.
- Corona, O. (2010) Wine-making with protection of must against oxidation in a warm, semi-arid terroir. *South African Journal of Enology and Viticulture* **31**, 58–63.
- Darriet, P., Tominaga, T., Lavigne, V., Boidron, J.N. and Dubourdieu, D. (1995) Identification of a powerful aromatic of *Vitis vinifera* L. var. sauvignon wines: 4-mercapto-4-methylpentan-2-one. *Flavour and Fragrance Journal* **10**, 385–392.
- De Lorenzis, G., Las Casas, G., Brancadoro, L. and Scienza, A. (2014) Genotyping of Sicilian grapevine germplasm resources (*V. vinifera* L.) and their relationships with Sangiovese. *Scientia Horticulturae* **169**, 189–198.
- Du Toit, W.J., Lisjak, K., Stander, M. and Prevo, D. (2007) Using LC-MSMS to assess glutathione levels in South African white grape juices and wines made with different levels of oxygen. *Journal of Agricultural and Food Chemistry* **55**, 2765–2769.
- Dubourdieu, D., Tominaga, T., Masneuf, L., des, Peyrot, Gachons, C. and Murat, M.L. (2006) The role of yeasts in grape flavor development during fermentation: the example of Sauvignon blanc. *American Journal of Enology and Viticulture* **57**, 81–88.
- Fedrizzi, B., Pardon, K.H., Sefton, M.A., Eley, G.M. and Jeffery, D. W. (2009) First identification of 4-S-glutathionyl-4-methylpentan-2-one, a potential precursor of 4-mercapto-4-methylpentan-2-one, in Sauvignon blanc juice. *Journal of Agricultural and Food Chemistry* **57**, 991–995.
- Fracassetti, D. and Tirelli, A. (2015) Monitoring of glutathione concentration during winemaking by a reliable high-performance liquid chromatography analytical method. *Australian Journal of Grape and Wine Research* **21**, 389–395.
- Grant-Preece, P.A., Pardon, K.H., Capone, D.L., Cordente, A.G., Sefton, M.A., Jeffery, D.W., Sefton, M.A. and Eley, G.M. (2010) Synthesis of wine thiol conjugates and labeled analogues: fermentation of the glutathione conjugate of 3-mercaptohexan-1-ol yields the corresponding cysteine conjugate and free thiol. *Journal of Agricultural and Food Chemistry* **58**, 1383–1389.
- Harsch, M.J., Benkwitz, F., Frost, A., Colonna-Ceccaldi, B., Gardner, R.C. and Salmon, J.M. (2013) New precursor of 3-mercaptohexan-1-ol in grape juice: thiol-forming potential and kinetics during early stages of must fermentation. *Journal of Agricultural and Food Chemistry* **61**, 3703–3713.



- Hatanaka, A., Kajiwaru, T. and Matsui, K. (1995) The biogenesis of green odour by green leaves and its physiological functions – past, present and future. *Zeitschrift für Naturforschung* **50c**, 467–472.
- Janés, L., Lisjak, K. and Vanzo, A. (2010) Determination of glutathione content in grape juice and wine by high-performance liquid chromatography with fluorescence detection. *Analytica Chimica Acta* **674**, 239–242.
- Kobayashi, H., Takase, H., Kaneko, K., Tanzawa, F., Takata, R., Suzuki, S. and Konno, T. (2010) Analysis of S-3-(hexan-1-ol)-glutathione and S-3-(hexan-1-ol)-l-cysteine in *Vitis vinifera* L. cv. Koshu for aromatic wines. *American Journal of Enology and Viticulture* **61**, 176–185.
- Kritzinger, E.C., Bauer, F.F. and Du Toit, W.J. (2013) Role of glutathione in winemaking: a review. *Journal of Agricultural and Food Chemistry* **61**, 269–277.
- Larcher, R., Tonidandel, L., Nicolini, G. and Fedrizzi, B. (2013a) First evidence of the presence of S-cysteinylated and S-glutathionylated precursors in tannins. *Food Chemistry* **141**, 1196–1202.
- Larcher, R., Nicolini, G., Tonidandel, L., Román Villegas, T., Malacarne, M. and Fedrizzi, B. (2013b) Influence of oxygen availability during skin-contact maceration on the formation of precursors of 3-mercaptohexan-1-ol in Müller-Thurgau and Sauvignon Blanc grapes. *Australian Journal of Grape and Wine Research* **19**, 342–348.
- Lund, C.M., Thompson, M.K., Benkwitz, F., Wohler, M.W., Triggs, C.M., Gardner, R., Heymann, H. and Nicolau, L. (2009) New Zealand Sauvignon blanc distinct flavor characteristics, sensory, chemical, and consumer aspects. *American Journal of Enology and Viticulture* **60**, 1–12.
- Mattivi, F., Fedrizzi, B., Zenato, A., Tiefenthaler, P., Tempesta, S., Perenzoni, D., Cantarella, P., Simeoni, F. and Vrhovsek, U. (2012) Development of reliable analytical tools for evaluating the influence of reductive wine making on the quality of Lugana wines. *Analytica Chimica Acta* **732**, 194–202.
- Murat, M.L., Masneuf, I., Darriet, P., Lavigne, V., Tominaga, T. and Dubourdieu, D. (2001) Effect of *Saccharomyces cerevisiae* strain on the liberation of volatile thiols in Sauvignon blanc. *American Journal of Enology and Viticulture* **52**, 136–139.
- Okuda, T. and Yokotsuka, K. (1999) Levels of glutathione and activities of related enzymes during ripening of Koshu and Cabernet Sauvignon grapes and during wine making. *American Journal of Enology and Viticulture* **50**, 264–270.
- Pardon, K.H., Graney, S.D., Capone, D.L., Swiegers, J.H., Sefton, M. A. and Elsey, G.M. (2008) Synthesis of the individual diastereomers of the cysteine conjugate of 3-mercaptohexanol (3-MH). *Journal of Agricultural and Food Chemistry* **56**, 3758–3763.
- Peña-Gallego, A., Hernández-Orte, P., Cacho, J. and Ferreira, V. (2012) S-Cysteinylated and S-glutathionylated thiol precursors in grapes. A review. *Food Chemistry* **131**, 1–13.
- Peyrot des Gachons, C., Tominaga, T. and Dubourdieu, D. (2002) Sulfur aroma precursor present in S-glutathione conjugate form: identification of S-3-(hexan-1-ol)-glutathione in must from *Vitis vinifera* L. cv. Sauvignon blanc. *Journal of Agricultural and Food Chemistry* **50**, 4076–4079.
- Peyrot des Gachons, C., Van Leeuwen, C., Tominaga, T., Sowyer, J. P., Gaudillère, J.P. and Dubourdieu, D. (2005) Influence of water and nitrogen deficit on fruit ripening and aroma potential of *Vitis vinifera* L. cv. Sauvignon blanc in field conditions. *Journal of the Science of Food and Agriculture* **85**, 73–85.
- Piano, F., Petrozziello, M., Vaudano, E., Bonello, F., Ferreira, V., Zapata, J. and Hernández-Orte, P. (2014) Aroma compounds and sensory characteristics of Arneis Terre Alfieri DOC wines: the concentration of polyfunctional thiols and their evolution in relation to different ageing conditions. *European Food Research and Technology* **239**, 267–277.
- Piano, F., Fracassetti, D., Buica, A., Stander, M., du Toit, W.J., Borsa, D. and Tirelli, A. (2015) Development of a novel liquid/liquid extraction and UPLC-MS/MS method for the assessment of thiols in South African Sauvignon blanc wines. *Australian Journal of Grape and Wine Research* **21**, 40–48.
- Pinu, F.R., Jouanneau, S., Nicolau, L., Gardner, R.C. and Villas-Boas, S.G. (2012) Concentrations of the volatile thiol 3-mercaptohexanol in Sauvignon blanc wines: no correlation with juice precursors. *American Journal of Enology and Viticulture* **63**, 407–412.
- Ribéreau-Gayon, P., Glories, Y., Maujean, A. and Dubourdieu, D. *Handbook of enology*. Vol. 1. (2006) The microbiology of wine and vinification. 2d ed. (John Wiley: Chichester, England).
- Robinson, J., Harding, J. and Vouillamoz, J. (2012) *Wine grapes: a complete guide to 1,368 vine varieties, including their origins and flavours* (Harper Collins: New York, NY, USA).
- Roland, A., Vialaret, J., Razungles, A., Rigou, P. and Schneider, R. (2010) Evolution of S-cysteinylated and S-glutathionylated thiol precursors during oxidation of Melon B. and Sauvignon blanc musts. *Journal of Agricultural and Food Chemistry* **58**, 4406–4413.
- Schneider, R., Charrier, F., Razungles, A. and Baumesb, R. (2006) Evidence for an alternative biogenetic pathway leading to 3-mercaptohexanol and 4-mercapto-4-methylpentan-2-one in wines. *Analytica Chimica Acta* **563**, 58–64.
- Subileau, M., Schneider, R., Salmon, J. and Degryse, E. (2008) New insights on 3-mercaptohexanol (3MH) biogenesis in Sauvignon blanc wines: Cys-3MH and (E)-hexen-2-al are not the major precursors. *Journal of Agricultural and Food Chemistry* **56**, 9230–9235.
- Thibon, C., Böcker, C., Shinkaruk, S., Moine, V., Darriet, P. and Dubourdieu, D. (2016) Identification of S-3-(hexanal)-glutathione and its bisulfite adduct in grape juice from *Vitis vinifera* L. cv. Sauvignon blanc as new potential precursors of 3SH. *Food Chemistry* **199**, 711–719.
- Tominaga, T., Furrer, A., Henry, R. and Dubourdieu, D. (1998a) Identification of new volatile thiols in the aroma of *Vitis vinifera* L. var. Sauvignon wines. *Flavour and Fragrance Journal* **13**, 159–162.
- Tominaga, T., Murat, M.L. and Dubourdieu, D. (1998b) Development of a method for analyzing the volatile thiols involved in the characteristic aroma of wines made from *Vitis vinifera* L. cv. Sauvignon blanc. *Journal of Agricultural and Food Chemistry* **46**, 1044–1048.
- Tominaga, T., Baltenweck-Guyot, R., des Peyrot, Gachons, C. and Dubourdieu, D. (2000) Contribution of volatile thiols to the aromas of white wines made from several *Vitis vinifera* grape varieties. *American Journal of Enology and Viticulture* **51**, 178–181.
- Winter, G., Van Der Westhuizen, T., Higgins, V.J., Curtin, C. and Ugliano, M. (2011) Contribution of cysteine and glutathione conjugates to the formation of the volatile thiols 3-mercaptohexan-1-ol (3MH) and 3-mercaptohexyl acetate (3MHA) during fermentation by *Saccharomyces cerevisiae*. *Australian Journal of Grape and Wine Research* **17**, 285–290.

Manuscript received: 23 March 2017

Revised manuscript received: 7 June 2017

Accepted: 8 June 2017

### Supporting information

Additional supporting information may be found in the online version of this article at the publisher's website: <http://onlinelibrary.wiley.com/doi/10.1111/ajgw.12311/abstract>.

**Figure S1.** NMR spectra of  $\gamma$ -L-glutamyl-S-[(1R/S)-1-(2-hydroxyethyl)butyl]-L-cysteinyl-glycine (GSH-3MH).

**Figure S2.** NMR spectra of  $\gamma$ -L-glutamyl-S-[(1R/S)-1-(2-hydroxyethyl)butyl-1d]-L-cysteinyl-glycine (d1-GSH-3MH).

**Figure S3.** NMR spectra of (R/S)-3-mercapto-1-hexan-3d-ol (d1-3MH).