

PAPER • OPEN ACCESS

Measurement of Solar pp-neutrino flux with Borexino: results and implications

To cite this article: O Yu Smirnov *et al* 2016 *J. Phys.: Conf. Ser.* **675** 012027

View the [article online](#) for updates and enhancements.

Related content

- [Going low: measurement of Solar pp-neutrino flux with liquid scintillator detector](#)
O Yu Smirnov and Borexino collaboration
- [Geochemical Determination of the Solar pp-Neutrino Flux with LOREX: A Progress Report](#)
M K Pavievi, W F Henning, F Bosch et al.
- [Physics news on the Internet \(based on electronic preprints\)](#)
Yu N Eroshenko

Measurement of Solar pp-neutrino flux with Borexino: results and implications

O Yu Smirnov^{1,29}, M Agostini², S Appel², G Bellini³, J Benziger⁴, D Bick⁵, G Bonfini⁶, D Bravo⁷, B Caccianiga³, F Calaprice⁸, A Caminata⁹, P Cavalcante⁶, A Chepurinov¹⁰, D D'Angelo³, S Davini¹¹, A Derbin¹², L Di Noto⁹, I Drachnev¹¹, A Etenko¹³, K Fomenko¹, D Franco¹⁴, F Gabriele⁶, C Galbiati⁸, C Ghiano⁹, M Giammarchi³, M Goeger-Neff², A Goretti⁸, M Gromov¹⁰, C Hagner⁵, E Hungerford¹⁵, Aldo Ianni⁶, Andrea Ianni⁸, K Jedrzejczak¹⁷, M Kaiser⁵, V Kobychew¹⁸, D Korablev¹, G Korga⁶, D Kryn¹⁴, M Laubenstein⁶, B Lehnert¹⁹, E Litvinovich^{13,20}, F Lombardi⁶, P Lombardi³, L Ludhova³, G Lukyanchenko^{13,20}, I Machulin^{13,20}, S Manecki⁷, W Maneschg²², S Marcocci¹¹, E Meroni³, M Meyer⁵, L Miramonti³, M Misiaszek^{17,6}, M Montuschi²³, P Mosteiro⁸, V Muratova¹², B Neumair², L Oberauer², M Obolensky¹⁴, F Ortica²⁴, M Pallavicini⁹, L Papp², L Perasso⁹, A Pocar²⁶, G Ranucci³, A Razeto⁶, A Re³, A Romani²⁴, R Roncin^{6,14}, N Rossi⁶, S Schönert², D Semenov¹², H Simgen²², M Skorokhvatov^{13,20}, A Sotnikov¹, S Sukhotin¹³, Y Suvorov^{27,13}, R Tartaglia⁶, G Testera⁹, J Thurn¹⁹, M Toropova¹³, E Unzhakov¹², A Vishneva¹, R B Vogelaar⁷, F von Feilitzsch², H Wang²⁷, S Weinz²⁸, J Winter²⁸, M Wojcik¹⁷, M Wurm²⁸, Z Yokley⁷, O Zaimidoroga¹, S Zavatarelli⁹, K Zuber¹⁹ and G Zuzel¹⁷ (Borexino Collaboration)

¹ Joint Institute for Nuclear Research, 141980 Dubna, Russia

² Physik-Department and Excellence Cluster Universe, Technische Universität München, 85748 Garching, Germany

³ Dipartimento di Fisica, Università degli Studi e INFN, 20133 Milano, Italy

⁴ Chemical Engineering Department, Princeton University, Princeton, NJ 08544, USA

⁵ Institut für Experimentalphysik, Universität, 22761 Hamburg, Germany

⁶ INFN Laboratori Nazionali del Gran Sasso, 67010 Assergi (AQ), Italy

⁷ Physics Department, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

⁸ Physics Department, Princeton University, Princeton, NJ 08544, USA

⁹ Dipartimento di Fisica, Università degli Studi e INFN, Genova 16146, Italy

¹⁰ Lomonosov Moscow State University Skobel'syn Institute of Nuclear Physics, 119234 Moscow, Russia

¹¹ Gran Sasso Science Institute (INFN), 67100 L'Aquila, Italy

¹² St. Petersburg Nuclear Physics Institute NRC Kurchatov Institute, 188350 Gatchina, Russia

¹³ NRC Kurchatov Institute, 123182 Moscow, Russia

¹⁴ AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, 75205 Paris Cedex 13, France

²⁹ Presenter. To whom any correspondence should be addressed.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

- ¹⁵ Department of Physics, University of Houston, Houston, TX 77204, USA
¹⁶ Institute for Theoretical and Experimental Physics, 117218 Moscow, Russia
¹⁷ M. Smoluchowski Institute of Physics, Jagiellonian University, 30059 Krakow, Poland
¹⁸ Kiev Institute for Nuclear Research, 06380 Kiev, Ukraine
¹⁹ Department of Physics, Technische Universität Dresden, 01062 Dresden, Germany
²⁰ National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Kashirskoe highway 31, Moscow 115409, Russia
²¹ Kepler Center for Astro and Particle Physics, Universität Tübingen, 72076 Tübingen, Germany
²² Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany
²³ Dipartimento di Fisica e Scienze della Terra Università degli Studi di Ferrara e INFN, Via Saragat 1-44122, Ferrara, Italy
²⁴ Dipartimento di Chimica, Università e INFN, 06123 Perugia, Italy
²⁵ Physics Department, Queen's University, Kingston ON K7L 3N6, Canada
²⁶ Amherst Center for Fundamental Interactions and Physics Department, University of Massachusetts, Amherst, MA 01003, USA
²⁷ Physics and Astronomy Department, University of California Los Angeles (UCLA), Los Angeles, California 90095, USA
²⁸ Institute of Physics and Excellence Cluster PRISMA, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

E-mail: osmirnov@jinr.ru

Abstract. Measurement of the Solar pp-neutrino flux completed the measurement of Solar neutrino fluxes from the pp-chain of reactions in Borexino experiment. The result is in agreement with the prediction of the Standard Solar Model and the MSW/LMA oscillation scenario. A comparison of the total neutrino flux from the Sun with Solar luminosity in photons provides a test of the stability of the Sun on the 10^5 years time scale, and sets a strong limit on the power production by the unknown energy sources in the Sun.

1. Introduction

The solar photon luminosity $L_{\odot} = 3.846 \times 10^{26}$ W is measured for a precision of 0.4% [1, 2]. The energy lost by neutrinos adds $L_{\nu} = 0.023 \cdot L_{\odot}$ to this value [3]. The solar luminosity constraint on the solar neutrino fluxes can be written as [4]:

$$\frac{L_{\odot}}{4\pi(1a.u.)^2} = \sum \alpha_i \Phi_i \quad (1)$$

where 1 a.u. is the average Earth-Sun distance, the coefficient α_i is the amount of energy provided to the star by nuclear fusion reactions associated with each of the important solar neutrino fluxes, Φ_i . The numerical values of the α 's are determined to an accuracy of 10^{-4} and better.

The estimated uncertainty in the luminosity of the Sun corresponds to less than 3% uncertainty in total solar neutrino flux.

Because of the relation (1) between the solar photon and neutrino luminosity, the measurement of the total neutrino luminosity will provide a test of the stability of the Sun at the time scale of 40000 years [5], the time needed for the radiation born at the center of the Sun to arrive to its surface. Finding a disagreement between L_{\odot} and L_{ν} would have significant long term environmental implications, and in the case of an agreement between the two measurements it would be possible to limit the unknown sources of the solar energy, different from the known thermonuclear fusion of light elements in the pp-chain and CNO-cycle.

The main neutrino sources in the Sun are the pp- and ${}^7\text{Be}$ reactions, providing roughly 91 and 7% of the total neutrino flux respectively. Borexino already measured the ${}^7\text{Be}$ neutrino

flux with 5% precision [6], but till recent time the pp-neutrino flux was derived in a differential measurement using solar detectors data.

A number of projects aiming to perform pp-neutrino detection have been put forward in past two decades, but with all the time passed since the proposals, none of them started the operation facing the technical problems with realization. A possibility to use ultrapure liquid organic scintillator as a low energy solar neutrino detector for a first time was discussed in [7, 8]. The authors come to the conclusion that a liquid scintillator detector with an active volume of 10 tons is an appropriate tool to register the solar pp-neutrino if operated at the target level of radiopurity for Borexino and good energy resolution (5% at 200 keV) is achieved.

2. Solar pp-neutrino flux measurement with Borexino detector

The low-energy range, namely 165-590 keV, of the Borexino experimental spectrum has been carefully analyzed with the purpose of the pp-neutrino flux extraction [9]. The data were acquired from January 2012 to May 2013 and correspond to 408 live days of data taking.

The main features of the experimental spectrum, as can be seen in in figure 1, are: a main contribution from the ^{14}C decays at low energies (below 200 keV) and the monoenergetic peak corresponding to 5.3 MeV α -particles from ^{210}Po decay. The statistics in the first bins used in the analysis is very high, of the order of 5×10^5 events, demanding the development of a very precise model for the studies - the allowed systematic precision at low energy part should be comparable to the statistical fluctuations of 0.14%. The development of such a precise model was a main goal of the analysis.

As one can see from figure 1 the shapes of contributions to the spectrum from ^{14}C and other background components are quite different from the contribution from the expected contribution of the electron recoil spectrum from the Solar pp-neutrino. This fact allows the extraction of the Solar pp-neutrino contribution from the data.

The stability and robustness of the measured pp neutrino interaction rate was verified by performing fits varying initial conditions, including fit energy range, method of pile-up construction, and energy estimator. The distribution of the central values for pp-neutrino interaction rates obtained for all these fit conditions was then used as an estimate of the maximal systematic error (partial correlations between different factors are not excluded).

3. Results and Implications

The solar pp neutrino interaction rate measured by Borexino is $pp = 144 \pm 13(stat) \pm 10(syst)$ cpd/100 t, compatible with the expected rate of $pp_{theor} = 131 \pm 2$ cpd/100 t. The corresponding total solar pp-neutrino flux is $\phi_{pp}(Borex) = (6.6 \pm 0.7) \times 10^{10} \text{ cm}^{-2}\text{s}^{-1}$, in a good agreement with the combined best fit value of the radiochemical and other solar experiments $\phi_{pp}(other) = (6.14 \pm 0.61) \times 10^{10} \text{ cm}^{-2}\text{s}^{-1}$ [10]. Both are in agreement with the expected value of $6.0 \times (1.000 \pm 0.006) \times 10^{10} \text{ cm}^{-2}\text{s}^{-1}$.

The survival probability for electron neutrino from pp-reaction is $P_{ee}(Borex) = 0.64 \pm 0.12$.

Taking into account that Borexino and other experiments measurements are independent, the results can be combined:

$$\phi_{pp} = (6.37 \pm 0.46) \times 10^{10} \text{ cm}^{-2}\text{s}^{-1}.$$

The electron neutrino survival probability measured in all solar but Borexino experiment is $P_{ee}(other) = 0.56 \pm 0.08$, combining it with Borexino one we obtain:

$$P_{ee} = 0.60 \pm 0.07,$$

well compatible with theoretical prediction of the MSW/LMA model $0.561^{+0.030}_{-0.042}$.




Figure 1. Borexino energy spectrum between 165 and 590 keV (in black). The pp neutrino component is shown in red, the ^{14}C spectrum in dark purple and the synthetic pile-up in light purple. The large green peak is ^{210}Po α -decays. ^7Be (dark blue), pep and CNO (light blue) solar neutrinos, and ^{210}Bi (orange) spectra are almost flat in this energy region.

Table 1. The Standard Solar Model predictions (for high metallicity and low metallicity abundances) and current experimental status of the Solar neutrino fluxes measurement. The limits are given for 90% C.L.. The corresponding energy release is calculated in the last column.

| Reaction | GS98 [11] | AGS09 [12] | Units $\text{cm}^{-2}\text{s}^{-1}$ | Measurement | MeV/ 1ν | L $\times 10^{26} \text{ W}\cdot\text{s}^{-1}$ |
|-----------------|------------------|------------------|--|---|-------------|---|
| pp | 5.98 ± 0.04 | 6.03 ± 0.04 | $\times 10^{10}$ | 6.14 ± 0.61 [10] 6.6 ± 0.7 [13] 6.37 ± 0.46 | 13.10 | 3.76 ± 0.28 |
| pep | 1.44 ± 0.012 | 1.47 ± 0.012 | $\times 10^8$ | 1.6 ± 0.3 [14] | 11.92 | 0.009 ± 0.002 |
| ^7Be | 5.00 ± 0.07 | 4.56 ± 0.07 | $\times 10^9$ | 4.87 ± 0.24 [6] | 12.60 | 0.276 ± 0.014 |
| ^8B | 5.58 ± 0.14 | 4.59 ± 0.14 | $\times 10^6$ | 5.25 ± 0.16 [15] | 6.63 | 1.57 ± 0.05 $\times 10^{-4}$ |
| hep | 8.0 ± 2.4 | 8.3 ± 2.5 | $\times 10^3$ | < 23 [16] | | |
| ^{13}N | 2.96 ± 0.14 | 2.17 ± 0.14 | $\times 10^8$ | CNO: | | |
| ^{15}O | 2.23 ± 0.15 | 1.56 ± 0.15 | $\times 10^8$ | < 7.4 [14] | | |
| ^{17}F | 5.52 ± 0.17 | 3.40 ± 0.16 | $\times 10^6$ | | | |

All available measurements of the solar neutrino fluxes are shown in table 1. The total energy production in the solar reactions observed till now (by detecting corresponding neutrino fluxes)

is $(4.04 \pm 0.28) \times 10^{26} \text{ W}\cdot\text{s}^{-1}$ in a good agreement with a total measured $L_{\odot} = 3.846 \times 10^{26} \text{ W}\cdot\text{s}^{-1}$. There is not much space left for the unknown energy sources, the 90% C.L. lower limit for the total energy production (conservatively assuming zero contribution from the not-observed reactions) is $L_{tot} = 3.68 \times 10^{26} \text{ W}\cdot\text{s}^{-1}$. If one assumes that such an unknown source exists, its total power with 90% probability can't exceed $0.15 \times 10^{26} \text{ W}\cdot\text{s}^{-1}$. In other words no more than 4% of the total energy production in the Sun is left for the unknown energy sources, confirming that the Sun shines due to the thermonuclear fusion reactions.

4. Conclusion

Borexino provided an independent measurement of the Solar pp-neutrino flux, which can be combined with measurements from other solar experiments. A comparison of the neutrino flux from the Sun with Solar luminosity in photons provides a test of the stability of the Sun on the 10^5 years time scale, and allows to set a limit of no more than 4% of the total power production for the unknown energy sources in the Sun at 90% C.L..

Acknowledgments

The Borexino program is made possible by funding from INFN (Italy), NSF (USA), BMBF, DFG, and MPG (Germany), RFBR: Grants 14-22-03031 and 13-02-12140, RFBR-ASPERA-13-02-92440 (Russia), and NCN Poland (UMO-2012/06/M/ST2/00426). We acknowledge the generous support and hospitality of the Laboratori Nazionali del Gran Sasso (LNGS). Russian colleagues from MEPhI acknowledge partial support from MEPhI Academic Excellence Project (contract No. 02.a03.21.0005, 27.08.2013).

References

- [1] Chapman G A 1997 *Encyclopedia of Planetary Science, Encyclopedia of Earth Science* (Springer Netherlands) 748
- [2] Fröhlich C and Lean J 1998 *Geophysical Research Letters* **25**(23) 4377–80
- [3] Bahcall J N 1989 *Neutrino Astrophysics* (Cambridge University Press)
- [4] Bahcall J N 2002 *Phys. Rev. C* **025801**
- [5] Fiorentini G and Ricci B 1999 *Comments on Astrophysics* **1** 49–51
- [6] Bellini G *et al.* 2011 *Physical Review Letters* **107**(14) 141302
- [7] Smirnov O Ju, Zaimidoroga O A and Derbin A V 2003 *Physics of Atomic Nuclei* **66**(4) 712–23
- [8] Derbin A V, Smirnov O.Yu. and Zaimidoroga O A 2004 *Physics of Atomic Nuclei* **67**(11) 2066–72
- [9] Bellini G *et al.* 2014 *Nature* **512** 383–6
- [10] Ianni A 2014 *Physics of the Dark Universe* **4** 44 – 9
- [11] Grevesse N and Sauval A J 1998 *Space Science Reviews* **85**(1-2) 161–74
- [12] Asplund M *et al.* 2009 *Annual Review of Astronomy and Astrophysics* **47**(1) 481–522
- [13] Bellini G *et al.* 2014 *Phys. Rev. D* **89** 112007
- [14] Bellini G *et al.* 2012 *Phys. Rev. Lett.* **108** 051302
- [15] Aharmim B *et al.* 2010 *Phys. Rev. C* **81** 055504
- [16] Aharmim B *et al.* 2006 *The Astrophysical Journal* **653**(2) 1545