

developmental therapeutics

375P A phase Ib dose-finding study of alpelisib (ALP; BYL719) and paclitaxel (PTX) in advanced solid tumors (aST)

J. Rodón¹, G. Curigliano², J.-P. Delord³, W. Harb⁴, A. Azaro¹, V. Donnet⁵, Y. Han⁶, L. Blumenstein⁷, C. Wilke⁷, J.T. Beck⁸

¹Medical Oncology, Vall d'Hebron University Hospital Institut d'Oncologia, Barcelona, Spain, ²Drug Development, European Institute of Oncology (IEO), Milan, Italy, ³Oncology, Institut Claudius Régaud, Toulouse, France, ⁴Unity Campus, Horizon Oncology Center, Lafayette, IN, USA, ⁵Oncology Global Development, Novartis Pharma S.A.S, Paris, France, ⁶Biostatistics, Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA, ⁷Oncology Global Development, Novartis Pharma AG, Basel, Switzerland, ⁸Oncology, Highlands Oncology Group, Lafayetteville, AZ, USA

Background: Aberrant activation of the phosphatidylinositol 3-kinase (PI3K)/mammalian target of rapamycin pathway due to alterations in *PIK3CA* (encoding PI3K α) frequently occurs in aST. We report safety findings from an ongoing, phase Ib dose-escalation study of ALP (PI3K α inhibitor) + PTX (NCT02051751).

Methods: Patients (pts) aged ≥ 18 years with aST (not amenable to resection/progressed on standard therapy), ECOG performance status ≤ 2 , adequate bone marrow/organ function, and no prior treatment with PI3K or AKT inhibitors were recruited. The primary objective was to determine the maximum tolerated dose (MTD) and/or recommended Phase II dose of ALP + PTX based on dose-limiting toxicities (DLTs) in Cycle 1. Dose escalation of ALP was guided by an adaptive Bayesian logistic regression model with escalation with overdose control principle.

Results: As of Dec 7, 2015, 19 pts received oral ALP (300 mg [n = 6], 250 mg [n = 4], or 150 mg [n = 9] once daily [QD]) and IV PTX (80 mg/m² once weekly [QW]). The most common primary sites of cancer were breast (n = 5) and rectum (n = 3). Treatment was discontinued in 18/19 pts due to disease progression (n = 12, 63%), pt decision (n = 3, 16%), adverse events (AEs; n = 2, 11%; 1 pt for grade [G]3 dehydration, G3 hyperglycemia, and G3 acute kidney injury; 1 pt for G4 neutropenia and G4 γ -glutamyltransferase increase), and physician decision (n = 1, 5%). DLTs occurred in 5/12 pts in the dose-determining set: 1/1 (100%) pt at 300 mg QD, 2/3 (67%) pts at 250 mg QD, and 2/8 (25%) pts at 150 mg QD. Six DLTs were reported: G2 hyperglycemia (n = 3), G4 hyperglycemia, G4 leukopenia, and G3 acute kidney injury (each n = 1). The MTD of ALP + PTX (80 mg/m² QW) was declared as 150 mg QD. All 19 pts had ≥ 1 treatment-emergent AE. Grade 3/4 AEs occurred in 11 (58%) pts, the most frequent being hyperglycemia (n = 6, 32%), diarrhea, anemia, lymphopenia, neutropenia, and leukopenia (each n = 2, 11%).

Conclusions: In pts with aST, the MTD of ALP + PTX (80 mg/m² QW) was 150 mg QD. Due to the challenging safety profile of the combination and lack of available data confirming the pharmacodynamics and/or clinical activity of ALP at 150 mg QD, planned dose expansion in pts with breast cancer and head and neck squamous cell carcinoma will not go forward.

Clinical trial identification: NCT02051751

Legal entity responsible for the study: Novartis

Funding: Novartis

Disclosure: J. Rodón: Advisory board for Novartis, Lilly, Servier, Leti, Oncompass, Orion Pharma. V. Donnet: Novartis Full-time Employee. Y. Han: I am an employee at Novartis and receive a salary from Novartis. L. Blumenstein: I hereby confirm to be a Novartis Pharma AG employee with stock ownership. C. Wilke: Employee of Novartis AG, sponsor of the study. All other authors have declared no conflicts of interest.