

to our study, only significant lesions were assessed.⁸

Finally, we are indebted to Sipahi *et al.* for the erratum found on the slopes of the regression lines of the second and third panels of *Figure 3*. Indeed, the value should be negative.

References

1. Jimenez-Quevedo P, Sabate M, Angiolillo D, Alfonso F, Hernandez-Antolin R, Banuelos C, Bernardo E, Ramirez C, Moreno R, Fernandez C, Escaned J, Macaya C. LDL-cholesterol predicts negative coronary artery remodeling in diabetic patients: an intravascular ultrasound study. *Eur Heart J* 2005;**26**:2307–2312.
2. Sabaté M, Costa MA, Kozuma K, Kay IP, van der Giessen WJ, Coen VLMA, Ligthart JMR, Serrano P, Levendag PC, Serruys PW. Geographic miss: a cause of treatment failure in radio-oncology applied to intracoronary radiation therapy. *Circulation* 2000;**101**:2467–2471.
3. Angiolillo DJ, Sabaté M, Alfonso F, Macaya C. 'Candy wrapper' effect after drug-eluting stent implantation: deja vu or stumbling over the same stone again? *Catheter Cardiovasc Interv* 2004;**61**:387–391.
4. Sabate M, Jimenez-Quevedo P, Angiolillo DJ, Gomez-Hospital JA, Alfonso F, Hernandez-Antolin R, Goicolea J, Banuelos C, Escaned J, Moreno R, Fernandez C, Fernandez-Aviles F, Macaya C, DIABETES Investigators. Randomized comparison of sirolimus-eluting stent versus standard stent for percutaneous coronary revascularization in diabetic patients: the diabetes and sirolimus-eluting stent (DIABETES) trial. *Circulation* 2005;**112**:2175–2183.
5. Mintz S, Kent KM, Pichard AD, Satler LF, Popma JJ, Leon MB. Contribution of inadequate arterial remodeling to the development of focal coronary artery stenoses. *Circulation* 1997;**95**:1791–1798.
6. Mintz GS, Nissen SE, Anderson WD, Bailey SR, Erbel R, Fitzgerald PJ, Pinto FJ, Rosenfield K, Siegel RJ, Tuzcu EM, Yock PG. American College of Cardiology Clinical Expert Consensus Document on Standards for Acquisition, Measurement and Reporting of Intravascular Ultrasound Studies (IVUS). A report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol* 2001;**37**:1478–1492.
7. Schoenhagen P, Ziada KM, Kapadia SR, Crowe TD, Nissen SE, Tuzcu EM. Extent and direction of arterial remodeling in stable versus unstable coronary syndromes: an intravascular ultrasound study. *Circulation* 2000;**101**:598–603.
8. Weissman NJ, Sheris SJ, Chari R, Mendelsohn FO, Anderson WD, Breall JA, Tanguay JF, Diver DJ. Intravascular ultrasonic analysis of plaque characteristics associated with coronary artery remodeling. *Am J Cardiol* 1999;**84**:37–40.

Pilar Jiménez-Quevedo
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Manel Sabaté
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain
Tel: +34913303283
E-mail address:
manelsabate1@telefonica.net

Dominick Angiolillo
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Fernando Alfonso
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Rosana Hernandez-Antolin
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Camino Bañuelos
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Esther Bernardo
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Celia Ramirez
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Raul Moreno
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Cristina Fernandez
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Javier Escaned
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

Carlos Macaya
Hospital Clínico San Carlos
Interventional Cardiology Department
Martin Lagos s/n
28040 Madrid
Spain

doi:10.1093/eurheartj/ehi839

Online publish-ahead-of-print 14 March 2006

Concurrent coronary and carotid artery surgery: an open debate

We read with great interest the article by Kolh *et al.*¹ dealing with the question about concurrent coronary artery bypass grafting (CABG) and carotid endarterectomy (CEA) surgery. Nevertheless, their report raises some concerns.

The article analysed factors influencing peri-operative outcomes, and ascending aortic calcification was found to be an independent predictor of both stroke and death. These data are considered a common finding in patients who underwent heart surgery.² However, in patients with concomitant severe carotid and coronary disease, the predictive role of aortic disease on stroke and death could be significantly overestimated. Carotid disease alone is a risk factor for stroke in patients who undergo surgical revascularization and stroke represents a known complication of CEA. In a recent large study, the adjusted stroke rate was 2.67-fold greater in the combined CEA-CABG group compared with CABG alone.³ By its nature, this study did not evaluate the impact of CEA on stroke incidence and the confounding effect of carotid disease on mortality and morbidity does not emerge. Hence the predictive effect of aortic disease could be significantly altered. Further studies on patients with and without concomitant CEA are needed to unmask the effect of carotid disease on stroke and to evaluate the real predictive role of aortic disease.

In addition, the multivariate model cannot predict precisely the odds ratio for variables associated with operative mortality and morbidity, as confidence intervals are large, although *P*-values are significant.

Some data about cardiopulmonary bypass and surgical techniques are lacking. We think that it could be important to explain where the arterial cannulation was made in

patients with ascending aorta disease, as the number of no-touch aorta technique and off-pump surgery was lower than expected. Moreover, no explanation was given about surgical approach to 17 patients with aortic dilatation, as it seems that dilated ascending aorta was not substituted.

This study confirms that concurrent CEA and CABG can be performed with acceptable mortality and morbidity, but it represents only one of the surgical option.⁴ We agree with the authors who consider treatment of symptomatic territory first reserving the combined simultaneous treatment for patients symptomatic in both territories.⁵

References

1. Kolh PH, Comte L, Tchana-Sato V, Honore C, Kerzmann A, Mauer M, Limet R. Concurrent coronary and carotid artery surgery: factors influencing perioperative outcome and long-term results. *Eur Heart J* 2006;**27**:49–56.
2. Kapetanakis EI, Stamou SC, Dullum MK, Hill PC, Haile E, Boyce SW, Bafi AS, Petro KR, Corso PJ. The impact of aortic manipulation on neurologic outcomes after coronary artery bypass surgery: a risk-adjusted study. *Ann Thorac Surg* 2004;**78**:1564–1571.
3. Hill MD, Shrive FM, Kennedy J, Feasby TE, Ghali WA. Simultaneous carotid endarterectomy and coronary artery bypass surgery in Canada. *Neurology* 2005;**64**:1435–1437.
4. Antunes PE, Anacleto G, de Oliveira JM, Eugenio L, Antunes MJ. Staged carotid and coronary surgery for concomitant carotid and coronary artery disease. *Eur J Cardiothorac Surg* 2002;**21**:181–186.
5. Huh J, Wall MJ Jr, Soltero ER. Treatment of combined coronary and carotid artery disease. *Curr Opin Cardiol* 2003;**18**:447–453.

Fabio Barili

Department of Cardiovascular Surgery
Centro Cardiologico Monzino
University of Milan
Via Parea 4
20138 Milan
Italy
Tel: +39 02 58002563
Fax: +39 02 58011194
E-mail address: fabarili@libero.it

Gianluca Polvani

Department of Cardiovascular Surgery
Centro Cardiologico Monzino
University of Milan
Via Parea 4
20138 Milan
Italy

Veli K. Topkara

Department of Cardiothoracic Surgery
Columbia University College of

Physicians and Surgeons
New York, NY
USA

Luca Dainese

Department of Cardiovascular Surgery
Centro Cardiologico Monzino
University of Milan
Via Parea 4
20138 Milan
Italy

Paolo Biglioli

Department of Cardiovascular Surgery
Centro Cardiologico Monzino
University of Milan
Via Parea 4
20138 Milan
Italy

doi:10.1093/eurheartj/ehi840

Online publish-ahead-of-print 14 March 2006

Concurrent coronary and carotid artery surgery: an open debate: reply

We read with interest the letter from Barili *et al.* on our work 'Concurrent coronary artery surgery: factors influencing perioperative outcome and long-term results'.¹ We thank Barili *et al.* for their comments that need our reply.

First, we share their doubts regarding the precise impact of asymptomatic carotid artery stenosis on cerebrovascular events occurring during coronary artery bypass grafting (CABG). This question could only be answered by randomizing patients with concurrent significant coronary and carotid artery lesions to either combined CABG and carotid endarterectomy (CEA) or CABG alone, with post-operative stroke rate being the primary endpoint. However, a very large number of patients would be required for such a study. Indeed, in the recent ACST trial,² a total of 3120 patients with asymptomatic carotid lesions were randomized to immediate CEA or indefinite deferral of any CEA to demonstrate a reduction of 5-year stroke risk from ~12 to ~6% in the immediate CEA group.

Therefore, to reach statistical significance, a randomized study comparing CABG and CEA with CABG alone would necessarily be multicentric (126 clinical sites from 30 countries participated in the ACST trial), with the inevitable differences concerning, for example, the use of a carotid shunt or the closure of the carotid arteriotomy with or without a patch.

Our policy has a more modest goal, which is to associate both surgical procedures during one anaesthesia when, in patients requiring CABG surgery, the carotid lesion meets the criteria accepted for surgical indication. We certainly are unsure that this policy significantly decreases the incidence of post-operative stroke. In our centre, the incidence of post-operative cerebrovascular events is higher after combined CEA–CABG when compared with CEA alone, and the incidence of post-operative myocardial infarction is higher after combined surgery than after CABG alone. This observation supports the notion that carotid artery lesions are a harbinger of severe atherosclerosis.

Secondly, we disagree with Barili *et al.* that our model was unable to accurately predict odds ratio for operative results. It is true that confidence intervals were large, but odds ratios were high (odds ratio for aortic calcifications was 6.3 for operative mortality and 5.2 for stroke).¹ Our results are in concordance with the observation of John *et al.*³ that aortic calcifications were the leading risk factor for stroke, with an odds ratio of 3.0, in CABG patients.

Thirdly, arterial cannulation was usually performed in the ascending or transverse aorta, at a site of normal palpation, especially in the early part of the series. When the aorta was very hostile, arterial cannulation was performed in the femoral artery or in the axillary artery. In the series reported,¹ a moderately dilated ascending aorta (>3.5 cm) was left in place. When the ascending aorta was more severely dilated (>5 cm), it was replaced with a prosthesis, but those patients were not included in this study.

Finally, we agree that patients with coronary and carotid symptomatic lesions should be treated with concurrent surgical procedures.

References

1. Kolh PH, Comte L, Tchana-Sato V, Honore C, Kerzmann A, Mauer M, Limet R. Concurrent coronary and carotid artery surgery: factors influencing perioperative outcome and long-term results. *Eur Heart J* 2006;**27**:49–56.
2. Halliday A, Mansfield A, Marro J, Peto C, Peto R, Potter J, Thomas D. Prevention of disabling and fatal strokes by successful carotid endarterectomy in patients without recent neurological symptoms: randomised controlled trial. *Lancet* 2004;**363**:1491–1502.
3. John R, Choudhri AF, Weinberg AD, Ting W, Rose EA, Smith CR, Oz MC. Multicenter review of preoperative risk factors for stroke after coronary artery bypass grafting. *Ann Thorac Surg* 2000;**69**:30–35.