

1 **Phase II Studies with Refametinib or Refametinib plus Sorafenib in**
2 **Patients with RAS-mutated Hepatocellular Carcinoma**

3 **Running title:** Refametinib or Refametinib + Sorafenib in RAS-mutated HCC

4 Ho Yeong Lim^{1,*}, Philippe Merle², Karl Heinz Weiss³, Thomas Yau⁴, Paul Ross⁵, Vincenzo
5 Mazzaferro⁶, Jean-Frédéric Blanc⁷, Yuk Ting Ma⁸, Chia Jui Yen⁹, Judit Kocsis¹⁰, Su Pin
6 Choo¹¹, Wattana Sukeepaisarnjaroen¹², René Gérolami¹³, Jean-François Dufour¹⁴, Edward J.
7 Gane¹⁵, Baek-Yeol Ryoo¹⁶, Markus Peck-Radosavljevic^{17,†}, Thong Dao¹⁸, Winnie Yeo¹⁹,
8 Wisut Lamlertthon²⁰, Satawat Thongsawat²¹, Michael Teufel²², Katrin Roth²³, Diego Reis²⁴,
9 Barrett H. Childs²², Heiko Krissel²³, and Josep M. Llovet^{25,26,27*}

10 ¹*Division of Hematology-Oncology, Samsung Medical Center, Sungkyunkwan University,*
11 *Seoul, Korea;* ²*Service of Hepato-Gastroenterology, Hepatology Unit, Croix-Rousse*
12 *Hospital, Lyon, France;* ³*Section of Transplant Hepatology, Liver Cancer Center Heidelberg,*
13 *Heidelberg, Germany;* ⁴*Department of Medicine, Queen Mary Hospital, Hong Kong;*
14 ⁵*Cancer Centre, Guy's & St Thomas' NHS Foundation Trust, London, UK;* ⁶*Gastrointestinal*
15 *Surgery and Liver Transplant Unit, The Fondazione IRCCS Istituto Nazionale Tumori*
16 *(National Cancer Institute) and University of Milan, Milan, Italy;* ⁷*Service of Hepato-*
17 *Gastroenterology and Digestive Oncology, Hôpital Haut-Lévêque, Bordeaux, France;*
18 ⁸*Department of Medical Oncology, University Hospitals Birmingham NHS Foundation Trust,*
19 *Birmingham, UK;* ⁹*Division of Hematology and Oncology, National Cheng Kung University*
20 *Hospital, Tainan, Taiwan;* ¹⁰*Oncology Department, Debrecen University Clinical Center,*
21 *Debrecen, Hungary;* ¹¹*Division of Medical Oncology, National Cancer Centre Singapore,*
22 *Singapore;* ¹²*Department of Medicine, Srinagarind Hospital, Khon Kaen, Thailand;* ¹³*Service*
23 *of Hepato-Gastroenterology, Aix-Marseille University, Marseille, France;* ¹⁴*Department of*
24 *Hepatology, University Clinic for Visceral Surgery and Medicine, University Hospital of*

25 *Bern, Bern, Switzerland;* ¹⁵*New Zealand Liver & Transplant Unit, Auckland City Hospital,*
26 *Auckland, New Zealand;* ¹⁶*Department of Oncology, Asan Medical Center, Seoul, Korea;*
27 ¹⁷*Department of Gastroenterology and Hepatology, Endocrinology, Rheumatology and*
28 *Nephrology, Medical University of Vienna, Vienna, Austria;* ¹⁸*Service of Hepato-*
29 *Gastroenterology and Nutrition, Caen University Hospital, Caen, France;* ¹⁹*Department of*
30 *Clinical Oncology, Chinese University of Hong Kong, Hong Kong;* ²⁰*Chulabhorn Hospital,*
31 *Bangkok, Thailand;* ²¹*Department of Internal Medicine, Maharaj Nakorn Chiang Mai*
32 *Hospital, Chiang Mai, Thailand;* ²²*Bayer HealthCare Pharmaceuticals, Inc., Whippany, NJ,*
33 *USA;* ²³*Bayer AG, Berlin, Germany;* ²⁴*Medical and Data Management, Bayer S.A., São*
34 *Paulo, Brazil;* ²⁵*Division of Liver Diseases, Icahn School of Medicine at Mount Sinai, New*
35 *York, NY, USA;* ²⁶*Liver Cancer Translational Research Laboratory, Barcelona Clinic Liver*
36 *Cancer Group (BCLC), IDIBAPS-Hospital Clínic de Barcelona, University of Barcelona,*
37 *Barcelona, Spain;* ²⁷*Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain*
38 [†]*Current affiliation: Department of Gastroenterology/Hepatology, Endocrinology and*
39 *Nephrology, Klinikum Klagenfurt am Wörthersee, Klagenfurt, Austria*

40 **Financial Support**

41 These studies were supported by Bayer AG. The study sponsor played a role in the design of
42 the study, data interpretation, writing of the manuscript, and the decision to submit for
43 publication. At each study center, the principal investigator was responsible for the study.
44 Medical writing support for the manuscript was funded by the study sponsor, based on
45 detailed direction and feedback from all authors. Medical writing assistance was provided by
46 Laura Badtke, PhD, at Complete HealthVizion, Chicago, IL, USA, based on detailed
47 discussion and feedback from all authors. Medical writing assistance was funded by
48 Bayer AG.

49 ***Corresponding Authors:**

50 Ho Yeong Lim, Division of Hematology-Oncology, Department of Medicine, Samsung
51 Medical Center, School of Medicine, Sungkyunkwan University, Seoul, Korea. Phone: 82-2-
52 3410-0914; Fax: 82-2-3410-1754; E-mail: hoylim@skku.edu

53 Josep M. Llovet, Director, Mount Sinai Liver Cancer Program, Division of Liver Diseases,
54 Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, 1425 Madison Ave., 11F-
55 70, Box 1123, New York, NY 10029, USA. Phone: 212-659-9503; Fax: 212-849-2574; E-
56 mail: josep.llovet@mssm.edu; Assistant: Melody Tam (melody.tam@mssm.edu)

57 **Disclosure of Potential Conflicts of Interest**

58 K.H. Weiss reports grants from Novartis and MSD, personal fees from Bayer, GMPO, and
59 Vivet Therapeutics, and grants and personal fees from Bristol-Myers Squibb, Univar, Wilson
60 Therapeutics, and Alexion. P. Ross reports grants from Sanofi Aventis, fees for educational
61 sessions from BMS, travel support from BMS, Sirtex, Bayer, Merck Serono, Servier, Amgen,
62 and Celgene, advisory board fees from BMS, Sirtex, Bayer, Merck Serono, Servier, and
63 Celgene, and speaking fees from Merck Serono, Sirtex, and Bayer. V. Mazzaferro reports
64 personal fees from Bayer and BTG. J.-F. Blanc reports personal fees from Bayer SP, Lilly,
65 and BMS. Y.T. Ma reports honoraria and travel support from Bayer and consultancy fees
66 from Baxalta UK, Ltd. S.P. Choo reports grants from Bristol-Myers Squibb, travel support
67 from Amgen, and honoraria from Bristol-Myers Squibb, Bayer, Novartis, and Sirtex. J.-F.
68 Dufour reports participating in advisory committees for AbbVie, Bayer, Bristol-Myers
69 Squibb, Falk, Genfit, Gilead Sciences, Intercept, Lilly, Merck, and Novartis, speaking and
70 teaching for AbbVie, Bayer, Bristol-Myers Squibb, Genfit, Gilead Science, and Novartis, and
71 an unrestricted research grant from Bayer. E.J. Gane reports participating in clinical advisory
72 committees or speaker bureaus for Alios, AbbVie, Arrowhead, Janssen, Merck, Gilead

73 Sciences, and Mylan. M. Peck-Radosavljevic reports being an investigator for AbbVie,
74 ArQule-Daiichi, Bayer, Bristol-Myers Squibb, Boehringer Ingelheim, Imclone, Lilly, MSD,
75 Novartis, and Roche, grant support from AbbVie, ArQule-Daiichi, Bayer, MSD, and Roche,
76 and acting as a speaker or advisor for Abbott, Bayer, Bristol-Myers Squibb, Boehringer
77 Ingelheim, Lilly, MSD, and Roche. T. Dao reports congress and travel support from Bayer,
78 Gilead Sciences, AbbVie, and MSD, and being an investigator for Lilly. M. Teufel and B.H.
79 Childs are employees of Bayer HealthCare Pharmaceuticals, Inc. K. Roth and H. Krissel are
80 employees of Bayer AG. H. Krissel also reports patent BHC 133035 EP 01 pending. D. Reis
81 is an employee of and has received personal fees from Bayer S.A. J.M. Llovet reports
82 research/education grants from Bayer, Blueprint Medicines, Boehringer Ingelheim, and
83 Incyte and consulting fees from Eli Lilly, Bayer, BMS, Blueprint Medicines, Eisai, Celsion,
84 and Boehringer Ingelheim. The remaining authors declare no potential conflicts of interest.

85 **Authors' Contributions**

86 All authors contributed to the collection and analysis of the data and to the writing of the
87 manuscript, and approved the final version before submission. H.Y. Lim and J.M. Llovet:
88 study design; coordinating investigators for the review; approved the clinical study reports;
89 had full access to all the data in the studies; vouch for the integrity of the data analyses; had
90 final responsibility for the decision to submit. K. Roth: statistical analysis. M. Teufel: study
91 design, interpretation of the biomarker results. D. Reis, B.H. Childs, and H. Krissel: study
92 design.

93 **ClinicalTrials.gov numbers:** NCT01915589, NCT01915602

94 **Keywords:** Hepatocellular carcinoma, *KRAS*, refametinib, sorafenib, BEAMing

95 **Word count:** 4717 (max 5000)

96 **Number of figures and tables:** 3 figures and 3 tables

97 **Translational Relevance**

98 The frequency of *RAS* mutations in hepatocellular carcinoma (HCC) is reported to be ~4%.
99 In this study, we report on the use of a liquid biopsy to prospectively screen patients with
100 HCC for *RAS* mutations using circulating tumor DNA for treatment with the MEK inhibitor
101 refametinib in monotherapy or in combination with sorafenib. The low prevalence of *RAS*
102 mutations in HCC was confirmed (4.4% of patients). *RAS* mutational status was confirmed
103 by next-generation sequencing using circulating tumor DNA, which allowed for the
104 determination of the mutational landscape in patients with HCC. The most frequently
105 detected mutations were in *TERT*, *TP53*, and β -catenin, confirming data reported in The
106 Cancer Genome Atlas. This is the first study using a liquid biopsy for large-scale mutational
107 testing, which offers the opportunity for comprehensive mutational analysis using a
108 non-invasive approach.

109 **Abstract**

110 **Purpose:** Refametinib, an oral MEK inhibitor, has demonstrated antitumor activity in
111 combination with sorafenib in patients with *RAS*-mutated hepatocellular carcinoma (HCC).
112 Two phase II studies evaluated the efficacy of refametinib monotherapy and refametinib plus
113 sorafenib in patients with *RAS*-mutant unresectable or metastatic HCC.

114 **Methods:** Eligible patients with *RAS* mutations of cell-free circulating tumor DNA (ctDNA)
115 determined by beads, emulsion, amplification, and magnetics technology received twice-daily
116 refametinib 50 mg ± sorafenib 400 mg. Potential biomarkers were assessed in ctDNA via
117 next-generation sequencing (NGS).

118 **Results:** Of 1318 patients screened, 59 (4.4%) had a *RAS* mutation, of whom 16 received
119 refametinib and 16 received refametinib plus sorafenib. With refametinib monotherapy, the
120 objective response rate (ORR) was 0%, the disease control rate (DCR) was 56.3%, overall
121 survival (OS) was 5.8 months, and progression-free survival (PFS) was 1.9 months. With
122 refametinib plus sorafenib, the ORR was 6.3%, the DCR was 43.8%, OS was 12.7 months,
123 and PFS was 1.5 months. In both studies, time to progression was 2.8 months. Treatment-
124 emergent toxicities included fatigue, hypertension, and acneiform rash. Twenty-seven
125 patients had ctDNA samples available for NGS. The most frequently detected mutations
126 were in *TERT* (63.0%), *TP53* (48.1%), and β -catenin (*CTNNB1*; 37.0%).

127 **Conclusions:** Prospective testing for *RAS* family mutations using ctDNA was a feasible, non-
128 invasive approach for large-scale mutational testing in HCC patients. A median OS of
129 12.7 months with refametinib plus sorafenib in this small population of *RAS*-mutant patients
130 may indicate a synergistic effect between sorafenib and refametinib – this preliminary finding
131 should be further explored.

132 **Word count:** 249

133 **Introduction**

134 Hepatocellular carcinoma (HCC) is a leading cause of cancer-related death worldwide (1,2),
135 and the prognosis for HCC remains extremely poor (2,3). The recommended standard of care
136 in advanced HCC is treatment with the multikinase inhibitor sorafenib (3-5). Lenvatinib has
137 been shown to be non-inferior to sorafenib in a recent phase III trial (6). In second-line, the
138 multikinase inhibitor regorafenib has been approved after showing significantly improved
139 survival versus placebo in patients who had disease progression on sorafenib (7). The kinase
140 inhibitor cabozantinib has also demonstrated promising survival improvements versus
141 placebo as second-line therapy in a phase III trial (8). Immunotherapy has shown promise in
142 HCC, with the immune checkpoint inhibitor nivolumab recently approved for the second-line
143 treatment of advanced HCC based on durable responses observed in a phase I/II trial (9).
144 Treatment with the monoclonal antibody ramucirumab has shown survival improvement
145 versus placebo in patients progressing to sorafenib with alpha-fetoprotein >400 ng/ml (10).
146 Poor prognosis and a lack of treatment options highlight a need for additional viable
147 treatment regimens in the advanced setting.

148 Refametinib (BAY 86-9766; Bayer AG, Berlin, Germany) is an oral, potent, non-adenosine
149 triphosphate competitive inhibitor targeting MEK 1 and 2 (11), which play a central role in
150 the RAS signal transduction cascade. RAS-MAPK signaling has been implicated in tumor
151 progression and dissemination in HCC (2). A phase I study of the combination of
152 refametinib with sorafenib in patients with advanced malignancies including HCC
153 demonstrated a favorable safety profile and pharmacokinetic profile at a maximum tolerated
154 dose of refametinib 50 mg twice daily in combination with sorafenib 400 mg twice daily (12).

155 The analysis of cell-free circulating tumor DNA (ctDNA) using beads, emulsion,
156 amplification, and magnetics technology (BEAMing; Sysmex Inostics GmbH, Hamburg,

157 Germany) enables tumor genotyping at the time of treatment and offers a viable, non-invasive
158 approach to identifying clinically relevant mutations (13,14). BEAMing may therefore be a
159 feasible tool to support the need for the identification of predictive biomarkers in HCC (3),
160 through proof-of-concept studies. Previous proof-of-concept studies of kinase inhibitors in
161 other cancer types have successfully detected predictive mutations, such as vemurafenib in
162 patients with inoperable melanoma with a *BRAF*^{V600} mutation (15) and crizotinib in patients
163 with non-small-cell lung cancer with *EML4-ALK* fusion (16).

164 A retrospective analysis in a phase II study evaluating refametinib plus sorafenib in Asian
165 patients with HCC found that patients with *RAS* mutations exhibited a robust clinical
166 response compared with patients with wild-type *RAS* (objective tumor response rate [ORR]:
167 3/4 patients [75.0%] compared with 1/65 patients [1.5%], respectively) (17). Here we
168 describe the first proof-of-concept studies based on mutations conducted in patients with
169 HCC. Two phase II studies prospectively evaluated the efficacy of refametinib monotherapy
170 (NCT01915589) and refametinib plus sorafenib (NCT01915602) in patients with
171 unresectable or metastatic HCC with mutated *RAS*, as determined by BEAMing of ctDNA.

172 **Patients and Methods**

173 **Study design**

174 These were two phase II, prospective, single-arm, multicenter, uncontrolled, open-label
175 studies. The primary objective was to evaluate the efficacy of refametinib alone or in
176 combination with sorafenib in patients with *RAS*- (*KRAS*- or *NRAS*-) mutated unresectable or
177 metastatic HCC. The primary efficacy variable was the central radiologic assessment of
178 ORR (complete response [CR] plus partial response [PR]) according to modified Response
179 Evaluation Criteria in Solid Tumors (mRECIST) (18). The secondary objective was safety,

180 and additional objectives included evaluation of biomarkers aiming to identify biomarkers or
181 biomarker signatures which could correspond to therapy response. Secondary efficacy
182 variables included centrally assessed ORR according to RECIST version 1.1, investigator-
183 assessed ORR according to mRECIST and RECIST version 1.1, overall survival, disease
184 control rate, time to radiographic tumor progression, duration of response, and progression-
185 free survival.

186 Fifteen patients with *RAS* mutations were planned to be included in the first stage of each
187 study. The second stage was to be initiated if five or more of these patients had a confirmed
188 objective response according to mRECIST.

189 **Important protocol amendments**

190 In the refametinib monotherapy study, the protocol was amended once, with changes
191 implemented globally. Prior cytotoxic chemotherapy was added as an exclusion criterion to
192 omit a population of overtreated patients who may have been different from patients
193 conventionally treated with sorafenib; this change affected eligibility criteria for *RAS*
194 mutation testing and treatment exclusion criteria. An exclusion criterion was also added
195 regarding women of childbearing potential to reduce the time gap between the pregnancy
196 evaluation and the beginning of treatment.

197 In the refametinib plus sorafenib study, the protocol was amended twice. Changes were
198 implemented globally and included the following amendments: patients with a corrected QT
199 interval >480 ms at the time of screening were excluded from the study because of the
200 potential for QT prolongation with sorafenib; the exclusion criterion regarding women of
201 childbearing potential was amended to reduce the time gap between the pregnancy evaluation
202 and the beginning of treatment; the exclusion criterion regarding systemic anticancer therapy
203 was clarified, as patients with prior systemic anticancer therapy were not eligible for this

204 study. In addition, a dose-modification scheme for hepatotoxic events was included, because
205 hepatotoxicity is an “identified risk” for the refametinib–sorafenib combination.

206 Amendments to the statistical analysis plan for the refametinib plus sorafenib study included
207 the collection of survival data to be continued until the last patient’s last visit instead of until
208 12 weeks after the last patient’s first treatment, or earlier if all patients had withdrawn from
209 the study. A data rule was also added regarding tumor assessment by centralized blinded
210 reading; for cases with missing adjudication for patients who had completed or withdrawn
211 from treatment at the time of primary analysis, the worst-case approach was to be applied.

212 **Patients**

213 Written, informed consent was obtained from all patients. The study protocol conforms to
214 the ethical guidelines of the 1975 Declaration of Helsinki as reflected in *a priori* approval by
215 the institutions’ human research committees.

216 Eligibility criteria for *RAS* mutational testing included: age ≥ 18 years with unresectable or
217 metastatic HCC, confirmed histologically (mandatory for non-cirrhotic patients) or by non-
218 invasive radiologic criteria; Eastern Cooperative Oncology Group performance status of 0 or
219 1; and life expectancy ≥ 12 weeks. Prior use of targeted agents, experimental therapy, or
220 systemic anticancer treatment was not allowed, although prior sorafenib treatment was
221 permitted in patients who received refametinib only.

222 Treatment eligibility criteria included: *KRAS* or *NRAS* mutation based on BEAMing plasma
223 test; Child-Pugh class A liver function status; at least one uni-dimensional measurable lesion
224 by computed tomography or magnetic resonance imaging; and Eastern Cooperative Oncology
225 Group performance status of 0 or 1. Treatment exclusion criteria included: any cancer
226 curatively treated less than 3 years before study entry (except cervical carcinoma *in situ*,

227 treated basal cell carcinoma, and superficial bladder tumors); eligibility for surgery, liver
228 transplantation, ablation, or transarterial chemoembolization for hepatocellular carcinoma;
229 renal failure requiring hemo- or peritoneal dialysis; a history of cardiac disease; or
230 uncontrolled hypertension.

231 **Treatment**

232 In both studies, eligible patients harboring *RAS* mutations received refametinib 50 mg twice
233 daily in 21-day cycles. In the refametinib plus sorafenib study, patients also received
234 standard sorafenib (400 mg twice daily), starting with a dose of 600 mg daily (200 mg in the
235 morning plus 400 mg in the evening) in cycle 1, escalating to the standard sorafenib dose in
236 cycle 2 if no hand-foot skin reaction, fatigue, or gastrointestinal toxicities of grade ≥ 2
237 occurred. Patients received treatment on a continuous basis until radiologic disease
238 progression, clinical progression, or other criteria for discontinuation of treatment were met.

239 **Assessments**

240 ctDNA from plasma samples collected in the pre-treatment period was centrally evaluated for
241 *RAS* mutational status using BEAMing technology (13), with a limit of detection at 0.02%
242 mutant allele. Tumor assessments were performed at screening and every 6 weeks.
243 Treatment response was centrally assessed according to mRECIST for the primary endpoint,
244 and was also investigator-assessed according to mRECIST (18). Safety, including adverse
245 events (AEs) and concomitant medications, was monitored throughout the studies. Creatine
246 phosphokinase (CPK) increase of grade ≥ 3 was considered an AE of special interest and was
247 to be reported as a serious AE (SAE). Plasma samples for biomarker analysis were collected
248 at screening, at cycle 1, days 1 and 15, and at cycle 2, day 15. Peripheral whole-blood
249 samples from patients with mutated *RAS* were analyzed for detection of genomic alterations
250 using FoundationACT[®] (Foundation Medicine[®], Cambridge, MA, USA), a targeted next-

251 generation sequencing (NGS)-based ctDNA assay (19). The detection limit of
252 FoundationACT[®] is specified at 0.1% mutant allele frequency, i.e. a lower sensitivity than
253 BEAMing. FoundationACT[®] is a hybrid-capture-based assay that is designed to interrogate
254 62 genes, identifying all classes of alterations including base substitutions, insertions and
255 deletions, copy number variations, and rearrangements/fusions through computational
256 algorithms (20).

257 **Statistical analysis**

258 In each study, it was estimated that approximately 350 patients were needed to be tested via
259 BEAMing to identify 15 patients with mutated *RAS* in stage 1, and that approximately
260 2300 patients would need to be tested via BEAMing to identify a sufficient number of
261 patients with mutated *RAS* to be treated in stage 2.

262 Descriptive statistics were calculated for the presented endpoints.

263 **Results**

264 **Patient disposition, demographics, and baseline characteristics**

265 In the refametinib monotherapy study, 498 patients were enrolled at 58 study centers in
266 17 countries across Asia, Europe, and the USA from September 2013 to June 2014. *RAS*
267 mutational testing was performed in 493 patients (Fig. 1A); 32 (6.5%) had a *RAS* mutation.
268 In the refametinib plus sorafenib study, 820 patients were enrolled at 80 study centers in
269 21 countries across Asia, Europe, and the USA from September 2013 to April 2015. *RAS*
270 mutational testing was performed in 815 patients (Fig. 1B); 27 (3.3%) had a *RAS* mutation.
271 Overall, 4.4% of HCC patients screened (59/1318) had a *RAS* mutation determined by
272 BEAMing. Of those, 32/59 patients received treatment, either refametinib monotherapy

273 ($n = 16$) or refametinib plus sorafenib ($n = 16$). Reasons for patients not receiving treatment
274 are summarized in Fig. 1.

275 In the refametinib monotherapy study, the median age was 69 years and the median time
276 since initial HCC diagnosis was 72.1 weeks (Table 1). Nine patients (56.3%) had received
277 prior first-line sorafenib treatment. In the refametinib plus sorafenib study, the median age
278 was 67 years and the median time since initial HCC diagnosis was 32.2 weeks (Table 1).
279 Demographics and baseline characteristics were similar between patients irrespective of *RAS*
280 mutational status in both studies.

281 **Efficacy**

282 Of the 16 patients treated with refametinib monotherapy, no patient had a CR or PR when
283 centrally assessed according to mRECIST, and the ORR was 0% (Table 2). One patient
284 (6.3%) achieved an unconfirmed PR and eight (50.0%) achieved stable disease; the disease
285 control rate was 56.3%. ORR was 0% by independent assessment according to RECIST
286 version 1.1: no patients had a CR or PR, 10 (62.5%) had stable disease, two (12.5%) had
287 disease progression, and four (25.0%) were not evaluable. The investigator-assessed ORR
288 was 0% according to mRECIST (Supplementary Table S1) and RECIST version 1.1:
289 no patients had a confirmed or unconfirmed CR or PR, 10 (62.5%) had stable disease,
290 two (12.5%) had disease progression, and four (25.0%) were not evaluable.

291 Of the 16 patients treated with refametinib plus sorafenib, one patient (6.3%) achieved a
292 confirmed PR when centrally assessed according to mRECIST, and the ORR was 6.3%
293 (Table 2). Two patients (12.5%) achieved unconfirmed PRs (confirmatory computed
294 tomography scan showed progression) and four (25.0%) had stable disease; the disease
295 control rate was 43.8%. Independent assessment according to RECIST version 1.1 reported
296 an ORR of 6.3%: one patient (6.3%) had a confirmed PR, six (37.5%) had stable disease,

297 five (31.3%) had disease progression, and four (25.0%) were not evaluable or had data
298 missing. Investigator-assessed ORR was 6.3% according to mRECIST (one patient [6.3%]
299 had a confirmed PR) (Supplementary Table S1). The investigator-assessed ORR was also
300 6.3% according to RECIST version 1.1: one patient (6.3%) had a PR, one (6.3%) had an
301 unconfirmed PR, five (31.3%) had stable disease, six (37.5%) had disease progression, and
302 three (18.8%) had missing data.

303 In the refametinib monotherapy study, four patients (25.0%) had radiologic progression and
304 the median time to progression was 2.8 months (Fig. 2A). Seven patients (43.8%) in the
305 refametinib plus sorafenib study had radiologic progression and the median time to
306 progression was 2.8 months (Fig. 2B).

307 Duration of response could not be calculated in the refametinib monotherapy study because
308 no patient achieved a CR or PR. Duration of response based on central assessment in the
309 refametinib plus sorafenib study was 1.4 months for the one patient who achieved a
310 confirmed PR; this patient had a *KRAS*^{G35A} point mutation. Duration of response was
311 2.7 months for the one patient who was investigator-assessed as achieving a confirmed PR;
312 this patient had a *KRAS*^{G38A} mutation.

313 In the refametinib monotherapy study, nine patients (56.3%) had disease progression or died
314 and median progression-free survival was 1.9 months (Fig. 2C). Eight patients (50.0%) died
315 and median overall survival was 5.8 months (Fig. 2E). In the refametinib plus sorafenib
316 study, 10 patients (62.5%) had disease progression or died and median progression-free
317 survival was 1.5 months (Fig. 2D). Nine patients (56.3%) died and median overall survival
318 was 12.7 months (Fig. 2F).

319 **Exposure and dose modifications**

320 With refametinib monotherapy, the median duration of treatment (including interruptions)
321 was 7.14 weeks. The mean (\pm standard deviation) daily dose of refametinib (excluding
322 interruptions) was 90.01 ± 13.88 mg.

323 With refametinib plus sorafenib, the median durations of treatment (including interruptions)
324 for refametinib and sorafenib were 8.21 weeks and 6.43 weeks, respectively. The mean
325 (\pm standard deviation) daily doses (excluding interruptions) of refametinib and sorafenib were
326 85.06 ± 16.07 mg and 514.24 ± 124.23 mg, respectively. The majority of patients
327 experienced treatment-emergent AEs (TEAEs) of hand-foot skin reaction, fatigue, or
328 gastrointestinal toxicities of grade ≥ 2 in cycle 1, so only three patients (18.8%) received the
329 full dose of sorafenib (800 mg/day) following cycle 1. One patient remained on treatment at
330 the time of data-cut off and has been ongoing for approximately 2 years.

331 TEAEs led to dose modification (interruption or reduction) in 14 patients (87.5%) receiving
332 refametinib monotherapy (Table 3) and were considered drug-related in 13 patients (81.3%).
333 Treatment was permanently discontinued because of TEAEs in four patients (25.0%) and
334 were considered drug-related in three patients (18.8%).

335 With refametinib plus sorafenib, dose modifications were reported in 11 patients (68.8%)
336 with refametinib and 11 patients (68.8%) with sorafenib. TEAEs led to dose modification in
337 15 patients (93.8%) (Table 3); events were considered refametinib-related in 13 patients
338 (81.3%) and sorafenib-related in 14 patients (87.5%).

339 **Safety**

340 At least one TEAE was reported in all 16 patients (100%) receiving refametinib monotherapy
341 (Table 3). The most common TEAEs of worst grade 3 were fatigue and increased CPK

342 (three patients each [18.8%]). Grade 5 TEAEs occurred in five patients (31.3%): sepsis,
343 death not otherwise specified, multi-organ failure, lung infection, and heart failure. The
344 causes of death were progressive disease (one patient) and AE associated with clinical
345 disease progression and AE not associated with clinical disease progression (two patients
346 each). Drug-related TEAEs occurred in 14 patients (87.5%) (Supplementary Table S2). In
347 most patients (75.0%), the worst grade of drug-related TEAEs was grade 3, and one patient
348 (6.3%) had a drug-related TEAE of grade 4 (increased serum amylase). Twelve patients
349 (75.0%) experienced SAEs (Supplementary Table S3), of which the most common worst
350 grade was grade 3 (43.8%). SAEs were refametinib-related in seven patients (43.8%), most
351 commonly increased CPK (three patients [18.8%]). All other refametinib-related SAEs were
352 reported in one patient each (6.3%) (Supplementary Table S3).

353 TEAEs occurred in all 16 patients (100%) receiving refametinib plus sorafenib (Table 3).
354 Hand-foot skin reaction was reported in two patients (12.5%). The most common TEAEs of
355 worst grade 3 were hypertension (10/16 [62.5%]) and increased aspartate aminotransferase
356 and increased CPK in five patients each (31.3%). Seven TEAEs of worst grade 4 were
357 reported in three patients (18.8%): increased aspartate aminotransferase, increased CPK,
358 decreased platelet count, investigations - other, hypophosphatemia, and hyperuricemia.
359 Grade 5 TEAEs included general disorders and administration site conditions - other and
360 dyspnea (one patient each); the cause of death was AE associated with clinical disease
361 progression and progressive disease (one patient each). Refametinib- and sorafenib-related
362 TEAEs were reported for all 16 patients (100%) (Supplementary Table S2). Nine patients
363 (56.3%) had refametinib-related TEAEs of grade 3 and three patients (18.8%) had
364 refametinib-related TEAEs of grade 4 (Supplementary Table S2). Twelve patients (75.0%)
365 had sorafenib-related TEAEs of worst grade 3 and three patients (18.8%) had sorafenib-
366 related TEAEs of worst grade 4. One patient (6.3%) had a grade 5 TEAE considered related

367 to both refametinib and sorafenib (general disorders and administration site conditions -
368 other). SAEs were reported in 13 patients (81.3%) (Supplementary Table S3), most
369 commonly increased CPK in six patients (37.5%; five grade 3, one grade 4). Refametinib-
370 related SAEs were experienced by 12 patients (75.0%), most frequently worst grade 3 (7/16
371 [43.8%]). Increased CPK was the most commonly reported refametinib-related SAE (5/16
372 [31.3%]; four grade 3, one grade 4). Sorafenib-related SAEs occurred in 10 patients (62.5%);
373 seven patients (43.8%) had events of worst grade 3 and one patient (6.3%) experienced worst
374 grade 4 (increased CPK). One SAE of grade 5 was considered refametinib-related and
375 sorafenib-related (general disorders and administration site conditions - other).

376 **Biomarker analyses**

377 To identify potential genomic biomarkers which might be associated with resistance to
378 refametinib monotherapy or combination therapy, NGS (FoundationACT[®]) was performed
379 on available ctDNA from 27 patients (refametinib monotherapy, $n = 15$; refametinib plus
380 sorafenib, $n = 12$). *RAS* mutations were not called by NGS in over 60% of the samples with a
381 mutant allele frequency of between 0.02% and 0.1% as determined by BEAMing. *RAS*
382 mutational status was confirmed by NGS in 12 patients (44.4%), all with a mutant allele
383 frequency above 0.1%. The *RAS* somatic aberration detected was concordant with BEAMing
384 results in 11 patients (91.7%). Excluding *RAS*, the most frequently detected mutation was in
385 the promoter region of telomerase reverse transcriptase (*TERT*; 17/27 [63.0%]), followed by
386 *TP53* (13/27 [48.1%]), and β -catenin (*CTNNB1*; 10/27 [37.0%]) (Fig. 3). Actionable
387 mutations were rare (<10%) and included oncogenes such as *EGFR*, *JAK2*, *BRAF*, *FLT3*,
388 *PIK3CA*, and *cKIT*.

389 Discussion

390 These two phase II proof-of-concept studies prospectively evaluated the efficacy and safety
391 of refametinib monotherapy or refametinib plus sorafenib in patients prospectively screened
392 for *RAS*-mutant unresectable or metastatic HCC based on evaluation of mutational status in
393 ctDNA. The previous phase II BASIL trial in a separate population of Asian patients with
394 HCC receiving refametinib plus sorafenib demonstrated that the majority of patients who
395 responded to this regimen had mutant *RAS* tumors, with an ORR of 75% in patients with
396 *RAS*-mutant HCC compared with 1.5% in HCC patients with no *RAS* mutation (17).

397 In these studies, prospective testing for *RAS* mutations using ctDNA isolated from plasma
398 was a feasible, non-invasive approach for large-scale mutational testing in HCC patients.

399 The current findings support a previous report of the use of ctDNA to detect *KRAS* mutations
400 via BEAMing in a small study of patients with refractory colorectal carcinoma treated with
401 regorafenib (21), although *KRAS* mutational frequency was notably higher in the colorectal
402 carcinoma population (~40%) compared with that reported here (~5%). Overall, 59/1318
403 (4.4%) of the HCC patients screened had a *RAS* mutation. The *RAS* mutation rates reported
404 here are consistent with previous reports in this patient population (~5%) (22-25). It should
405 therefore be noted that the low *RAS* mutational frequency in this population suggests that
406 identifying *RAS*-mutant patients may be challenging in practice.

407 The primary efficacy variable was not met in the refametinib monotherapy study, with no
408 patient with mutated *RAS* achieving a CR or PR. In the refametinib plus sorafenib study, one
409 patient with mutated *RAS* achieved a PR, resulting in an ORR of 6.3%, which is broadly
410 similar to the ORR of 6.9% reported in the BASIL trial (17).

411 The target for the first stage of the trials ($\geq 5/15$ patients with a CR or PR) was not reached;
412 therefore, these studies did not proceed to the planned evaluation of refametinib monotherapy
413 or combination therapy in a larger number of patients. Further exploration would be required
414 to understand the lower ORR with refametinib plus sorafenib in this study compared with
415 previous reports (17). These results suggest that the use of *RAS* mutational status as a
416 prognostic biomarker for treatment response to refametinib monotherapy or in combination
417 with sorafenib was unsuccessful, and targeting MEK with refametinib in this *RAS*-mutant
418 patient population did not lead to a significant proportion of objective responses. However,
419 the low number of patients treated should be taken into account, and the low proportion of
420 responses observed may reflect random error – these results should therefore be interpreted
421 with caution. Additional molecular events may explain the limited responses seen using
422 mutated *RAS* as a prognostic biomarker for targeted MEK inhibition in these studies. It is
423 possible that with intra-tumor heterogeneity, mutations occurring in low-frequency subclonal
424 tumor cell populations may have acquired mutations that conferred resistance to refametinib,
425 which was targeted to progenitor cells expressing truncal driver mutations in *RAS*, negatively
426 affecting clinical outcomes (26). Evaluation of non-truncal mutations, together with longer-
427 term evaluations of changes in allele frequency, were not planned in this study, although may
428 provide useful insights into the development of resistance to refametinib in patients with
429 HCC.

430 Median overall survival was 5.8 months with refametinib monotherapy and 12.7 months with
431 refametinib plus sorafenib, with over half of events occurring during the study period. *KRAS*
432 mutation is generally associated with poorer outcomes in most cancers, although there are no
433 established data in HCC due to the lack of robust testing in large studies of advanced disease
434 (23). In our study, the effect of refametinib monotherapy on overall survival can be
435 considered insignificant, since the expected outcome of placebo at first or second line is 7–8

436 months (27,28). In fact, this survival of under 6 months might indicate that advanced *RAS*-
437 positive HCC tumors have a poor natural history. It should also be noted that 56% of patients
438 in the monotherapy arm had received prior sorafenib, possibly contributing to the poor
439 survival seen. However, the approximately 13-month survival outcome with refametinib plus
440 sorafenib treatment is more intriguing, considering the expected median survival with first-
441 line sorafenib monotherapy alone is 11 months (29). This result may indicate a synergistic
442 effect between sorafenib and refametinib, which is relevant as tumors harboring *RAS*
443 mutations remain some of the most challenging to treat because of the paucity of successful
444 drugs targeting the *RAS* pathway (30). However, this finding should be interpreted with
445 caution because of the heterogeneity in baseline liver function and tumor factors, which could
446 affect response to treatment. Also, patients in the refametinib plus sorafenib study had a
447 much shorter median time from initial diagnosis to study treatment compared with patients in
448 the monotherapy study (32.1 weeks vs. 72.1 weeks, respectively). The overall survival
449 findings in the combination study support those described for patients receiving refametinib
450 plus sorafenib in the BASIL trial (17), although median overall survival was increased in our
451 study (12.7 months vs. 9.5 months, respectively).

452 Median time to progression was the same across the refametinib monotherapy and
453 refametinib plus sorafenib studies (2.8 months), with similar median progression-free
454 survival observed between the studies (1.9 months and 1.5 months, respectively). However,
455 progression-free survival times were lower than previous reports (17).

456 Drug exposure was similar between both studies and similar to the median refametinib dose
457 observed in the BASIL study (17). The majority of patients in the refametinib plus sorafenib
458 study experienced AEs during cycle 1 that prevented sorafenib dose escalation to 800 mg per
459 day, which was also observed in the BASIL study (17). The majority of patients across both

460 studies experienced AEs leading to dose modifications, which may have caused insufficient
461 drug exposure, potentially leading to reduced efficacy of both refametinib regimens. Overall,
462 median duration of treatment was relatively short in both trials (7 weeks and 8 weeks,
463 respectively), similar to that reported for the BASIL study (8 weeks) (17).

464 Overall, refametinib was tolerated as monotherapy and combination therapy, and the majority
465 of TEAEs were manageable in both studies. In patients receiving refametinib monotherapy,
466 the most common TEAEs of limb edema, fatigue, nausea, and vomiting were consistent with
467 the safety profile previously reported in a phase I study of refametinib (31). The high overall
468 incidence of grade 3 TEAEs irrespective of causality in both studies (68.8%) was similar to
469 that reported in the BASIL trial (60.0%) (17). Generally, the observed incidence and severity
470 of refametinib-related TEAEs observed with refametinib monotherapy were comparable with
471 data from the previous refametinib phase I study (31). Refametinib-related SAEs were less
472 frequent with refametinib monotherapy than with refametinib plus sorafenib (43.8% vs.
473 75.0%, respectively). Increased CPK grade ≥ 3 was the most common refametinib-related
474 SAE reported in both studies, consistent with reports of increased CPK as a class effect of
475 MEK inhibitors (32-34).

476 Compared with the known safety profile of sorafenib monotherapy (29,35), a higher
477 incidence of liver and gastrointestinal toxicities and rash was observed in patients who
478 received refametinib plus sorafenib. However, alopecia and hand-foot skin reaction were less
479 common compared with those reported for sorafenib monotherapy (14% vs. 12.5% and 21%
480 vs. 6.3%, respectively) (29), possibly due to the reduced exposure to sorafenib in the majority
481 of patients in our study.

482 Biomarker analysis of ctDNA analyzed by NGS showed the observed mutational landscape
483 to be consistent with published data for HCC (36). The most common mutation was in the

484 promoter region of *TERT*, supporting previous observations in patients with HCC and
485 combined HCC-cholangiocarcinoma (37,38). Few actionable mutations were found, with
486 none appearing to explain the resistance to refametinib alone or in combination with
487 sorafenib, and few of the detected mutations are feasible for targeting with existing drugs. It
488 therefore remains inconclusive from our results as to whether somatic mutations in oncogenes
489 affect the efficacy of refametinib in monotherapy or combination therapy. Although analyses
490 were planned to evaluate the role of biomarkers in the response to treatment, due to limited
491 sample size and early study termination it was not possible to fully address the role of intra-
492 tumor heterogeneity (26). In addition, although the two study populations included only
493 Child-Pugh A patients, these patients were heterogeneous for various factors that may be
494 prognostic for treatment response, such as a history of ascites (in four patients overall
495 [12.5%]) (39), alpha-fetoprotein (>400 µg/L in 12 patients [37.5%]) (40), microvascular
496 invasion (in 11 patients [34%]) (41), extrahepatic spread (in 16 patients [50%]) (41), and
497 hepatitis C (in seven patients [21.9%]) (41). However, no formal analysis of lung function
498 status and tumor factors as prognostic markers for treatment response was planned in these
499 studies.

500 In these studies, *RAS* mutational status as determined by BEAMing was confirmed in 44% of
501 samples using NGS, all with mutant allele frequencies of 0.1% or higher. Although
502 BEAMing technology is highly sensitive (42), the newly developed NGS from ctDNA
503 approach has demonstrated high concordance, confirming nearly all mutations identified by
504 BEAMing, and offers the additional advantage of providing the mutational landscape based
505 on ctDNA. A comparison of sensitivity between both assays is difficult due to the different
506 detection limit of each method (0.02% for BEAMing vs. 0.1% for NGS), which did not allow
507 for the detection of *RAS* mutational status by NGS in over 60% of samples with mutant allele
508 frequency between 0.02% and 0.1%. Nonetheless, our results demonstrated that NGS

509 appears to be a promising non-invasive approach to determine the landscape of somatic
510 mutations, particularly for patients in whom a biopsy is not an option (19).

511 Despite the poor ORR in patients with *RAS* mutations, a median overall survival of
512 13 months in the small population included in the refametinib plus sorafenib study may
513 indicate a synergistic effect between refametinib and sorafenib that should be further
514 explored in a larger patient population that is not stratified by *RAS* mutational status, taking
515 into account other prognostic factors based on patient heterogeneity and intra-tumor
516 heterogeneity. The analysis of mutational status using ctDNA isolated from plasma as a
517 liquid biopsy was a feasible, non-invasive technique in patients with unresectable or
518 metastatic HCC, although *RAS* mutational frequency was low. Further analysis of this
519 technique is warranted for discovery of predictive biomarkers in HCC and other cancers.

520 **Acknowledgments**

521 The authors wish to thank the patients and their families. These studies were supported by
522 Bayer AG. Medical writing assistance was provided by Laura Badtke, PhD, at Complete
523 HealthVizion, Chicago, IL, USA, based on detailed discussion and feedback from all authors.
524 Medical writing assistance was funded by Bayer AG.

525 **References**

- 526 1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer
527 statistics, 2012. *CA Cancer J Clin* **2015**;65(2):87-108.
- 528 2. Llovet JM, Zucman-Rossi J, Pikarsky E, Sangro B, Schwartz M, Sherman M, *et al.*
529 Hepatocellular carcinoma. *Nat Rev Dis Primers* **2016**;2:16018.
- 530 3. European Association for the Study of the Liver, European Organisation for Research
531 and Treatment of Cancer. EASL-EORTC clinical practice guidelines: management of
532 hepatocellular carcinoma. *J Hepatol* **2012**;56(4):908-43.
- 533 4. Omata M, Lesmana LA, Tateishi R, Chen PJ, Lin SM, Yoshida H, *et al.* Asian Pacific
534 Association for the Study of the Liver consensus recommendations on hepatocellular
535 carcinoma. *Hepatol Int* **2010**;4(2):439-74.
- 536 5. Bruix J, Sherman M. Management of hepatocellular carcinoma: an update.
537 *Hepatology* **2011**;53(3):1020-2.
- 538 6. Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, *et al.* Lenvatinib versus
539 sorafenib in first-line treatment of patients with unresectable hepatocellular
540 carcinoma: a randomised phase 3 non-inferiority trial. *Lancet* **2018**:[Epub ahead of
541 print] doi 10.1016/S0140-6736(18)30207-1.
- 542 7. Bruix J, Qin S, Merle P, Granito A, Huang YH, Bodoky G, *et al.* Regorafenib for
543 patients with hepatocellular carcinoma who progressed on sorafenib treatment
544 (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*
545 **2017**;389(10064):56-66 doi 10.1016/S0140-6736(16)32453-9.

- 546 8. Abou-Alfa GK, Meyer T, Cheng AL, El-Khoueiry AB, Rimassa L, Ryoo BY, *et al.*
547 Cabozantinib (C) versus placebo (P) in patients (pts) with advanced hepatocellular
548 carcinoma (HCC) who have received prior sorafenib: results from the randomized
549 phase III CELESTIAL trial. *J Clin Oncol* **2018**;36 (Suppl 4S):abstr 207.
- 550 9. El-Khoueiry AB, Sangro B, Yau T, Crocenzi TS, Kudo M, Hsu C, *et al.* Nivolumab in
551 patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label,
552 non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet*
553 **2017**;389(10088):2492-502 doi 10.1016/S0140-6736(17)31046-2.
- 554 10. Zhu AX, Park JO, Ryoo BY, Yen CJ, Poon R, Pastorelli D, *et al.* Ramucirumab
555 versus placebo as second-line treatment in patients with advanced hepatocellular
556 carcinoma following first-line therapy with sorafenib (REACH): a randomised,
557 double-blind, multicentre, phase 3 trial. *Lancet Oncol* **2015**;16(7):859-70 doi
558 10.1016/S1470-2045(15)00050-9.
- 559 11. Iverson C, Larson G, Lai C, Yeh LT, Dadson C, Weingarten P, *et al.* RDEA119/BAY
560 869766: a potent, selective, allosteric inhibitor of MEK1/2 for the treatment of cancer.
561 *Cancer Res* **2009**;69(17):6839-47.
- 562 12. Adjei AA, Richards DA, El-Khoueiry A, Braiteh F, Becerra CHR, Stephenson JJ, *et*
563 *al.* A Phase I study of the safety, pharmacokinetics, and pharmacodynamics of
564 combination therapy with refametinib plus sorafenib in patients with advanced cancer.
565 *Clin Cancer Res* **2016**;22(10):2368-76.
- 566 13. Tabernero J, Lenz HJ, Siena S, Sobrero A, Falcone A, Ychou M, *et al.* Analysis of
567 circulating DNA and protein biomarkers to predict the clinical activity of regorafenib

- 568 and assess prognosis in patients with metastatic colorectal cancer: a retrospective,
569 exploratory analysis of the CORRECT trial. *Lancet Oncol* **2015**;16(8):937-48.
- 570 14. Thierry AR, Mouliere F, El Messaoudi S, Mollevi C, Lopez-Crapez E, Rolet F, *et al.*
571 Clinical validation of the detection of *KRAS* and *BRAF* mutations from circulating
572 tumor DNA. *Nat Med* **2014**;20(4):430-5.
- 573 15. Sosman JA, Kim KB, Schuchter L, Gonzalez R, Pavlick AC, Weber JS, *et al.*
574 Survival in BRAF V600-mutant advanced melanoma treated with vemurafenib. *N*
575 *Engl J Med* **2012**;366(8):707-14.
- 576 16. Solomon BJ, Mok T, Kim DW, Wu YL, Nakagawa K, Mekhail T, *et al.* First-line
577 crizotinib versus chemotherapy in *ALK*-positive lung cancer. *N Engl J Med*
578 **2014**;371(23):2167-77.
- 579 17. Lim HY, Heo J, Choi HJ, Lin CY, Yoon JH, Hsu C, *et al.* A phase II study of the
580 efficacy and safety of the combination therapy of the MEK inhibitor refametinib
581 (BAY 86-9766) plus sorafenib for Asian patients with unresectable hepatocellular
582 carcinoma. *Clin Cancer Res* **2014**;20(23):5976-85.
- 583 18. Lencioni R, Llovet JM. Modified RECIST (mRECIST) assessment for hepatocellular
584 carcinoma. *Semin Liver Dis* **2010**;30(1):52-60.
- 585 19. Clark TA, Kennedy M, Young L, Zhao M, Coyne M, Breese V, *et al.* Rigorous
586 validation of clinical circulating tumor DNA for cancer molecular profiling. 107th
587 Annual Meeting of the American Association for Cancer Research, New Orleans, LA,
588 USA, April 16-20. 3965 ed2016.
- 589 20. Foundation Medicine Inc. FoundationACT[®] Technical Information. 2016.

- 590 21. Wong ALA, Lim JSJ, Sinha A, Gopinathan A, Lim R, Tan CS, *et al.* Tumour
591 pharmacodynamics and circulating cell free DNA in patients with refractory
592 colorectal carcinoma treated with regorafenib. *J Transl Med* **2015**;13:57.
- 593 22. Hou W, Liu J, Chen P, Wang H, Ye BC, Qiang F. Mutation analysis of key genes in
594 RAS/RAF and PI3K/PTEN pathways in Chinese patients with hepatocellular
595 carcinoma. *Oncol Lett* **2014**;8(3):1249-54.
- 596 23. Turhal NS, Savaş B, Çoşkun Ö, Baş E, Karabulut B, Nart D, *et al.* Prevalence of K-
597 Ras mutations in hepatocellular carcinoma: a Turkish Oncology Group pilot study.
598 *Mol Clin Oncol* **2015**;3(6):1275-9.
- 599 24. Tornesello ML, Buonaguro L, Izzo F, Buonaguro FM. Molecular alterations in
600 hepatocellular carcinoma associated with hepatitis B and hepatitis C infections.
601 *Oncotarget* **2016**;7(18):25087-102.
- 602 25. Villanueva A, Llovet JM. Liver cancer in 2013: mutational landscape of HCC - the
603 end of the beginning. *Nat Rev Clin Oncol* **2014**;11(2):73-4.
- 604 26. Fisher R, Puzstai L, Swanton C. Cancer heterogeneity: implications for targeted
605 therapeutics. *Br J Cancer* **2013**;108(3):479-85 doi 10.1038/bjc.2012.581.
- 606 27. Llovet JM, Decaens T, Raoul JL, Boucher E, Kudo M, Chang C, *et al.* Brivanib in
607 patients with advanced hepatocellular carcinoma who were intolerant to sorafenib or
608 for whom sorafenib failed: results from the randomized Phase III BRISK-PS study. *J*
609 *Clin Oncol* **2013**;31(28):3509-16.
- 610 28. Bruix J, Merle P, Granito A, Huang YH, Bodoky G, Yokosuka O, *et al.* Efficacy and
611 safety of regorafenib versus placebo in patients with hepatocellular carcinoma (HCC)

- 612 progressing on sorafenib: results of the international, randomized phase 3 RESORCE
613 trial. *Ann Oncol* **2016**;27 (Suppl 2):ii140-ii1 doi 10.1093/annonc/mdw237.03.
- 614 29. Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, *et al.* Sorafenib in
615 advanced hepatocellular carcinoma. *N Engl J Med* **2008**;359(4):378-90.
- 616 30. Gysin S, Salt M, Young A, McCormick F. Therapeutic strategies for targeting ras
617 proteins. *Genes Cancer* **2011**;2(3):359-72.
- 618 31. Gore L, Lewis K, Von Hoff DD, Weiss GJ, Ramanathan RK, Adjei AA, *et al.* Safety,
619 pharmacokinetics, and pharmacodynamics results from a phase I trial of BAY 86-
620 9766 (RDEA119), a MEK inhibitor, in patients with advanced cancer. *J Clin Oncol*
621 (Meeting Abstracts). Volume 29 (Suppl)2011.
- 622 32. Akinleye A, Furqan M, Mukhi N, Ravella P, Liu D. MEK and the inhibitors: from
623 bench to bedside. *J Hematol Oncol* **2013**;6:27.
- 624 33. Zhao Y, Adjei AA. The clinical development of MEK inhibitors. *Nat Rev Clin Oncol*
625 **2014**;11(7):385-400.
- 626 34. Martinez-Garcia M, Banerji U, Albanell J, Bahleda R, Dolly S, Kraeber-Bodéré F, *et*
627 *al.* First-in-human, Phase I dose-escalation study of the safety, pharmacokinetics, and
628 pharmacodynamics of RO5126766, a first-in-class dual MEK/RAF inhibitor in
629 patients with solid tumors. *Clin Cancer Res* **2012**;18(17):4806-19.
- 630 35. Ha Y, Lee D, Shim JH, Lim YS, Lee HC, Chung YH, *et al.* Role of transarterial
631 chemoembolization in relation with sorafenib for patients with advanced
632 hepatocellular carcinoma. *Oncotarget* **2016**;7:74303-13.

- 633 36. Schulze K, Imbeaud S, Letouzé E, Alexandrov LB, Calderaro J, Rebouissou S, *et al.*
634 Exome sequencing of hepatocellular carcinomas identifies new mutational signatures
635 and potential therapeutic targets. *Nat Genet* **2015**;47(5):505-11.
- 636 37. Sasaki M, Sato Y, Nakanuma Y. Mutational landscape of combined hepatocellular
637 carcinoma and cholangiocarcinoma, and its clinicopathological significance.
638 *Histopathology* **2016**;70:423-34.
- 639 38. Lee SE, Chang SH, Kim WY, Lim SD, Kim WS, Hwang TS, *et al.* Frequent somatic
640 *TERT* promoter mutations and *CTNNB1* mutations in hepatocellular carcinoma.
641 *Oncotarget* **2016**;7:69267-75.
- 642 39. Kim HY, Park JW, Joo J, Kim H, Woo SM, Lee WJ, *et al.* Worse outcome of
643 sorafenib therapy associated with ascites and Child-Pugh score in advanced
644 hepatocellular carcinoma. *J Gastroenterol Hepatol* **2013**;28(11):1756-61 doi
645 10.1111/jgh.12310.
- 646 40. Liu L, Zhao Y, Jia J, Chen H, Bai W, Yang M, *et al.* The prognostic value of alpha-
647 fetoprotein response for advanced-stage hepatocellular carcinoma treated with
648 sorafenib combined with transarterial chemoembolization. *Sci Rep* **2016**;6:19851 doi
649 10.1038/srep19851.
- 650 41. Bruix J, Cheng AL, Meinhardt G, Nakajima K, De Sanctis Y, Llovet J. Prognostic
651 factors and predictors of sorafenib benefit in patients with hepatocellular carcinoma:
652 analysis of two phase III studies. *J Hepatol* **2017**;67(5):999-1008 doi
653 10.1016/j.jhep.2017.06.026.
- 654 42. Li M, Diehl F, Dressman D, Vogelstein B, Kinzler KW. BEAMing up for detection
655 and quantification of rare sequence variants. *Nat Methods* **2006**;3(2):95-7.

Table 1. Demographics and baseline characteristics of patients receiving refametinib monotherapy or refametinib plus sorafenib

| | Refametinib monotherapy (n = 16) | Refametinib plus sorafenib (n = 16) |
|---|---|--|
| Male, <i>n</i> (%) | 13 (81.3) | 12 (75.0) |
| Race, <i>n</i> (%) | | |
| White | 9 (56.3) | 9 (56.3) |
| Asian | 7 (43.8) | 6 (37.5) |
| Black or African American | 0 | 1 (6.3) |
| Median age, years (range) | 69 (37–84) | 67 (53–82) |
| Median body mass index, kg/m ² (range) | 23.7 (20.5–31.8) | 23.6 (16.4–34.8) |
| Baseline ECOG PS, <i>n</i> (%) | | |
| 0 | 7 (43.8) | 10 (62.5) |
| 1 | 9 (56.3) | 6 (37.5) |
| Medical history, <i>n</i> (%) ^a | | |
| Hepatic cirrhosis | 14 (87.5) | 11 (68.8) |
| Ascites | 3 (18.8) | 1 (6.3) |
| Gastroesophageal reflux disease | 3 (18.8) | 1 (6.3) |
| Abdominal pain ^b | 2 (12.5) | 2 (12.5) |
| Esophageal varices | 2 (12.5) | 1 (6.3) |
| Portal hypertension | 1 (6.3) | 2 (12.5) |
| Confirmation of liver cirrhosis, <i>n</i> (%) | | |
| Histologic | 4 (25.0) | 3 (18.8) |
| Clinical | 10 (62.5) | 5 (31.3) |
| Histologic and clinical | 0 | 3 (18.8) |
| Missing | 2 (12.5) | 5 (31.3) |

Etiology of HCC, *n* (%)

| | | |
|--|-----------------------|------------------|
| Alcohol use | 6 (37.5) | 3 (18.8) |
| Alcohol use/genetic/metabolic | 0 | 1 (6.3) |
| Alcohol use/hepatitis B | 2 (12.5) | 0 |
| Alcohol use/hepatitis C | 0 | 1 (6.3) |
| Hepatitis B | 2 (12.5) | 5 (31.3) |
| Hepatitis C | 3 (18.8) | 4 (25.0) |
| Non-alcoholic steatohepatitis | 1 (6.3) | 1 (6.3) |
| Unknown | 2 (12.5) | 1 (6.3) |
| Overall Child-Pugh A score, <i>n</i> (%) | | |
| 5 | 8 (50.0) | 10 (62.5) |
| 6 | 8 (50.0) | 6 (37.5) |
| BCLC stage, <i>n</i> (%) | | |
| A (early) | 0 | 2 (12.5) |
| B (intermediate) | 2 (12.5) | 2 (12.5) |
| C (advanced) | 14 (87.5) | 12 (75.0) |
| Presence of macrovascular invasion, <i>n</i> (%) | 4 (25.0) | 7 (43.8) |
| Presence of extrahepatic spread, <i>n</i> (%) | 10 (62.5) | 6 (37.5) |
| Alpha-fetoprotein >400 µg/L, <i>n</i> (%) | 9 (56.3) ^c | 3 (18.8) |
| Bilirubin, mg/dL, median (range) | 0.9 (0.4–2.3) | 0.7 (0.3–1110.9) |
| Albumin, g/dL, median (range) | 3.9 (2.8–4.3) | 3.9 (3.2–38.0) |
| Prothrombin INR, median (range) | 1.1 (0.9–1.3) | 1.1 (0.9–1.2) |
| Median time since initial diagnosis, weeks (range) | 72.1 (5.9–262.3) | 32.2 (8.1–342.7) |
| Median time since most recent progression, weeks (range) | 8.6 (1.1–57.0) | 8.6 (3.1–21.0) |
| Prior anticancer therapies and procedures, <i>n</i> (%) | | |

| | | |
|--|-----------|-----------|
| Surgical therapeutic procedure | 6 (37.5) | 9 (56.3) |
| Systemic anticancer therapy (sorafenib) | 9 (56.3) | 0 |
| Local anticancer therapy | 6 (37.5) | 8 (50.0) |
| Number of target lesions (mRECIST), <i>n</i> (%) | | |
| 1 | 2 (12.5) | 4 (25.0) |
| 2 | 10 (62.5) | 10 (62.5) |
| 3 | 4 (25.0) | 1 (6.3) |
| 4 | 0 | 1 (6.3) |
| Number of non-target lesions (mRECIST), <i>n</i> (%) | | |
| 0 | 4 (25.0) | 6 (37.5) |
| 1 | 9 (56.3) | 8 (50.0) |
| 2 | 2 (12.5) | 0 |
| 3 | 1 (6.3) | 1 (6.3) |
| 4 | 0 | 1 (6.3) |

^aIn two or more patients overall; ^bIncludes upper and lower abdominal pain in one patient each in the combination study; ^cBaseline data missing for one patient.

Abbreviation: BCLC, Barcelona Clinic Liver Cancer; ECOG PS, Eastern Cooperative Oncology Group performance status; INR, international normalized ratio.

Table 2. Response evaluation by central assessment using mRECIST in patients receiving refametinib monotherapy or refametinib plus sorafenib

| <i>n</i> (%) [95% CI] | Refametinib monotherapy (<i>n</i> = 16) | Refametinib plus sorafenib (<i>n</i> = 16) |
|-----------------------------------|--|---|
| Best overall response | | |
| Complete response | 0 | 0 |
| Confirmed partial response | 0 | 1 (6.3) [0.16–30.23] |
| Unconfirmed partial response | 1 (6.3) [0.16–30.23] | 2 (12.5) [7.27–52.38] |
| Stable disease | 8 (50.0) [24.65–75.35] | 4 (25.0) [7.27–52.38] |
| Disease progression | 3 (18.8) [4.05–45.65] | 5 (31.3) [11.02–58.66] |
| Not evaluable | 0 | 1 (6.3) [0.16–30.23] |
| Missing | 4 (25.0) [7.27–52.38] | 3 (18.8) [4.05–45.65] |
| Objective tumor response rate | 0 | 1 (6.3) [0.16–30.23] |
| Disease control rate ^a | 9 (56.3) [29.88–80.25] | 7 (43.8) [19.75–70.12] |

^aIncludes unconfirmed complete and partial responses ≥ 6 weeks from baseline assessment.

Abbreviation: CI, confidence interval.

Table 3. Summary of safety and incidence of treatment-emergent adverse events (by worst CTCAE grade) occurring in three or more patients receiving refametinib monotherapy or refametinib plus sorafenib

| <i>n</i> (%) | Refametinib monotherapy (<i>n</i> = 16) | Refametinib plus sorafenib (<i>n</i> = 16) |
|---|--|---|
| Any TEAE ^a | 16 (100) | 16 (100) |
| Worst grade | | |
| 3 | 11 (68.8) | 11 (68.8) |
| 4 | 0 | 3 (18.8) |
| 5 (death) | 5 (31.3) | 2 (12.5) |
| Serious adverse events | 12 (75.0) | 13 (81.3) |
| Led to dose modification | 14 (87.5) | 15 (93.8) |
| Led to permanent discontinuation | 4 (25.0) | 5 (31.3) |
| Incidence of TEAEs (any grade) occurring in $\geq 10\%$ of the total population | | |
| Limb edema | 7 (43.8) | 3 (18.8) |
| Fatigue | 6 (37.5) | 12 (75.0) |
| Nausea | 6 (37.5) | 2 (12.5) |
| Vomiting | 6 (37.5) | 5 (31.3) |
| Increased creatine phosphokinase | 5 (31.3) | 8 (50.0) |
| Diarrhea | 5 (31.3) | 10 (62.5) |
| Acneiform rash | 5 (31.3) | 8 (50.0) |
| Increased aspartate aminotransferase | 4 (25.0) | 8 (50.0) |
| Maculo-papular rash | 4 (25.0) | 6 (37.5) |
| Hypertension | 4 (25.0) | 13 (81.3) |
| Anemia | 3 (18.8) | 2 (12.5) |

| | | |
|---|----------|----------|
| Abdominal pain | 3 (18.8) | 2 (12.5) |
| Ascites | 3 (18.8) | 3 (18.8) |
| Anorexia | 3 (18.8) | 4 (25.0) |
| Hypoglycemia | 3 (18.8) | 0 |
| Back pain | 3 (18.8) | 1 (6.3) |
| Dyspnea | 3 (18.8) | 2 (12.5) |
| Dry skin | 3 (18.8) | 2 (12.5) |
| Skin and subcutaneous tissue disorders - other, specify | 3 (18.8) | 0 |
| Oral mucositis | 2 (12.5) | 5 (31.3) |
| Hypoalbuminemia | 2 (12.5) | 4 (25.0) |
| Decreased platelet count | 2 (12.5) | 4 (25.0) |
| Constipation | 2 (12.5) | 3 (18.8) |
| Investigations - other, specify | 2 (12.5) | 3 (18.8) |
| Hyperglycemia | 2 (12.5) | 3 (18.8) |
| Increased alanine aminotransferase | 1 (6.3) | 3 (18.8) |
| Malaise | 1 (6.3) | 3 (18.8) |
| Skin infection | 1 (6.3) | 3 (18.8) |
| Headache | 0 | 3 (18.8) |
| Increased lipase | 0 | 3 (18.8) |

^aNumber (%) of patients with the specified event starting or worsening between the start of treatment and 30 days after the end of treatment.

Abbreviation: CTCAE, Common Terminology Criteria for Adverse Events.


Figure legends

Figure 1. Patient disposition in the two phase II studies. (A) Refametinib monotherapy study. (B) Refametinib plus sorafenib study.


Figure 2. Kaplan-Meier curves of TTP, PFS, and OS in the two phase II studies. (A) TTP in patients who received refametinib monotherapy. (B) TTP in patients who received refametinib plus sorafenib. (C) PFS in patients who received refametinib monotherapy. (D) PFS in patients who received refametinib plus sorafenib. (E) OS in patients who received refametinib monotherapy. (F) OS in patients who received refametinib plus sorafenib. Abbreviations: CI, confidence interval; NE, not estimable due to censored data; OS, overall survival; PFS, progression-free survival; TTP, time to progression.



Figure 3. Somatic aberrations of patients with *RAS* mutations as detected in circulating tumor DNA. Abbreviations: r, rearrangement; s, short variant.



A



B












Figure 3



Clinical Cancer Research

Phase II Studies with Refametinib or Refametinib plus Sorafenib in Patients with *RAS*-mutated Hepatocellular Carcinoma

Ho Yeong Lim, Philippe Merle, Karl Heinz Weiss, et al.

Clin Cancer Res Published OnlineFirst June 27, 2018.

| | |
|--------------------------|--|
| Updated version | Access the most recent version of this article at: doi: 10.1158/1078-0432.CCR-17-3588 |
| Author Manuscript | Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited. |

| | |
|-----------------------------------|--|
| E-mail alerts | Sign up to receive free email-alerts related to this article or journal. |
| Reprints and Subscriptions | To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org . |
| Permissions | To request permission to re-use all or part of this article, use this link http://clincancerres.aacrjournals.org/content/early/2018/06/27/1078-0432.CCR-17-3588 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site. |