

17 **Abstract.** Flow cytometry (FC) is widely applied to characterize and stage nodal lymphomas in
18 dogs because it has a short turnaround time, requires minimally invasive sampling, and allows
19 contemporary evaluation of neoplastic cells in the primary lesion and of blood and marrow
20 involvement. We investigated advantages and limitations of FC in suspected extranodal
21 lymphomas in dogs. The likelihood of obtaining a suitable FC sample was significantly lower for
22 aspirates of extranodal lesions than for lymph node aspirates. However, we noted no differences
23 among different extranodal lesion sites. We also describe FC results for 39 samples compatible
24 with extranodal lymphoma. A dominant population of large cells was easily identified on
25 morphologic FC scattergrams in many cases. Phenotypic aberrancies were frequently present,
26 mainly in T-cell lymphomas. Lymphoma cells were distinguishable from normal residual
27 lymphocytes in >85% of cases, facilitating the quantification of putative blood and marrow
28 involvement by FC. Despite the high percentage of non-diagnostic samples (32 of 73, >40%), we
29 support the inclusion of FC in the diagnostic workup of suspected extranodal lymphomas in
30 dogs, in conjunction with histopathology. Histopathology is the gold standard for diagnosing
31 lymphoma, provides relevant information, including tissue invasion and epitheliotropism, but has
32 a longer turnaround time.

33

34 **Key words:** Dogs; extranodal lymphoma; flow cytometry.

35

Introduction

Diagnostic flow cytometry (FC) was introduced in veterinary medicine at the end of the 20th century³⁵; the primary use of FC is in characterizing canine hematopoietic neoplasms.^{6,27} FC provides useful information in dogs with lymphoma, including phenotype of neoplastic cells,¹² lymphoma subtype,³⁶ expression of specific markers,^{25,26} stage,^{20,21,28,29} and presence of minimal residual disease.¹

Most FC studies in dogs have focused on nodal lymphoma; only a few reports have characterized primary extranodal lymphomas.^{9,11,15,31} Conversely, in human medicine, FC is routinely included in the diagnostic workup of extranodal lymphomas.^{3,19,32,34,37,38} In addition, FC is a safe and minimally invasive technique to confirm the diagnosis of lymphoma in cats, a species with a high prevalence of extranodal forms.^{14,23}

Given these premises, the aim of our retrospective study was to describe FC results in a case series of suspected canine extranodal lymphomas. We initially questioned whether the likelihood of obtaining a diagnostic sample varied between nodal and extranodal forms, and among different extranodal sites. Then, we describe the FC features of a series of cases compatible with extranodal lymphomas, including detailed phenotype and putative tumor burden in blood and bone marrow quantified by FC. For this second aim, we added cases obtained from a second FC database in order to consolidate the dataset.

Materials and methods

Case selection

To assess the likelihood of obtaining diagnostic samples, the FC database of the Department of Veterinary Medicine, University of Milan (Milan, Italy) was examined retrospectively from January 2009 to September 2017, and canine cases were extracted. Cases were included if FC immunophenotyping was requested for a fine-needle aspirate obtained from lesions suspected to

60 be lymphoma at extranodal sites, excluding effusions and lymphoid organs (thymus, spleen, and
61 tonsils). Cases were subsequently excluded if generalized lymphadenopathy was identified or if
62 the nodal status was not reported. For each case, lesion site and FC results were retrieved.

63 FC results were considered consistent with lymphoma if cases fulfilled the following
64 criteria: 1) cytologic evaluation performed prior to FC was compatible or suggestive of round-
65 cell neoplasia, likely of lymphoid origin; 2) well-preserved FC sample, with minimal debris and
66 disrupted cells and minimal hemocontamination; and 3) presence of a dominant lymphoid
67 population sharing a unique morphologic and phenotypic pattern.⁵ Hepatic samples were
68 diagnosed as lymphoma only if infiltrating leukemia was unambiguously ruled out. Acute
69 leukemia was suspected if CD34+ cells were detected, whereas chronic lymphocytic leukemia
70 (CLL) was included in the differential if neoplastic cells had a mature morphologic appearance,
71 were small, CD34-, and CD21+, CD4+, or CD8+ on FC. One dog had a proliferation of small
72 mature CD4+ lymphocytes in the liver and was retained in the study. CLL was excluded in this
73 dog based on the following: slight peripheral blood (PB) lymphocytosis, mild enlargement of
74 hepatic lymph nodes (LNs) with all other LNs within normal limits, and/or death from
75 progressive disease within 15 d of the diagnosis, despite corticosteroid therapy.

76 Control canine cases consisted of suspected nodal lymphomas cases extracted from the
77 same FC database, according to the following inclusion criteria: 1) generalized
78 lymphadenopathy; 2) no extranodal lesions; and 3) LN aspirate sent to the laboratory for FC.

79 In a second step, to consolidate the dataset of samples with FC results compatible with
80 extranodal lymphomas, additional cases were included from the FC database of the Veterinary
81 Teaching Hospital, University of Turin (Grugliasco, Italy). For all cases included in this dataset,
82 the referring veterinarians were contacted to retrieve follow-up data.

83 All of the dogs were privately owned and sampled for diagnostic purposes with the
84 informed consent of the owners. Thus, specific formal approval by the authors' Institution
85 Committee for Animal Care was not required (protocol 1965-2017, Ethical Committee,
86 University of Turin).

87 **Flow cytometry**

88 FC was performed on tissue aspirates collected in tubes containing 1 mL of a liquid medium
89 (either RPMI-1640 or saline solution). Prior to labeling, sample cellularity was assessed with an
90 automated hematology analyzer (XT-2000iV, Sysmex, Kobe, Japan; or ADVIA 120, Siemens
91 Healthcare Diagnostics, Milan, Italy). Also, a visual inspection was done by the operator, in
92 order to assess the gross quality of the sample and decide whether to perform FC labeling.²³

93 Processing for FC was performed as described previously,¹² using different combinations
94 of antibodies (Table 1), with a multicolor approach. For intracellular staining, a permeabilization
95 procedure was performed using either Leucoperm reagents (Serotec, Oxford, UK) following the
96 manufacturer's instructions, or FACS permeabilizing solution 2 (Becton Dickinson, San José,
97 CA) as described.¹² Samples were acquired with a FACSCalibur or a Accuri C6 flow cytometer
98 (Becton Dickinson) and analyzed with Cell Quest or CFlow Plus software (Becton Dickinson).

99 Cells were classified as "small" when showing the same FC morphologic properties
100 (forward-scatter [FSC] and side-scatter [SSC]) of residual normal lymphocytes, and as "large"
101 when 2 distinct lymphoid populations were identifiable based on FSC and SSC (normal small
102 lymphocytes and large neoplastic cells). Cases compatible with lymphoma were classified as B
103 cell if cells from the dominant population expressed CD21, CD79a, or CD79b, and did not
104 express any T-cell marker. T-cell lymphomas were identified when cells expressed CD3, CD5,
105 CD4, CD8, or CytD3, and did not express any B-cell marker. Cases were classified as T cell

106 when the dominant population expressed concomitantly any T-cell marker and CD21 but not
107 CD45.²²

108 When available, PB and/or bone marrow (BM) samples collected into EDTA tubes were
109 also analyzed by FC to quantify the infiltration by putative neoplastic cells. Red blood cells were
110 lysed prior to labeling by means of an erythrocyte lysis buffer containing 8% ammonium
111 chloride. Putative infiltration degree was defined as the percentage of nucleated cells showing
112 the same FC morphologic and phenotypic characteristics of the dominant population identified in
113 the tissue aspirate.

114 **Statistical analysis**

115 For statistical analysis, extranodal lesions were grouped into skin, liver, head and neck, and other
116 sites. Contingency tables were prepared, and the Pearson chi-square test was performed to assess
117 possible different likelihoods of obtaining a sample suitable for FC assessment by comparing
118 nodal and extranodal samples, and extranodal samples among different sites. Analyses were then
119 performed (Statistics v.20.0, SPSS, Chicago, IL). Significance was set at $p \leq 0.05$.

120 **Results**

121 **Diagnostic yield of nodal and extranodal samples from different sites**

122 Seventy-three canine extranodal samples were retrieved from the FC database of the Department
123 of Veterinary Medicine (Milan). Poor cellularity, excessive hemodilution, and high content of
124 debris and disrupted cells were the principal limitations affecting the quality of the samples and
125 the suitability for FC assessment. Lesions were located in the skin in 33 cases (45%), liver in 13
126 (18%), head and neck in 8 (11%), and elsewhere in 19 (26%), including kidney ($n = 5$),
127 cerebrospinal fluid ($n = 4$), bowel ($n = 2$), lung ($n = 2$), and 1 each of the following: anal sac,

128 joint, pancreas, stomach, peritoneum, and adrenal gland. Forty-one samples (56%) were
129 considered suitable for FC assessment.

130 Considering only the acceptable samples, FC results were compatible with histiocytic
131 proliferative disease in 3 cases (7%), and 1 case (2%) was suspect for infiltration by CLL
132 because a unique population of small mature CD8+ lymphocytes was identified invading liver
133 and bone marrow. A mixed population of small lymphocytes was present in 9 cases (22%), with
134 large cells accounting for <5% of the population in all instances; because none of the lymphoid
135 population was dominant, these samples were considered negative for hematopoietic neoplasia.
136 In the remaining 28 (68%) cases, a dominant cell population was identifiable; these samples
137 were considered consistent with extranodal lymphoma (Table 2).

138 The control population consisted of 894 dogs in which nodal lymphoma was suspected. A
139 total of 820 cases (92%) were suitable for FC assessment, and 743 cases (91%) were consistent
140 with lymphoma (Table 2).

141 The proportion of samples suitable for FC assessment varied significantly between
142 extranodal and nodal samples ($p < 0.001$), with the latter being more likely acceptable (odds
143 ratio: 8.64; 95% confidence interval: 5.14–14.5). The difference was not significant among
144 different extranodal sites ($p > 0.05$).

145 **FC features of extranodal lymphomas**

146 To consolidate the dataset, 11 additional samples consistent with extranodal lymphoma obtained
147 from the Veterinary Teaching Hospital (Turin) were included. Thus, the FC features of 39 cases
148 were described (Table 3). The skin was involved in 16 (41%) dogs, the liver in 9 (23%), and the
149 head and neck in 5 (13%). The remaining 9 (23%) cases involved kidneys ($n = 3$), bowel ($n = 2$),
150 and 1 each in urinary bladder, tibia, lung, and peritoneum. A round cell tumor was suspected in

151 all cases based on cytologic examination performed prior to FC; histopathology and
152 immunohistochemistry were performed in 9 cases, with a diagnosis of lymphoma.

153 A dominant population of cells was easily identified based on morphologic features (large
154 cells) in 25 (64%) cases. Phenotypic aberrancies were commonly found in T-cell lymphomas (20
155 cases, 80%), including lack of expression of both CD4 and CD8, discordant expression of pan-
156 T-cell markers, expression of CD8 but not of CD3 or CD5, lack of expression of CD45, and
157 expression of CD21. In particular, 10 (50%) of these cases had more than 1 phenotypic
158 aberrancy. Among suspected B-cell lymphomas, only 1 case (7%) was CD21-CD79+.
159 Interestingly, KIT expression was detected in 2 B-cell lymphomas (14%). In 5 (13%) cases, the
160 dominant population was cells with FC properties overlapping those of normal small
161 lymphocytes; the dominant cells were also small and showed no phenotypic aberrancies, with the
162 exception of KIT expression in 1 dog.

163 Concerning phenotype distribution, the skin was more commonly affected by
164 proliferation of CD45+CD4-CD8- T cells ($n = 7$, 44%), the liver by B cells ($n = 5$, 56%), and
165 the head and neck by CD45- T-cell lymphomas ($n = 3$, 60%). B-cell proliferation was more
166 common in the miscellaneous group ($n = 5$, 56%).

167 PB and BM samples were available for FC analysis in 25 and 22 cases, respectively.
168 Putative neoplastic lymphoid cells were present in 15 (60%) PB samples, with a mean infiltration
169 level of $20 \pm 24\%$ (median: 8%, min.-max.: 1-79%) and in 9 (41%) BM samples, with a mean
170 infiltration level of $14 \pm 17\%$ (median: 4%, min.-max.: 1-37%).

171 Follow-up data were retrieved for 14 dogs. Lesions involved the skin in 8 cases; head and
172 neck in 3; liver, bowel, and kidney in 1 each. Ten (71%) dogs were treated with corticosteroids
173 or different chemotherapy protocols and died because of progressive disease, with a median

174 survival of 77 d (range: 15–249 d). Two (14%) dogs with cutaneous lymphoma were treated with
175 multi-agent chemotherapy obtaining partial and complete remission and died as a result of
176 lymphoma-unrelated causes after 113 and 603 d, respectively. Two (14%) dogs with tongue
177 lesions received chemotherapy and obtained complete clinical remission; one was still alive after
178 579 d, the second one died because of lymphoma-unrelated causes after 427 d.

179 **Discussion**

180 Approximately half of the suspected extranodal lymphomas included in our study had an intra-
181 abdominal location, and most of them involved the liver, which caused the dominant health issue
182 in these dogs. This was unexpected, considering the low frequency reported in the literature for
183 this lymphoma presentation in dogs.¹⁶ Conversely, cutaneous lymphomas are common, with the
184 epitheliotropic subtype alone accounting for 3–8% of all canine lymphomas.⁴ This result can be
185 explained by the preferential use of less invasive sampling when intra-abdominal lesions are
186 present, whereas the skin is more likely to be biopsied. Nevertheless, the presence of mild
187 peripheral lymphadenomegaly may have gone unnoticed by the clinicians in some cases, thereby
188 raising the number of dogs with suspected primary hepatic lymphoma.

189 The phenotypes of the dominant cell population identified by FC in our study showed a
190 different distribution between nodal and extranodal samples. As described previously, B cell is
191 the most common phenotype in nodal lymphomas,^{2,36} and CD4+ phenotype is the most frequent
192 in multicentric T-cell lymphomas.⁹ Cutaneous lymphomas are more frequently composed of
193 CD4– T cells with variable expression of CD8,^{4,24} whereas cutaneous B-cell lymphomas are less
194 common.^{7,8}

195 Interestingly, the prevalence of a CD45– T-cell phenotype did not vary between nodal
196 and suspected extranodal lymphomas. This phenotype is considered highly suggestive of T-zone

197 lymphoma (TZL) in dogs.^{22,30,33} Three of 4 CD45– T-cell extranodal lymphomas in our case
198 series were located within the oral cavity, and 2 of them involved the tongue. This has been
199 described as a novel presentation of canine TZL.¹⁵ The remaining CD45– T-cell lymphoma was
200 a peritoneal mass near the spleen. The neoplastic cells in the mass were large based on the FC
201 morphologic scattergram, and expressed CD3 and CD8. Unfortunately, CD5 and CD21
202 expression was not tested, and cytologic specimens were not available for review. Because cells
203 were large, it is likely that the final diagnosis in this case was peripheral T-cell lymphoma rather
204 than TZL.

205 We found that <60% of submitted extranodal samples were suitable for FC assessment,
206 which is considerably lower than the results obtained for nodal samples (>90%). Several
207 possibilities might explain this difference, including discomfort in the sampling procedure given
208 the intra-abdominal location of the lesions, different size and characteristics of the lesion
209 (extranodal tissue thickening vs. gross LN enlargement), different cellular exfoliation from
210 tissues other than nodes, and higher content of debris and parenchymal cells in the extranodal
211 lesions. Further, a greater variety of lesions can affect extranodal sites compared to LN, which is
212 confirmed by the relatively high frequency (>20%) of non-hematopoietic tumors that were
213 encountered among extranodal samples. Finally, other pre-analytical factors may have influenced
214 the diagnostic yield of the samples, as has been demonstrated in canine LNs⁵ and nodal and
215 extranodal lesions in cats.²³ Regardless of the underlying causes, the risk of obtaining samples
216 not suitable for FC assessment should be taken into account in the diagnostic workup of
217 extranodal lesions.

218 Phenotypic aberrancies were commonly found in the cases consistent with extranodal
219 lymphomas. Phenotypic aberrancies are defined as gross antigen deletions, expression of

220 antigens that are not normally present on the cells, and co-expression or loss of both CD4 and
221 CD8.¹⁸ In our case series, phenotypic aberrancies were detected in 20 (80%) samples consistent
222 with T-cell lymphoma and in 1 (7%) consistent with B-cell lymphoma. Phenotypic aberrancies
223 are generally more common in T-cell than B-cell nodal lymphomas,¹² likely because a larger
224 panel of T-cell antigens is tested routinely, allowing more detailed phenotypic definition.
225 Conversely, only a few canine B-cell antigens can be tested by FC in dogs. CD79 labeling
226 requires permeabilization, and is expensive, time-consuming, and is generally omitted if the cells
227 of interest are CD21+. This approach further reduces the number of B-cell antigens tested.

228 We recorded KIT expression in 2 cases. KIT expression has been investigated in canine
229 nodal lymphomas,¹³ and the activity of masitinib, a tyrosine kinase inhibitor, has been
230 documented in 10 dogs with epitheliotropic lymphoma, although none of them expressed KIT by
231 immunohistochemistry.¹⁷ Further studies are needed to assess the possible clinical and
232 therapeutic significance of KIT expression in canine extranodal lymphomas.

233 We found high percentages of putative neoplastic cells in PB and BM of dogs with
234 suspected extranodal lymphoma, which supports the utility of assessing infiltration of these 2
235 tissues in extranodal lymphomas. We did not consider clinical and prognostic significance in our
236 study. FC has optimal diagnostic performance in staging canine large B-cell lymphoma,²⁸ but
237 validation in other lymphoma subtypes is still lacking. However, large cells or phenotypic
238 aberrancies are not present in PB from healthy dogs.¹⁰ Thus, we have confidence in the
239 specificity of the results obtained for FC staging for most of the cases included in our study. A
240 small subset of cases had a dominant population of small cells without phenotypic aberrancies in
241 the primary extranodal lesion; thus, the cells with the same morphologic and phenotypic
242 properties detected in PB and BM may represent either infiltrating neoplastic cells or

265 The authors declared no potential conflicts of interest with respect to the research, authorship,
266 and/or publication of this article.

267 **Funding**

268 The authors received no financial support for the research, authorship, and/or publication of this
269 article.

270 **References**

- 271 1. Aresu L, et al. Minimal residual disease detection by flow cytometry and PARR in lymph
272 node, peripheral blood and bone marrow, following treatment of dogs with diffuse large
273 B-cell lymphoma. *Vet J* 2014;200:318–324.
- 274 2. Aresu L, et al. Canine indolent and aggressive lymphoma: clinical spectrum with histologic
275 correlation. *Vet Comp Oncol* 2015;13:348–362.
- 276 3. Cannon CR, Richardson D. Value of flow cytometry with fine needle aspiration biopsy in
277 patients with head and neck lymphoma. *Otolaryngol Head Neck Surg* 2000;123:696–699.
- 278 4. Chan CM, et al. Clinical outcome and prognosis of dogs with histopathological features
279 consistent with epitheliotropic lymphoma: a retrospective study of 148 cases (2003–
280 2015). *Vet Dermatol* 2018;29:154–e59.
- 281 5. Comazzi S, et al. Effects of pre-analytical variables on flow-cytometric diagnosis of canine
282 lymphoma: a retrospective study (2009–2015). *Vet J* 2018;232:65–69.
- 283 6. Comazzi S, Gelain ME. Use of flow cytometric immunophenotyping to refine the cytological
284 diagnosis of canine lymphoma. *Vet J* 2011;188:149–155.
- 285 7. Day MJ. Immunophenotypic characterization of cutaneous lymphoid neoplasia in the dog and
286 cat. *J Comp Pathol* 1995;112:79–96.

- 287 8. De Mello Souza CH, et al. Immunohistochemical detection of retinoid receptors in tumors
288 from 30 dogs diagnosed with cutaneous lymphoma. *J Vet Intern Med* 2010;24:1112–
289 1117.
- 290 9. Deravi N, et al. Specific immunotypes of canine T-cell lymphoma are associated with
291 different outcomes. *Vet Immunol Immunopathol* 2017;191:5–13.
- 292 10. Faldyna M, et al. Lymphocyte subsets in peripheral blood of dogs—a flow cytometric study.
293 *Vet Immunol Immunopathol* 2001;82:23–37.
- 294 11. Foglia Manzillo V, et al. Extranodal gamma-delta T-cell lymphoma in a dog with
295 leishmaniasis. *Vet Clin Pathol* 2008;37:298–301.
- 296 12. Gelain ME, et al. Aberrant phenotypes and quantitative antigen expression in different
297 subtypes of canine lymphoma by flow cytometry. *Vet Immunol Immunopathol*
298 2008;121:179–188.
- 299 13. Giantin M, et al. Evaluation of tyrosine-kinase receptor c-kit mutations, mRNA and protein
300 expression in canine lymphoma: might c-kit represent a therapeutic target? *Vet Immunol*
301 *Immunopathol* 2013;154:153–159.
- 302 14. Guzera M, et al. The use of flow cytometry for immunophenotyping lymphoproliferative
303 disorders in cats: a retrospective study of 19 cases. *Vet Comp Oncol* 2016;14:40–51.
- 304 15. Harris LJ, et al. Clinicopathologic features of lingual canine T-zone lymphoma. *Vet Comp*
305 *Oncol* 2018;16:131–139.
- 306 16. Hirose N, et al. A retrospective histopathological survey on canine and feline liver diseases at
307 the University of Tokyo between 2006 and 2012. *J Vet Med Sci* 2014;76:1015–1020.
- 308 17. Holtermann N, et al. Masitinib monotherapy in canine epitheliotropic lymphoma. *Vet Comp*
309 *Oncol* 2016;14:127–135.

310 18. Jamal S, et al. Immunophenotypic analysis of peripheral T-cell neoplasms. A multiparameter
311 flow cytometric approach. *Am J Clin Pathol* 2001;116:512–526.

312 19. Jaso J, et al. CD5-positive mucosa-associated lymphoid tissue (MALT) lymphoma: a
313 clinicopathologic study of 14 cases. *Hum Pathol* 2012;43:1436–1443.

314 20. Joetzke AE, et al. Flow cytometric evaluation of peripheral blood and bone marrow and fine-
315 needle aspirate samples from multiple sites in dogs with multicentric lymphoma. *Am J*
316 *Vet Res* 2012;73:884–893.

317 21. Marconato L, et al. Assessment of bone marrow infiltration diagnosed by flow cytometry in
318 canine large B-cell lymphoma: prognostic significance and proposal of a cut-off value.
319 *Vet J* 2013;197:776–781.

320 22. Martini V, et al. Flow-cytometric detection of phenotypic aberrancies in canine small clear
321 cell lymphoma. *Vet Comp Oncol* 2015;13:281–287.

322 23. Martini V, et al. Flow cytometry for feline lymphoma: a retrospective study regarding pre-
323 analytical factors possibly affecting the quality of samples. *J Feline Med Surg*
324 2018;20:494–501.

325 24. Moore PF, et al. Canine epitheliotropic cutaneous T-cell lymphoma: an investigation of T-
326 cell receptor immunophenotype, lesion topography and molecular clonality. *Vet*
327 *Dermatol* 2009;20:569–576.

328 25. Poggi A, et al. Prognostic significance of Ki67 evaluated by flow cytometry in dogs with
329 high grade B-cell lymphoma. *Vet Comp Oncol* 2017;15:431–440.

330 26. Rao S, et al. Class II major histocompatibility complex expression and cell size
331 independently predict survival in canine B-cell lymphoma. *J Vet Intern Med*
332 2011;25:1097–1105.

- 333 27. Reggeti F, Bienzle D. Flow cytometry in veterinary oncology. *Vet Pathol* 2011;48:223–235.
- 334 28. Riondato F, et al. Analytical and diagnostic validation of a flow cytometric strategy to
335 quantify blood and marrow infiltration in dogs with large B-cell lymphoma. *Cytometry B*
336 *Clin Cytom* 2016;90:525–530.
- 337 29. Riondato F, et al. Identification of peripheral blood involvement in dogs with large B-cell
338 lymphoma: comparison of different methods. *Res Vet Sci* 2017;115:288–293.
- 339 30. Rout ED, Avery PR. Lymphoid neoplasia. Correlation between morphology and flow
340 cytometry. *Vet Clin North Am Small Anim Pract* 2017;47:53–70.
- 341 31. Rütgen B, et al. Cutaneous T-cell lymphoma—Sézary syndrome in a Boxer. *Vet Clin Pathol*
342 2016;45:172–178.
- 343 32. Schniederjan SD, Osunkoya AO. Lymphoid neoplasms of the urinary tract and male genital
344 organs: a clinicopathological study of 40 cases. *Mod Pathol* 2009;22:1057–1065.
- 345 33. Seelig DM, et al. Canine T-zone lymphoma: unique immunophenotypic features, outcome
346 and population characteristics. *J Vet Intern Med* 2014;28:878–886.
- 347 34. Stacchini A, et al. Extranodal lymphoproliferative processes and flow cytometry. *Acta Cytol*
348 2016;60:315–325.
- 349 35. Tarrant JM. The role of flow cytometry in companion animal diagnostic medicine. *Vet J*
350 2005;170:278–288.
- 351 36. Valli VE, et al. Canine lymphomas: association of classification type, disease stage, tumor
352 subtype, mitotic rate, and treatment with survival. *Vet Pathol* 2013;50:738–748.
- 353 37. Yu JB, et al. Identification of immunophenotypic subtypes with different prognoses in
354 extranodal natural killer/T-cell lymphoma, nasal type. *Hum Pathol* 2014;45:2255–2262.

355 38. Zeppa P, et al. Fine needle aspiration cytology and flow cytometry immunophenotyping of
356 non-Hodgkin lymphoma: can we do better? *Cytopathology* 2010;21:300–310.
357

358 **Table 1.** Antibodies used for flow cytometric immunophenotyping of suspected canine
 359 lymphoma samples.

Target molecule	Antibody clone	Source	Specificity
CD45	YKIX716.13	Serotec, Oxford, UK	All leukocytes
CD3	CA17.2A12	Serotec	T cells
CD5	YKIX322.3	Serotec	T cells
CycD3	CD3-12	Serotec	T cells
CD4	YKIX302.9	Serotec	T-helper cells and neutrophils
CD8	YCATE55.9	Serotec	T-cytotoxic cells
CD21	CA2.1D6	Serotec	Mature B cells
CD79a	HM57	Serotec	B cells
CD79b	AT107-2	Serotec	B cells
CD34	1H6	BD Pharmingen, San José, CA	Precursors
MHC II	YKIX334.2	Serotec	Lymphocytes, monocytes
CD117	ACK45	BD Pharmingen	Precursors and mast cells

360

361 **Table 2.** Flow cytometric immunophenotype of 771 dogs with nodal or suspected extranodal lymphoma.

	Lymphoma phenotype						Total
	B cell	T cell					
		CD45+CD4+CD8-	CD45+CD4-CD8+	CD45+CD4+CD8+	CD45+CD4-CD8-	CD45-	
Nodal	550 (74)	82 (11)	21 (3)	5 (1)	22 (3)	63 (9)	743 (100)
Extranodal	8 (29)	1 (4)	9 (32)	0 (0)	7 (25)	3 (11)	28 (100)
Total	558 (72)	83 (11)	30 (4)	5 (1)	29 (4)	66 (9)	771 (100)

362 Numbers in parentheses are percentages.

363

364 **Table 3.** Flow cytometric features of the dominant lymphoid population in extranodal lesions compatible with lymphoma from 39
365 dogs.

Case ID	Lesion site	Cell size	Dominant population (%)	CD45	CD3	CD5	CD4	CD8	CD21	CD79 (a or b)	CD34	MHC II	KIT	CycD3	PB lymphocyte count ($\times 10^9/L$)	Putative infiltration (%)	
																PB	BM
1	Skin	Large	63	+	+	-	-	-	-	-	ND	ND	ND	ND	4.3	-	-
2*	Skin	Large	89	+	-	+	-	-	-	ND	-	ND	-	ND	1.1	-	-
3*	Skin	Small	69	+	-	+	-	-	-	ND	ND	ND	ND	ND	0.8	4	-
4	Skin	Small	98	+	+	+	-	-	-	ND	ND	ND	-	ND	7.6†	30	45
5	Skin	Large	80	+	+	-	-	-	-	ND	-	ND	ND	ND	ND	ND	ND
6	Skin	Large	93	+	+	-	-	-	-	ND	-	+	-	ND	ND	ND	ND
7	Skin	Large	70	+	-	-	-	-	-	-	-	+	ND	+	ND	ND	ND
8	Skin	Small	87	+	-	-	-	+	-	ND	ND	ND	-	ND	1.2	-	ND
9*	Skin	Large	91	+	+	+	-	+	-	ND	ND	ND	ND	ND	0.7	5	1
10	Skin	Small	64	+	ND	+	-	+	-	ND	ND	ND	ND	ND	ND	ND	ND
11*	Skin	Large	87	+	ND	-	-	+	-	ND	ND	ND	-	ND	1.3	-	-
12*	Skin	Small	64	+	ND	-	-	+	-	ND	ND	ND	ND	ND	2.1	-	-
13	Skin	Small	71	+	-	-	-	-	+	+	-	ND	+	ND	50.4†	30	ND
14	Skin	Small	81	+	ND	-	-	-	+	+	-	ND	ND	-	0.7	4	ND
15	Skin	Large	99	+	ND	-	ND	ND	+	ND	-	+	ND	ND	ND	ND	ND
16	Skin	Large	99	+	-	-	-	-	+	ND	-	ND	ND	ND	1.7	2	3
17	Liver	Large	82	+	ND	-	-	-	+	ND	-	ND	ND	ND	85.2†	79	ND
18	Liver	Large	88	+	ND	ND	-	ND	+	ND	-	+	ND	ND	0.9	-	ND
19	Liver	Large	81	+	-	-	-	-	+	+	-	ND	ND	ND	1.1	-	-
20	Liver	Large	86	+	ND	-	ND	ND	+	ND	-	ND	ND	ND	1.9	1	2
21	Liver	Large	94	+	-	-	-	-	+	+	-	ND	ND	ND	ND	ND	ND
22	Liver	Large	80	+	+	-	-	+	ND	ND	-	ND	ND	ND	ND	ND	ND
23	Liver	Large	85	+	-	-	-	+	-	-	ND	ND	ND	ND	0.8	17	36
24	Liver	Small	76	+	+	+	+	-	-	ND	-	ND	ND	ND	6.2†	23	ND
25	Liver	Large	78	+	-	+	-	-	-	-	-	ND	ND	ND	1.5	-	-
26*	Tongue	Small	87	-	+	+	+	-	+	-	-	ND	ND	ND	7.9†	70	ND
27	Tongue	Small	78	-	+	+	-	+	+	ND	-	ND	-	ND	1.3	4	-
28	Oral mucosa	Small	64	-	ND	+	ND	ND	-	ND	-	ND	ND	ND	1.1	16	8
29*	Lip	Small	97	+	ND	+	-	-	-	ND	ND	ND	ND	ND	ND	ND	-
30	Gum	Small	70	+	+	+	-	+	-	ND	-	ND	ND	ND	2.3	2	2
31	Kidney	Large	89	+	-	-	-	-	+	+	-	ND	+	ND	ND	ND	28
32	Kidney	Large	96	+	-	-	-	-	+	+	-	ND	ND	-	1.5	-	-
33	Kidney	Large	62	+	ND	-	ND	+	-	ND	ND	ND	ND	ND	2.2	-	-

34*	Bowel	Large	97	+	-	ND	ND	ND	+	ND	ND	ND	ND	ND	ND	ND	ND
35	Bowel	Large	66	+	+	-	-	-	-	-	ND	ND	ND	ND	ND	ND	ND
36	Urinary bladder	Large	98	+	ND	ND	ND	ND	+	+	-	ND	ND	ND	ND	ND	ND
37	Tibia	Small	65	+	-	-	-	-	-	+	-	ND	-	ND	ND	ND	-
38*	Lung	Large	97	+	+	-	-	+	-	ND	-	ND	-	ND	ND	ND	-
39	Peritoneum	Large	86	-	+	ND	ND	+	ND	ND	-	ND	ND	ND	1.6	8	4

366 - = negative; BM = bone marrow; ND = not done; PB = peripheral blood.

367 * Histopathology performed.

368 † Exceeding laboratory upper reference limit.