

**The assessment of the carbonaceous component in black crusts damaging the  
stone surfaces of historical monuments  
(from Trevi fountain Roma to square San Marco Venice)**

V.Comite<sup>a</sup>, P. Fermo<sup>a</sup>, M.F. La Russa<sup>b</sup>, S.A Ruffolo<sup>b</sup>

<sup>a</sup> *Dipartimento di Chimica, Università degli Studi di Milano, Via Golgi 19, 20133 Milano, Italy;*

<sup>b</sup> *Università della Calabria, Dipartimento di Biologia, Ecologia e Scienze della Terra (DiBEST),  
Via Pietro Bucci 87036, Arcavacata di Rende, CS, Italy*

*E-mail of the presenting author: [valeria.comite@gmail.com](mailto:valeria.comite@gmail.com)*

The issue of conservation of the monumental heritage is mainly related to atmospheric pollution that causes the degradation of stone surfaces. Black crusts can be formed as a result of different chemical and physical reactions between the stone surface and environmental factors (such as gaseous pollutants and aerosol particulate matter, PM). These black layers present on the stone monuments reflect the composition of the aerosol particulate matter to which the surfaces are exposed. In particular elemental carbon (EC, also known as black carbon, typically emitted by combustion processes) is the PM component responsible for the characteristic black color of the crusts where it is embedded together with calcium sulphate due to the conversion of calcium carbonate, the main constituent of the stone. Organic carbon (OC) represents the other carbonaceous component of PM and it is present in the black crusts, too. It is of both primary or secondary origin and is linked to numerous sources (traffic, heating plants, biomass burning, etc.).

A deep knowledge of the crust composition in terms of OC and EC optical properties is mandatory in order to get information on the sources responsible for the surface darkening.

OC/EC in PM samples are generally quantified by a reference method (TOT, Thermal Optical Transmittance) not suitable for the analysis of these components in the crusts. A new approach for OC/EC quantification based on a thermal protocol and including CHN and TGA analyses, has been set-up. The method validation has been performed analyzing suitable reference standard samples prepared by mixing different chemical species in order to simulate the composition of the black crusts present on the monument surfaces. Real samples of black crusts coming from historical monuments placed in Rome, Milan and Venice have been analyzed.