



5th - 7th November 2018

Milan, Italy

[HOME](#)

[PROGRAM](#)

[REGISTRATION](#)

[MORE](#)

Spread the meeting



Duccio Cavalieri (University of Firenze, Italy)
Jin-ichi Inokuchi (Tohoku Medical and Pharmaceutical University, Japan)
Hiroshi Kiyono (University of Tokyo, Japan)
Jerry M. Wells (Wageningen, The Netherlands)

Speakers

This international event is organized as the final event within the European MSCA-ETN project TOLLerant (www.tollerant.eu) and it will focus on molecular aspects of microbiota-host interaction in health and disease.



Partner

BEST4FOOD

Bicocca cEnter of Science and Technology for FOOD



Con il patrocinio

UNIMIB

Università degli Studi di Milano – Bicocca

EuChemS

European Chemical Society

ANSISA

Associazione Nazionale Scienze dell'Alimento



P38 – Soil microbiome before and after fertilization with bovine manure

Penati M.¹, Grilli G.¹, Carraro L.², Cagnardi P.³, Piccirillo A.²

¹Università degli Studi di Milano, Dept. Veterinary Medicine, Milan, Italy.

²Università degli Studi di Padova, Dept. Comparative Biomedicine and Food Science, Legnaro, Padova, Italy

³Università degli Studi di Milano, Dept. Health, Animal Sciences and Food Safety, Milan, Italy.

martinapenati@icloud.com

Background: Dairy manure is commonly used for the fertilization of agricultural soils, and it includes the intestinal microbial community influenced by the feed and possible drug residues used for therapy. The excessive use of antimicrobials in animal husbandry and subsequent land application of animal wastes may introduce massive quantities of antimicrobials and resistant bacteria into the soil environment.

Objective: The objective of this paper is to evaluate the possible change of the soil microbiome after fertilization with manure coming from intensive dairy farms.

Methodology: Samples of manure and soil (before and after 30 days from fertilization) were collected from intensive dairy farms.

Microbioma of samples was determined by a culture independent approach based on NGS sequencing. All samples were also screened for detecting and quantifying the most commonly used antimicrobials (i.e. beta-lactams, fluoroquinolones, polymyxins and macrolides) by HPLC methods.

Results: All samples were negative for the antimicrobials screened. The manure microbioma was homogenous among different farms and regions of Northern Italy, but very different from the soil one. In manure, a higher presence of *Porphyromonadaceae*, *Lachnospiraceae*, *Clostridiaceae*, *Pseudomonadaceae* and *Ruminococcaceae* families was observed, whereas in soil the *Chitinophagaceae* family was the most commonly detected. After fertilization, the microbioma composition of soil was not changed, however the absence of drugs concentration in manure may have had a role in this.

Conclusions / Implications for practice: This is one of the first study evaluating the modifications of the microbial communities of dairy manure and agricultural soil before and after fertilization in Northern Italy.

References

- Hodkinson B. P., Grice E. A. (2014), *Next-Generation Sequencing: a review of technologies and tools for wound microbiome research*. *Advances In Wound Care*. 4: 50-58.
- Hu H.-W., Han X.-M., Shi X.Z.n, Wang J.T., Ha2 L.Li, Chen D., He J.Z. (2016), *Temporal changes of antibiotic resistance genes and bacterial communities in two contrasting soils treated with cattle manure*. *FEMS Microbiology Ecology*, Vol. 92, No. 2.
- Jechalke S., Heuer H., Siemens J., Amelung W., Smalla K. (2014), *Fate and effects of veterinary antibiotics in soil*. *Trends in Microbiology*, Vol. 22, No. 9.