

1 **Membrane restructuring following in situ sialidase digestion of gangliosides: complex model**  
2 **bilayers by synchrotron radiation reflectivity.**

3

4 Valeria Rondelli<sup>a</sup>, Paola Brocca<sup>a</sup>, Giovanna Fragneto<sup>b</sup>, Jean Daillant<sup>c</sup>, Cristina Tringali<sup>a</sup>, Laura  
5 Cantu<sup>'a\*</sup>, Elena Del Favero<sup>a</sup>

6 <sup>a</sup> Dept. of Medical Biotechnologies and Traslational Medicine, University of Milano, LITA, Via  
7 F.lli Cervi, 93. 20090 Segrate, Italy.

8 <sup>b</sup> Institut Laue-Langevin, 71 avenue des Martyrs, BP 156, 38000 Grenoble Cedex, France

9 <sup>c</sup> Synchrotron SOLEIL, L'Orme des Merisiers, Saint-Aubin - BP 48, 91192 Gif-sur-Yvette Cedex,  
10 France

11 Corresponding author:

12 Laura Cantu'

13 Dept. of Medical Biotechnologies and Traslational Medicine,

14 University of Milano,

15 LITA, Via F.lli Cervi 93

16 20090 Segrate,

17 Italy

18 +39 0250330362

19 laura.cantu@unimi.it

20

21 **Keywords:** Gangliosides; NEU2; X-Ray reflectometry; single model membrane; biomembrane;

22 GEMs

23

1

## 2 **Abstract**

3 Synchrotron radiation reflectometry was used to access the transverse structure of model  
4 membranes under the action of the human sialidase NEU2, down to the Ångström length scale.  
5 Model membranes were designed to mimic the lipid composition of so-called Glycosphingolipids  
6 Enriched Microdomains (GEMs), which are membrane platforms specifically enriched in  
7 cholesterol and sphingolipids, where also typical signalling molecules are hosted. Gangliosides,  
8 glycosphingolipids containing one or more sialic acid residues, are asymmetrically embedded in  
9 GEMs, in the outer membrane leaflet. There, gangliosides are claimed to directly interact with  
10 growth-factor receptors, modulating their activation and then the downstream intracellular  
11 signalling pathways. Thus, membrane dynamics and signalling could be strongly influenced by the  
12 activity of enzymes regulating the membrane ganglioside composition, including sialidases. Our  
13 results, concerning the structure of single membranes undergoing *in-situ* enzymatic digestion, show  
14 that the outcome of the sialidase action is not limited to the emergence of lower-sialylated  
15 ganglioside species. In fact, membrane reshaping occurs, involving a novel arrangement of the  
16 headgroups on its surface. Thus, sialidase activity reveals to be a potential tool to dynamically  
17 control the structural properties of the membrane external leaflet of living cells, influencing both the  
18 morphology of the close environment and the extent of interaction among active molecules  
19 belonging to signalling platforms.

20

21

## 22 **1. Introduction**

1 Signal transduction, cell migration, endocytosis and exocytosis of proteins and lipids are largely  
2 influenced by the dynamic structure of the plasma membrane and by the local aggregation of  
3 specific lipids and proteins [1-4]. The assembly of receptors and associated transducing elements  
4 typically occur in the lipid bilayer matrix of membranes. Specific lipid classes, like sphingolipids,  
5 are driving agents for membrane partition into functional micro-domains [5,6]. Membrane micro-  
6 domains enriched in cholesterol and sphingolipids, including glycosphingolipids, leading to a liquid  
7 ordered phase, are usually identified by the term GEMs (Glycosphingolipid Enriched Membrane  
8 micro-domains) [7]. Gangliosides and cholesterol have been found to act synergistically, forcing a  
9 preferential redistribution of components across the membrane and contributing to the super-  
10 structuring of their environment [8-10].

11 Moreover, specific lipid mixtures have been found in association with particular types of signalling  
12 molecules. To this regard, the FAS/CD95 receptor is typically found in association with the  
13 ceramide/sphingomyelin/gangliosides complexes, while it has been reported that the epidermal  
14 growth factor receptor (EGFR) is mostly associated with the  
15 cholesterol/sphingomyelin/gangliosides ones [11]. Based on recent studies, the emerging concept is  
16 that the local environment of any protein within membrane microdomains may modulate its  
17 propensity to interact with other proteins [12]. Within GEMs, gangliosides appear to play a  
18 multifaceted role, entangling structural and functional properties. In particular, the interactions  
19 between gangliosides and receptors have been demonstrated to be crucial to modulate downstream  
20 signalling pathways. In fact, the interaction between ganglioside GM3 and multi N-  
21 acetylglucosamine (GlcNAc) residues of N-linked glycans belonging to EGFR down-regulates the  
22 activation of the receptor [13]. Moreover, the interaction with gangliosides GM3 and GM2 has  
23 shown to inhibit the activation of the hepatocyte growth factor receptor, cMet [14].

24 In this view, the remodelling of gangliosides in plasma membranes could deeply alter cell  
25 signalling, and enzymes regulating ganglioside metabolism could play a key role in this context.

1 Sialidases (EC 3.2.1.18), which catalytically remove sialic acid from different sialo-glyco-  
2 conjugates, including gangliosides [15], have been demonstrated to be involved in these  
3 mechanisms.


4 In mammals, four sialidases (NEU1, NEU2, NEU3, and NEU4) are present. They are involved in  
5 several key physiological events [16] and their deregulation appears to be related to cancer  
6 transformation [17-23]. The typical plasma membrane-associated sialidase NEU3 shows strict  
7 substrate specificity for gangliosides [16] and is prevalently associated with GEMs [24]. Through  
8 the regulation of membrane ganglioside composition, NEU3 modulates the activation of membrane  
9 receptors including EGFR [19,25], androgen receptor [26], and  $\beta$ 1 integrin trafficking [23].  
10 Nonetheless, recently, it has been demonstrated that NEU3 is not the unique sialidase present in the  
11 plasma membrane but also other sialidases, typically located in other cellular districts, can move to  
12 the cell surface and may be involved in many events. Lysosomal sialidase, NEU1, can migrate to  
13 the plasma membrane in different cells, like T lymphocytes [27], macrophages [28], and  
14 erythrocytes [29], there accomplishing important physiological tasks. Sialidase NEU2 is cytosolic  
15 and occurred to be found associated to mouse thymus cells [30]. Thus, plasma membrane proteins,  
16 lipids and enzymes involved in their metabolism appear to be dynamic entities that can move,  
17 entering and exiting from dynamical structures. Alterations occurring in such dynamics could be  
18 related to the deregulation of signal transmission or protein trafficking.

19 The study of portions of plasma membrane undergoing enzymatic action is then a very interesting  
20 topic. In particular, nanoscale modifications of the membrane structural properties induced by the  
21 punctual action of an enzyme on one of its components are now made possible by innovative  
22 experimental techniques on model systems. Among them, the synchrotron radiation reflectometry  
23 technique [31] is a powerful tool to access the transverse structure of a membrane down to the  
24 Ångström length scale.

1 In the work presented here, we applied synchrotron radiation reflectometry to perform a structural  
2 study of single ganglioside-containing model membranes under the action of sialidase NEU2. In  
3 particular, our aim was to observe the changes induced in the cross structure of a biomimetic  
4 membrane by the *in situ* chemical conversion of one component, namely GD1a ganglioside, in its  
5 metabolic product. It follows previous works we performed concerning the sialidase digestion of  
6 GD1a substrate in the form of colloidal aggregates in solution, either as pure GD1a micelles, or as  
7 mixed micelles or vesicles containing GD1a. [32]. The chemical formula of GD1a ganglioside is  
8 reported in Figure 1 (MW=1849 Da). Among others, the human sialidase NEU2 (HsNEU2) was  
9 chosen because it is water soluble, and therefore it could be added directly in the measuring cell to  
10 the buffer in contact with a pre-existing and characterized target membrane, without the help of any  
11 solubilizing agent. In addition, HsNEU2 is well characterized as a globular protein of ~42 kDa, it  
12 has been successfully purified and its structure and kinetic properties toward many substrates  
13 (gangliosides, oligosaccharides, and sialoglycoproteins) have been determined [33,34]. Within the  
14 family of neuraminidases, HsNEU2 has been found to be the most specific in substrate [35], being  
15 very efficient in removing the  $\alpha$ 2-3 linked- external sialic acid residues, as in GD1a, GD1b, GT1b,  
16 GM3, sialyllactose, and sialoglycoproteins [34], while showing no activity on *internal* sialic acid  
17 residues, like in GM1 or in GM2. Some residual activity has been detected on monomeric  
18 dispersions of GM1 [34], Moreover, we observed [32] that, while chemically undergoing the  
19 enzymatic action, biomimetic mixed GD1a-phospholipid vesicles in solution display a transient  
20 response to sialidase, followed by full recovery.

21 Structural studies often require simplified model membranes. Still, some important features of  
22 biological membranes should be maintained. When mimicking GEMs, the presence of cholesterol  
23 and the asymmetric disposition of gangliosides are of primary importance. By a combination of  
24 Langmuir-Blodgett and Langmuir-Schaefer techniques [36], we deposited single asymmetric  
25 membranes composed by DPPC, cholesterol and gangliosides (only in the outer layer), on  
26 macroscopic silicon supports, with biomimetic molar ratio and disposition [37].

1 The reflectometry experiment is illustrated in Figure 1. Two replicas of a GD1a containing  
 2 membrane have been built and accurately characterized. Then the target biomimetic membranes  
 3 were submitted to the enzymatic treatment, and, finally, the structural properties of the digested  
 4 membrane were assessed. Moreover, a much larger experimental landscape has been designed, as  
 5 reported in the scheme of Figure 1. On this wider basis, we could finally gain novel and important  
 6 additional suggestions on membrane reshaping following sialidase action.



7  
 8 **Figure 1** Scheme of the experimental design. The GD1a-containing membranes *mA* and *mB* are  
 9 compared to their own *in situ* digested membrane and with other putative-product membranes.  
 10 Membrane C and Membrane D have GM1 and AsialoGM1 gangliosides respectively embedded,  
 11 instead of GD1a ganglioside. The lightning symbol stands for enzyme intervention. Nomenclature  
 12 is according to Chester [38]. At the bottom: structure of ganglioside GD1a.

13  
 14


15 **2. Materials and Methods**

16 **2.1. Membrane preparation**

1 DPPC (1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine) was purchased from Avanti Polar Lipids.  
2 Cholesterol was from Sigma-Aldrich Co. GD1a ( $\alpha$ -Neu5Ac-(2-3)- $\beta$ -Gal-(1-3)- $\beta$ -GalNAc-(1-4)-[ $\alpha$ -  
3 Neu5Ac-(2-3)]- $\beta$ -Gal-(1-4)- $\beta$ -Glc-(1-1)-Cer), GM1 ( $\beta$ -Gal-(1-3)- $\beta$ -GalNAc-(1-4)-[ $\alpha$ -Neu5Ac-(2-  
4 3)]- $\beta$ -Gal-(1-4)- $\beta$ -Glc-(1-1)-Cer) and AsialoGM1 ( $\beta$ -Gal-(1-3)- $\beta$ -GalNAc-(1-4)- $\beta$ -Gal-(1-4)- $\beta$ -Glc-  
5 (1-1)-Cer) were extracted and purified as described by Tettamanti et al. [39] and obtained as sodium  
6 salt powder. HsNEU2 sialidase expression and purification were conducted following the procedure  
7 described by Tringali et al. [33]. In order to obtain stable, single asymmetric membranes deposition,  
8 we used the Langmuir-Blodgett/Langmuir-Schaefer techniques [40]. A brief description of the  
9 technique is reported in *SI Materials and Methods*.

10 After the Langmuir-Schaefer deposition of the planned layers on a UV-Ozone treated [41] single  
11 crystal of silicon ( $5 \times 5 \times 1.5 \text{ cm}^3$ ), the sample were closed on a homemade Teflon holder [42],  
12 directly mounted on the measuring station of ID10B beamline. The sample was mounted  
13 horizontally, face down, as schematically reported in Figure 2.

14



15

16 **Figure 2** Schematic representation of the experimental set-up mounted on ID10B beamline.

17

18

## 19 2.2. Synchrotron Radiation Reflectivity

20 A reflectivity experiment allows to get information about the internal structure of stratified samples

1 [43-46]. Details of the technique are given in *SI* Materials and Methods.

2 Reflectivity measurements were performed on the horizontal reflectometer ID10B at the European  
3 Synchrotron Radiation Facility in Grenoble, France. Data were analyzed using the software Motofit  
4 [47] with a 7 layers model (hydrophilic out, 3 hydrophobic, hydrophilic in, a water layer and the  
5 silicon oxide). Electron densities were calculated starting from Nagle and Tristram-Nagle [48],  
6 Boretta et al. [49] and Greenwood et al. [50].

7

### 8 **3. Results**

9 The scheme of the performed reflectivity experiment is reported in Figure 1. We investigated the  
10 changes induced in model membranes by *in situ* enzymatic removal of sialic acid from parent  
11 gangliosides. Afterwards, we compared the structure of the so-obtained final membrane with that of  
12 a membrane originally built up with the lower-sialylated ganglioside. Moreover, we submitted also  
13 these last membranes to sialidase, in order to test for residual activity on *internal* sialic acid residues  
14 and for possible non specific interaction with the membrane itself.

15 The contrast of a material for synchrotron radiation is related to its electron density, which is higher  
16 for the glycolipids than for other lipids (see *SI*, Table S2) in the region of the sugar heads. Then,  
17 reflectometry allows detecting structural changes of the saccharidic substrates of HsNEU2, within  
18 the membrane, with high sensitivity. To our scope, we prepared and studied four supported  
19 membranes, containing each a different ganglioside species: GD1a (two replicas), GM1 and  
20 AsialoGM1, differing in the number of sialic acid residues in their polar region (see Figure 1). The  
21 lipid matrix consisted of di-palmitoyl-phosphatidyl-choline (DPPC), with a distinct asymmetric  
22 disposition of cholesterol and gangliosides in the two leaflets. The overall components molar ratio  
23 was DPPC: cholesterol: ganglioside = 10: 1.25: 0.5. Gangliosides were deposited in the outer leaflet  
24 of the membranes, exposed to the bulk water, together with 30% of cholesterol. The inner leaflet  
25 composition was therefore DPPC: cholesterol = 10: 1.75 mol, whereas the outer leaflet composition  
26 was DPPC: cholesterol: ganglioside = 10: 0.75: 1 mol. Saturated lipids were used, as they are best




1 suited to prepare full-coverage bilayers. Samples were submitted to annealing (see *SI*) and then  
2 reflectivity spectra were collected at 22°C.

3 The GD1a-containing membrane (sample *mA*) was measured both in pure water and in 150 mM  
4 NaCl aqueous solution as bulk solution. The obtained results indicated that the membrane was  
5 stable and resistant against changes in the ionic strength of the solvent. This feature is highly  
6 desirable for a model membrane, suitable for the study of the interactions between the membrane  
7 and approaching macromolecules, usually dispersed in buffer solutions, or for multistep processes,  
8 requiring buffer replacing. Upon salt addition, a slight increase in membrane thickness was detected  
9 (5.7 nm instead of 5.4 nm in pure water). This is consistent with the screening of electrostatic  
10 repulsive interactions among charged ganglioside headgroups, allowing tighter packing of lipids  
11 within the membrane. Then, for all other membranes, experiments were made in the presence of  
12 150 mM NaCl solution, safely flushed before reflectivity measurement.

13 After characterization of the target membrane *mA*, the solution containing HsNEU2 sialidase was  
14 injected directly in the measuring cell and, after 0.5 h, reflectivity was measured again. Based on  
15 the known kinetics of HsNEU2 [34], such a delay is suitable for the enzymatic reaction to be fully  
16 completed. In Figure 3A, the reflectivity spectra of the membrane *mA* before and after the action of  
17 HsNEU2 are shown, together with the best fitting curve. The density profiles, corresponding to the  
18 best fit of the reflectivity curves and describing the cross structure of the membrane, are reported in  
19 Figure 3B. Vertical dashed lines are drawn to guide the eye and approximately identify different  
20 regions of the deposited raft-mime membrane. Profiles differ mainly in the outer hydrophilic layer  
21 (region *b*), where ganglioside headgroups are exposed to the enzyme. In particular, the action of  
22 HsNEU2 resulted in a significant increase of the contrast, and therefore of the electron density, of  
23 this region.

24 The reproducibility of this result was tested on a second replica of GD1a-containing membrane *mA*,  
25 namely membrane *mB*. The two membranes (*mA* and *mB*) have been independently prepared in  
26 different days, following the same protocol, and are then independent replicas of the same


1 membrane system. The measured reflectivity curves are very nicely superimposed, as can be seen in  
 2 *SI*, Figure S1. After exposition to sialidase, results obtained on the membrane transverse structure of  
 3 the replica membrane *mB* confirmed that the enzymatic digestion produced a pronounced increase  
 4 of the electron density of the external polar region, from  $416 \text{ e}^-/\text{nm}^3$  to  $480 \text{ e}^-/\text{nm}^3$  (as reported in *SI*,  
 5 Table S1), the hydrophobic core being nearly unaffected.



7  
 8 **Figure 3** Reflectivity curves (A) and density profiles (B) of membrane *mA* before (blue dots) and  
 9 after (red triangles) the action of HsNEU2 sialidase. The solvent is a 150 mM NaCl solution.  $T=$   
 10  $22^\circ\text{C}$ . In A) symbols correspond to the experimental data, lines to the best fits, corresponding to the  
 11 profiles in B). In B) vertical dashed lines are drawn to guide the eye to approximately identify 9  
 12 regions, referring to different portions of the interfacial system: bulk solution (*a*), outer hydrophilic  
 13 layer (*b*), outer  $\text{CH}_2$  groups layer (*c*),  $\text{CH}_3$  groups layer (*d*), inner  $\text{CH}_2$  groups layer (*e*), inner  
 14 hydrophilic layer (*f*), water layer (*g*), silicon oxide (*h*), silicon (*i*).

15  
 16  
 17 Then, we wondered whether the “digested” membrane, resulting from the *in situ* action of the  
 18 sialidase on the matrix membrane containing GD1a, had the same structure of a membrane prepared  
 19 by mixing the matrix with the digestion product, *i.e.* GM1. We therefore prepared an asymmetric  
 20 membrane, *mC*, containing GM1 instead of GD1a and measured its reflectivity spectrum. The  
 21 density profile is reported in Figure 4, in comparison with that of the digested-*mB* membrane.

1 Remarkably, *mB* membrane (and replica *mA*) obtained after HsNEU2 digestion of GD1a was  
2 different from *mC* membrane, prepared by directly mixing GM1 to the matrix. In particular, the  
3 electron density of the hydrophilic layer was higher in the digested-*mA* and digested-*mB*  
4 membranes ( $480 \text{ e}^-/\text{nm}^3$ ) than in the *mC* membrane ( $419 \text{ e}^-/\text{nm}^3$ ).




5  
6 **Figure 4** Contrast profiles of the *in situ* digested-*mB* membrane (after the action of HsNEU2  
7 sialidase, red) and of membrane *mC* (green). The solvent is 150 mM aqueous solution. T= 22°C.  
8 Vertical dashed lines are drawn to guide the eye, as in Figure 3.

9  
10 Why digested-GD1a-containing membranes could be different from *ab initio* GM1-containing ones  
11 is not trivial. In principle, because of the reported activity of HsNEU2 on monomeric GM1 [34], a  
12 weak activity of sialidase on GM1 produced by GD1a hydrolysis in model membranes could not be  
13 excluded. Thus, to address this point, we exposed the GM1-containing model membrane, *mC*, to  
14 HsNEU2.

15 The reflectivity profiles of the *mC* membrane before and after the action of sialidase were almost  
16 superimposable, as seen in Figure 5. This finding also excludes that non specific interaction occurs  
17 between the enzyme and the membrane. A minor effect could be related to the mentioned weak  
18 action of the sialidase on dispersed GM1 molecules.

19



1


2 **Figure 5** Reflectivity curves from membrane *mC* before (green dots) and after (violet triangles) the  
 3 action of HsNEU2 sialidase. The solvent is 150 mM NaCl aqueous solution.  $T = 22^\circ\text{C}$ . Symbols  
 4 correspond to the experimental data, line to the best fit, corresponding to the profile in Figure 4,  
 5 green curve.

6

7 We performed reflectivity experiments on a membrane containing Asialo-GM1, *mD*. The  
 8 corresponding spectrum, together with the best fit line, and the fitted cross profile are reported in  
 9 Figure 6. The transverse structure of membrane *mD* was different from membrane *mA* both before  
 10 and after HsNEU2 digestion.

11

12




13

14 **Figure 6** Reflectivity curve from membrane *mD* (A) and contrast profile (B). The solvent is 150  
 15 mM NaCl aqueous solution.  $T = 22^\circ\text{C}$ . Symbols correspond to the experimental data, line to the best

1 fit, corresponding to the profiles in B). In B) vertical dashed lines are drawn to guide the eye, as in  
2 Figure 3.

3

4 Finally, in Figure 7, we compare all the recovered electron density profiles, restricted to the external  
5 headgroup region. It is easily seen that all of them appear very similar, except for the one  
6 corresponding to the digested-GD1a model membrane, remarkably different from the others. In  
7 fact, its external polar region has a much higher contrast than all of the other model membranes,  
8 including the chemically-identical GM1-containing membrane, *mC*.



9

10 **Figure 7** Contrast profiles of the external head-group region of the different membranes in 150 mM  
11 NaCl solution: *mA* before (blue) and after (red) HsNEU2 digestion, *mC* before (green) and after  
12 (violet) HsNEU2 digestion and *mD* (black). T=22°C. The external polar region GD1a-digested  
13 membrane *mA* (red) has a remarkably higher contrast. The horizontal axis refers to distances from  
14 the mid-plane ( $z = 0$ ) of the external head-group regions.

15

## 16 4. Discussion

17 In biological membranes, gangliosides are transformed into lower-sialylated species by sialidases,  
18 while being already packed in the membrane aggregate. From this start point, interesting issues  
19 emerge. Gangliosides disposition and relative concentration within the lipid structure could affect  
20 enzymatic activity and kinetics. Also, and more intriguing, the enzymatic action could not only

1 produce a new molecule in the membrane, but also induce changes in the membrane organization,  
2 such as lipid lateral segregation, membrane core fluidity and roughness, maybe reminiscent of its  
3 chemical and structural history.

4 In the experiment presented here, our aim was to study the transverse structure of ganglioside  
5 containing membranes submitted to sialidase digestion. So we applied synchrotron radiation  
6 reflectometry to four supported model membranes containing three different ganglioside species:  
7 GD1a, GM1 and AsialoGM1. We investigated the changes induced by sialic acid removal by *in-situ*  
8 enzymatic digestion, and we compared the final membrane with a membrane originally built up  
9 with the lower sialylated ganglioside. The successful deposition of single macroscopic membranes  
10 containing GD1a and AsialoGM1 was itself a first remarkable result, following the previous  
11 successful preparations with GM1 and cholesterol [8,51] and opens the way to the construction of  
12 model membranes with increasing bio-similarity.

13 The occurrence of a change in the cross structure of the GD1a-containing membrane, *mA*, upon  
14 enzymatic action is clearly visible in Figure 3. Then, the technique proves to be extremely sensitive  
15 in following this kind of membrane-enzyme interaction and constitutes an experimental and  
16 technical achievement. In fact, the technique is able to reveal structural changes in the sugar-  
17 containing hydrophilic layer of the membrane, where sialidase is acting. However, before the  
18 experiment had been performed, it was difficult to foresee the extent of experimental sensitivity to  
19 these structural changes. In fact, the mole fraction of GD1a with respect to the hosting  
20 phospholipids of the outer layer is as low as 1:10. In addition, the enzyme removes only the external  
21 sialic acid residue of GD1a (one sugar unit out of 6), likely located quite far from the membrane  
22 surface (about 1.5 nm distance), while the phospholipid headgroups of the matrix are confined in a  
23 closer layer (some 1 nm thick). Remarkably, despite the visibility of a clear effect was not obvious  
24 to anticipate, an unambiguous variation of the average electron density of the hydrophilic layer as a  
25 whole was observed. We note that the outer hydrophilic layer (region *b*) accounts for the *dense*  
26 hydrophilic region of the membrane, where DPPC headgroups (the majority compound) are

1 confined, contributing to space filling with the bound hydration water. Reversely, the protruding  
2 portions of the ganglioside headgroups, containing the external, terminal, sialic acid unit to be  
3 removed by the enzyme (see Figure 1), sparingly populate a far-layer that is not dense enough to  
4 effectively contrast against bulk water. In this far layer one can estimate that the headgroup surface  
5 occupancy is in the range 10-20% of the total available area.

6 The density profiles reported in Figure 3B differ mainly in the outer hydrophilic layer (region *b*),  
7 where ganglioside headgroups are exposed to the enzyme. Upon HsNEU2 digestion, a significant  
8 increase of the contrast of this region occurred, *i.e.*, its electron density increased. We remark that  
9 the same result was obtained on the replica GD1a-containing membrane, *mB*; thus, the  
10 reproducibility stands as a strong proof of the reliability of the outcome.

11 This result is of great importance and adds interesting information on the cross local structure of the  
12 ganglioside-containing membrane, integrating the compositional result of GD1a turned to GM1 by  
13 biochemical catalysis. In fact, a decrease in the average specific volume of the layer, corresponding  
14 to increased lipid layer compactness, cannot be invoked to explain the observed large increase in the  
15 average electron density of the hydrophilic close-layer, about 1 nm thick. In fact, no corresponding  
16 change is seen in the electron density of the hydrophobic region of the outer layer (see *SI*, Table  
17 S1). A more tricky effect should occur.

18 We observe that an increase in the electron density profile of the hydrophilic close-layer, without  
19 affecting the hydrophobic region, can be obtained by replacing a fraction of the hydrophilic close-  
20 layer with an equal volume with higher electron density. This suggests that the headgroups of the  
21 GM1 molecules emerging from *in situ* digestion are not only shortened by one charged external  
22 unit, but also retracted closer to the membrane surface with respect to the parent-GD1a headgroups.  
23 It seems that they lay down on the membrane surface. We recall that the lying conformation,  
24 requiring the ganglioside headgroups to tilt with respect to the axis of the hydrophobic chains  
25 embedded in the membrane core, has often been hypothesized in the literature. The interconversion  
26 between the standing and lying conformations of headgroups has been observed to occur in

1 ganglioside aggregates as a consequence of the action of external parameters like temperature,  
2 dehydration, or salinity [9,52].

3 By integration of the  $\rho(z)$  profiles in the region of the external polar heads of membrane *mA* before  
4 and after digestion (blue and red curves of Figure 7), the number density of additional electrons in  
5 the close-layer can be calculated. A 30% increase in the electron density excess with respect to  
6 water has occurred, corresponding to a 1.3% GM1 head retraction per unit volume. Very  
7 interestingly, this is consistent with a glycolipid head volume redistribution increasing its lateral  
8 hindrance by 4%, the same value estimated for the interfacial area increase upon GM1 tilting in  
9 micelles [53]. The corresponding variation in the low-contrast hydrophobic moiety, 4% each for the  
10 10% ganglioside molecules in the layer, is not visible.

11 The comparison between the “enzyme digested” membrane and a membrane prepared by mixing  
12 the matrix with the digestion product, reported in Figure 4, is extremely interesting. In particular,  
13 the electron density of the hydrophilic layer is higher for the digested-*mA* (digested-*mB*) membrane  
14 than for *mC*.

15 This is an interesting point. Why should digested-GD1a-containing membranes be different from *ab*  
16 *initio* GM1-containing ones is not obvious. An incomplete digestion of GD1a into GM1 or further-  
17 proceeding digestion leading to AsialoGM1 can be excluded. In fact, despite direct biochemical  
18 assessment on the deposited membranes is unfeasible, enzymatic activity has been previously  
19 studied in different ganglioside-containing systems: incomplete digestion of GD1a is unlikely since  
20 HsNEU2, similarly to other sialidases, has been proved to go through full GD1a-to-GM1  
21 conversion in different aggregated structures in solution, as micelles or mixed vesicles [32,54].  
22 Moreover, the amount of added enzyme was intentionally in large excess with respect to the  
23 substrate. Moreover, being the electron density of the hydrophilic outer layer higher than both  
24 GD1a- and GM1-containing membranes, so out of any weighted average of the two, trivial  
25 incomplete digestion is unable to provide straightforward explanation of the experimental result. On  
26 the other hand, a weak activity of sialidase on GM1 inserted in model membranes, producing the



1 Asialo-GM1, could not be excluded [34] in principle. This was ruled out by testing that the  
2 reflectivity spectra of the GM1-containing *mC* membrane before and after exposure to sialidase  
3 were almost superimposable, as seen in Figure 5. Furthermore, the transverse structure of the  
4 Asialo-GM1-containing *mD* membrane is different from that of GD1a-containing membranes *mA*  
5 and *mB*, both before and after HsNEU2 digestion (Figures 6A and 6B).  
6 From the comparison of the electron density profiles restricted to the external headgroups region,  
7 reported in Figure 7, we see that the lying conformation, corresponding to high close-shell contrast,  
8 is not the only one admitted for GM1, in the membrane. As well, GM1 headgroups can stand more  
9 perpendicular to the membrane (as in *mC*), displacing their center of mass farther from the surface.  
10 This arrangement appears to be similar to the one adopted by both GD1a and AsialoGM1 in  
11 membranes *mA* (and its replica *mB*) and *mD*. Notably, when the same protocol was followed for  
12 sample preparation, the cross profiles of the external close-layers at 40mN/m were nicely the same.  
13 Although the polar heads of the different gangliosides are different, their initial portion, included in  
14 the close-layer, is the same and, at the given surface dilution, behaves the same way. The obtained  
15 results (Figure 7) indicate that in this condition of surface concentration and pressure, the  
16 ganglioside headgroups assume the standing conformation with respect to the membrane plane. But,  
17 upon specific and proper stimulation, they can turned to lying, seemingly modulating the  
18 mechanical properties of their local environment.

19

## 20 **5. Conclusion**

21 The outcome of this work is that the action of the enzyme HsNEU2 has a deeper effect on  
22 biomimetic membranes than just turning GD1a to GM1 by external sialic acid removal. The *in-situ*  
23 produced GM1 glycosidic headgroup is retracted to the membrane surface, storing additional  
24 packing surface. Stored surface could eventually be used to promote or allow for mechanical  
25 deformations of the membrane, providing structural support to biological functions. In fact,

1 ganglioside-enriched domains are easily associated with membrane protrusion or caveolae. The  
2 present observations, then, support the hypothesis that the sialidase action drives the membrane to a  
3 structural turning point, where it can be readdressed to different final fates.

4 Finally, it is sometimes argued that the packing properties of aggregating molecules are likely to be  
5 washed out once in a large structure, like a membrane, where they happen to be mixed with other  
6 different species. In this experiment, as in many others, it comes out that, instead, molecular  
7 packing and membrane properties respond to each other, contributing to membrane structural and  
8 functional evolution.

9

10


11

## 12 **Acknowledgments**


13 The authors wish to acknowledge the ESRF (Grenoble, FR) Synchrotron Radiation Facility for  
14 beamtime allocation, and to thank Oleg Konovalov for assistance on ID10B beamline. The authors  
15 also thank the Institut Laue-Langevin in Grenoble (FR) for the use of the Partnership for Soft  
16 Condensed Matter (PSCM) facilities.

1 **Graphical Abstract:**

2



3



4

5

6 **References**

7 [1] K. Simons, M. J. Gerl, Revitalizing membrane rafts: new tools and insights, Nature  
8 Reviews - Molecular Cell Biology 11 (2010) 688-699.

9 [2] A. Kusumi, T.K. Fujiwara, R. Chadda, M. Xie, T.A. Tsunoyama, Z. Kalay, R.S. Kasai,  
10 K.G.Suzuki, Dynamic Organizing Principles of the Plasma Membrane that Regulate Signal  
11 Transduction: Commemorating the Fortieth Anniversary of Singer and Nicolson's Fluid-Mosaic  
12 Model, Ann Rev Cell Dev Biol 28 (2012) 215-250.

13 [3] H.E. Grecco, M. Schmick, P.I.H. Bastiaens, Signaling from the Living Plasma Membrane,  
14 Cell 144 (2011) 897-909.

15 [4] B.N. Kholodenko, J.F. Hancock, W. Kolch, Signalling ballet in space and time, Nat Rev  
16 Mol Cell Biol 11 (2010) 414-426.

- 1 [5] V. Mueller, A. Honigmann, C. Ringemann, R. Medda, G. Schwarzmann, C. Eggeling, FCS  
2 in STED microscopy: studying the nanoscale of lipid membrane dynamics, *Methods Enzymol* 519  
3 (2013) 1-38.
- 4 [6] J.M. Kwiatek, D. M. Owen, A. Abu-Siniyeh, P. Yan, L.M. Loew, K. Gaus, Characterization  
5 of a New Series of Fluorescent Probes for Imaging Membrane Order, *Plos One* 8 (2013) 7-13.
- 6 [7] P. Quinn, Sphingolipid Symmetry Governs Membrane Lipid Raft Structure, *BBA* 1838  
7 (2014) 1922-1930.
- 8 [8] V. Rondelli, G. Fragneto, S. Motta, E. Del Favero, P. Brocca, S. Sonnino, L. Cantù,  
9 Ganglioside GM1 forces the redistribution of cholesterol in a biomimetic membrane, *BBA* 1818  
10 (2012) 2860–2867.
- 11 [9] L. Cantù, E. Del Favero, P. Brocca, M. Corti, Multilevel structuring of ganglioside-  
12 containing aggregates: From simple micelles to complex biomimetic membranes, *Adv Coll Interf Sci*  
13 205 (2014) 177-186.
- 14 [10] L. Cantù, E. Del Favero, S. Sonnino, A. Prinetti, Gangliosides and the multiscale modulation  
15 of membrane structure, *Chem Phys Lip* 164 (2011) 796-810.
- 16 [11] S. Staubach, F.G. Hanisch, Lipid rafts: signaling and sorting platforms of cells and their  
17 roles in cancer, *Expert Rev Proteom* 8 (2011) 263-277.
- 18 [12] F. Hullin-Matsuda, T. Kobayashi, Monitoring the distribution and dynamics of signaling  
19 microdomains in living cells with lipid-specific probes, *Cell Mol Life Sci* 64 (2007) 2492-2504.
- 20 [13] K. Handa, S.I. Hakomori, Carbohydrate to carbohydrate interaction in development process  
21 and cancer progression, *Glycoconj J* 29 (2012) 627-637.
- 22 [14] A.R. Todeschini, J.N. Dos Santos, K. Handa, S.I. Hakomori, Ganglioside GM2/GM3  
23 complex affixed on silica nanospheres strongly inhibits cell motility through CD82/cMet-mediated  
24 pathway, *Proc Natl Acad Sci* 105 (2008) 1925-1930.

- 1 [15] E. Monti, E. Bonten, A. D'Azzo, R. Bresciani, B. Venerando, G. Borsani, R. Schauer, G.  
2 Tettamanti, Sialidases in vertebrates: a family of enzymes tailored for several cell functions, *Adv*  
3 *Carbohydr Chem Biochem* 64 (2010) 403-479.
- 4 [16] E. Monti, A. Preti, B. Venerando, G. Borsani, Recent development in mammalian sialidase  
5 molecular biology, *Neurochem Res* 27 (2002) 649-663.
- 6 [17] T. Miyagi, T. Wada, K. Yamaguchi, K. Hata, Sialidase and malignancy: a minireview,  
7 *Glycoconj J* 20 (2004) 189-198.
- 8 [18] T. Miyagi, T. Wada, K. Yamaguchi, K. Shiozaki, I. Sato, Y. Kakugawa, H. Yamanami, T.  
9 Fujiya, Human sialidase as a cancer marker, *Proteomics* 8 (2008)3303-3311.
- 10 [19] Wada T, K. Hata, K. Yamaguchi, K. Shiozaki, K. Koseki, S. Moriya, T. Miyagi, A crucial  
11 role of plasma membrane-associated sialidase in the survival of human cancer cells ,*Oncogene* 26  
12 (2007) 2483-2490.
- 13 [20] C. Tringali, B. Lupo, L. Anastasia, N. Papini, E. Monti, R. Bresciani, G. Tettamanti, B.  
14 Venerando, Expression of Sialidase Neu2 in Leukemic K562 Cells Induces Apoptosis by Impairing  
15 Bcr-Abl/Src Kinases Signaling, *J Biol Chem* 282(2007) 14364–14372.
- 16 [21] C. Tringali, B. Lupo, F. Cirillo, N. Papini, L. Anastasia, G. Lamorte, P. Colombi, R.  
17 Bresciani, E. Monti, G. Tettamanti, B..Venerando, Silencing of membrane-associated sialidase Neu3  
18 diminishes apoptosis resistance and triggers megakaryocytic differentiation of chronic myeloid  
19 leukemic cells K562 through the increase of ganglioside GM3, *Cell Death & Differ* 16 (2009) 164-  
20 174.
- 21 [22] C. Tringali, F. Cirillo, G. Lamorte, N. Papini, L. Anastasia, B. Lupo, I. Silvestri, G.  
22 Tettamanti, B. Venerando, NEU4L sialidase overexpression promotes  $\beta$ -catenin signaling in  
23 neuroblastoma cells enhancing stem-like malignant cell growth, *Int J Cancer* 31 (2012) 1768-1778.
- 24 [23] C. Tringali, I. Silvestri, N. Papini, L. Anastasia, G. Tettamanti, B. Venerando, The Plasma  
25 Membrane Sialidase NEU3 Regulates the Malignancy of Renal Carcinoma Cells by Controlling  $\beta$ 1  
26 Integrin Internalization and Recycling, *J Biol Chem* 287 (2012) 42835-42845.

- 1 [24] X. Wang, K. Yamaguchi, T. Wada, K. Hata, X. Zhao, T. Fujimoto, T. Miyagi, A Close  
2 Association of the Ganglioside-specific Sialidase Neu3 with Caveolin in Membrane Microdomains, J  
3 Biol Chem 277 (2002) 26252–26259.
- 4 [25] R. Scaringi, M. Piccoli, N. Papini, F. Cirillo, E. Conforti, S. Bergante, C. Tringali, A.  
5 Garatti, C. Gelfi, B. Venerando, L. Menicanti, G. Tettamanti, L. Anastasia, NEU3 Sialidase Is  
6 Activated under Hypoxia and Protects Skeletal Muscle Cells from Apoptosis through the Activation  
7 of the Epidermal Growth Factor Receptor Signaling Pathway and the Hypoxia-inducible Factor  
8 (HIF)-1 $\alpha$ , J Biol Chem 288 (2013) 3153-3162.
- 9 [26] S. Kawamura, I. Sato, T. Wada, K. Yamaguchi, Y. Li, D. Li, X. Zhao, S. Ueno, H. Aoki, T.  
10 Tochigi, M. Kuwahara, T. Kitamura, K. Takahashi, S. Moriya, T. Miyagi, Plasma membrane-  
11 associated sialidase (NEU3) regulates progression of prostate cancer to androgen-independent growth  
12 through modulation of androgen receptor signalling, Cell Death & Diff 19 (2012) 170-179.
- 13 [27] X. Nan, I. Carubelli, N.M. Stamatou, Sialidase expression in activated human T lymphocytes  
14 influences production of IFN-gamma, J Leukoc Biol 81 (2007) 284-296.
- 15 [28] F. Liang, V. Seyrantepe, K. Landry, R. Ahmad, A. Ahmad, N.M. Stamatou, A.V.  
16 Pshezhetsky, Monocyte differentiation up-regulates the expression of the lysosomal sialidase neu1  
17 and triggers its targeting to the plasma membrane via major histocompatibility complex class ii-  
18 positive compartments, J Biol Chem 281 (2006) 27526-27538.
- 19 [29] F. D'Avila, C. Tringali, N. Papini, L. Anastasia, G. Croci, L. Massaccesi, E. Monti, G.  
20 Tettamanti, B. Venerando, Identification of Lysosomal Sialidase NEU1 and Plasma Membrane  
21 Sialidase NEU3 in Human Erythrocytes, J Cell Biochem 114 (2013) 204-211.
- 22 [30] S. Kijimoto-Ochiai, N. Doi, M. Fujii, S. Go, K. Kabayama, S. Moriya, T. Miyagi, T. Kodama,  
23 Possible association of Neu2 with plasma membrane fraction from mouse thymus exhibited sialidase  
24 activity with fetuin at pH 7.0 but not at pH 4.5, Microbiol Immunol 57 (2013) 569-582.
- 25 [31] J. Daillant, A. Gibaud, X-ray and Neutron Reflectivity: Principles and Applications,  
26 Springer, 1999.

- 1 [32] E. Del Favero, P. Brocca, S. Motta, V. Rondelli, S. Sonnino, L. Cantù, Nanoscale structural  
2 response of ganglioside-containing aggregates to the interaction with sialidase, *J Neurochem* 116  
3 (2011) 833-839. M. Leonard, Crystal Structure of the Human Cytosolic Sialidase Neu2, *JBC* 280  
4 (2005) 469-475.
- 5 [33] C. Tringali, N. Papini, P. Fusi, G. Croci, G. Borsani, A. Preti, P. Tortora, G. Tettamanti, B.  
6 Venerando, E. Monti, Properties of Recombinant Human Cytosolic Sialidase HsNEU2, *JBC* 279  
7 (2004) 3169–3179.
- 8 [34] V. Smutova, A. Albohy, X. Pan, E. Korchagina, T. Miyagi, N. Bovin, C. W. Cairo, A. V.  
9 Pshezhetsky, Structural Basis for Substrate Specificity of Mammalian Neuraminidases, *PLoS One* 9  
10 (2014) e106320.
- 11 [35] T. Charitat, E. Bellet-Amalric, G. Fragneto, F. Graner, Absorbed and free bilayers at the  
12 solid liquid interface, *Eur Phys J B* 8 (1999) 583-593.
- 13 [36] V. Rondelli, E. Del Favero, S. Motta, L. Cantù, G. Fragneto, P. Brocca, Neutrons for rafts,  
14 rafts for neutrons, *Eur Phys J E* 36 (2013) 73-80. M.A. Chester UPAC-IUB Joint Commission on  
15 Biochemical Nomenclature (JCBN) Nomenclature of glycolipids, *Eur J Biochem* 257 (1998) 293-  
16 298.
- 17 [37] G. Tettamanti, F. Bonali, S. Marchesini, V. Zambotti, A new procedure for the extraction  
18 purification and fractionation of brain gangliosides, *BBA* 296 (1973) 160-170.
- 19 [38] A.P. Girard-Egrot, L.J. Blum, *Langmuir-Blodgett Technique for Synthesis of Biomimetic*  
20 *Lipid Membranes, Nanobiotech of Biomim Mem*, Ed D.K. Martin, Springer, 2007 .
- 21 [39] J.R. Vig, UV/Ozone cleaning of surfaces, *Vac Sci Technol A* 3 (1985) 1027-1034.
- 22 [40] J. Daillant, E. Bellet-Amalric, A. Braslau, T. Charitat, G. Fragneto, F. Graner, S. Mora, F.  
23 Rieutord, B. Stidder, Structure and fluctuations of a single floating lipid bilayer, *Proc Natl Acad Sci*  
24 102 (2005) 11639-11644.
- 25 [41] M. Born, E. Wolf, *Principles of Optics*, Pergamon Press Ltd, London, 1959.

- 1 [42] J. Penfold, R.K. Thomas, The application of the specular reflection of neutrons to the study  
2 of surfaces and interfaces, *JPCM* 2 (1990) 1369-1412.
- 3 [43] L.G. Parratt, Surface Studies of Solids by Total Reflection of X-Rays, *Phys Rev* 95 (1954)  
4 359-369.
- 5 [44] F. Abeles, Les applications des couches minces en polarimétrie, *J Phys Radium* 11 (1950)  
6 403-406.
- 7 [45] A. Nelson, Co-refinement of multiple-contrast neutron/X-ray reflectivity data using  
8 MOTOFIT, *J Appl Cryst* 39 (2006) 273-276.
- 9 [46] J.F. Nagle, S. Tristram-Nagle, Structure of lipid bilayers, *BBA* 1469 (2000) 159–195.
- 10 [47] M. Boretta, L. Cantù, M. Corti, E. Del Favero, Cubic phases of gangliosides in water:  
11 possible role of the conformational bistability of the headgroup, *Physica A* 236 (1997) 162–176.
- 12 [48] A.I. Greenwood, S. Tristram-Nagle, J.F. Nagle, Partial molecular volumes of lipids and  
13 cholesterol, *Chem Phys Lip* 143 (2006) 1–10.
- 14 [49] V. Rondelli, G. Fragneto, S. Motta, E. Del Favero, L. Cantù, Reflectivity from floating  
15 bilayers: can we keep the structural asymmetry?, *J Phys Conf Ser* 340 (2012) 012083.
- 16 [50] E. Del Favero, P. Brocca, V. Rondelli, S. Motta, A. Raudino, L. Cantù, Optimizing the  
17 crowding strategy: sugar-based ionic micelles in the dilute-to-condensed regime, *Langmuir* 30 (2014)  
18 9157–9164.
- 19 [51] L. Cantù, M. Corti, E. Del Favero, E. Digirolamo, S. Sonnino, G. Tettamanti, Experimental  
20 evidence of a temperature-related conformational change of the hydrophilic portion of gangliosides,  
21 *Chem Phys Lip* 79 (1996) 137-145.
- 22 [52] E. Del Favero, A. Raudino, M. Pannuzzo, P. Brocca, S. Motta, L. Cantù, Transient step-like  
23 kinetics of enzyme reaction on fragmented-condensed micellar substrates, *J Phys ChemB* 116 (2012)  
24 9570–9579.