

Spermidine Associated to Non-Surgical Treatment of Periodontal Disease: Split Mouth Randomized Controlled Clinical Trial

Daniela Carmagnola¹, Carlo Angelo Ghisalberti², Gaia Pellegrini^{1*}, Lucrezia Cinquanta³, Marilisa Toma¹ & Claudia Dellavia¹

¹*Department of Biomedical, Surgical and Dental Sciences, Università degli Studi di Milano, via Mangiagalli, Milan, Italy*

²*Department of Biomedical Science for Health, Università degli Studi di Milano, via Mangiagalli, Milan, Italy*

***Correspondence to:** Dr. Gaia Pellegrini, Department of Biomedical, Surgical and Dental Sciences, Università degli Studi di Milano, via Mangiagalli, Milan, Italy.

Copyright

© 2019 Dr. Gaia Pellegrini, *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 13 May 2019

Published: 04 July 2019

Keywords: *Spermidine; Non-Surgical Therapy; Periodontal Disease Treatment*

Abstract

Objectives

Spermidine is an endogenous polyamine whose increase induces the autocrine remodeling into targeted cells. The aim of the present study was to evaluate the adjunctive beneficial effects of the local delivery of spermidine to periodontal non-surgical therapy on reduction of periodontal pocket.

Methods

In this split mouth, blind, randomized controlled clinical trial, 20 patients with severe chronic generalized periodontal disease scheduled for cause related non-surgical periodontal treatment were enrolled. In three quadrants for each patients, three experimental teeth with probing pocket depth (PPD) ≥ 6 mm were selected and randomly assigned to one of the following treatments: scaling and root planning (SRP)+ spermidine gel (spermidine + alginate) (group a), SRP+ placebo

gel (only alginate) (positive control, group b) or SRP (negative control, group c). PPD at baseline, 3 and 6 months after treatment was assessed. Descriptive and inferential statistics was done.

Results

After periodontal treatment, a reduction of PPD was observed in all sites. No differences emerged between groups at each time point. In all groups, a significant PPD reduction was observed at T1 and T2 compared to baseline ($p < 0.05$ and $p < 0.001$). PPD reduction from T1 to T2 was significant only in group a ($p < 0.001$).

Conclusions

After non-surgical treatment, local delivery of spermidine may induce prolonged improvement of clinical outcome.

Abbreviations

Probing Pocket Depth (PPD);
Full Mouth Plaque Scores (FMPS)
Full Mouth Bleeding Scores (FMBS)
Scaling and Root Planning (SRP)
Percentage of PPD reduction (%PPDr)

Introduction

Periodontitis is a wide range of bacterially induced inflammatory diseases that cause the destruction of tooth supporting tissues (gingiva, periodontal ligament and bone) and the formation of periodontal pocket [1]. Extension and severity of periodontal tissue damage depends on different factors, such as the quality and quantity of bacteria, the host response and, for example, smoking or some systemic conditions [2]. The ratio between pro (bacteria, systemic diseases, risk factors..) and anti inflammatory (immune system) mechanisms will determine the story of the disease, that is usually slow and progressive [2].

The treatment of periodontal disease consists in cause related therapy, the aim of which is to remove all local etiological factors and irritants such as plaque and calculus by powered or manual instruments, to obtain adequate compliance by the patient in the maintenance of a satisfying level of oral hygiene and to modify any modifiable negative habit. In most cases, such approach is effective at arresting the progression of the disease, bringing the periodontal tissue back to healthy conditions and eliminating periodontal pockets. However, cases in which infection is particularly aggressive and/or pockets defects are particularly deep or difficult to cleanse, further local or systemic therapeutic strategies are needed to reduce the bacteria load or to support the immune system [3]. Subsequently, any remaining periodontal pockets may be eliminated by a surgical approach that may contemplate the use of regenerative substances or biomaterials [4,5]. A large variety of drugs or substances that inhibit or eliminate periodontopathogenic microorganisms as well as modulate the inflammatory response of tissues has been proposed with the aim to make the cause-related therapy more effective, however so far only limited beneficial effects have been proved [3].

Polyamines, namely putrescine, spermidine and spermine, are low molecular weight cationic molecules synthesized by many types of cells, especially those in tissues with high turnover, and also introduced with the common diet. Spermidine acts as conformation modifier on most biologically active, anionic biopolymers [6,7]. Such electrostatic, non-pharmacologic mode of action seems to end-up with the promotion of physiologic events including autophagy induction, immunomodulation, extracellular matrix turnover, cardioprotection, neuroprotection and stem cells activation [6,8]. Spermidine also has demonstrated *in vitro* to suppress the synthesis of pro-inflammatory cytokines [9-11] and to downregulate chemotaxis of polymorphonuclear leucocytes when administered at concentrations similar to those found in gingival crevicular fluid [12]. Higher concentration of spermidine has also been found in tissues during inflammatory disease, suggesting a modulating activity mediated by the electrostatic interaction with the anionic mediators of the inflammatory patterns. The tissue content of spermidine was found to decrease with age and its higher levels in blood of nona/centenarians seemingly indicating a protective activity on tissue aging, thus improving human longevity [13].

Due to its role as endogenous regulator, the delivery of spermidine on periodontal pockets may be proposed to improve/accelerate the periodontal tissue response and healing after non-surgical therapy of periodontitis. However the effect of this polyamine on periodontal status has not been assessed yet.

The aim of the present study was to assess if the local delivery of spermidine brings adjunctive beneficial effects to periodontal non-surgical therapy in terms of reduction of periodontal pocket.

Study Population and Methodology

Study Design

This is a split mouth, blind, randomized controlled clinical trial. The study was approved by the ethical committee of Università degli Studi di Milano (n. 655, 3rd May 2016). All clinical procedures were performed in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards for studies involving human participants.

Participants

20 patients with periodontal disease of Stage 3 Grade B scheduled for cause related non-surgical periodontal treatment were enrolled.

The included patients presented the following inclusion criteria: age range: 25-80 years old; at least 3 monoradicular teeth with probing pocket depth (PPD) ≥ 6 mm distributed in three different quadrants; full mouth plaque (FMPS) and bleeding (FMBS) scores $> 25\%$ at the beginning of the study (baseline).

They all agreed to sign a written informed consent to take part to the study.

The following exclusion criteria were applied: current smokers or former smokers who had smoked in the previous 6 months; patients with clinically significant or unstable organic diseases, immune compromised

subjects, subjects taking steroid medications or chronically treated (i.e., two weeks or more) with any medication known to affect periodontal status (i.e., antibiotics or NSAIDS), patients who took antibiotics within 90 days of baseline; patients displaying a compromised healing potential such as those affected by connective tissue disorders or bone metabolic diseases; patients with conditions requiring antibiotic prophylaxis; pregnant or lactating women, or women who were of childbearing potential and not using birth control or abstinence.

Randomization

In each patient, three pockets with ≥ 6 mm were assigned by a balanced random permuted block randomization approach to one of the following treatments: a) SRP + spermidine gel, b) SRP + placebo gel, c) SRP. In order to reduce the chance of unfavorable splits between treatment groups in terms of key prognostic factors, randomization process of the treatment assignment (a, b and c) of 3 experimental sites for each patient was done by the central registrar (DC) and took into account the mean PPD at baseline (PPD \leq of 7mm). The treatment assignment was communicated to the operator (MT) immediately after scaling and root planning of all teeth.

Treatment Procedures

At baseline, a full periodontal charting was completed for each enrolled patient. Intraoral radiographs of the experimental quadrants were also taken using A Rinn's RING and a long cone parallel technique. In three quadrants for each patients, three experimental teeth with PPD ≥ 6 mm were selected and assigned to one of the following treatments: scaling and root planning (SRP)+ spermidine gel (spermidine + alginate) (group a), SRP+ placebo gel (only alginate) (positive control, group b) or SRP (negative control, group c). After inclusion into the study, each patient was thereafter scheduled the next appointments for non-surgical periodontal treatments (T0), charting at 3 (T1) and 6 (T2) months of follow up. Briefly, all pockets were cleaned under local anesthesia by means of ultrasonic scalers and hand instrumentation (Gracey standard and mini five curettes), and the roots were planned. Non-surgical treatment was performed by the same operator (MT) following a quadrant by quadrant approach at Dental Building at San Paolo Hospital (Milan). Further mechanical instrumentation was not performed during the following 3 months.

Immediately after mechanical instrumentation, for each experimental pocket, one of the following treatments was performed: a) injection of gel with spermidine, b) injection of gel with placebo, c) no adjunctive treatment (Figure 1). Intraoral photographs of the experimental sites were taken before SRP, after 3 and 6 months.



Figure 1: Local delivery of spermidine gel into the periodontal pocket.

Clinical Measurements

The following clinical measurements were taken at baseline, 3 (T1) and 6 (T2) months after SRP (T0) by a further blind operator (LC): FMPS and FMBS at four sites for all teeth; probing pocket depth (PPD) assessed at six sites of each tooth treated. All measurements were taken with a UNC periodontal probe (Hu Friedy Manufacturing Company Inc., Chicago, IL, USA).

Statistical Analysis

The primary outcome of the study was PPD reduction. Sample size calculation was performed using $\alpha = 0.05$ (5%) and the power of sample 80%. For the variability the value of 0.8mm (ng/mL) (standard deviation of PPD) obtained in previous papers was considered [14]. The minimum significant value considered was 1mm [14]. On the basis of these data, the needed number of patients to be enrolled resulted 17. Three more patients were enrolled considering a drop out of 15%.

In each treatment group, mean and standard deviation were calculated for probing pocket depth (PPD) (mm) at baseline, 3 months after treatment, and 6 months after treatment. Furthermore the percentage of PPD reduction at T1 and T2 compared to baseline was computed.

To study the trend of PPD over time within each treatment group the Friedman test was applied for PPD. To compare data between treatment groups at each time point, data analysis was performed by means of Friedman test. In case of emerging significant differences, a paired t test was applied to calculate differences in PPD levels between the two groups for each time point. A paired t test was used to study differences between all time points in the same group. Furthermore a paired t test was used to assess differences on percentages of PPD reduction between treatment groups.

Data were analyzed with SAS PROC MIXED (SAS Institute, Inc. 2008. SAS/STATVR 9.2 User's guide. Cary, NC: SAS Institute, Inc). Level of significance was set for 5% ($p < 0.05$).

Gaia Pellegrini, *et al.* (2019). Spermidine Associated to Non-Surgical Treatment of Periodontal Disease: Split Mouth Randomized Controlled Clinical Trial. *CPQ Dentistry*, 1(4), 01-12.

Results

Study Population

A total of 20 patients (9 males and 11 females, mean age 54 years) non smokers and systemically healthy were enrolled in the present study. Patients were enrolled from July 2016 until March 2017 and follow-ups at 6 months were completed by September 2017.

At baseline, bleeding on probing was 33.55% (± 18.65) and significantly reduced at T1 ($p < 0.05$) and T2 ($p < 0.05$) compared to baseline (paired *t* test, one tail). The plaque index was 48.52% (± 26.28) at baseline and significantly reduced at T1 ($p < 0.05$) and at T2 ($p < 0.05$) compared to baseline (paired *t* test, one tail).

Clinical Data

A total of 20 pockets for each group were treated and analyzed. Data on PPD for each group are reported in table 1. PPD at baseline was:

- in group a: 6mm in 12 sites, 7mm in 7 sites and 8mm in 1 site;
- in group b: 6mm in 14 sites, 7mm in 4 sites and 8mm in 2 sites;
- in group c: 6mm in 14 sites, 7mm in 5 sites and 8mm in 1 site.

Table 1: Data on periodontal pocket depth at each time point for groups. Significance of differences between time points (paired *t* test) and groups (Friedman test) has been indicated

	Baseline	T1	T2	T1 vs baseline (p)	T2 vs baseline	T2 vs T1
Group a	6.5 (± 0.7)	4.4 (± 0.69)	3.8 (± 0.7)	$< .0005$	$< .0001$	$< .001$
Group b	6.4 (± 1.3)	4.5 (± 0.9)	3.9 (± 0.9)	$< .0001$	$< .0001$	n.s.
Group c	6.4 (± 1.1)	4.1 (± 1.3)	4 (± 0.9)	$< .0001$	$< .0001$	n.s.
p between groups (Friedman test)	n.s.	n.s.	n.s.			

After periodontal treatment, a reduction of PPD was observed in all sites. No significant differences emerged between groups at each time point, while significant differences resulted within each treatment group between time points (see table 1) (Friedman test). In all groups, a statistically significant PPD reduction was observed at T1 and T2 compared to baseline ($p < 0.05$ and $p < 0.001$). PPD reduction from T1 to T2 resulted significant only in group a ($p < 0.001$).

The mean percentage of PPD reduction (%PPDr) compared to baseline resulted 31.3% at T1 and 40.4% at T2 in group a, 29.9% at T1 and 38% at T2 in group b, 35.8% at T1 and 38.8% at T2 in group c. No significant differences were found between groups. The percentage of PPD reduction was also computed considering PPD at baseline 6mm or >6 mm and is reported in table 2. No significant differences were found in %PPDr between sites with PPD 6mm at baseline and those with PPD > 6 mm.

Table 2: Mean percentage of periodontal pocket depth reduction for each group, separately indicated considering PPD at baseline 6mm or > 6mm

PPD at baseline	T1 vs baseline		T2 vs baseline	
	6mm	> 6mm	6mm	> 6mm
Group a	27.8%	36.6%	33.3%	50.9%
Group b	26.2%	38.7%	35.7%	43.4%
Group c	33.3%	41.7%	35.7%	44.3%



Figure 2a: Pictures before treatment with placebo



Figure 2b: Pictures after treatment with placebo. Gingival tissue appears pink without edema.



Figure 3a: Pictures before treatment with spermidine. The gingival inflammation due to plaque and calculus is evident




Figure 3b: Pictures after treatment with spermidine. The gingiva appears healed. The opening of the interdental spaces may indicate the resolution of the edema.

Discussion

In the present study the beneficial effects of spermidine locally delivered into periodontal pockets as adjunctive treatment to non-surgical therapy of periodontal disease were investigated. A split mouth randomized controlled study was drawn in order to eliminate interindividual differences concerning the response to the treatment at patient and operator level. From the analysis of data collected 3 months after treatment, it resulted that all the treatment options determined a significant improvement of clinical outcomes without differences between groups. However at the analysis of the reduction trend, only sites treated with spermidine demonstrated a prolonged (6 months) improvement of periodontal pocket depth compared to placebo and traditional non-surgical mechanical treatment. As reported above, polyamines and spermidine in particular, proven to play a non-pharmacologic, electrostatic micro-modification of some key protein involved in immune regulation [15]. Through this primary mode of action, spermidine exerts the synthesis inhibition of pro-inflammatory mediators including nitric oxide, prostaglandin E2 and cytokines such as tumor necrosis factor α and interleukin 1 β [16]. In a rodent model of Parkinson disease it showed

antioxidant and anti-inflammatory properties [17]. Spermidine was also reported to favor the formation of CD8+ memory T cells via induction of autophagy [18]. CD8+ memory T cells were found to play a protective role on bone tissue, by promoting the secretion of anti-inflammatory cytokines (IL 10), the suppression of osteoclastogenesis, thus promoting bone regeneration [19]. Furthermore, CD8 T cells synthesize amphiregulin, a cytokine member of the epidermal growth factor family with anti-inflammatory properties [20]. This protein plays a crucial role in orchestrating the repair tissue activity of lungs, muscles and skin [21]. In periodontal tissue, inflammatory and reparative processes that follow non-surgical treatment may involve the activation of the immune system and induce the secretion of amphiregulin. Indeed, an *in vitro* study observed that epithelial cells rests of Malassez express amphiregulin [22]. In adults, such cells appear to play a role in maintaining the normal periodontal cellular elements and function and also seem an important source of stem cells that might play a pivotal role in periodontal regeneration [23]. In consideration of what is reported in literature, it could be supposed that the prolonged improvement of reduction of PPD that was found at 6 months in sites treated with spermidine led to a sustained tissue repair activity over time. To confirm this hypothesis, further studies assessing the immune response of periodontal tissue to spermidine should be designed. In the treatment of periodontal disease, the strategy of local delivery has been proposed with the aim to expose the infected/inflamed gingival tissue to high doses, under controlled release, of medications limiting the systemic involvement. Several drugs have been released in such way as an adjunct to scaling and root planning with the aim to reduce the pathogenetic biofilm, improve bone formation or contrast its resorption and reduce tissue inflammation. Bone stimulatory and anti inflammatory agents such as boric acid gel, simvastatin, rosuvastatin and atorvastatin have been clinically tested with encouraging results on the improvement of clinical parameters including pocket depth reduction and percentage of bone fill up to 6 months after delivery [24-26]. A recently published review observed that local antibiotics/antimicrobials can offer clinical improvements, in particular in persistent or recurrent localized deep sites, mostly when using vehicles with proven sustained release of the antimicrobial agent [27]. Local delivery of gels containing bisphosphonates were also investigated and resulted in a significant increase in PPD reduction and percentage of bone fill [14,28]. In the present study, non-surgical treatment was done following a quadrant-by-quadrant approach to limit contamination of control sites with spermidine during the initial healing phase. However this study design limited the blinding of the operator who performed the therapy [29].

Conclusion

The local delivery of spermidine in addition to a non-surgical approach may improve the clinical outcomes of periodontal therapy by stimulating a prolonged repair capacity within periodontal cells. Further studies should be designed to investigate the effects of this polyamine on cells of periodontal tissues. It would be also interesting to assess the clinical effects of repeated deliveries of spermidine in periodontal pockets after non-surgical treatment in the long term.

Bibliography

1. Bosshardt, D. D. (2018). The periodontal pocket: pathogenesis, histopathology and consequences. *Periodontol 2000.*, 76(1), 43-50.

2. Kinane, D. F., Stathopoulou, P. G. & Papapanou, P. N. (2017). Periodontal diseases. *Nat Rev Dis Primers.*, 3, 17038.
3. Da Rocha, H. A., Silva, C. F., Santiago, F. L., Martins, L. G., Dias, P. C. & De Magalhães, D. (2015). Local drug delivery systems in the treatment of periodontitis: a literature review. *J Int Acad Periodontol.*, 17(3), 82-90.
4. Carmagnola, D., Pellegrini, G., Dellavia, C., Rimondini, L. & Varoni, E. (2018). Tissue engineering in periodontology: biological mediators for periodontal regeneration. *Int J Artif Organs.*, 42(5), 241-257.
5. Dellavia, C., Canciani, E., Rasperini, G., Pagni, G., Malvezzi, M. & Pellegrini, G. (2019). CEMP-1 Levels in Periodontal Wound Fluid during the Early Phase of Healing: Prospective Clinical Trial. *Mediators Inflamm.*, 2019(1737306).
6. Yang, J. & Rau, D. C. (2005). Incomplete ion dissociation underlies the weakened attraction between DNA helices at high spermidine concentrations. *Biophys J.*, 89(3), 1932-1940.
7. Berwanger, A., Eyrisch, S., Schuster, I., Helms, V. & Bernhardt, R. (2010). Polyamines: naturally occurring small molecule modulators of electrostatic protein-protein interactions. *J Inorg Biochem.*, 104(2), 118-125.
8. Stabellini, G., Moscheni, C., Gagliano, N., Dellavia, C., Calastrini, C., Ferioli, M. E. & Gioia, M. (2005). Depletion of polyamines and increase of transforming growth factor-beta1, c-myc, collagen-type I, matrix metalloproteinase-1, and metalloproteinase-2 mRNA in primary human gingival fibroblasts. *J Periodontol.*, 76(3), 443-449.
9. Southan, G. J., Szabo, C. & Thiernemann, C. (1994). Inhibition of the induction of nitric oxide synthase by spermine is modulated by aldehyde dehydrogenase. *Biochem Biophys Res Comm.*, 203(3), 1638-1644.
10. Szabó, C., Southan, G. J., Wood, E., Thiernemann, C. & Vane, J. R. (1994). Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor. *Br J Pharmacol.*, 112(2), 355-356.
11. Zhang, M., Caragine, T., Wang, H., Cohen, P.S., Botchkina, G., Soda, K., *et al.* (1997). Spermine inhibits proinflammatory cytokine synthesis in human mononuclear cells: a counterregulatory mechanism that restrains the immune response. *J Exp Med.*, 185(10), 1759-1768.
12. Walters, J. D., Miller, T. J., Cario, A. C., Beck, F. M. & Marucha, P. T. (1995). Polyamines found in gingival fluid inhibit chemotaxis by human polymorphonuclear leukocytes *in vitro*. *J Periodontol.*, 66(4), 274-278.
13. Pucciarelli, S., Moreschini, B., Micozzi, D., De Fronzo, G. S., Carpi, F. M., Polzonetti, V., *et al.* (2012). Spermidine and spermine are enriched in whole blood of nona/centenarians. *Rejuvenation Res.*, 15(6), 590-595.

14. Dutra, B. C., Oliveira, A. M. S. D., Oliveira, P. A. D., Manzi, F. R., Cortelli, S. C., Cota, L. O. M. & Costa, F. O. (2017). Effect of 1% sodium alendronate in the non-surgical treatment of periodontal intraosseous defects: a 6 month clinical trial. *J Appl Oral Sci.*, 25(3), 310-317.
15. Ylilauri, M., Mattila, E., Nurminen, E. M., Kämpylä, J., Niinivehmas, S. P., Määttä, J. A., *et al.* (2013). Molecular mechanism of T-cell protein tyrosine phosphatase (TCPTP) activation by mitoxantrone. *Biochim Biophys Acta.*, 1834(10), 1988-1997.
16. Jeong, J. W., Cha, H. J., Han, M. H., Hwang, S. J., Lee, D. S., Yoo, J. S., *et al.* (2018). Spermidine protects against oxidative stress in inflammation models using macrophages and zebrafish. *Biomol Ther (Seoul)*, 26(2), 146-156.
17. Sharma, S., Kumar, P. & Deshmukh, R. (2018). Neuroprotective potential of spermidine against rotenone induced Parkinson's disease in rats. *Neurochem Int.*, 116, 104-111.
18. Puleston, D. J. & Simon, A. K. (2015). New roles for autophagy and spermidine in T cells. *Microb Cell.*, 2(3), 91-93.
19. Cardoso, E. M. & Arosa, F. A. (2017). CD8+ T cells in chronic periodontitis: roles and rules. *Front Immunol.*, 8, 145.
20. Zaiss, D. M. W., Gause, W. C., Osborne, L. C. & Artis, D. (2015). Emerging functions of amphiregulin in orchestrating immunity, inflammation, and tissue repair. *Immunity*, 42(2), 216-226.
21. Vasanthakumar, A. & Kallies, A. (2015). The regulatory T cell: jack of all trades. *Trends Immunol.*, 36(12), 756-758.
22. Ohshima, M., Yamaguchi, Y., Micke, P., Abiko, Y. & Otsuka, K. (2008). *In vitro* characterization of the cytokine profile of the epithelial cell rests of Malassez. *J Periodontol.*, 79(5), 912-919.
23. Xiong, J., Mrozik, K., Gronthos, S. & Bartold, P. M. (2012). Epithelial cell rests of Malassez contain unique stem cell populations capable of undergoing epithelial mesenchymal transition. *Stem Cells Dev.*, 21(11), 2012-2025.
24. Priyanka, N., Abhilash, A., Saquib, S., Malgaonkar, N., Kudiyar, N., Gupta, A., *et al.* (2017). Clinical efficacy of subgingivally delivered 1.2mg simvastatin in the treatment of patients with aggressive periodontitis: a randomized controlled clinical trial. *Int J Periodontics Restorative Dent.*, 37(2), e135-e141.
25. Muniz, F. W. M. G., Taminski, K., Cavagni, J., Celeste, R. K., Weidlich, P. & Rösing, C. K. (2018). The effect of statins on periodontal treatment a systematic review with meta analyses and meta regression. *Clin Oral Investig.*, 22(2), 671-687.
26. Singhal, S., Pradeep, A. R., Kanoriya, D., Garg, S. & Garg, V. (2018). Boric acid gel as local drug delivery in the treatment of class II furcation defects in chronic periodontitis: a randomized, controlled clinical trial. *J Investig Clin Dent.*, 9(1).

-
27. Jepsen, K. & Jepsen, S. (2016). Antibiotics/antimicrobials: systemic and local administration in the therapy of mild to moderately advanced periodontitis. *Periodontol*, 71(1), 82-112.
28. Sharma, A., Raman, A. & Pradeep, A. R. (2017). Role of 1% alendronate gel as adjunct to mechanical therapy in the treatment of chronic periodontitis among smokers. *J Appl Oral Sci.*, 25(3), 243-249.
29. Madeo, F., Eisenberg, T., Pietrocola, F. & Kroemer, G. (2018). Spermidine in health and disease. *Science*, 26, 359(6374).