

# Contribution of respiratory syncytial virus (RSV) among patients <15 years hospitalized with severe acute respiratory infection (SARI) in Milan, 2014-2017.



22<sup>nd</sup> ESCV,  
European Society for Clinical  
Virology

Laura Pellegrinelli<sup>1</sup>, Sara Colonia Uceda Renteria<sup>2</sup>, Cristina Galli<sup>1</sup>, Letizia Greco<sup>2</sup>, Annarosa Orlandi<sup>2</sup>, Sandro Binda<sup>1</sup>, Elena Pariani<sup>1</sup>, Giovanna Lunghi<sup>2</sup>

1. Department of Biomedical Sciences for Health, University of Milan, Milan, Italy

2. Virology Unit, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

Contact details: Laura Pellegrinelli; Laura.pellegrinelli@unimi.it, +390250325071

**Keywords:** Respiratory syncytial virus; severe acute respiratory infection; pediatric hospitalization; molecular diagnostic.

## Background

Respiratory syncytial virus (RSV) infections can be asymptomatic or associated with symptoms ranging from mild cold to severe acute respiratory infection (SARI)<sup>1</sup> and are responsible for substantial global morbidity in young children and elderly individuals<sup>2</sup>. To implement preventive and control measures and to inform vaccination strategies, it is critical to characterize the epidemiological patterns of circulating RSV.

## Objectives

This study aimed at describing the results of RSV molecular detection in respiratory samples collected from children <15 years hospitalized with SARI in Milan (Italy) during four consecutive years (2014-2017) and to estimate the prevalence of RSV, the risk of infection from RSV and the incidence of hospitalization of RSV-positive SARI stratified by age.

## Methods

From January 1<sup>st</sup>, 2014, to December 31<sup>st</sup>, 2017, 3013 respiratory samples (2826 upper respiratory tract [URT] and 187 lower respiratory tract [LRT] specimens) collected from as many children <15 years hospitalized with SARI at a university hospital in Milan were analysed. After nucleic acids extraction, samples were tested by a multiplex real-time PCR to detect RSV<sup>3</sup>.

**SARI case definition:** the standard SARI case definition is: acute respiratory infection with history of high fever or measured high fever ( $\geq 38^\circ\text{C}$ ) and cough with onset within the last 10 days and requiring hospitalization<sup>4</sup>.

**Incidence rate of hospitalization of RSV-positive SARI:** incidence were calculated using the average number of population <15 years of age for each year of study as denominator.

## Results

### 1 - Prevalence of RSV-positive SARI cases

During the study period, 571 out of 3013 (19%) respiratory samples tested positive to RSV (Table 1). In this SARI series, RSV positivity rate identified in LRT samples (27/187; 14.4%) and URT samples (544/2826; 19.2%) was similar ( $p=0.09$ ). The median age of RSV-positive SARI cases was 6.6 months (inter quartile range; 17.2 months) with a male-to-female ratio of 1:1.12. Overall, 22% (355/1613) and 19% (125/658) of children aged less than 1 year and those aged 1-3 years, respectively, tested positive to RSV (Table 1). Cumulatively, 62.2% (355/571) of RSV were identified in children <1 year (Fig. 1) and 12.3% (70/571) in children <1 month.

### 2 - Risk of infection from RSV by age-groups

The risk of RSV infection in children aged 0-1 year was 1.5-fold (95% CI: 1.2-1.9) higher with respect to all the other age groups.

### 3 - Incidence of hospitalization of RSV-positive SARI


The overall incidence of hospitalization RSV-positive SARI was 40.5 per 100'000 children aged <15 years with the highest incidence rate observed in children less than 1 year of age (426/100'000) (Fig. 2).

### 4 - Temporal distribution of RSV


RSV circulated along the entire study period with the highest positivity rate (59.2%; 338/571) identified during seasonal peaks occurring from December to February each year.

**Table 1.** Prevalence of RSV detection in children with SARI by age.

Age group (year)	No. of patients	No. of RSV-positive patients	% of RSV-positive patients
0-1	1613	355	22.0%
2-3	658	125	19.0%
4-6	386	55	14.2%
7-11	210	22	10.5%
12-15	146	14	9.6%
<b>Total</b>	<b>3013</b>	<b>571</b>	<b>19.0%</b>



**Figure 1.** Distribution of RSV-positive SARI by age.



**Figure 2.** Incidence of hospitalization of RSV-positive SARI per 100'000 children <15 years of age.

## Discussion and Conclusions

Accordingly to other studies<sup>1-2</sup>, RSV was detected in 19% of hospitalized SARI cases <15 years, mainly in children <1 year in which the risk of RSV infection was 1.5-fold higher with respect to all the other age groups. As expected<sup>1-2</sup>, circulation of RSV peaked from December to February each year. In this series, sampling of upper or lower respiratory tract resulted in similar RSV-positivity rate. RSV is recognized as a major cause of hospital admissions in young children; in this study, the highest incidence of hospitalization of children less than 15 year with RSV-positive SARI was 426 per 100'000 children aged <1 years. Routine molecular testing to detect RSV and epidemiological study are warranted to implement preventive and control measures and to drive further vaccination strategies.

## References

1. Ogra PL. Paediatr Respir Rev. 2004;5 Suppl A:S119-26

2. Shi T, et al. Lancet. 2017;2;390(10098):946-958

3. Anyplex™ II, RV16-Detection, Seegene Inc.

4. World Health Organization (WHO). [https://www.who.int/influenza/surveillance\\_monitoring/ili\\_sari\\_surveillance\\_case\\_definition/en/](https://www.who.int/influenza/surveillance_monitoring/ili_sari_surveillance_case_definition/en/)