

## Unlocking pre-1850 instrumental meteorological records: A global inventory

Stefan Brönnimann, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, stefan.broennimann@giub.unibe.ch

Rob Allan, International ACRE Project Manager, Climate Monitoring and Attribution Group, Met Office Hadley Centre, UK, rob.allan@metoffice.gov.uk

Linden Ashcroft, School of Earth Sciences, University of Melbourne, Australia, linden.ashcroft@unimelb.edu.au.

Saba Baer, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, saba.baer@giub.unibe.ch

Mariano Barriandos, Department of History and Archaeology, University of Barcelona, Spain, barriandos@telefonica.net

Rudolf Brázdil, Institute of Geography, Masaryk University, Brno, and Global Change Research Institute, Czech Academy of Sciences, Brno, Czech Republic, brazdil@sci.muni.cz

Yuri Brugnara, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, yuri.brugnara@giub.unibe.ch

Manola Brunet, Centre for Climate Change, Department of Geography, University Rovira i Virgili, Tarragona, Spain, manola.brunet@urv.cat

Michele Brunetti, Istituto di Scienze dell'Atmosfera e del Clima (ISAC-CNR), Bologna, Italy, m.brunetti@isac.cnr.it

Barbara Chimani, Zentralanstalt für Meteorologie und Geodynamik, Vienna, Austria, barbara.chimani@zamg.ac.at

Richard Cornes, Koninklijk Nederlands Meteorologisch Instituut, De Bilt, Netherlands, now at: National Oceanography Centre, Southampton, UK, richard.cornes@noc.ac.uk

Fernando Domínguez-Castro, Instituto Pirenaico de Ecología (IPE-CSIC), Zaragoza, Spain, f.dominguez@ipe.csic.es

Janusz Filipiak, Department of Climatology and Meteorology, Institute of Geography, University of Gdansk, Gdansk, Poland, filipiak@ug.edu.pl

Dimitra Founda, Institute for Environmental Research & Sustainable Development, National Observatory of Athens, Athens, Greece, founda@noa.gr

Joelle Gergis, Fenner School of Environment and Society, ARC Centre of Excellence for Climate Extremes, Australian National University, Australia, joelle.gergis@anu.edu.au

Stefan Grab, School of Geography, Archaeology & Environmental Studies, University of the Witwatersrand, Johannesburg, South Africa, stefan.grab@wits.ac.za

Lisa Hannak, Deutscher Wetterdienst DWD, Lisa.Hannak@dwd.de

Ricardo García Herrera, Departamento Física de la Tierra y Astrofísica, Facultad de Ciencias Físicas, Universidad Complutense de Madrid and IGEO, Instituto de Geociencias (CSIC, UCM), Madrid, Spain, rgarciah@fis.ucm.es

Heli Huhtamaa, Institute of History and Oeschger Centre, University of Bern, Switzerland, heli.huhtamaa@hist.unibe.ch

Kim S. Jacobsen, Royal Museum for Central Africa, Leuvensesteenweg 13, Tervuren, Belgium, kim.jacobsen@africamuseum.be

Phil Jones, Climatic Research Unit, University of East Anglia, Norwich, UK (ORCID of PD: 0000-0001-5032-5493), P.Jones@uea.ac.uk

Sylvie Jourdain, Météo-France, Direction de la Climatologie et des Service Climatiques, Toulouse, France, sylvie.jourdain@meteo.fr

Andrea Kiss, Institute of Hydraulic Engineering and Water Resources Management, TU Vienna, Vienna, Austria, kiss@hydro.tuwien.ac.at

Kuanhui Elaine Lin, Taiwan, Research Center for Environmental Changes, Academia Sinica, Taipei, khelin@gate.sinica.edu.tw

Andrew Lorrey, National Institute of Water and Atmospheric Research, Auckland 1010, New Zealand, Andrew.Lorrey@niwa.co.nz

Elin Lundstad, Norwegian Meteorological Institute, Oslo, Norway, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, [elin.lundstad@giub.unibe.ch](mailto:elin.lundstad@giub.unibe.ch)

Jürg Luterbacher, Department of Geography, Climatology, Climate Dynamics and Climate Change, Justus-Liebig-University Giessen, Giessen, Germany and; Centre for International Development and Environmental Research, Justus Liebig University Giessen, Giessen, Germany, [juerg.luterbacher@geogr.uni-giessen.de](mailto:juerg.luterbacher@geogr.uni-giessen.de)

Anders Moberg, Department of Physical Geography, Stockholm University, Sweden, [anders.moberg@natgeo.su.se](mailto:anders.moberg@natgeo.su.se)

Franz Mauelshagen, Centre for Global Cooperation Research, University of Duisburg-Essen, Germany, [mauelshagen@gcr21.uni-due.de](mailto:mauelshagen@gcr21.uni-due.de)

Maurizio Maugeri, Università degli Studi di Milano, Department of Environmental Science and Policy, Milan, Italy, [maurizio.maugeri@unimi.it](mailto:maurizio.maugeri@unimi.it)

Nicolas Maughan, Aix-Marseille University, I2M UMR-CNRS 7373, Marseille, France, [nicolas.maughan@gmail.com](mailto:nicolas.maughan@gmail.com)

Raphael Neukom, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, [neukom@giub.unibe.ch](mailto:neukom@giub.unibe.ch)

Sharon Nicholson, Florida State University, Tallahassee FL, USA, [snicholson@fsu.edu](mailto:snicholson@fsu.edu)

Simon Noone, Irish Climate Analysis and Research Units (ICARUS), Department of Geography, Maynooth University, Maynooth, Ireland, [simon.noone@mu.ie](mailto:simon.noone@mu.ie)

Øyvind Nordli, Norwegian Meteorological Institute, Oslo, Norway, [oyvind.nordli@met.no](mailto:oyvind.nordli@met.no)

Kristín Björg Ólafsdóttir, The Icelandic Meteorological Office, Reykjavík, Iceland, [kbo@vedur.is](mailto:kbo@vedur.is)

Petra R. Pearce, National Institute of Water and Atmospheric Research, Auckland 1010, New Zealand, [petra.pearce@niwa.co.nz](mailto:petra.pearce@niwa.co.nz)

Lucas Pfister, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, [lucas.pfister@giub.unibe.ch](mailto:lucas.pfister@giub.unibe.ch)

Kathleen Pribyl, Climatic Research Unit, University of East Anglia, UK, [k.pribyl@uea.ac.uk](mailto:k.pribyl@uea.ac.uk)

Rajmund Przybylak, Department of Meteorology and Climatology, Faculty of Earth Sciences, Nicolaus Copernicus University, Toruń, Poland, [rp11@umk.pl](mailto:rp11@umk.pl)

Christa Pudmenzky, Centre for Applied Climate Sciences (CACS), University of Southern Queensland, Christa.Pudmenzky@usq.edu.au

Dubravka Rasol, Meteorological and Hydrological Service of Croatia, Zagreb, Croatia, [rasol@cirus.dhz.hr](mailto:rasol@cirus.dhz.hr)

Delia Reichenbach, Institute of Geography and Oeschger Centre, University of Bern, Switzerland, [delia.reichenbach@giub.unibe.ch](mailto:delia.reichenbach@giub.unibe.ch)

Ladislava Řezníčková, Institute of Geography, Masaryk University, Brno, and Global Change Research Institute, Czech Academy of Sciences, Brno, Czech Republic, [ladkar@sci.muni.cz](mailto:ladkar@sci.muni.cz)

Fernando S. Rodrigo, Department of Chemistry and Physics, University of Almería, Spain, [frodrigo@ual.es](mailto:frodrigo@ual.es)

Robert Rohde, Berkeley Earth Berkeley, CA, USA, [robert@berkeleyearth.org](mailto:robert@berkeleyearth.org)

Christian Rohr, Institute of History and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland, [christian.rohr@hist.unibe.ch](mailto:christian.rohr@hist.unibe.ch)

Oleg Skrynyk, Ukrainian Hydrometeorological Institute, Kyiv, Ukraine, [skrynyk@uhmi.org.ua](mailto:skrynyk@uhmi.org.ua)

Victoria Slonosky, McGill University, Montreal/Montreal, Canada, [victoria.slonosky@mail.mcgill.ca](mailto:victoria.slonosky@mail.mcgill.ca)

Peter Thorne, Irish Climate Analysis and Research Units (ICARUS), Department of Geography, Maynooth University, Maynooth, Ireland, [peter.thorne@mu.ie](mailto:peter.thorne@mu.ie)

Maria Antónia Valente, Instituto Dom Luiz, University of Lisbon, Lisbon, Portugal, [mavalente@fc.ul.pt](mailto:mavalente@fc.ul.pt)

José M. Vaquero, Centro Universitario de Mérida, Universidad de Extremadura, Spain, [jvaquero@unex.es](mailto:jvaquero@unex.es)

Nancy E. Westcott, Midwestern Regional Climate Center, Illinois State Water Survey, University of Illinois, IL, USA, [nan@illinois.edu](mailto:nan@illinois.edu)

Fiona Williamson, School of Social Sciences, Singapore Management University, Singapore, [fwilliamson@smu.edu.sg](mailto:fwilliamson@smu.edu.sg)

Przemysław Wyszynski, Department of Meteorology and Climatology, Faculty of Earth Sciences, Nicolaus Copernicus University, Toruń, Poland, [Przemyslaw.Wyszynski@umk.pl](mailto:Przemyslaw.Wyszynski@umk.pl)

## **Abstract**

Instrumental meteorological measurements from periods prior to the start of national weather services are designated “early instrumental data”. They have played an important role in climate research as they allow daily-to-decadal variability and changes of temperature, pressure, and precipitation, including extremes, to be addressed. Early instrumental data can also help place 21<sup>st</sup> century climatic changes into a historical context such as to define pre-industrial climate and its variability. Until recently, the focus was on long, high-quality series, while the large number of shorter series (which together also cover long periods) received little to no attention. The shift in climate and climate impact research from mean climate characteristics towards weather variability and extremes, as well as the success of historical reanalyses which make use of short series, generates a need for locating and exploring further early instrumental measurements. However, information on early instrumental series has never been electronically compiled on a global scale. Here we attempt a worldwide compilation of metadata on early instrumental meteorological records prior to 1850 (1890 for Africa and the Arctic). Our global inventory comprises information on several thousand records, about half of which have not yet been digitized (not even as monthly means), and only approximately 20% of which have made it to global repositories. The inventory will help to prioritize data rescue efforts and can be used to analyze the potential feasibility of historical weather data products. The inventory will be maintained as a living document and is a first, critical, step towards the systematic rescue and re-evaluation of these highly valuable early records. Additions to the inventory are welcomed.

## **1. Introduction**

As Enlightenment scientists initiated regular meteorological measurements in the 17<sup>th</sup> and 18<sup>th</sup> centuries (Wolf 1962), scientists today have access to a wealth of early weather and climate information from across Europe (Jones 2001), and also other parts of the world. Very long series such as Central England temperatures (start date 1659; Manley 1974, Parker et al. 1992) or Paris temperatures (1658; Rousseau 2015) are widely used as a baseline for current temperature changes and to study past climatic variations at regional scale. The same holds for long rainfall series such as those for Paris (1688; Slonosky 2002), Ireland (1716; Murphy et al. 2018), and Seoul (1770; Arakawa 1956). A considerable number of other long, mostly European, instrumental records have been published (*e.g.*, Wheeler 1995, Moberg and Bergström 1997, Jones and Lister 2002, Moberg et al. 2002, Bergström and Moberg 2002, Maugeri et al. 2002ab, Camuffo et al. 2006, Bryś and Bryś 2010ab, Cornes et al. 2011ab, 2012, Brázdil et al. 2012). While these records shed light on past climatic variations, they constitute only a subset of all measurements taken. Here we aim at providing an inventory, still far from comprehensive, of where, when, and by whom meteorological measurements were made prior to ca. 1850.

Numerous efforts by individuals, weather services, international projects, and efforts coordinated by the Atmospheric Circulation Reconstructions over the Earth Initiative (ACRE, Allan et al. 2011) form the basis of our inventory. A selection of long European daily records (from Italy, Spain, Sweden,

Belgium, Russia) of air temperature and air pressure, some reaching far back into the 18th century, were compiled within the project IMPROVE (Camuffo and Jones 2002). The Austrian-led dataset HISTALP (Auer et al. 2007) collected and processed long instrumental series for temperature, pressure, precipitation, cloudiness, sunshine duration, water vapor pressure and relative humidity back to 1760. The project ADVICE compiled monthly mean air pressure data for several sites in Europe back to 1780 (Jones et al. 1999). Early instrumental series were compiled for Portugal and its colonies (Alcoforado et al. 2012), Spain (Barriendos et al. 2002, Domínguez-Castro et al. 2014, Prohom et al. 2016, Sanchez-Rodrigo 2019), Italy and the Western Mediterranean (Cantù and Narducci 1967, Brunetti et al. 2001, 2006, Camuffo and Bertolin 2012, Camuffo et al. 2013, 2017), the Mediterranean North Africa and the Middle East (Brunet et al. 2014, Ashcroft et al. 2018), the Czech Republic (Brázdil et al. 2005, 2012), and Poland (Przybylak and Pospieszyńska 2010, Przybylak et al. 2014). Canadian data were published by Slonosky (2003, 2014) and long series compiled in the US by Burnette et al. (2010). Historical USA daily weather data were archived and imaged by NOAA under the Climate Data Modernization Program, with more than 15,000 station months (more than 140 stations) digitized for the period prior to 1850 (Dupigny-Giroux et al. 2007, Westcott et al. 2011). Domínguez-Castro et al. (2017) compiled early instrumental series for Latin America, while Ashcroft et al. (2014, 2016) and Gergis and Ashcroft (2013) provide data for southeastern Australia and Williamson et al. (2018) for southeastern Asia. For an overview of data in Africa see Nash and Adamson (2014). A global inventory of these records, however, does not exist yet.

Interest in such historical weather data is not new. Scientists in the early 18<sup>th</sup> century compiled meteorological data in their efforts to study and understand weather and climate in different parts of the world (Maraldi 1709, Derham 1735, Hadley 1741, 1744, Wargentin 1758, Kirwan 1787). In the 19<sup>th</sup> and early 20<sup>th</sup> century, numerous inventories were compiled (*e.g.*, Schouw 1839, Dove 1838, 1839, 1841, 1842, 1845, 1852, Blodget 1857, Schott 1876ab, 1881, Angot 1897, Hellmann 1883, 1901ab, 1927, Supan 1898, Eredia 1912, 1919). Data were used for regional climate descriptions (*e.g.*, Kreil 1865, Strzelecki 1845) or for the construction of isothermal maps (Humboldt 1817, Berghaus 2004, Dove 1852). Although attempts were made again in the 1970s through 2000s to systematically compile 18<sup>th</sup> century records (Kington et al. 1988), most historical inventories are nowadays largely forgotten, mainly because the metadata from these inventories have never been digitized.

Dove's articles did not just include summaries of the data and source details, but monthly averages for over a thousand series. These were used in Northern Hemisphere temperature estimates produced in the early 1980s (Bradley et al. 1985; Jones et al. 1985, 1986). Almost 100 years earlier, these same Dove sources, together with data published in European meteorological journals were used by Köppen (1873, 1881) to produce the first estimate of average temperatures in the Northern Hemisphere.

Large fractions of the data listed in these inventories, particularly the shorter series, have not been digitized. The value of such shorter series is well recognized today. Historical reanalyses such as the “Twentieth Century Reanalysis” (20CR, Compo et al. 2011) are able to generate useful weather reconstructions from records of only a few years, as demonstrated for the “Year Without a Summer” of 1816 (Brugnara et al. 2015; Brohan et al. 2016). A backward extension not just of climate data, but also weather data contributes towards a better understanding of extreme, and thus rare, events. It allows important climate processes to be addressed such as the transit of the climate system out of Little Ice Age climate, the impacts of volcanic eruptions or natural, interannual-to-decadal, variability. However, this requires rescuing additional historical weather data (Brönnimann et al. 2018a). Together with other sources, they also allow climate data to be connected with climatic impacts and associated societal responses (*e.g.*, resilience, adaptation).

Here we provide an initial, systematic global compilation of information on early instrumental data. The paper documents efforts within ACRE, the International Surface Temperature Initiative (Thorne et al. 2011) as well as results from a workshop held in Bern, Switzerland, 18-21 June 2018. Experts from all parts of the world are involved in this initiative (the authors of this paper represent 26 different countries) and have contributed to making the inventory as comprehensive as possible.

The focus is on instrumental series with regular (daily or more frequent) measurements over at least one year prior to 1850, even if some of the data have so far not been found. The inventory contains the relevant meta information when and where available (station name, coordinates, altitude, observer, source) as well as information on availability or state of data rescue. Though far from complete, our inventory, along with data rescue services (Brönnimann et al. 2018b), aims to support future data rescue efforts. It should also help to counteract the danger of losing data. Climate data are societal products (Brönnimann and Wintzer, 2019) and form part of our cultural heritage. The inventoried information may also interest historians and experts of related disciplines.

The paper is organized as follows. Section 2 describes the generation of the inventory. Section 3 provides an overview of the stations inventoried, starting with a global overview and then focusing on World Meteorological Organization (WMO) regions. Conclusions follow in Section 4.

## **2. Methods**

### *2.1. Criteria for data collection*

Before inventorying the data, we defined criteria for collection. We defined 1850 as a cut-off year because this approximately reflects the start of national weather services (*e.g.*, Prussian Meteorological Institute 1847, Smithsonian Institution network 1849, weather service of the Austrian empire 1851, French meteorological service at the Observatoire de Paris 1855); this makes 1850 also

broadly the beginning of international standards. For the same reason, many global data sets reach back to around that time. For instance, CRUTEM4 (Jones et al. 2012) and 20CRv2c go back to 1850 and 1836, respectively.

However, for some regions the 1850 cut-off is too early. Africa has almost no measurements prior to 1850, while much data from the second half of the century also have never been systematically compiled. The same is true for the Arctic (Przybylak et al. 2010). For these regions, we therefore set a later cut-off at 1890. Series which are listed in inventories published prior to 1850, but for which measurement years were not given, are also included (in the figures they are attributed to the years immediately before the inventories were published).

A second criterion is that only metadata on instrumental measurements was searched. We did not consider wind observations (only wind measurements) or the number of rainy days, even though these are quantitative data that may be of high quality. This information could be compiled at a later stage in an inventory for non-instrumental observations. The same holds for inches of rain and snow in China (*Yu-xue-feng-cun*, Pei et al. 2018). Unlike the measurement of depth in a snow/rainfall gauge, these numbers refer to the depth of soil seeped through by rainwater for example (Ge et al. 2005, Wang and Zhang 1988). Although widely analyzed for climate reconstructions (Hao et al 2012, Ding et al. 2014) we do not consider such data in the present inventory.

A third criterion concerned a minimum length. As a rule of thumb, we compiled only data series with a length of at least one year, although this lower limit was sometimes disregarded for very early data or shorter records from remote regions. If the length of the record was not known, the information was collected nevertheless (in the figures they appear as one year long).

Marine data are not considered in this inventory, except for the “stationary ships” of the European powers moored in the harbors of their various colonies around the world. To date, most of the ships identified are British vessels, serving as regional command posts, hospitals, coaling stations, guard ships, convict ships, receiving ships, store ships/depots, mooring ships, and hulks. Other European powers are highly likely to have had similar vessels stationed in their major colonial ports.

## 2.2. Sources used

Our procedure consisted of (1) searching existing digital data repositories, (2) searching readily available national inventories, (3) compiling community knowledge through ACRE, through the Bern workshop, and through further involvement with the community, (4) compiling early (non-digital) inventories and (5) compiling early collections and networks.

- (1) We searched 5 global repositories (Table 1). These contained the series as well as some metadata (here we only compile the metadata). Many of the series were found in several of the repositories. In total, 741 unique records were found.

- (2) We next searched regional and national data repositories or metadata inventories (Table 1). These comprised around 1500 records, but with a large overlap with (1).
- (3) The community knowledge compiled was particularly important for those regions for which no repositories exist, and for time periods before the foundation of national agencies. The inventory for Europe and North America was greatly extended by the authors of this paper.
- (4) A number of early inventories were consulted where known and recoverable, some covering the globe, some covering certain regions or countries. Typically these inventories contain only metadata, often in condensed form (Fig. 1). Note that the inventories may not be free of errors or ambiguities, and many errors were corrected by the authors.
- (5) From the earliest days of meteorology, scientists collected and published data (Table 2); some of which, such as the Societas Meteorologica Palatina (SMP) Network, can be considered coordinated networks.

### *2.3. Metadata compiled*

The inventory should be able to store comprehensive meta information, but should also collect sparse information (*e.g.*, a reference pointing to the existence of a measurement series, even without measurement years, observers, or exact location). The following information was compiled:

- (1) existing station identifications (WMO, national, or other repositories);
- (2) the station location: continent, modern country, location, other names, location details, coordinates, elevation, WMO region;
- (3) observer name and context (*e.g.*, profession of observer, missionary or military contexts, etc.), variables, measurement frequency;
- (4) inventory in which it is contained, source of metadata and data (see also Table 3);
- (5) start and end year and availability in present repositories;
- (6) status on digitizing and availability; and
- (7) comments.

Comments are used, for instance, to mark data which has already been investigated and discovered to be unusable, or data which have been irretrievably lost.

### *2.4. Quality control of metadata*

After compiling the metadata, a quality control of the metadata was performed. Coordinates were tested against station names, countries, and a land-sea mask. Errors found in this procedure were corrected. Duplicates were identified, suspicious series were checked in the original sources (and sometimes identified as non-instrumental) and series starting after 1850 (or 1890 for Africa and the Arctic) or too short series were flagged. During this procedure, no information was removed. In case of duplicates, the unique information was combined into one record and the duplicates were flagged, with a cross-reference to the primary record.

Duplicate removal proved difficult. Often duplication was only partial; both entries were then left in. Some series were made up of multiple stations if within a certain distance, and some have overlapping periods. Some series have unique entries per variable (which may have different start and end dates), some have a unique entry per observer, or per source. We did not attempt to unify this. “Entry” in the following may therefore mean one long multi-variable record of a station, or only a segment, or just one variable.

### 3. Results

#### 3.1. Global overview

The inventory currently has 10,280 entries, the majority of which is flagged as duplicates, post-1850, too short, or non-instrumental. The remaining 4,626 entries are from 2,250 locations. There are still numerous partial duplicates. Plotted as a map (Fig. 2), duplicates lie on top of each other and do not distort the result. Almost all early and long series are from Europe. Long records, though reaching less far back, also exist for New England and Canada. Early records from other WMO regions are sporadic and short. These data are mostly from expeditions and colonial endeavors. A close to global (though sparse) coverage, with continuous series in all WMO regions except Africa, is reached only after 1800. For Africa and the Arctic, good coverage emerges only in the period 1851-90. No entry exists for Antarctica (first measurements were taken during the *S.Y. Belgica* expedition 1897-99; Arctowski 1904; see also Jones 1990).

The first series start in the 1650s (Fig. 3). From the 1680s onward at least 10 records are inventoried in any year (but some are known to be lost). The number increases to approximately 50 in 1720, reaching the hundreds by 1800. The increases in the early 18<sup>th</sup> century as well as the peak in the late 18<sup>th</sup> century indicate coordinated activities, such as those of Johann Kanold and James Jurin in the 1710s and 1720s, and the SMP network in the 1780s (Table 2). The 30% drop in the late 18<sup>th</sup> century corresponds to the politically unstable Napoleonic period in Europe. The inventory lists the status (*e.g.*, imaged, transcribed), although not systematically. A brief analysis shows that ca. 25% of entry years are fully transcribed another 25% partly so (*e.g.*, as monthly means). Only around 20% of entry years are available from global repositories (Table 1), indicating that a considerable fraction of the transcribed data has not yet made it into these repositories. Table 3 provides URL's of important data collections. In the following results are discussed by WMO regions (plus Arctic).

#### 3.2. WMO Region 1: Africa

For Africa, the start of instrumental meteorological measurements varies greatly by country. Rainfall records generally go back further than temperature records. The inventory map (Fig. 4) shows that the



starting year is mostly after 1800. A good fraction of the series has already been digitized but due to the sparse coverage, additional series are particularly valuable. Africa's oldest instrumental records come from *South Africa*. Although Robert Jacob Gordon refers to barometric pressure records going back to at least 1737 in Cape Town, the original records have yet to be found. Four short records are known from Cape Town in the following decades (1751-52 by explorer Nicolas-Louis de Lacaille, 1766-68 by Paulus Henricus Eksteen, 1779 by the Bataviaasch Genootschap and 1789-92 by R. J. Gordon). The first regional network was set up by Colonial district offices and covers the period 1818-27. However, there are large data gaps for several of these stations. In 1829, the Cape Town Port Office began meteorological measurements. Daily records the Royal Astronomical Observatory in Cape Town began in 1834 and continue to this day (stationary ships recorded weather at Cape of Good Hope, 1834-1912).

Outside of South Africa, instrumental records concern mostly rainfall, which relates to the importance of the amount of rain in the rainy seasons on all aspects of life, an emphasis that continues to this day. Most measurements were taken by colonial administrations. In some cases, regular measurements were made by missionaries, such as in *Malawi*, 1876-80 (Nash et al. 2018), or by individuals, such as the Urquiola sisters in *Equatorial Guinea* in 1875 (Gallego et al. 2011). In some cases, lengthy series of measurements were made during scientific expeditions, such as the Loango expedition of 1873-76 (Danckelman 1878, 1884). In general, the longest records are available for stations near the coast, with the far inland being nearly devoid of measurements until the 20<sup>th</sup> century.

In *Algeria*, the first two rain gauges were installed in 1837 in Algiers by the Administration of Drying and in Constantine in the military and colonial hospital. Before 1890, measurements were made by engineers from the National Road and Bridge Administration, by military doctors at the colonial hospitals and by military engineers. In 1865, 34 meteorological stations had been created in the main military hospitals. Rainfall data for Algeria were compiled by Raulin (1875b).

In *Senegal*, records begin in the 1830s (all available pre-1900 measurements have been compiled by Nicholson et al. 2012a). Meteorological measurements, made by military pharmacists at the military colonial hospitals, started before 1860 in Bakel, Podor, Saint-Louis and Gorée (Raulin 1875a).

In *Madagascar*, measurements were taken at the military and colonial hospitals on Nosy Be (Nossi Be) Island in 1855 and on Nosy Bohara (Sainte-Marie) Island in 1863. Colin (1889) compiled the measurements in Antananarivo starting with Laborde, the first French consul in Madagascar, 1872-78, and followed by the Fathers of the Catholic mission, 1879-89 (a short record from 1829 is known).

Less extensive records are available for *Egypt*, *Morocco*, *Tunisia*, and *Namibia*, with numerous stations in these countries starting in the 1870s or 1880s (in Egypt in the 1860s). In the same period, stations were established by the Germans in *Togo*, *Tanzania*, and *Cameroon*; by the British in *Sudan*, *Ghana*, *Uganda*, *Kenya*, *Zimbabwe*, *Lesotho*, and *Malawi*; and by the Spanish in the *Canary Islands*.

Isolated measurements from this time were also made in the Portuguese colonies of *Angola*, *Mozambique* and the *Cape Verde Islands*, and in the Italian colonies of *Eritrea* and *Libya*. Rainfall records from these and other sites are described in Nicholson (2001) and Nicholson et al. (2012ab). Several earlier records are also noteworthy. Rainfall was recorded at the Danish Fort at Christianborg (today Accra, *Ghana*), 1829-34 and 1839-42. Records exist for Freetown, *Sierra Leone*, 1793-95 (rainfall, temperature, pressure), 1819, 1828, and 1847-51 (rainfall only) and since 1849 (temperature; plus measurements onboard stationary ships 1832-41 and 1861-67). Short records are available for *Zanzibar* since 1850, with continuous data since 1874. Measurements started in Luanda, *Angola*, in 1858. Sporadic records in the late 1850s and early 1860s exist for Bukoba, *Uganda*; Natete, *Mozambique*; Libreville and Sibange Farm, *Gabon*; Bathurst, *Gambia*; Grand Bassam, *Ivory Coast*; and Lagos, *Nigeria*.

For the *Congo Basin*, the frequency of measurements follows colonial history. Shortly after King Leopold II of Belgium declared the Congo Free State his personal territory, a publication was issued containing meteorological measurements for the coastal village of Banana (Etienne 1892). More commonly however, early measurements remained unpublished, and mostly in personal archives of explorers. Often measurements are merely sporadic, such as those by Cyriaque Gillian and Francis Dhanis noted in their travel diaries for 1889 and 1890-91, respectively.

The earliest measurements for *Mauritius* were made by French colonial residents and administrators, Jean-Nicolas Céré, Director of the Botanical Gardens at Pamplemousses (1770-90), the astronomer, botanist and cartographer Jean-Baptiste Lislet-Geoffroy at Port Louis (1784-1834) and Mr Labutte at Yemen (1812-47). A mix of French and English colonials followed, such as Lt.-Col. John Augustus Lloyd, Surveyor-General of Mauritius (1831-49) at the ‘Government Observatory’ at Port Louis, and Julien Desjardins, the first Secretary of the Natural History Society of Mauritius, at Flacq (1836-38). Most of these measurements are lost or only survive in summary form. For *Réunion*, various sources suggested that Jean Nicolas Céré took, or compiled, instrumental records from the late 18<sup>th</sup> century (but these have never been found). Stationary ships provide measurements for *Ascension Island* (1844-71) and *St. Helena* (1819-31).

### 3.3. WMO Region 2: Asia

The first measurements in Asia date back to the late 17<sup>th</sup> century, when Scottish surgeon and naturalist James Cunningham made meteorological measurements on his journey to China. Early records are also available for India (Fig. 5 left). The longest continuous series are rainfall in Seoul as well as temperature, pressure, and rainfall in Chennai, both reaching back to the 18<sup>th</sup> century. Unlike Africa, the records concern not only rainfall, but also pressure and temperature. Most records start after 1800. Various data rescue activities are underway. However, many records, including many shorter fragments, have not yet been digitized.

*China* holds a wealth of documentary climate data (compilation digitized in the REACHES data set, Wang et al. 2018), but measurements are scarce. The earliest known measurements were made by Cunningham at Xiamen (Amoy) from October 1698 to January 1699. Between 1700 and 1702, a continuous record was also made by Cunningham at Zhoushan, Zhejiang Province. Beijing temperatures were measured by P. Antoine Gaubil in 1743. Missionary Joseph Amiot recorded temperature, pressure, and rainfall in Beijing, 1757-62. The Russian observatory network also made measurements at their Consulate in Beijing from 1841.

Guangzhou (Canton) has records for 1771-74, 1785, 1789, 1804, 1829-31, 1830-39, 1836-38, and then periods in the 1840s. Some of these data can be found in old newspapers such as the “Canton Register”, including measurements made in Hong Kong (Tsukahara 2013). An early dataset was recovered for *Macau* in 1780. Measurements were also made in 1787 by the French orientalist Joseph de Guignes. For the period 1830-37 temperature, pressure and rainfall measurements exist in printed journals for Hong Kong and Macao. A couple of years for daily measurements of temperature and pressure at Macao (1840-41) were also published in the “Canton Register”.

*Japan* holds exceptionally early instrumental datasets (Zaiki et al. 2006). Early measurements were made by foreigners such as Carl Peter Thunberg in 1775/76 and 1779 (Demarée et al. 2013), J. Cock Blomhoff, 1819-23, and von Siebold, 1825-28, all in Nagasaki (Können et al. 2003). Other station records include Osaka, 1828-71, and Tokyo, 1825-75 (all digitized; Table 3). In the Asiatic part of *Russia*, the longest series, Irkutsk and Jakutsk, reach back to the 1820s (digitized; Table 3).

The earliest measurements made on the Indian sub-continent were by European missionaries in Puducherry and Chandannagar in the 1730s and 1740s. More regular measurements were later carried out at Chennai and Calcutta by British colonial officers. At Fort Saint George in Chennai, daily meteorological measurements were recorded in 1776-78, by respectively the assistant surgeon Dr William Roxburgh and Colonel James Capper. The next instrumental record from Chennai is from William Petrie in 1787 (India Meteorological Department 1976) and from missionaries in 1789-91 (Walsh et al. 1999). From 1791 onward rainfall, temperature and pressure were measured at an observatory set up by the East India Company. The records from John Goldingham in 1793 (India Meteorological Department 1976) and successive astronomers until 1843 (Goldingham 1826, Goldingham and Taylor 1844) are gradually being digitized under ACRE. Monthly mean air pressures taken at the Chennai Observatory from 1796-2000 were recovered (Allan et al. 2002).

In Calcutta (Kolkata), the earliest known measurements are those made by British colonial officers, such as Henry Traill, Treasurer of the Asiatic Society of Bengal, 1784/85 (Traill 1790) and Colonel Thomas Dean Pearce, 1785/86 (Pearce, 1788). Temperature data from 1816 onward (Hardwicke 1830) are now in global repositories.

Pressure time series prior to 1850 have been recovered and digitized for Benares/Varanasi (1823-27) (Prinsep 1829ab), Bangalore (1830-35) (Kingsford 1843) and others (see also Adamson and Nash 2014) as well as for locations in *Nepal* (including Kathmandu) in 1802/3 (Hamilton 1819). Long runs of such data tend to begin from the 1840s onwards, such as for Hoshungabad (from 1849). Data from stationary ships exist from Trincomalee (Sri Lanka) (1819-64) and Hong Kong (1842-1920).

#### 3.4. WMO Region 3: South America

The longest instrumental record from South America is temperature from Lima, *Peru*, which reaches back to 1754 but is only available as annual minimum and maximum temperatures. Another noteworthy record is from Rio de Janeiro, 1781-88 (Farrona et al. 2012). Almost all other records inventoried start after 1800 (Fig. 5 right). Fewer records are available than for Asia, mostly less than 10 records at any time. Most records include at least temperature. No systematic inventories are currently available and data collected prior to the foundation of the National Meteorological Services is dispersed among a variety of archives. The most important effort to collect measurements prior to 1900 was made recently by Domínguez-Castro et al. (2017) who retrieved more than 300,000 meteorological measurements from 20 countries in South and Central America (Table 3).

This effort allowed the identification of some of the most relevant sources for the region. A large number of the measurements made during the colonial period were recorded with the objective of characterizing the climate of the region to evaluate possibilities to improve agricultural production as well as the influence of climate on health. The main observers in the region were scientists, explorers and military personnel.

Explorers recorded meteorological data, but their series are short because they changed their observation locations frequently. They used portable instruments, sometimes with poor calibration processes. Measurements by explorers are usually recorded in printed books that describe the explored territories. Unfortunately, as the meteorological tables took up many pages, the measurements were usually summarized as monthly means (*e.g.*, Boussingault 1849). A compilation of Spanish explorers in the region is given in Lucerna Giraldo (2008).

The longest series were recorded by researchers with the intention of starting a national observatory (generally astronomical and meteorological) or involved in academic initiatives, as José Celestino Mutis and José de Caldas in *Colombia* or Francisco Aguilar in *Ecuador*. The continuity of the series depends on the support they could obtain. Generally, these measurements were performed at sub-daily scale but the manuscripts of the measurements were frequently lost or remain undiscovered.

During the early 19<sup>th</sup> century most of the countries suffered from wars of independence. Most of the series have gaps during this period and it is likely that some original records were lost during the wars because many observatories were used for military proposes. Stationary ship data are listed in the inventory for Rio de Janeiro (1840-75) and Valparaiso (1843-79).

### 3.5. WMO Region 4: North America, Central America and the Caribbean

Sporadic instrumental measurements from North America reach back to 1697. The first actual series started in the 1730s and 1740s in Charleston and in various places in Massachusetts, Quebec, and Philadelphia (Fig. 6). Havens (1958) inventories the early instrumental data (see also van der Schrier and Jones 2008, for an example of use of long early US records). Many of the early (prior to ca. 1820) series listed in our inventory have not yet been digitized, so there is potential for data rescue activities in North America. One of the most important collections of historical climatic data for North America originates from the efforts of the Smithsonian Institute, led by Joseph Henry, to collect meteorological measurements from across North America to compile statistics on the climate for agricultural purposes, and later to issue storm warnings (beginning in the 1840s). Although the active exchange of contemporary meteorological records by the Smithsonian started in the 1850s, the Smithsonian also became a repository for older climatic records. These records were compiled and analyzed in several publications over the course of the 19<sup>th</sup> century, including publications by Blodget (1857), Hough (1872), and Schott (1876ab). The “Forts” dataset, with information from military and other early measurements sites across the US, has 142 stations beginning before 1850 (Dupigny-Giroux et al. 2007, Westcott et al. 2011).

Among the earliest systematic measurements taken in North America are those kept by the medical establishment, including the *Médécins du Roi* under the French colonial regime, the US Army Medical Department in the War of 1812, and monthly and quarterly reports sent to the US Army Surgeon General, 1820-59. Other military units which kept systematic records include the Royal Artillery and the Royal Engineers. Observers at colleges and universities made measurements for the purposes of agricultural planning, and individual volunteers also made important contributions.

Many early records from eastern Canada have been digitized with the help of citizen science efforts. A nearly continuous daily temperature series has been developed for the St Lawrence Valley region back to 1798, and more sporadically to 1742 (Slonosky 2015). Missionaries, explorers and Hudson’s Bay Company (HBC) factors provided important measurements from the north and northwestern interior regions of the continent. Over 40 station records were found in the HBC and Royal Society archives for northern Canada, 1771-1840, in pioneering work by climatologist Cynthia Wilson in the 1980s.

The earliest instrumental series in *Mexico* were recorded by Jose Antonio Alzate in Mexico City in 1769. In the early 19<sup>th</sup> century some short series have been recorded mainly in the central region. The digital archives of the Spanish National Library contain records included in the Mexican newspapers “*Gazeta del Gobierno de Mexico*” “*Diario del Gobierno de la Republica Mexicana*”. They include around four years of monthly data (1827-30) from Orizaba, one year (1831) from Veracruz and daily data from Mexico City for some period of 1833 and 1842-43.

In Central America and the Caribbean, some records reach back to the 18<sup>th</sup> century. The Royal Engineers network had stations in *Bahamas, Jamaica, Honduras* and *Barbados*. Records exceeding 10 years are listed for *Savanna-la-Mar, Jamaica* (1764-76), *Saint-Domingue* (1772-84), *St. Barthélemy* (1786-96), and *Havana, Cuba* (starting in 1791). In *Guadeloupe* measurements started in the 1780s; from the 1820s they were taken in the military hospitals by pharmacists or doctors. Charles Saint-Claire Deville, geologist and meteorologist, co-founder of the Société Météorologique de France and founder of the Paris-Montsouris observatory compiled several short temperature and pressure series measured in West Indies, 1840-50. Stationary ships are a source of meteorological measurements for *Bermuda* (1824-1904), *Havana* (1837-45), and *Jamaica* (1823-1903).

### 3.6. WMO Region 5: South-West Pacific

In *Malaysia*, measurements were made at Prince of Wales Island in Penang/George Town between 1815-16 and 1820-23 and at Malacca in 1809. The Royal Society in London holds a set of subdaily pressure, temperature and rainfall measurements kept at Penang, 1843-45. At *Singapore*, measurements were made from at least 1820 virtually continuously through until 1845, at which point there is currently a data gap until 1861. No datasets longer than one year are available for *Indonesia* (Dutch East Indies) pre-1850 except for West Java at Buitenzorg (Bogor), 1841-55 and for Padang, 1850-53 (both digitized by KNMI).

Pre-1850s land-based measurements for *Australia* date back to European settlement on the continent in 1788 (Fig. 7). They were taken primarily at the main colonial centers of population in southern Australia such as Sydney, Hobart, Melbourne, Adelaide and Perth. However, other locations such as Albany (Frederick Town), the oldest colonial settlement in Western Australia, Launceston, Port Arthur (then a major penal settlement) and various properties of the Van Diemen's Land Company (such as in the Hampshire Hills) in Tasmania also kept records starting in the 1830-40s. Records should also exist for Brisbane, taken at the General Hospital, 1825-44 (Bartkey 2008, Ashcroft et al. 2016). Consequently, the spatial and temporal coverage of the pre-1850 data map (Fig. 8) closely relates to the expansion of English settlements, with early records from coastal regions and penal colonies in the southeast, before moving inland and westward with explorers and farming communities.

The earliest land-based measurement of temperature and pressure were recorded by Lieutenant William Dawes, 1788-91, from his observatory in Sydney Cove (Gergis et al. 2009). A detailed collection of 39 pre-1860 instrumental data sources for south-eastern Australia is provided by Ashcroft et al. (2014). They show that only sporadic measurements have so far been uncovered for the 1790-1820 period, with an increase in the number of records available from the mid-1820s.

Measurements have also been found in newspapers, explorers' journals, colonial diaries, convict settlement documents, official observatory and government publications, doctors' records and private

letters. A short-lived network of stations existed at penal colonies along the New South Wales and Tasmanian coast in 1822. More continuous records start later, such as temperature measurements for Sydney, 1826-48, and measurements made in Melbourne, Sydney and Port Macquarie, 1840-51 under Government order. Instrumental measurements were made on various expeditions, e.g., by John Oxley to Bathurst, New South Wales in 1823, or by Sir Thomas Livingstone Mitchell in south-eastern Australia during 1831-32, 1835, 1836 and 1845-46 (Mitchell 1838), and are being digitized under ACRE Australia.

Recent efforts are currently underway to develop near-continuous daily records dating back to the early 1830s for the cities of Adelaide and Perth. The former consists of measurements made by Dr William Wyatt in Adelaide from 1838–43 (Ashcroft et al. 2014), newly imaged data from 1843-56 held by the National Archives of Australia, and available data from the Adelaide Observatory from 1856. Measurements of temperature and pressure from the Swan River settlement area of Perth, 1830–74, have recently been digitized. These records, in addition to the extensive data available for Sydney (Ashcroft et al. 2014), improve the possibility of developing long instrumentals records back to 1830 for the underrepresented Southern Hemisphere. In Hobart, a stationary ship provided measurements for 1844-51. Additional pre-1850 instrumental data and supporting metadata around the scientific practices used in colonial Australia continue to be recovered around the country, many through citizen science activities (Ashcroft et al. 2016).

Limited land-based instrumental climate data are available before 1850 for *New Zealand*. The earliest known multi-year measurement record is the diary of Reverend Richard Davis, an English missionary stationed at Waimate North and Kaikohe, Northland, in the 1830s-1850s. Davis' weather diary covers nine years within the 1839-51 timespan (Lorrey and Chappell 2015). In addition, pre-1839 instrumental measurements are provided by Davis from 1836 onward in his personal diary, but these are sporadic.

Old newspapers also contain printed measurements. Barometric pressure and temperature have been found for 1840-43 in the “New Zealand Gazette” and “Wellington Spectator” for Wellington. Similar data for 1842-1843 have been found in the “Nelson Examiner” and “New Zealand Chronicle”, however no metadata has been found about the location of these measurements.

### 3.7. WMO Region 6: Europe

The first European measurements reach back to the 17<sup>th</sup> century. A good coverage of Central Europe is reached by the first half of the 18<sup>th</sup> century (Fig. 9). The records usually comprise temperature and pressure, less often precipitation. A particularly dense coverage is reached in the 1780s, when many networks were active (Table 2). Although many of these series have been digitized, an even greater coverage could be reached for this period and for the first half of the 18<sup>th</sup> century.

*Italy* was home to the “Medici Network”, the first international network of meteorological measurements established within the sphere of the Accademia del Cimento in Tuscany by the Grand Duke Ferdinand II de’ Medici and his brother Prince Leopold (Camuffo and Bertolin 2012). The scientists of the Accademia del Cimento invented instruments and performed well-coordinated measurements with identical instruments. Observational activities were extended beyond the borders of Florence, with stations in Vallombrosa, Pisa, Cutigliano, Bologna, Parma, Milan, Paris, Innsbruck, Osnabrück and Warsaw. The data were digitized and found to be of useable quality (Camuffo and Bertolin 2012). Italy also has several other long records (Brunetti et al. 2006).

In mainland *France*, instrumental measurements were carried out by astronomers, doctors and Jesuits during the late-17<sup>th</sup> and the 18<sup>th</sup> century. Paris holds the longest temperature (back to 1658), pressure (1670) and precipitation records (1688), and several other stations started in the first half of the 18<sup>th</sup> century. An early collection was compiled by Louis Cotte (Table 2). Monthly weather reports, stored in the library of the Académie de Médecine in Paris, were sent to him by 173 doctors during the period 1776-1793, until the French revolution put an end to this network. Météo-France’s inventory shows the gap during the period 1795-1805 with only 14 French stations, including the two astronomical observatories Paris and Marseille. For the Southeast of the country, the history of instrumental measurements was compiled by Pichard (1988).

Northern Europe is relatively well covered. For *Iceland* our inventory lists 49 series prior to 1850. The earliest available measurements from *Sweden* (see Moberg 1998) are from 1719-23, though from an unknown site. Measurements in Uppsala started in 1722, arranged by astronomers at the Society of Science in the city and apparently in connection with Jurin’s invitation (Jurin 1723). This became the start of what is now the longest instrumental record in Sweden, as measurements are still made in Uppsala. Two other Swedish cities with long continuous meteorological records are Lund and Stockholm, starting in 1740 and 1754. An effort to establish a national meteorological network was made by the Royal Academy of Sciences in 1786. At some sites, this led to continuous measurements being made for several decades, but none of their records stretch longer than to 1840s.

Long instrumental records in present-day *Finland* encompass Turku (starting in 1748) and Oulu (1776). Data from Copenhagen (*Denmark*) reach back to 1786. In *Norway*, Trondheim has the longest record (1762) followed by a continuous record from Oslo (1816) (Hestmark and Nordli 2016). Long records reaching back to the 18<sup>th</sup> century are also available for Vilnius (*Lithuania*, 1777) and Riga (*Latvia*, 1795). The series from Tallinn (*Estonia*) starts in 1806.

First measurements were taken in *Germany* in the 17<sup>th</sup> century. The longest temperature record is that of Berlin, reaching back to 1701, but many other series start prior to 1750. The SMP network based in Mannheim comprised 14 stations in Germany, among them the mountain station Hohenpeissenberg. Another important record is that of Karlsruhe, reaching back to the 1770s. Hellmann’s inventory



(Table 2) of German instrumental series lists more than 450 stations prior to 1850, many have not been digitized.

Long records are also available for the *Netherlands*. A composite series for De Bilt back to 1706 has been composed, but there are also other long series (Table 3). A Central *Belgium* daily temperature series was compiled from different records back to 1769 (Demarée et al. 2002).

The *United Kingdom* has some of the earliest meteorological measurement series. The Central England temperature record, which is not an individual series but a composite, reaches back to 1659. Early pressure records are available from Oxford and later London (Cornes et al. 2011a). The Royal Society and its journal, particularly through the invitation of James Jurin (1723), played an important role in the collection and dissemination of data. A composite precipitation record has been constructed for *Ireland* back to 1716 (Murphy et al. 2018).

*Spain* hosts several long instrumental measurement series such as Barcelona (back to 1780), Cadiz and Madrid (1786) and Valencia (1790); earlier short records from Madrid and Granada date back to the 1720s and 1730s (Rodrigo 2019). The first meteorological measurements in the Iberian Peninsula were taken in *Portugal* between 1 November 1724 and 11 January 1725 (Domínguez-Castro et al., 2013). The series for Lisbon reaches back to 1781. The Balkan Peninsula has only poor coverage prior to 1850. Our inventory lists two series from the 1780s to 1800s (Piran, *Slovenia*, and Timișoara, *Romania*). Data from Corfu (*Greece*) reach back to the early 19<sup>th</sup> century, but there are only few other records and they only reach back to the 1830s (Hvar, *Croatia*; Cluj and Stansilav, *Romania*; Athens, *Greece*).

Instrumental measurements in *Poland* began in the 17<sup>th</sup> century. Apart from Warsaw (in the Medici network), measurements took place in Gdansk and Wroclaw, for the latter including the series by David von Grebner (1710-21), Johann Kanold (1717-26) and Elias Büchner (1727-30). Although not all data have been found, Warsaw, Gdansk, Wroclaw, and Cracow all have long records (Przybylak 2010). At Gdansk, observations were conducted 1655-1701, 1721-1786 (Filipiak et al. 2019) and from 1739 onward. In the *Czech Republic*, the series from Prague-Klementinum reaches back to the 1770s, but other long series (e.g., Brno) exist. The Budapest series in *Hungary* reaches back to 1780. The first instrumental measurement in *Austria* date back to the early 18<sup>th</sup> century, continuous records started around 1770 (Kremsmünster, Vienna, Innsbruck). The longest *Swiss* records are those from Basel and Geneva, both reaching back to the mid-18<sup>th</sup> century, but a recent inventory (Pfister et al. 2019) found data from several other stations for which continuous series back to the 18<sup>th</sup> century can be produced.

The earliest measurements in today's *Ukraine* date back to 1738 (Kharkiv) and 1770 (Kyiv), however, they were probably lost. Currently, the archive of the Ukrainian weather service contains eight pre-1850 stations; the earliest measurements date to 1808 (Kherson).

Measurements in *Russia* started in 1724 in St. Petersburg. Only monthly minimum and maximum pressure were published during the first 20 years; from the 1740s onward, temperature data are available from global repositories. The Moscow series reaches back to the 1770s, other series then followed in the 1810s (for instance Old Archangel) and particularly in the 1830s from the “Annuaire magnétique et météorologique du corps des ingénieurs des mines de Russie” (1837-46), the “Annales de l'observatoire physique central de Russie” (1847-61), and “Observations météorologiques faites à Nijné-Taguisk et Vicimo-Outkink (Monts Oural), Gouvernement de Perm” (1839/40–1865/1866) (Weselowksij, 1857; Wild 1881, 1887, Leyst 1887, Bergman 1892).

### 3.8. Arctic

The first instrumental observations in the Arctic (defined after Atlas Arktiki, see Przybylak 2016), at least of duration of three months or longer, were initiated by the “Moravian Brethren” in 1767 in Neu-Herrnhut near Godthåb (present-day Nuuk), Greenland, and four years later Nain (Labrador) (Demarée and Ogilvie 2008). Demarée and Ogilvie (2008) proposed to distinguish four distinct periods in the timeframe of the measurements in Labrador, from which three cover the period of interest in the present paper: 1771-90, 1801-1883 and 1883-beginning of the 20<sup>th</sup> century. Early instrumental meteorological data for Greenland are available from 1873 onward for the four regular stations run by the Danish Meteorological Institute and published in yearbooks until 1960: Ilulissat (Jakobshavn), Upernavik, Nuuk (Godthåb), and Ivittuut (Ivigut) (Cappelen 2018). However, measurements began much earlier, *i.e.*, in 1807 for Ilulissat and 1784 for Nuuk. All available series (see Table 1 and Fig. 2 in Vinther et al. 2006), however, contain many gaps.

Another region covered by early measurements is the Canadian Arctic. Meteorological observations began here in 1819 when the Royal Navy sent the first expedition in search of the Northwest Passage. Later expeditions included those looking for the lost expedition of Sir John Franklin in 1848-59. A large number of those data (monthly and annual means for fixed hours) is available in the publications of Strachan (1879-88), while all source data with sub-daily resolution (hourly, two-hourly, four-hourly, or six-hourly) are in ship logbooks (Przybylak and Vizi 2005).

A third region of the Arctic with important sets of early meteorological measurements is Novaya Zemlya. Seven series with sub-daily resolution are available, usually of 1-year duration, from periods 1832-39 and 1872-83 (Przybylak and Wyszynski 2017). The best coverage for the entire Arctic exists for the 1<sup>st</sup> International Polar Year, 1882/83, when nine stations were operating in the high Arctic, of which two (Sagastyr and Lady Franklin Bay) continued until 1884 (Przybylak et al. 2010; Wyszynski and Przybylak 2014).

## 4. Conclusions

This article describes a global inventory of terrestrial meteorological measurements prior to 1850 (1890 for Africa and the Arctic) which will support data compilation and data rescue efforts. The inventory comprises 4,626 unique entries from 2,250 locations. This is more than anticipated and suggests that climate or weather reconstruction (*e.g.*, by means of reanalyses) based on instrumental data might be extended back well into the 18<sup>th</sup> century. Such data sets would allow new insights into the transition of the climate system from the Little Ice Age climate into the present climate, longer samples to learn from past extreme events, and new opportunities to analyze the climate-society interface.

However, the data are not readily available. Roughly half of the series (in terms of entry years) have not yet been transcribed, and of those that have been partly or fully transcribed, only half is represented in global inventories. Extending the data series backward thus requires further efforts on various aspects, including metadata cataloguing, current data holdings inventorying and updating, maintaining and expanding data compilations and enforcing data standards (see Thorne et al. 2017). Activities currently undertaken within Copernicus Climate Change Services (C3S) (Brönnimann et al. 2018b) can support this process with broader contributions from the communities. The inventory will be maintained as a living document at the C3S Climate Data Store, additions to the inventory are welcomed.

## **Acknowledgements.**

SB acknowledges funding from the European Research Council (787574), Copernicus Climate Change Services (311a Lot 1) and from the Swiss National Science Foundation (IZSEZ0\_180328, 205121\_169676). The research work of RP and PW was supported by the National Science Centre, Poland (Grants No. DEC-2012/07/B/ST10/04002 and 2015/19/B/ST10/02933). JL and RA acknowledge the ongoing support of CSSP China under the BEIS UK–China Research & Innovation Partnership Fund through the Met Office Climate Science for Service Partnership (CSSP) China as part of the Newton Fund. J. Luterbacher also acknowledges the DAAD project “The Mediterranean Hot-Spot” and the JPI-Climate/Belmont Forum collaborative Research Action “INTEGRATE, An integrated data-model study of interactions between tropical monsoons and extratropical climate variability and extremes”. SN and PWT were funded by the European Commission via the Copernicus Climate Change Service contract C3S 311a Lot 2. RB and LR acknowledge the support by the project SustES-Adaptation strategies for sustainable ecosystem services and food security under adverse environmental conditions project no. CZ.02.1.01/0.0/0.0/16\_019/0000797. VS is grateful to support from Environment and Climate Change Canada. JMV was supported by the Junta of Extremadura (grant GR18097) and by the Spanish Government (CGL2017-87917-P). HH was supported by the Swiss National Science Foundation (P2BEP1\_175214)

## **References**

- Adamson, G. C. D., and D. J. Nash, 2014: Documentary reconstruction of monsoon rainfall variability over western India, 1781–1860. *Clim. Dyn.*, **42**, 749–769, <https://doi.org/10.1007/s00382-013-1825-6>.
- Alcoforado, M. J., J. M. Vaquero, R. M. Trigo, and J. P. Taborada, 2012: Early Portuguese meteorological measurements (18th century). *Clim. Past*, **8**, 353–371, <https://doi.org/10.5194/cp-8-353-2012>.

- Allan, R. J., C. J. C. Reason, P. Carroll, and P. D. Jones, 2002: A reconstruction of Madras (Chennai) mean sea-level pressure using instrumental records from the late 18th and early 19th centuries. *Int. J. Climatol.*, **22**, 1119–1142, <https://doi.org/10.1002/joc.678>.
- Allan, R., P. Brohan, G. P. Compo, R. Stone, J. Luterbacher, and S. Brönnimann, 2011: The international Atmospheric Circulation Reconstructions over the Earth (ACRE) initiative. *Bull. Amer. Meteorol. Soc.*, **92**, 1421–1425, <https://doi.org/10.1175/2011BAMS3218.1>.
- Angot A. 1897: Premier catalogue des observations météorologiques, faites en France, depuis l'origine jusqu'en 1850, Annales du Bureau central météorologique de France Année 1895, Mémoires, t.1, B91-B144, Gauthier-Villars et fils, Paris. Retrieved on May 16, 2019 from <https://gallica.bnf.fr/ark:/12148/bpt6k65116714?rk=64378;0>.
- Arakawa, H., 1956: On the secular variation of annual totals of rainfall at Seoul from 1770 to 1944. *Arch. Met. Geoph. Biokl. B.*, **7**, 205-211.
- Arctowski, H., 1904: *Rapports scientifiques, Expédition Antarctique Belge, Res. voy. S.Y. Belgica. Météorologie. Rapport sur les observations météorologiques horaires faites pendant l'hivernage antarctique de la "Belgica"*, J. E. Buschmann, Anvers.
- Ashcroft, L., J. Gergis, and D. J. Karoly, 2014: A historical climate dataset for southeastern Australia, 1788-1859. *Geosci. Data J.*, **1**, 2, 158–178, <https://doi.org/10.1002/gdj3.19>.
- Ashcroft, L., R. Allan, H. Bridgman, J. Gergis, C. Pudmenzky, and K. Thornton, 2016: Current climate data rescue activities in Australia. *Adv. Atmos. Sci.*, Science Press **33**, 12, 1323–1324, <https://doi.org/10.1007/s00376-016-6189-5>.
- Ashcroft, L., J. R. Coll, A. Gilabert, P. Domonkos, M. Brunet, E. Aguilar, M. Castella, J. Sigro, I. Harris, P. Unden, and P. Jones, 2018: A rescued dataset of sub-daily meteorological observations for Europe and the southern Mediterranean region, 1877–2012, *Earth Syst. Sci. Data*, **10**, 1613-1635, <https://doi.org/10.5194/essd-10-1613-2018>.
- Auer, I., and Coauthors, 2007: HISTALP - historical instrumental climatological surface time series of the Greater Alpine Region. *Int. J. Climatol.*, **27**, 17-46, <https://doi.org/10.1002/joc.1377>.
- Barriendos, M., J. Martín-Vide, J. C. Peña, and R. Rodríguez, 2002: Daily Meteorological Observations in Cádiz - San Fernando. Analysis of the Documentary Sources and the Instrumental Data Content (1786-1996). *Clim. Change*, **53**, 151-170.
- Bartkey, I. R., 2008: The (Almost) Unseen Total Eclipse of 1831. *J. Astron. Hist. Herit.*, **11**, 1, 55-62.
- Berghaus, H., 2004: *Physikalischer Atlas*. H. M. Enzensberger, Ed., Frankfurt a.M.: Eichborn
- Bergman, R., 1892: *O raspriedeleniji o diejatielnosti meteorologiczeskich stancji w Rossijskoj imperii s naczala ich wozniknowienija do 1889*, St. Petersburg 1892
- Bergström, H., and A. Moberg, 2002: Daily Air Temperature and Pressure Series for Uppsala (1722–1998). *Clim. Change*, **53**, 213-252.
- Blodget, L., 1857: *Climatology of the United States and temperate latitudes of the North American continent*, Lippincott, Philadelphia.
- Boussingault, 1849: *Viajes científicos a los Andes Ecuatoriales*, Libreria Castellana, Paris.
- Bradley, R.S., P. M. Kelly, P. D. Jones, C. M. Goodess, and H. F. Diaz, 1985: A Climatic Data Bank for Northern Hemisphere Land Areas, 1851-1980. U.S. Dept. of Energy, Carbon Dioxide Research Division, *Technical Report TRO17*, 335 pp.
- Brázdil, R., H. Valášek, and J. Macková, 2005: *Meteorological observations in Brno in the first half of the nineteenth century (History of weather and hydrometeorological extremes)* (in Czech). Brno: Archiv města Brna.
- Brázdil, R., A. Kiss., J. Luterbacher, and H. Valášek, 2008: Weather patterns in eastern Slovakia 1717–1730, based on records from Breslau meteorological network. *Int. J. Climatol.*, **28**, 12, 1639–1651, <https://doi.org/10.1002/joc.1667>.
- Brázdil, R., M. Bělinová, P. Dobrovolný, J. Mikšovský, P. Pišof, L. Řezníčková, P. Štěpánek, H. Valášek, and P. Zahradníček, 2012: *Temperature and Precipitation Fluctuations in the Czech Lands During the Instrumental Period*. Brno: Masaryk University.
- Brohan, P., G. P. Compo, S. Brönnimann, R. J. Allan, R. Auchmann, Y. Brugnara, P. D. Sardeshmukh, and J. S. Whitaker 2016: The 1816 ‘year without a summer’ in an atmospheric reanalysis, *Clim. Past Discuss.*, <https://doi.org/10.5194/cp-2016-78>.
- Brönnimann, S., and Coauthors, 2018a: Observations for Reanalyses. *Bull. Amer. Meteorol. Soc.*, **99**, 1851–1866, <https://doi.org/10.1175/BAMS-D-17-0229.1>.
- Brönnimann, S., and Coauthors, 2018b: A roadmap to climate data rescue services. *Geosci. Data J.*, **5**, 8–39, <https://doi.org/10.1002/gdj3.56>.
- Brönnimann, S., and J. Wintzer, 2019: Climate Data Empathy. *WIREs Clim. Change*, **10**, e559, <https://doi.org/10.1002/wcc.559>.
- Brugnara, Y., and Coauthors, 2015: A collection of sub-daily pressure and temperature observations for the early instrumental period with a focus on the “year without a summer” 1816. *Clim. Past*, **11**, 1027–1047, <https://doi.org/10.5194/cp-11-1027-2015>.

- Brunet, M., P. Jones, S. Jourdain, D. Efthymiadis, M. Kerrouche, and C. Boroneant 2013: Data sources for rescuing the rich heritage of Mediterranean historical surface climate data. *Geosc. Data J.*, **1**, 61-73.
- Brunet, M., A. Gilabert, and P. Jones, 2014: A historical surface climate dataset from station observations in Mediterranean North Africa and Middle East areas. *Geosci. Data J.*, **1**, 121-128.
- Brunetti, M., L. Buffoni, G. Lo Vecchio, M. Maugeri, and T. Nanni, 2001: *Tre Secoli di Meteorologia a Bologna*. Edizioni Cusl: Milano.
- Brunetti, M., M. Maugeri, F. Monti, and T. Nanni, 2006: Temperature and precipitation variability in Italy in the last two centuries from homogenised instrumental time series. *Int. J. Climatol.*, **26**, 345-381, <https://doi.org/10.1002/joc.1251>.
- Burnette, D. J., D. W. Stahle, and C. J. Mock, 2010: Daily-mean temperature reconstruction for Kansas from early instrumental and modern observations. *J. Clim.*, **23**, 1308-1333, <https://doi.org/10.1175/2009JCLI2445.1>.
- Bryś, K., and T. Bryś, 2010a: The first one hundred year (1791-1890) of the Wrocław air temperature series. *The Polish Climate in the European Context: An Historical Overview*. R. Przybylak, J. Majorowicz, R. Brázdil, and M. Kejna, Eds., Springer, Dordrecht, 485-524.
- Bryś, K., and T. Bryś, 2010b: Reconstruction of the 217-year (1791-2007) Wrocław air temperature and precipitation series. *Bulletin of Geography. Physical Geography Series*, **3**, 121-171.
- Camuffo, D., and C. Bertolin, 2012: The earliest temperature observations in the world: the Medici Network (1654-1670). *Clim. Change*, **111**, 335-363, <https://doi.org/10.1007/s10584-011-0142-5>.
- Camuffo, D., and P. Jones, 2002: Improved Understanding of Past Climatic Variability from Early Daily European Instrumental Sources. *Clim. Change*, **53**, 1-4, <http://doi.org/10.1023/A:1014902904197>.
- Camuffo, D., C. Cocheo, and G. Sturaro, 2006: Corrections of systematic errors, data homogenisation and climatic analysis of the Padova pressure series (1725-1999). *Clim. Change*, **78**, 493-514, <http://doi.org/10.1007/s10584-006-9052-3>.
- Camuffo, D., C. Bertolin, N. Diodato, C. Cocheo, M. Barriendos, F. Dominguez-Castro, E. Garnier, M. J. Alcoforado, and M. F. Nunes, 2013: Western Mediterranean precipitation over the last 300 years from instrumental observations. *Clim. Change*, **117**, 85-101, <https://doi.org/10.1007/s10584-012-0539-9>.
- Camuffo, D., A. della Valle, C. Bertolin, and E. Santorelli, 2017: Temperature observations in Bologna, Italy, from 1715 to 1815: a comparison with other contemporary series and an overview of three centuries of changing climate. *Clim. Change*, **142**, 7-22, <https://doi.org/10.1007/s10584-017-1931-2>.
- Cantù, V., and P. Narducci, 1967: Lunghie serie di osservazioni meteorologiche. *Rivista di Meteorologia Aeronautica* XXVII, 71-79. I
- Cappelen, J., (ed.), 2018: Greenland - DMI Historical Climate Data Collection 1784-2017, DMI Report 18-04, pp. 118.
- Colin, E., 1889: *Résumé des observations météorologiques faites à Tananarive*, Imprimerie de la mission catholique, Tananarive.
- Compo G. P., and Coauthors, 2011: The Twentieth Century Reanalysis project. *Q. J. R. Meteorol. Soc.*, **137**, 1-28, <https://doi.org/10.1002/qj.776>.
- Cornes, R. C., P. D. Jones, K. R. Briffa, and T. J. Osborn, 2012a: A daily series of mean sea-level pressure for London, 1692-2007. *Int. J. Climatol.*, **32**, 641-656, <https://doi.org/10.1002/joc.2301>.
- Cornes, R. C., P. D. Jones, K. R. Briffa, and T. J. Osborn, 2012b: A Daily Series of Mean Sea-Level Pressure for Paris, 1670-2007. *Int. J. Climatol.*, **32**, 1135-1150, <https://doi.org/10.1002/joc.2349>.
- Cotte, L., 1788: Mémoires sur la météorologie pour servir de suite et de supplément au traité de météorologie publié en 1774, tome II. Imprimerie royale, Paris. Retrieved on May 22, 2019 from <https://gallica.bnf.fr/ark:/12148/bpt6k94862j?rk=107296;4>
- Cram, T. A., and Coauthors, 2015: The International Surface Pressure Databank version 2. *Geosci. Data J.*, **2**, 31-46, <https://doi.org/10.1002/gdj3.25>.
- Danckelman, A. v., 1878: *Die meteorologischen Beobachtungen der Güßfeldtschen Loango-Expedition*, Froberg, Leipzig.
- Danckelman, A. v., 1884: *Mémoire sur les observations météorologiques faites à Vivi et sur la climatologie de la côte Sud-Ouest d'Afrique en general*, Asher and Co., Berlin.
- Demarée, G. R., and A. E. J. Ogilvie, 2008: The Moravian missionaries at the Labrador coast and their centuries-long contribution to instrumental meteorological observations. *Clim. Change*, **91**, 3-4, 423-450, <http://doi.org/10.1007/s10584-008-9420-2>.
- Demarée, G. R., P. J. Lachaert, T. Verhoeve, and E. Thoen, 2002: The Long-Term Daily Central Belgium Temperature (CBT) Series (1767-1998) and Early Instrumental Meteorological Observations in Belgium. *Improved understanding of past climatic variability from early daily European instrumental sources*, D. Camuffo, and P. Jones, Ed., Kluwer Academic Publishers, Dordrecht.
- Demarée, G. R., T. Mikami, T. Tsukahara, and M. Zaiki, 2013: The meteorological observation of the "Vereenigde Oost-indische Compagnie (VOC)" - What can be learned from them? *Historical Geographers in Japan*, **55**, 99-106.
- Derham, W., 1735: An abstract of the meteorological diaries, communicated to the Royal Society; with Remarks upon them *Phil. Trans. Roy. Soc.*, 1733-34, 38, 334-344 and 458-467.

- Ding, L., Q. Ge, J. Zheng, and Z. Hao, 2014: Variations in the starting date of the pre-summer rainy season in South China, 1736–2010, *J. Geogr. Sci.*, **24**, 5, 845–857, <https://doi.org/10.1007/s11442-014-1124-0>.
- Domínguez-Castro, F., R. M. Trigo, and J. M. Vaquero, 2013: The first meteorological measurements in the Iberian Peninsula: evaluating the storm of November 1724, *Climatic Change*, **118**, 443–455
- Domínguez-Castro, F., J. M. Vaquero, F. S. Rodrigo, A. M. M. Farrona, M. C. Gallego, R. García-Herrera, M. Barriendos, and A. Sanchez-Lorenzo, 2014: Early Spanish meteorological records (1780–1850). *Int. J. Climatol.*, **34**, 3, 593–603, <https://doi.org/10.1002/joc.3709>.
- Domínguez-Castro, F., and Coauthors, 2017: Early meteorological records from Latin-America and the Caribbean during the 18th and 19th centuries. *Sci. Data*, **4**, 170169, <https://doi.org/10.1038/sdata.2017.169>.
- Dove, H. W., 1838: *Über die nichtperiodischen Änderungen der Temperaturverteilung auf der Oberfläche der Erde. 1. Teil.* Reimer, Berlin
- Dove, H. W., 1839: *Über die nichtperiodischen Änderungen der Temperaturverteilung auf der Oberfläche der Erde. 2. Teil.* Reimer, Berlin
- Dove, H. W., (Ed.) 1841: *Repertorium der Physik.* Vol. 4, Verlag von Veit & Comp, Berlin
- Dove, H. W., 1842: *Über die nichtperiodischen Änderungen der Temperaturverteilung auf der Oberfläche der Erde. 3. Teil.* Reimer, Berlin
- Dove, H. W., 1845: *Über die nichtperiodischen Änderungen der Temperaturverteilung auf der Oberfläche der Erde. 4. Teil.* Reimer, Berlin
- Dove, H. W., 1852: Die Verbreitung der Wärme auf der Oberfläche der Erde erläutert durch Isothermen, thermische Isanomalien und Temperaturcurven. 2. Aufl., Reimann, Berlin.
- Dupigny-Giroux, L.-A., T. F. Ross, J. D. Elms, R. Truesdell, and S. Doty, 2007: NOAA's Climate Database Modernization Program: Rescuing, Archiving, and Digitizing History. *Bull. Amer. Meteorol. Soc.*, **88**, 7, 1015–1017,
- Eredia, F., 1912: La temperatura in Italia. *Annali dell'Ufficio Centrale di Meteorologia.* Serie II, vol. XXXI, anno 1909. UCM, Roma.
- Eredia, F., 1919: *Osservazioni pluviometriche raccolte a tutto l'anno 1915 dal R. Ufficio Centrale di Meteorologia e Geodinamica.* Ministero dei Lavori Pubblici. UCM, Roma.
- Etienne, E., 1892: *Le Climat de Banana en 1890, faites du 1er décembre 1889-16 mai 1891*, Imprimerie Typographique Jules Vanderauwera, Bruxelles.
- Farrona, A. M. M., J. M. Vaquero, M. C. Gallego and R. M. Trigo, 2012: The meteorological observations of Bento Sanches Dorta, Rio de Janeiro, Brazil: 1781-1788. *Clim. Change*, **115**, 579–595.
- Filipiak, J., R. Przybylak, and P. Oliński, 2019: The longest one-man weather chronicle (1721–1786) by Gottfried Reyger for Gdańsk, Poland as a source for improved understanding of past climate variability. *Int. J. Climatol.*, **39**, 828–842, <http://doi.org/10.1002/joc.5845>.
- Gallego, M.C., F. Dominguez-Castro, J. M. Vaquero, and R. Garcia-Herrera, 2011: The hidden role of women in monitoring nineteenth-century African weather: instrumental observations in equatorial Guinea. *Bull. Amer. Meteor. Soc.*, **92**, 3, 315–324, <https://doi.org/10.1175/2010BAMS2807.1> <https://doi.org/10.1175/2010BAMS2807.1>.
- Gazina, E. A., and V. V. Klimenko, 2008: Climatic changes of the Eastern Europe during the last 250 years by instrumental data. *Moscow University Vestnik.* Series 5: Geography, 3, 60–66 (in Russian).
- Ge, Q.-S., J. Y. Zheng, Z. X. Hao, P. Y. Zhang, and W.-C. Wang, 2005: Reconstruction of historical climate in China: high-resolution precipitation data from Qing Dynasty archives. *Bull. Amer. Meteorol. Soc.*, **86**, 5, 671–679.
- Gergis, J., and L. Ashcroft, 2013: A rainfall history of south-eastern Australia Part 2: a comparison of documentary, early instrumental and palaeoclimate records, 1788–2008. *Int. J. Climatol.*, DOI: 10.1002/joc.3639.
- Gergis, J., D. J. Karoly, and R. Allan, 2009: A climate reconstruction of Sydney Cove, New South Wales, using weather journal and documentary data, 1788–1791. *Australian Meteorological and Oceanographic Journal*, **58**, 2, 83–98,
- Goldingham, J., 1826: Tables containing the results of the meteorological observations taken at the Madras Observatory. Microfilm from the Oriental and India Office Collections (OIOC) of the British Library, London.
- Goldingham, J., and T. G. Taylor, 1844: Meteorological Register kept at the Honorable The British East India Company's Observatory at the Madras for the years 1822-1843. Madras Government. Microfilm from the Oriental and India Office Collections (OIOC) of the British Library, London.
- Hadley, G., 1741: An account and abstract of the meteorological diaries, communicated to the Royal Society, for the years 1729 and 1730. *Phil. Trans. Roy. Soc.*, **40**, 154–175.
- Hadley, G., 1744: An account and abstract of the meteorological diaries, communicated to the Royal Society, for the years 1731, 1732, 1733, 1734 and 1735. *Phil. Trans. Roy. Soc.*, **42**, 243–263.
- Hamilton, F., 1819: *An account of the Kingdom of Nepal: and of the territories annexed to this Dominion by the House of Gorkha.* Archibald Constable and Company, Edinburgh [Meteorological Observations 318-345], 376 pp.
- Hao, Z., J. Zheng, Q. Ge and W. Wang, 2012: Winter temperature variations over the middle and lower reaches of the Yangtze River since 1736 AD. *Clim Past*. **8**:1023-1030
- Hardwicke, T., 1830: Appendix No 1. Meteorological Registers. *Transactions of the Royal Asiatic Society*, **2**, i–xix.

- Havens, J. M., 1958: An annotated bibliography of meteorological observations in the United States, 1715-1718. U.S. Department of Commerce, Weather Bureau, Washington D.C.
- Hellmann, G., 1883: *Repertorium der deutschen Meteorologie*, Wilhelm Engelmann, Leipzig.
- Hellmann, G., 1901a: Die Entwicklung der meteorologischen Beobachtungen bis zum Ende des XVII. Jahrhunderts. *Meteorol. Z.*, **18**, 145–157.
- Hellmann, G., 1901b: *Meteorologische Beobachtungen vom XIV. bis XVII. Jahrhundert*, A. Aher, Berlin.
- Hellmann, G., 1927: Die Entwicklung der meteorologischen Beobachtungen bis zum Ende des XVIII. Jahrhunderts. Abhandlungen der Preußischen Akademie der Wissenschaften, Physikalisch-Mathematische Klasse.
- Hestmark G. and Nordli Ø. 2016: Jens Esmark's Christiania (Oslo) meteorological observations 1816–1838: the first long-term continuous temperature record from the Norwegian capital homogenized and analysed. *Clim Past*, **12**, 2087-2106.
- Hough, F., 1872: *Results of a series of Meteorological Observations made under Instructions from the Regents of the University*. Weed, Parsons and Company, Albany, NY.
- Humboldt, A. v. 1817: Des lignes isothermes et de la distribution de la chaleur sur le globe. *Annales de Chimie et de Physique*, **5**, 102–112.
- India Meteorological Department, 1976: *Hundred Years of Weather Service (1875–1975)*. Dy. Director General of Observatories: Poona; 207 pp.
- Jones, P. D., 1990: Antarctic temperatures over the present century - a study of the early expedition record. *J. Clim.*, **3**, 1193-1203.
- Jones, P. D., 2001: Early European Instrumental Records. In: Jones P.D., Ogilvie A.E.J., Davies T.D., Briffa K.R., Eds., *History and Climate*. Springer, Boston, MA.
- Jones, P. and D. Lister, 2002: The daily temperature record for St. Petersburg (1743–1996). *Clim. Change*, **53**, 253–267.
- Jones, P. D., and Coauthors, 1985: A Grid Point Surface Air Temperature Data Set for the Northern Hemisphere, U.S. Dept. of Energy, Carbon Dioxide Research Division, *Technical Report TRO22*, 251 pp.
- Jones, P. D., S. C. B Raper, R. S. Bradley, H. F. Diaz, P. M. Kelly, and T. M. L. Wigley, 1986: Northern Hemisphere surface air temperature variations: 1851-1984. *J. Clim. Appl. Meteorol.*, **25**, 161–179.
- Jones, P. D., and Coauthors, 1999: Monthly Mean Pressure Reconstructions for Europe for the 1780–1995 Period. *Int. J. Clim.*, **19**, 347–364.
- Jones, P. D., D. H. Lister, T. J. Osborn, C. Harpham, M. Salmon, and C. P. Morice, 2012: Hemispheric and large-scale land-surface air temperature variations: An extensive revision and an update to 2010. *J. Geophys. Res.*, **117**, D05127.
- Jurin, J., 1723: Invitatio ad Observationes Meteorologicae communi consilio instituendas. *Phil. Trans. Roy. Soc.*, **32**, 422–427.
- Kingsford, S., 1843: Summary of meteorological observations of barometer and thermometer made at Bangalore from 1820 to 1835 inclusive. *The Quarterly Journal of Meteorological Physical Science*, **1**, 36 pp.
- Kington, J. A., 1974: The Societas Meteorologica Palatina: An Eighteenth-century meteorological society. *Weather*, Nov. 1974, 416-426.
- Kington, J., 1988: *The weather of the 1780s over Europe*. Cambridge University Press, Cambridge.
- Kirwan, R., 1787: *An Estimate of the Temperature of Different Latitudes*. London.
- Können, G.P., M. Zaiki, A. P. M. Baede, TY. Mikami, P. D. Jones, and T. Tsukahara, 2003: Pre-1872 extension of the Japanese instrumental meteorological observation series back to 1819. *J. Clim.*, **16**, 118-131
- Köppen, W., 1873: Über mehrjährige Perioden der Witterung, insbesondere über die 11-jährige Periode der Temperatur. *Z. Österr. Ges. Meteor.* **8**, 241–248, 257–267.
- Köppen, W., 1881: Über mehrjährige Perioden der Witterung–III. Mehrjährige Änderungen der Temperatur 1841 bis 1875 in den Tropen der nördlichen und südlichen gemässigten Zone, an den Jahresmitteln untersucht. *Z. Österr. Ges. Meteor.*, 141–150.
- Kreil, C., 1865: *Klimatologie von Böhmen*. Carl Gerold's Sohn, Wien.
- Lawrimore, J. H., M. J. Menne, B. E. Gleason, C. N. Williams, D. B. Wuertz, R. S. Vose, and J. Rennie, 2011: An overview of the Global Historical Climatology Network monthly mean temperature dataset, version 3. *J. Geophys. Res.*, **116**, D19121.
- Leyst, E., 1887: *Katalog der meteorologischen Beobachtungen in Russland und Finnland, vierter Supplementband zum Repertorium für Meteorologie*, Commissionäre der Kaiserlichen Akademie der Wissenschaften, St. Petersburg.
- Lorrey, A. M., and P. R. Chappell, 2015: The “dirty weather” diaries of Reverend Richard Davis: insights about early colonial-era meteorology and climate variability for northern New Zealand, 1839-1851, *Clim. Past*, **12**, 553–573.
- Lucerna Giraldo, M. (ed.), 2009: *Atlas de los exploradores españoles*, GeoPlaneta, Sociedad Geográfica Española, Barcelona.
- Lüdecke, C., 2010: Von der Kanoldsammlung (1717-1730) zu den Ephemeriden der Societas Meteorologica Palatina (1781-1792). Meteorologische Quellen zur Umweltgeschichte des 18. Jahrhunderts. *Landschaften agrarisch-ökonomischen Wissens: Strategien innovativer Ressourcennutzung in Zeitschriften und Sozietäten des 18. Jahrhunderts*. M. Popplow, Ed., Waxmann, Münster, 97–119.

- Manley, G., 1974: Central England temperatures: monthly means 1659 to 1973. *Quart. J. R. Met. Soc.*, **100**, 389–405.
- Maraldi, J.-P., 1709: Comparaison des Observations de Barometre faites en differens lieux. *Memoires de l'Academie Royale des Sciences*, 233–245.
- Maugeri, M., L. Buffoni, and F. Chlistovsky, 2002a: Daily Milan temperature and pressure series (1763-1998): history of the observations and data and metadata recovery. *Clim. Change*, **53**, 101–117.
- Maugeri, M., L. Buffoni, B. Delmonte, and A. Fassina, 2002b: Daily Milan Temperature and Pressure Series (1763–1998): Completing and Homogenising the Data. *Clim. Change*, **53**, 119–149.
- Maugeri, M., M. Brunetti, L. Buffoni, G. Lentini, F. Mangianti, F. Monti, T. Nanni, and R. Pastorelli, 2006: Variabilità e cambiamenti climatici in Italia nel corso degli ultimi due secoli documentati da serie storiche secolari omogeneizzate. In: *Progetto di Ricerca CLIMAGRI-Cambiamenti climatici e agricoltura (finanziato dal Ministero delle Politiche Agricole, Alimentari e Forestali) – Risultati conclusivi*. S. Esposito, C. Epifani, C. Serra, Eds., CRA-Ufficio Centrale di Ecologia Agraria, Roma, 15–29
- Menne, M. J., I. Durre, R. S. Vose, B. E. Gleason, and T. G. Houston, 2012: An overview of the global historical climatology network-daily database. *J. Atmospheric Ocean. Technol.*, **29**, 897–910.
- Mitchell, T. L., 1838: *Three Expeditions into the Interior of Eastern Australia: With Descriptions of the Recently Explored Region of Australia Felix, and of the Present Colony of New South Wales*. Published by London, T. & W. Boone, Vol. 1, p. 355.
- Moberg, A., 1998: Meteorological observations in Sweden made before A.D. 1860. *Paläoklimaforschung/Palaeoclimate Research*, **23**, 99–119.
- Moberg, A. and H. Bergström, 1997: Homogenization of Swedish temperature data. Part III: the long temperature records from Uppsala and Stockholm. *Int. J. Climatol.*, **17**, 667–699.
- Moberg, A., H. Bergström, J. R. Kringsman, and O. Svanered, 2002: Daily air temperature and pressure series for Stockholm (1756–1998). *Clim. Change*, **53**, 171–212.
- Murphy, C., and Coauthors, 2018: A 305-year continuous monthly rainfall series for the island of Ireland (1711–2016). *Clim. Past*, **14**, 413–440.
- Nash, D. J., and G. C. D. Adamson, 2014: Recent Advances in the Historical Climatology of the Tropics and Subtropics. *Bull. Amer. Meteorol. Soc.*, **95**, 131–146.
- Nash, D. J., K. Pribyl, G. H. Endfield, J. Klein, and G. C. D. Adamson, 2018: Rainfall variability over Malawi during the late 19th century. *Int. J. Climatol.*, **38**, e629-e642.
- Nicholson, S. E., D. Klotter, and A. K. Dezfuli, 2012a: Spatial reconstruction of semi-quantitative precipitation fields over Africa during the nineteenth century from documentary evidence and gauge data. *Quat. Res.*, **78**, 13–23.
- Nicholson, S.E., 2001; A semi-quantitative, regional precipitation data set for studying African climates of the nineteenth century, part I. Overview of the data set. *Clim. Change*, **50**, 317–353.
- Nicholson, S.E., A. K. Dezfuli, and D. Klotter, 2012b: A two-century precipitation dataset for the continent of Africa. *Bull. Amer. Meteorol. Soc.*, **93**, 1219–1231.
- Parker, D. E., T. P. Legg, and C. K. Folland, 1992: A new daily Central England temperature series. *Int. J. Climatol.*, **12**, 317–342.
- Pearce, T. D., 1788: A meteorological journal kept by Colonel T.D. Pearce, from 1 March 1785, to 28<sup>th</sup> February 1786. In *Appendix to Asiatic Researches*, **1**, 441–465.
- Pei, Q., P. Forêt, and M. Hall, 2018: Introduction to the Climate Records of Imperial China. *Environmental History*, 3–24, DOI: 10.1093/envhis/emv052.
- Pfister, L., F. Hupfer, Y. Brugnara, L. Munz, L. Villiger, L. Meyer, M. Schwander, F. A. Isotta, C. Rohr, and S. Brönnimann, 2019: Swiss Early Instrumental Meteorological Measurements, *Clim. Past Discuss.*, <https://doi.org/10.5194/cp-2019-26>.
- Peterson, T.C. and R. S. Vose, 1997: An overview of the Global Historical Climatology Network temperature data base. *Bull. Amer. Meteorol. Soc.*, **78**, 2837–2849.
- Pichard, G. 1988: Les météorologistes provençaux aux XVIIe et XVIIIe siècles. *Provence Historique*, **153**, 249-284.
- Prinsep, J., 1829a: Abstract of a meteorological journal kept at Benares, during the years 1824, 1825, and 1826. *Gleanings in Science*, **5**, 157–158.
- Prinsep, J., 1829b: Abstract of a meteorological journal kept at Benares, during the year 1827. *Phil. Trans. Roy. Soc.*, **118**, 251-255
- Prohom, M., M. Barriendos, and A. Sanchez-Lorenzo, 2016: Reconstruction and homogenization of the longest instrumental precipitation series in the Iberian Peninsula (Barcelona, 1786-2014). *Int. J. Climatol.*, **36**, 8, 3072–3087, DOI: 10.1002/joc.4537.
- Przybylak R., 2010: The Climate of Poland in Recent Centuries: A Synthesis of Current Knowledge: Instrumental observations. *The Polish Climate in the European Context: An Historical Overview*. R. Przybylak, J. Majorowicz, R. Brázdil, and M. Kejna, Eds., Springer, Dordrecht, 129–166.



- Przybylak, R., 2016: *The Climate of the Arctic, Second edition*, Atmospheric and Oceanographic Sciences Library 52, Springer, Dordrecht.
- Przybylak, R., and A. Pospieszńska, 2010: Air temperature in Wrocław (Breslau) in the period 1710-1721 based on measurements made by David Von Grebner. *Acta Agrophysica*, **184**, 35–43.
- Przybylak, R., and Z. Vizi, 2005: Air temperature changes in the Canadian Arctic from the early instrumental period to modern times. *Int. J. Climatol.*, **25**, 1507–1522, DOI: 10.1002/joc.1213.
- Przybylak, R., and P. Wyszynski, 2017: Air temperature in Novaya Zemlya Archipelago and Vaygach Island from 1832 to 1920 in the light of early instrumental data. *Int. J. Climatol.*, **37**, 8, 3491–3508, DOI: 10.1002/joc.4934.
- Przybylak, R., Z. Vizi, and P. Wyszynski, 2010: Air temperature changes in the arctic from 1801 to 1920. *Int. J. Climatol.*, **30**, 791–812, DOI:10.1002/joc.1918.
- Przybylak, R., A. Pospieszńska, P. Wyszynski, and M. Nowakowski, 2014: Air temperature changes in Żagan (Poland) in the period from 1781 to 1792. *Int. J. Climatol.*, **34**, 2408–2426, DOI: 10.1002/joc.3847.
- Raulin, V. 1875a: Observations pluviométriques faites dans les colonies françaises (anciennes et actuelles) de la zone torride de 1731 à 1870. Actes de l'Académie nationale des sciences, belles lettres et arts de Bordeaux, France, Paris, 1874, 36<sup>e</sup> année, 3e série, trimestres 1 et 2 France, 451-527. Retrieved on May 22, 2019 from <https://gallica.bnf.fr/ark:/12148/bpt6k33972n/f352.item>
- Raulin, V. 1875b: Régime pluvial de l'Algérie d'après les observations de l'administration des Ponts -et-Chaussées. Actes de l'Académie nationale des sciences, belles lettres et arts de Bordeaux, France, Paris, 36<sup>e</sup> année, 1874, 3e série, trimestre 1 et 2, 351-450. Retrieved on May 22, 2019 from <https://gallica.bnf.fr/ark:/12148/bpt6k33972n/f352.item>.
- Raulin, V. 1876: Observations pluviométriques faites dans la France méridionale (Sud-Ouest, Centre et Sud-Est) de 1704 à 1870 avec les grandes séries de Paris, Genève et le Grand-St-Bernard, Extraits des Actes de l'Académie des Sciences, belles lettres et arts de Bordeaux, France, Bordeaux et Paris, 1083pp. Retrieved on May 22, 2019 from [http://doc-meteo.meteo.fr/exl-php/docs/mf\\_-\\_bibliotheque\\_numerique\\_-\\_collections\\_patrimoniales/12310/iso00009683\\_\\_PDF.txt](http://doc-meteo.meteo.fr/exl-php/docs/mf_-_bibliotheque_numerique_-_collections_patrimoniales/12310/iso00009683__PDF.txt).
- Raulin, V. 1881: Observations pluviométriques faites dans la France septentrionale (Est, Neustrie et Bretagne) de 1688 à 1870 Actes de l'Académie nationale des sciences, belles lettres et arts de Bordeaux, France, Paris, 1878, 40e année (supplément), 3e série, trimestres 1 et 2, 812pp. Retrieved on May 22, 2019 from <https://gallica.bnf.fr/ark:/12148/bpt6k339761>.
- Rennie, J. J., and Coauthors, 2014: The international surface temperature initiative global land surface databank: monthly temperature data release description and methods. *Geosci. Data J.*, **1**, 75–102, DOI: 10.1002/gdj3.8.
- Rodrigo, F. S. 2019: The climate of Granada (southern Spain) during the first third of the 18th century (1706–1730) according to documentary sources, *Clim. Past*, **15**, 647-659.
- Rohde, R., and Coauthors, 2013: A New Estimate of the Average Earth Surface Land Temperature Spanning 1753 to 2011. Geoinfor Geostat: An Overview 1:1
- Rousseau, D., 2015: Variabilité des températures mensuelles à Paris de 1658 à 2014. *XXVIIIe Colloque de l'Association Internationale de Climatologie*, Liège 2015. 597-602.
- Sanchez-Rodrigo, F., 2019: Early Meteorological Data in Southern Spain during the Dalton Minimum. *Int J Climatol*. Accepted Author Manuscript. doi:10.1002/joc.6041
- Schott, C. A., 1876a: *Tables, distribution, and variations of the atmospheric temperature in the United States*, Smithsonian Institution Washington DC.
- Schott, C. A., 1876b: *Tables, distribution and variations of the atmospheric temperatures in the US and adjacent parts of America*, Smithsonian Institution, Washington DC.
- Schott, C. A., 1881: *Tables and results of the precipitation, in rain and snow, in the United States: and at some stations in adjacent parts of North America, and in Central and South America*, Smithsonian Institution, Washington DC.
- Schouw, J. F., 1839: *Tableau du climat et de la végétation de l'Italie. Résultat de deux voyages en ce pays dans les années 1817-1819 et 1829-1830*, Librairie Gyldendal, Copenhagen.
- Slonosky, V. C., 2002: Wet winters, dry summers? Three centuries of precipitation data from Paris. *Geophys. Res. Lett.*, **29**, 18, 1887, DOI: 10.1029/2001GL014302.
- Slonosky, V. C., 2003: The Meteorological Observations of Jean-François Gaultier, Quebec, Canada: 1742–56. *J. Clim.*, **16**, 2232–2247.
- Slonosky, V. C., 2014: Historical climate observations in Canada: 18th and 19th century daily temperature from the St. Lawrence Valley, Quebec. *Geosci. Data J.*, **1**, 103–120, DOI: 10.1002/gdj3.11.
- Slonosky, V. C., 2015: Daily Minimum and Maximum Temperature in the St-Lawrence Valley, Quebec: Two Centuries of Climatic Observations from Canada. *Int. J. Climatol.*, **35**, 1662–81.
- Strachan M., 1879-1888: Contributions to Our Knowledge of the Meteorology of the Arctic Regions. Authority of the Meteorology: London; Part I (1879), Part II (1880), Part III (1882), Part IV (1885), Part V (1888).
- Strzelecki, P. E., 1845: *Physical description of New South Wales and Van Diemen's Land*. Longman; Brown; Green; and Longmans; London.
- Supan, A., 1898: *Die Verteilung des Niederschlags auf der festen Erdoberfläche*. *Petermanns Mitt., Ergänzungsheft* **124**, 103 pp.


- Thorne, P. W., and Coauthors, 2011: Guiding the creation of a comprehensive surface temperature resource for twenty - first century climate science. *Bull. Amer. Meteorol. Soc.*, **92**, ES40– ES47, doi: 10.1175/2011BAMS3124.1
- Thorne, P. W., and Coauthors, 2017: Towards an integrated set of surface meteorological observations for climate science and applications. *Bull. Amer. Meteorol. Soc.*, **98**, 2689–2702, DOI: 10.1175/BAMS-D-16-0165.1.
- Traill, H., 1790: A meteorological diary kept at Calcutta by Henry Traill, Esq. from 1<sup>st</sup> February 1784, to 31<sup>st</sup> December 1785. In Appendix to *Asiatic Researches*, **2**, 419–471.
- Tsukahara, T., 2013: 19th century Chinese Meteorology: Climate reconstruction based on historical record in English newspapers. *Historical Geographers in Japan*, **55**, 69–81.
- van der Schrier, G., and P. D. Jones, 2008: Storminess and cold air outbreaks in NE America during AD 1790-1820. *Geophys. Res. Lett.*, **35**, L02713, DOI: 10.1029/2007GL032259.
- Vinther, B. M., K. K. Andersen, P. D. Jones, K. R. Briffa, and J. Cappelén, 2006: Extending Greenland temperature records into the late-18th century. *J. Geophys. Res.*, **111**, D11105, DOI: 10.1029/2005JD006810.
- Walsh, R., R. Glaser, and S. Miltzer, 1999: The climate of Madras in the eighteenth century. *Int. J. Climatol.*, **19**, 1025–1047.
- Wang, P. K. and D. Zhang, 1988: Introduction to some historical governmental weather records of China. *Bulletin American Meteorological Society* 69:753-758
- Wang, P. K., and Coauthors, 2018: Construction of the REACHES climate database based on historical documents of China. *Sci. Data*, **8**, 1–14.
- Wargentin, P., 1758: Jämförelse imellan Svenska och Franska Climaterna, samt tvänne andra sydligare. *Kungliga Vetenskapsakademiens Handlingar*, **19**, 1–15.
- Weselowksij, K. S., 1857: On the climate of Russia” (О климате России)  
<https://books.google.ch/books?id=DOdPAAAАсAAJ>; 770 pages
- Westcott, N. E., and Coauthors, 2011: Quality Control of 19th Century Weather Data. Prepared for the Midwestern Regional Climate Center, CR 11-02. Illinois State Water Survey, Illinois, 48p.
- Wheeler, D., 1995: Early instrumental weather data from Cadiz: A study of late eighteenth and early nineteenth century records. *Int. J. Climatol.*, **15**, 801–810, DOI: 10.1002/joc.3370150707
- Wild, H., 1881: *Die Temperatur-Verhältnisse des Russischen Reiches, Supplementband zum Repertorium für Meteorologie*, Buchdruckerei der Kaiserlichen Academie der Wissenschaften, St. Petersburg 1881.
- Wild, H., 1887: *Die Temperatur-Verhältnisse des Russischen Reiches, Supplementband zum Repertorium für Meteorologie, Erste Hälfte*, Buchdruckerei der Kaiserlichen Academie der Wissenschaften, St. Petersburg 1887.
- Williamson, F., and Coauthors, 2018: Collating Historic Weather Observations for the East Asian Region: Challenges, Solutions, and Reanalyses. *Adv. Atmos. Sci.*, **35**, 899–904.
- Wolf, A., 1962: A history of science, philosophy and technology in the eighteenth century, Vol. 1. George Allen & Unwin, London, 274–305.
- Wyszynski, P., and R. Przybylak, 2014: Variability of humidity conditions in the Arctic during the first International Polar Year, 1882–83. *Polar Res.*, **33**, 23896, DOI: 10.3402/polar.v33.23896.
- Zaiki, M., G. P. Konnen, T. Tsukahara, P. D. Jones, T. Mikami, and K. Matsumoto, 2006: Recovery of nineteenth-century Tokyo/Osaka meteorological data in Japan. *Int. J. Climatol.*, **26**, 399–423.

# Figures


Darwar	16.28 15.32?	72.50	2250	23.7	<p>Jan.—Oct. 1827, obs. Turnbull, Christie 10 u. 10<sup>h</sup> u. 2-stdl. (Nov. u. Dec. eingeschalt.); James N. Ph. J. 1828.            23 J. 1796—1807 u. 1813—25 nach stdl. Beob. (monatl. 3 Tage) auf wahre Med. red. von Goldingham, Madras Observ. Pap. 1827 p. 859 sq. <math>\frac{w. 24.6}{s. 30.3}</math>; <math>\frac{k. M. 24.0}{w. M. 31.2}</math> (a. 21 J.)            — Roxburgh's Beob. geb. nur 26°.9 Phil. Tr. In einer kalten Ebene.            7 J. Febr. 1820—Dec. 25, 1830 u. 35, obs. Monat Sid. Med. der ersten 6 J. 23°.7.            D. einz. Jahresmed. zeigen eine Diff. von 3°.6. Calc. Tr. v. VI; 2-stündl. Bb. 1835 v. Monat geben 23°.56; 10<sup>h</sup> Mg. 0°.3 weniger. Beng. J. V. 296.            1 J. 1827 Std. Ebdas.            Arnee: Sid. etc. ebf. unbek.; 1828, 1 J.: 29°.0.            2 J. 1814 u. 16, obs. Seemann i. Zimmer, Oaig. u. 3—4<sup>h</sup>. Brewst. Ed. J. Sc. V. 259. Kämtz rechnet 25°.2            (1 J., 6 u. 10<sup>h</sup> Morg.; a. d. 2-stdl. Bb. in Bangalore ist die Corr. auf 10<sup>h</sup> Morg. angewendet, daher das Mittel höher als Baikie's Angabe (18°.5); vielleicht zu hoch, Madr. J. of Science, Oct. 1836 (Bibl. univ.).            A. mangelh. Ang. geschätzt; Not. statist. — 29°.6 v. Humb. l. is., gewiss zu hoch. N. Cossigny's Bb. (vgl. Port Louis!) u. Le Gentil Voy. mers de l'Inde I. 474.            4 J. 1736—9, 68 u. 69. K. M. 24.5; w. M. 33.0? Sandige Küste.            J. Mg. u. Mittg.-Bb. Not. stat. Col. Fr. III. 23.            (Auf d. Vorhöh. d. Ghats bei Tellichery; 10 J. Std.? (1810—13, 18—23), obs. Murdoch Browne. Tr. Lit. Soc. Madr. P. I. 1827. p. 89. <math>\frac{k. M. 23.7}{w. M. 29.7}</math>            3 J.: 1 (?) J. Beob. aus 6, 3 u. 9 (13°.6) und 24 J. (1831—33) aus 6 u. 3<sup>h</sup> v. Baikie. Ph. S. Calc. Tr. v. IV. 77, J. Beng. III. 653, B's Topogr. of Neilgher., Martin Br. Col. I.            Jackanary: 4700' h. (1 J.?) a. Beob. um 6, 3 u. 9<sup>h</sup>: 15°.6. Zu niedrig?            Dimhutti: 5800' h.: 17°.8; nach Ritter's Asien IV.</p>
Madras	13.5	77.57	—	27.8	
Bangalore	12.58	75.17	2730	23.7	
Arcot	12.54	77.8	590	27.7	
Seringapatam	12.45	74.21	2260	25.0	
Mascara	12.26	73.30	4200	20.8	
Pondichéry	11.56	77.32	—	28.6?	
Mahé	11.42	73.12	...	26.1	
Anjarakandy	11.40	78.20	40	27.2	
Ootacamund	11.25	74.30	6900	14.0	

Fig. 1. Excerpt taken from the inventory of Dove (1841). It contains a lot of useful information in condensed form.


Start year




Length (yrs)




**Fig. 2.** Coverage of entries in the inventory as a function of (top) start years and (bottom) record length, i.e., the number of years prior to 1850 (1890 for Africa and Arctic).




**Fig. 3.** Number of entries as a function of time and region (note the logarithmic scaling of the y-axis).



**Fig. 4.** Series inventoried for Africa.



**Fig. 5.** Series inventoried for (left) Asia and (right) South America. The thick grey line denotes the Arctic region.



**Fig. 6.** Series inventoried for North and Central America and the Caribbean. The thick grey line denotes the Arctic region.




Fig. 7. Series inventoried for the Southwest Pacific.





Fig. 8. Series inventoried for Europe. The thick grey line denotes the Arctic region.



**Fig. 9.** Series inventoried for the Arctic (definition shown by the thick grey line).



## Tables

**Table 1.** Global (italics) and national repositories searched (see also Table 3), number of pre-1850 records, and reference.

<b>Repository</b>	<b>Abbr.</b>	<b>N</b>	<b>Reference</b>
<i>Global Historical Climatology Network</i>	GHCN	596	Lawrimore et al. 2011, Menne et al. 2012
<i>International Surface Temperature Initiative</i>	ISTI	710	Rennie et al. 2014
<i>Climatic Research Unit Temperature</i>	CRUTEM3/4	476	Jones et al. 2012
<i>Berkeley Earth</i>	BEST	358	Rohde et al. 2013
<i>International Surface Pressure Databank</i>	ISPD	193	Cram et al. 2015
Historical Instrumental Climate Series of the Greater Alpine Region	HISTALP	85	Auer et al. 2007
Canada (incl. Hudson Bay Company)		369	Slonosky 2014; Slonosky pers. comm.
US Army Signal Service and other 19th Century Voluntary Observations	CDMP Forts	142	Dupigny-Giroux et al. 2007, Westcott et al. 2011
MétéoFrance	MétéoFrance	236	Brunet et al. 2013
French National Archives	FNA	53	Brunet et al. 2013
Swiss Metadata Inventory	CHIMES	200	Pfister et al. 2019
German Weather Service	DWD	138	
Austria	ZAMG	66	
Sweden	Moberg	ca. 100	Moberg 1998
National Library of Iceland	ICELAND	65	
Cambiamenti climatici e agricoltura	CLIMAGRI	28	Maugeri et al. 2006
Early meteorological records from Latin-America and the Caribbean	EMERLAC	33	Domínguez-Castro et al. 2017
Russia	RIHMI	28	Gazoina and Klimenko 2008

**Table 2.** Historical inventories (italic) and collections considered, number of pre-1850 records, and reference.

<b>Inventory/collection</b>	<b>Title/Source</b>	<b>Region</b>	<b>N</b>	<b>Reference</b>
<i>Berghaus</i>	<i>Physikalischer Atlas</i>	<i>Global</i>	304	<i>Berghaus 2004</i>
<i>Dove</i>	<i>Repertorium/Isotherme Linien</i>	<i>Global</i>	1246	<i>Dove 1841, 1852</i>
<i>Hellmann Germany</i>	<i>Repertorium</i>	<i>Germany</i>	457	<i>Hellmann 1883</i>
<i>Hellmann global</i>		<i>Global</i>	197	<i>Hellmann 1901, 1927</i>
<i>Schott</i>	<i>Tables, distribution and variations of the atmospheric temperatures in the US and adjacent parts of America</i>	<i>North America</i>	586	<i>Schott 1876</i>
<i>Schouw</i>	<i>Tableau du climat et de la végétation de l'Italie</i>	<i>Italy</i>	49	<i>Schouw 1839</i>
<i>Angot</i>	<i>Premier catalogue des observations météorologiques faites avant 1850</i>	<i>France</i>		<i>Angot. 1897</i>
<i>Raulin</i>	<i>Observations pluviométriques faites en France et dans les colonies françaises</i>	<i>France/Colonies</i>	100	<i>Raulin .1875ab, 1876, 1881</i>
<i>Havens</i>	<i>An annotated bibliography of meteorological observations in the United States</i>	<i>USA</i>	89	<i>Havens 1958</i>
Kanold	Kanold Colletion	Europe	30	Brázdil et al. 2008, Lüdecke 2010
Jurin	Royal Society	Europe North America Asia	28 1 1	Jurin 1723, Derham 1735, Hadley 1741, 1744
Cotte	Mémoires sur la météorologie /Bibliothèque de l'Académie nationale de médecine	Europe	75	Cotte, 1788
Palatina	Ephemerides	Europe/North America	37	Kington 1974, 1988

**Table 3.** Publicly accessible data repositories (*italics*) and with pre-1850 measurements as well as larger data collections published in data journals.

<b>Repository</b>	<b>Abbr.</b>	<b>URL</b>
<i>CRU Temperature</i>	CRUTEM	<a href="https://crudata.uea.ac.uk/cru/data/temperature/crutem4/station-data.htm">https://crudata.uea.ac.uk/cru/data/temperature/crutem4/station-data.htm</a>
<i>Global Historical Climatology Network</i>	GHCN	<a href="http://www.ncdc.noaa.gov/data-access/land-based-station-data/land-based-datasets/global-historical-climatology-network-ghcn">//www.ncdc.noaa.gov/data-access/land-based-station-data/land-based-datasets/global-historical-climatology-network-ghcn</a>
<i>International Surface Pressure Database</i>	ISPD	<a href="https://reanalyses.org/observations/international-surface-pressure-databank">https://reanalyses.org/observations/international-surface-pressure-databank</a>
<i>International Surface Temperature Initiative</i>	ISTI	<a href="ftp://ftp.ncdc.noaa.gov/pub/data/globaldatabank/monthly/stage3/">ftp://ftp.ncdc.noaa.gov/pub/data/globaldatabank/monthly/stage3/</a>
<i>BerkeleyEarth Surface Temperature</i>	BEST	<a href="http://berkeleyearth.org/source-files/">http://berkeleyearth.org/source-files/</a>
<i>Royal Dutch Weather Service</i>	KNMI	<a href="http://projects.knmi.nl/klimatologie/daggegevens/antieke_wrn/index.html">http://projects.knmi.nl/klimatologie/daggegevens/antieke_wrn/index.html</a>
<i>German Weather Service</i>	DWD	<a href="ftp://ftp-cdc.dwd.de/pub/CDC/observations_germany/climate/daily/kl/historical/">ftp://ftp-cdc.dwd.de/pub/CDC/observations_germany/climate/daily/kl/historical/</a> <a href="ftp://ftp-cdc.dwd.de/pub/CDC/observations_germany/climate/monthly/kl/historical/">ftp://ftp-cdc.dwd.de/pub/CDC/observations_germany/climate/monthly/kl/historical/</a>
<i>German Weather Service, overseas data</i>	DWD	<a href="https://www.dwd.de/DE/leistungen/ueberseestationen/ueberseestationen.html">https://www.dwd.de/DE/leistungen/ueberseestationen/ueberseestationen.html</a>
<i>Russian Hydrometeorological Service</i>	RIHMI	<a href="http://meteo.ru/english/climate/cl_data.php">http://meteo.ru/english/climate/cl_data.php</a>
<i>Norwegian Weather Services</i>	MetNo	<a href="http://eklima.met.no">http://eklima.met.no</a>
<i>Japanese Data</i>	JCDP	<a href="https://jcdp.jp/instrumental-meteorological-data/">https://jcdp.jp/instrumental-meteorological-data/</a>
<i>Historical Arctic Database</i>	NCU	<a href="http://www.hardv2.prac.umk.pl/">http://www.hardv2.prac.umk.pl/</a>
<i>Mediterranean Data Rescue Project</i>	MEDARE	<a href="http://app.omm.urv.cat/urv/accessdata/">http://app.omm.urv.cat/urv/accessdata/</a>
<i>Climate Database Modernization Programme</i>	CDMP	<a href="https://www.ncdc.noaa.gov/IPS/">https://www.ncdc.noaa.gov/IPS/</a> and <a href="https://mrcc.illinois.edu/data_serv/cdmp/cdmp.jsp">https://mrcc.illinois.edu/data_serv/cdmp/cdmp.jsp</a>
<i>Early (pre-1900) rainfall records for Africa</i>	NIC131	<a href="https://www1.ncdc.noaa.gov/pub/data/paleo/historical/africa/africa2001precip.txt">https://www1.ncdc.noaa.gov/pub/data/paleo/historical/africa/africa2001precip.txt</a>
Early meteorological records from Latin-America and the Caribbean during the 18th and 19th centuries	EMERLAC	Domínguez-Castro et al. (2017), doi:10.1038/sdata.2017.169
Southeastern Australian rescued observational climate network, 1788–1859	SEARCH	Ashcroft et al. (2014), doi:10.5281/zenodo.7598
A historical surface climate dataset from station observations in Mediterranean North Africa and Middle East areas	EURO4M	Brunet et al. (2014), doi:10.5281/zenodo.7531